



Swiech, W. M., Hamerton, I., Zeng, H., Watson, D. J., Mason, E., & Taylor, S. E. (2017). Water-based fractionation of a commercial humic acid. Solid-state and colloidal characterization of the solubility fractions. *Journal of Colloid and Interface Science*, 508, 28-38.
<https://doi.org/10.1016/j.jcis.2017.08.031>

Peer reviewed version

Link to published version (if available):
[10.1016/j.jcis.2017.08.031](https://doi.org/10.1016/j.jcis.2017.08.031)

[Link to publication record in Explore Bristol Research](#)
PDF-document

This is the author accepted manuscript (AAM). The final published version (version of record) is available online via Elsevier at <http://www.sciencedirect.com/science/article/pii/S0021979717309372?via%3Dihub>. Please refer to any applicable terms of use of the publisher.

University of Bristol - Explore Bristol Research

General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:
<http://www.bristol.ac.uk/pure/about/ebr-terms>

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21

Water-based fractionation of a commercial humic acid. Solid-state and colloidal characterization of the solubility fractions.

Weronika M. Swiech^a, Ian Hamerton^{b,1}, Huang Zeng^c, David J. Watson^b,

Eleonore Mason^d, and Spencer E. Taylor^{a*}

^a Centre for Petroleum and Surface Chemistry, Department of Chemistry, Faculty of Engineering and Physical Sciences, University of Surrey, Guildford, Surrey GU2 7XH, UK

^b Department of Chemistry, Faculty of Engineering and Physical Sciences, University of Surrey, Guildford, Surrey GU2 7XH, UK

^c BP America, Upstream Technology, 501 Westlake Blvd., Houston TX 77079, USA

^d Department of Chemistry, University of Reading, Reading, RG6 6AD

Email: s.taylor@surrey.ac.uk

Abstract

Background and hypothesis. Humic acid (HA) is of considerable environmental significance, being a major component of soil, as well as being considered for application in other technological areas. However, its structure and colloidal properties continue to be the subject of debate, largely owing to its molecular complexity and association with other humic

¹ Current address: Bristol Composites Institute (ACCIS), Department of Aerospace Engineering, Queen's Building, University of Bristol, University Walk, Bristol, BS8 1TR

1 substances and mineral matter. As a class, HA is considered to comprise supramolecular
2 assemblies of heterogeneous species, and herein we consider a simple route for the separation
3 of some HA sub-fractions.

4 *Experiments.* A commercial HA sample from Sigma-Aldrich has been fractionated into two
5 soluble (**S1**, **S2**) and two insoluble (**I1**, **I2**) fractions by successive dissolution in deionized
6 water at near-neutral pH. These sub-fractions have been characterized by solution and solid-
7 state approaches.

8 *Findings.* Using this simple approach, the HA has been shown to contain non-covalently
9 bonded species with different polarity and water solubility. The soluble and insoluble
10 fractions have very different chemical structures, as revealed particularly by their solid-state
11 properties (¹³C NMR and IR spectroscopy, and TGA); in particular, **S1** and **S2** are
12 characterized by higher carbonyl and aromatic contents, compared with **I1** and **I2**. As shown
13 by solution SAXS measurements and AFM, the soluble fractions behave as hydrophilic
14 colloidal aggregates of at least 50 nm diameter.

15

16 **Keywords**

17 AFM, colloidal properties, humic acid fractionation, IR spectroscopy, solid-state NMR,
18 surface tension, SAXS, TGA, zeta potential

19

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23

1. Introduction

The present paper considers the aqueous solubility and composition of a commercial sample of humic acid (HA). This has been chosen as an example of one of the classes of “humic substances” produced by the environmental decomposition of natural organic matter, and which are loosely defined in terms of the procedures used in their extraction. Humic substances are present in soils, rivers and other aquifers, and HAs have been shown to exist as aggregates of colloidal size at sufficiently high concentration [1,2]. Our original interest in these materials stems from their role in bitumen extraction from oil sand ores [3,4], where HA is mostly complexed with clays which, depending on the geological origin, can have different consequences for bitumen recovery [5,6]. However, the present work has a more general context.

On a molecular level, HAs are difficult to characterize. They are known to be (or contain) polyelectrolytes which interact strongly with certain cations to form insoluble complexes [7,8], and have been shown to be active in removing metal contaminants from water [9]. Other studies have demonstrated an improved ability of humic substances over conventional surfactants, such as Triton X-100 or sodium dodecyl sulfate, to aid soil bioremediation through the removal of pollutants such as polyaromatic hydrocarbons, thiophenes, sulfones or biphenyls [10].

As has recently been discussed [11], humic substances have been subdivided into three separate classes based on their respective solubility characteristics. Thus, humins are insoluble under alkaline conditions, fulvic acids are soluble throughout the pH range, and HAs are insoluble under acidic conditions. These materials have intrinsic surfactant-like

1 tendencies, e.g. reducing the surface tension of aqueous solutions [2,12-14] and solubilizing
2 organic molecules in colloidal aggregates [15].

3 The separation of humic substances has been the subject of a number of studies, as recently
4 reviewed [16]. The analysis of different fractions taken from within individual classes has
5 only served to highlight the molecular complexity of these substances, and detailed molecular
6 structures still remain elusive. Indeed, this may not be surprising based on the arguments that
7 humic substances are more likely to be a “continuum of progressively decomposing organic
8 compounds” than “inherently stable and chemically unique compounds” [11].

9 In this paper, we are particularly interested in the composition and colloidal properties of
10 HA component species. In the majority of previous studies, chemical and molecular size
11 differences have been exploited in order to effect fractionation. For example, the recent
12 procedure used by Chilom et al. [13] involved alkali- and solvent-based extractions, enabling
13 the isolation of one lipid-like and two humic-like fractions. Powell and Town solubilized
14 HAs using a combination of pH and different ionic media which were then fractionated based
15 on molecular size using gel permeation chromatography combined with equilibrium dialysis
16 [17]. Conte et al. used preparative size exclusion chromatography to produce seven fractions
17 from a lignite HA which were shown by different spectroscopic analyses to vary in aromatic-
18 aliphatic character and oxidized state [18]. Using ultrafiltration, Carlsen et al. produced six
19 molecular size fractions ranging from <1 kDa to <100,000 kDa from Aldrich HA and
20 examined their respective interactions with Eu^{3+} ions [19]. Similarly, Francioso et al.
21 produced six fractions from Na humate solution ranging from 5-10 kDa to >300 kDa using
22 tangential ultrafiltration, which were characterized by vibrational and NMR spectroscopic
23 techniques [20].

1 However, Pitois et al. [21] used a different approach to produce “chemical adsorptive”
2 fractions from commercial HA by adsorption on quartz sand. They found that adsorption of
3 HA was a two-stage process, one occurring faster (~3 h) than the other (~45 h). The HA
4 solutions remaining after each adsorption period were analyzed using asymmetric flow-field
5 flow fractionation coupled with UV/visible absorption spectrophotometry, which showed that
6 adsorption of lower molecular weight components (<4800 Da) occurs initially, followed by
7 higher molecular weight components (1400-9200) [21]. Fractions produced by this method
8 are clearly towards the lower molecular weight end of the ultrafiltration fraction ranges.

9 A contemporary model for the structure of HAs proposes supramolecular assemblies of
10 smaller heterogeneous molecules held together by hydrophobic interactions and hydrogen
11 bonding [22,23]. The composition of humic substances, including HA, are naturally
12 heterogeneous and source-dependent, since they originate from the breakdown of plant and
13 animal matter by microbiological and abiotic transformation.

14 Herein, we present results from an experimental study on a commercial HA sample from
15 Sigma-Aldrich which has been the subject of several reports in the literature over the years
16 [1,7,9,12,21]. We report that it is possible to isolate “solubility fractions” based on
17 differential solubility in deionized water under near-neutral pH conditions. Klučáková and
18 Pekař [24,25] have previously noted different dissolution/solubility characteristics of solid
19 HAs, including a commercial sample from Fluka, which on the basis of its pH-concentration
20 behavior is considered similar to the product used herein (see later). These workers proposed
21 a multi-step dissolution mechanism upon interaction with water, involving dissolution, with
22 or without acid dissociation [24,25]. We were therefore interested to investigate further the
23 water solubility of the commercial HA and the nature of the solubility fractions produced.

24
25

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25

2. Experimental

2.1. Materials and HA fractionation in water

All reagents, including neutral HA, the corresponding sodium salt (NaHA), the inorganic salts NaCl, CaCl₂.2H₂O and LaCl₃.7H₂O, and cetyltrimethylammonium bromide (CTAB) were the highest purity grades available from Sigma-Aldrich, UK and were used as received. Aqueous solutions were prepared using deionized water (resistivity 18.2 MΩ.cm) from a Millipore Direct-Q system.


Inorganic matter is known to be an impurity in HA and this is confirmed by the lower elemental compositions in the present sample of HA compared with the literature values of a similar product produced from NaHA: found (lit. [26]) (%): C, 36.9 (49.7); H, 3.10 (4.49); C/H ratio 0.99 (0.92). On the other hand, the similar C/H ratio found for the present unpurified product is consistent with an impurity devoid of carbon and hydrogen. Thermal analysis data are reported later in this paper. Since the compositions of HA and NaHA could potentially vary from sample to sample, experiments were conducted as much as possible using a single commercial batch (identified by lot number).

HA and NaHA solutions were prepared by agitating the appropriate quantities of the respective dry solids in deionized water using an orbital shaker at ambient temperature (20 ± 2 °C). Solutions of the desired concentration of the sodium salt were prepared by dissolving the exact weighed amount of NaHA in deionized water. However, in the case of HA, it was discovered that a proportion of the HA was soluble in deionized water, the concentration of the soluble fraction being dependent on the initial mass of HA, and to a lesser extent, time (from 4-48 hours used). The solutions produced in this way were then filtered to remove

1 undissolved solid (termed “insoluble 1”, **I1**), and the concentrations of the soluble fraction
2 (termed “soluble 1”, **S1**) in the filtrates were determined gravimetrically after drying. Lower
3 concentration solutions were prepared by dilution.

4 This procedure was extended by removing the undissolved **I1** from the first dissolution
5 stage and adding this to fresh deionized water. This resulted in further, but also incomplete,
6 dissolution and, again, the undissolved solid (termed “insoluble 2”, **I2**) was removed by
7 filtration, the filtrate containing the second soluble component (“soluble 2”, **S2**). This
8 procedure is shown in Scheme 1. The corresponding “soluble” components (**S1** and **S2**) were
9 recovered by carefully drying aliquots of the respective solutions in preparation for further
10 analysis.

11



12

13 **Scheme 1.** Sequence of humic acid dissolution in water to produce soluble (**S1**, **S2**) and insoluble (**I1**, **I2**)
14 fractions.

15

16 **2.2. Characterization Methods**

17 *2.2.1. Colloidal and solution analysis*

18 Zeta potentials of **S1** aqueous solutions (prepared by dilution of a stock solution as
19 described above to produce solutions in the range 2.6–1320 mg/L) were determined using a

1 ZetaMaster (Malvern Instruments, Malvern, UK). Surface tensions of the aqueous solutions
2 of NaHA and **S1** were determined as a function of concentration by the du Noüy ring method
3 with a Krüss K10 tensiometer (Krüss GmbH, Hamburg, Germany). The Pt-Ir ring was rinsed
4 thoroughly with deionized water and flamed before making each measurement, which were
5 made at least in triplicate. Both sets of measurements were made at ambient temperature (20
6 ± 2 °C).

7 Small-angle X-ray scattering (SAXS) measurements on aqueous **S1** and **S2** solutions were
8 made using the B21 beamline at the Diamond Light Source of the UK National Synchrotron
9 Science Facility (Harwell Science and Innovation Campus, UK). Solutions were manually
10 injected via a coupled syringe into a quartz capillary (1.8 mm internal diameter) in the X-ray
11 beam. The quartz capillary was enclosed in a vacuum chamber, in order to reduce
12 background scatter. Experiments were carried out with a camera length of 4.01 m and fixed
13 energy (12.4 keV, corresponding to a wavelength of 0.999 Å), providing a q range of 0.022 to
14 4.2 nm^{-1} . Scattering patterns were acquired using a Pilatus 2M detector. Background (water)
15 subtraction and radial averaging were performed using the dedicated beamline software
16 ScÅtter.

17 Concentrations of acid groups in **S1** and **S2** were determined using a modification of the
18 cationic surfactant-based precipitation method described by Nobili et al. [27]. In the present
19 work, typically 5-10 mg of solid **S1** or **S2** was dissolved in deionized water (40 mL) and
20 titrated with a solution of CTAB (i.e. $\text{CTA}^+ \text{Br}^-$) of known concentration ($\sim 5 \times 10^{-3} \text{ M}$) at
21 either pH 7 or pH 9.8. Reaction with CTA^+ at the former pH provides a measure of acid
22 groups dissociated at pH 7 (mainly, but not exclusively, carboxylic), whereas at the higher
23 pH CTA^+ reacts additionally with phenolic groups [27]. The original method identifies the
24 most optically-clear supernatant, after centrifuging a range of mixtures with different
25 HA/CTAB ratios. Here, we detect the end-point visually by the appearance of large flocs,

1 assisted by “spotting” one drop of the titration mixture onto a filter paper (Whatman #1). The
2 presence of large flocs is apparent as dark particles (see Fig. S1, Supporting Material). We
3 also continuously monitored the solution pH during the titration, which decreases as the end-
4 point is approached, in a single stage at the lower pH, and in two stages at the higher pH, as
5 also shown in Fig. S1. We suggest that the two stages indicate the types of acid groups
6 involved [27].

7 The effect of ionic strength and cation valency on the dissolution of HA, and the colloid
8 stability of **S1**, were analyzed spectrophotometrically (Thermo-Scientific Evolution 2000
9 UV-Vis spectrophotometer). In the former experiments, HA was suspended in NaCl, CaCl₂
10 and LaCl₃ solutions and agitated using an orbital shaker (4 h; 200 rpm). After filtration
11 (Whatman #1 filter paper), the absorbance at 450 nm of each filtrate was compared with the
12 corresponding value for a suspension of HA in deionized water. For the colloid stability
13 measurements, aliquots (2.5 mL) of a stock **S1** solution (0.62 g/L) were mixed with equal
14 volumes of known concentration salt solutions and also agitated using an orbital shaker (4 h;
15 200 rpm). In these latter experiments, any precipitate formed was removed by centrifugation
16 (Rotina 380 Hettich benchtop centrifuge at 5000 rpm for 10 minutes), and the residual **S1**
17 concentrations determined by comparing absorbance values at 450 nm with a deionized water
18 reference (i.e. equivalent to 0.31 g/L).

19

20 2.2.1. *Solid-state analysis*

21 Thermogravimetric analysis (TGA) was conducted using a TA Instruments Q500 TGA
22 analyzer. Powdered samples (~5 mg) were placed in a shallow platinum crucible and heated
23 in static air at a rate of 10 K/min from room temperature to 950 °C. The results are expressed
24 as thermogravimetric (TG) and first derivative (DTG) plots. Elemental analyses (CHN) of the

1 humic substances were determined by combustion analysis using an Exeter Analytical CE440
2 elemental analyzer. Infrared spectroscopic analysis was carried out using a Brüker Alpha
3 instrument in ATR mode (2 cm^{-1} spectral resolution).

4 Solid-state NMR experiments were conducted on HA, NaHA, **S** and **I** samples at the
5 former EPSRC National Solid-State NMR Service facility at the University of Durham. ^{13}C
6 Multiple cross-polarization magic angle spinning (CPMAS) spectra [28] were obtained at
7 100.56 MHz at ambient probe temperature using a Varian VNMRs spectrometer based on a
8 9.4 T Oxford Instruments superconducting magnet.

9 Imaging of mica surfaces containing humic acid fraction **S1** by atomic force microscopy
10 (AFM) were obtained under ambient conditions in air using a Brüker Innova atomic force
11 microscope in tapping mode using 256 scan lines and a scan rate of 4 Hz. The images were
12 post-processed using Brüker NanoScope software, which was used primarily to level the
13 images.

14

15 **3. RESULTS AND DISCUSSION**

16 **3.1. Aqueous fractionation of HA**

17 The dissolution behavior of the commercial HA powder is very different from that of
18 NaHA. Unsurprisingly, aqueous NaHA solutions are easily prepared simply by stirring the
19 solute into water. On the other hand, as previously also found by Klučáková and Pekař, HA is
20 only partially soluble, even when very small quantities are added to water under neutral pH
21 conditions [24,25]. Here, we have found that proportionately more concentrated solutions
22 were produced by increasing the initial quantities of HA powder, and even after prolonged


1 (~48 h) agitation the water/HA mixtures contained undissolved (organic and inorganic)
2 components.

3 As an example, the addition of as-received HA powder (0.125 g) to deionized water (100
4 mL) at ambient temperature consistently produced a concentration of 0.39 ± 0.03 g/L in
5 solution, suggesting the presence of ~30% of a soluble component (**S1**) under these
6 conditions. The concentration increased to 0.50 and 0.54 g/L at 50° and 80° C, respectively.
7 Extending the number of ambient temperature experiments produced the data shown in Fig.
8 1(a). During dissolution, the pH was neutral to slightly acidic, and changed little with time, as
9 noted elsewhere for another commercial HA sample [25]. Over the course of a 4 h
10 experiment in which 2 g HA was suspended in deionized water (100 mL), the pH of the
11 solution decreased slightly from 6.11 to 6.02, while for an initial suspension of 6 g HA in the
12 same volume of deionized water, the pH remained at 5.74 throughout. The removal of excess
13 undissolved solid (i.e. **I1**, Scheme 1) by filtration or centrifugation enabled **S1** solutions of
14 relatively high concentration to be obtained.

15 The data in Fig. 1(a) indicate that a steady linear increase in the **S1** concentration is seen as
16 a function of the initially suspended HA, which indicates a limiting solubility for the HA
17 components comprising **S1** under these conditions. **S1** could potentially exist in a dissolved
18 state or in colloidal suspension, and coupled with the solubility behavior, led us to consider
19 the compositional and solution properties of the soluble and insoluble fractions.

20

21



1

2 **Fig. 1.** (a) Dissolution of HA powder in deionized water to produce **S1** in solution, leaving insoluble **I1**. The
 3 two highlighted points (*a* and *b*) identify the **S1** solutions used to produce samples for AFM. (b) Plot showing
 4 the solubility of excess powdered **I1** deionized water to yield **S2**. The curves are drawn to guide the eye.

5

6 Upon isolating the separated insoluble HA fraction **I1** (see Scheme 1), it was also found to
 7 yield ~30% of a second soluble fraction **S2** when added to further (100 mL) deionized water;

1 filtration also produced another residual insoluble fraction **I2**. The **S2** concentration is plotted
2 in Fig. 1(b) as a function of the original suspended **I1** concentration, from which it can be
3 seen by comparison with Fig. 1(a) that dissolution is generally lower than for HA. This is
4 indicative of a selective dissolution process, and the differences in the fractions from
5 successive dissolution steps would be expected to reflect the respective hydrophilicity of the
6 different fractions.

7 **3.2 Compositional and functional analysis of HA and fractions**

8 Thermogravimetric analysis (TGA) has been shown to be a useful technique for
9 characterizing HAs and other humic substances in past studies [29,30]. The pattern of HA
10 decomposition of a standard Suwannee River humic acid, for example, shows a small mass
11 loss in the region of 300–400 °C followed by a larger mass loss between 450–550 °C [29].
12 Below ~400 °C, mass loss is generally a result of decomposition or elimination of oxygen-
13 containing functional groups, whereas above 400 °C it reflects the high aromatic content [29].
14 Here, TGA in air has been performed on the fractions **S1**, **S2**, **I1** and **I2** as well as the
15 precursor HA. As can be seen in Fig. 2, the TG and DTG profiles for the two soluble
16 fractions are similar, as are those for the two insoluble fractions, although the respective
17 behaviors of soluble and insoluble fractions are distinctly different; the profile for the original
18 HA is intermediate.

19 All the samples show ~10-15% mass loss up to ~300 °C, most probably a result of
20 dehydration. However, true thermal degradation first appears as a small peak in the derivative
21 plots at ~300-325 °C for HA and the insoluble fractions, each amounting to ~10% mass loss.
22 The main degradation peaks for **I1** and **I2** follow at 423 and 390 °C, respectively, compared
23 with 555 °C for HA. The widths of these peaks decrease in the order HA > **I1** > **I2**, which
24 may reflect decreasing sample heterogeneity.

1 On the other hand, degradation of **S1** and **S2** is entirely different, exhibiting much higher
2 thermal stability. In these fractions, a steady mass loss totaling ~15% occurs up to 600 °C.
3 This is followed by the most significant mass loss (~45%) in the range 600-900 °C, which
4 contains two main degradation peaks at 784 and 890 °C for **S1**, and 754 and 832 °C for **S2**,
5 indicative of char formation from aromatic structures; oxidation subsequently occurs at much
6 higher temperatures.

7 HA, **I1** and **I2** each produce ~25-33% residue (note that the HA supplier quotes ~20 %
8 ash), whereas the corresponding yields from **S1** and **S2** are approximately half these values
9 (see Table 1). A more detailed comparison of the DTG plots in Fig. 2(b) reveals that
10 degradation of the original HA contains minor contributions from each of the individual
11 fractions. A broad peak at ~800 °C, for instance, is an indication of the soluble fractions,
12 whereas the insoluble fractions are more apparent in the temperature region leading up to the
13 main HA degradation at ~550 °C. However, it is interesting to note that the responses of the
14 respective fractions are not reflected in the overall behavior of HA. By considering the
15 TG/DTG profiles for the HA/**S1** and **I1/S2** solubility pairs, it appears that char formation in
16 the soluble fractions is suppressed by the presence of the respective insoluble fractions – for
17 example, mass loss from HA at >600 °C is 5.8% whereas it is 50.3% in **S1**, and yet HA
18 contains ~30% **S1**.

19

20

21

22

23

1 **Table 1. Microanalytical and TGA residue data and calculated atomic ratios for HA and its fractions.**

	C (wt%)	H (wt%)	N (wt%)	C/H atomic ratio	O/C atomic ratio ^a	Residue from TGA (%)
HA ^b						
Batch 1 (Lot # BCBG7429V)	40.33	3.69	0.80	1.10	0.40	33.9
Batch 2 (Lot # BCBK5107V)	42.75	3.61	0.99	1.01 (1.06) ^c	0.47	25.9
S1	30.5	1.64	<0.3	0.61	1.28	12.7
S2	36.9	2.20	<0.3	0.72	0.91	16.1
I1	40.5	2.90	<0.3	0.86	0.58	25.5
I2	41.8	3.03	<0.3	0.86	0.55	24.4

2

3 ^a %O calculated by difference, including residue content.


4 ^b Supplier's data.

5 ^c From ref. [31] for an earlier product dating from 1974.

6

7 The microanalytical results, summarized in Table 1, also indicate that higher TGA residue
 8 yields correspond to higher H/C atomic ratios (nitrogen levels were found to be below the
 9 0.3% level of uncertainty of the analytical procedure). It is therefore likely that the residues
 10 contain different amounts of ash originating from the original HA, which would be retained
 11 in the insoluble fractions. Fig. 2 and Table 1 show that residue yields for HA, **I1** and **I2** are
 12 very similar, whereas the corresponding **S1** and **S2** values are significantly lower. The O/C
 13 atomic ratios, estimated by taking residue levels into account, also reveal significant polarity
 14 differences between the soluble and insoluble fractions. Being the first fraction to dissolve, it

- 1 is not surprising that **S1** has a higher relative oxygen content than **S2**, and more than double
- 2 that of the insoluble fractions.



3
4 **Fig. 2.** (a) TG and (b) DTG profiles obtained in air for HA and the solubility fractions **S1**, **S2**, **I1** and **I2**.

5
6 Differences in oxygen content would also be expected to be evident from infrared
7 spectroscopic analysis. Thus, Figs. 3 and S2 (Supporting Material) reveals the present HA
8 sample and its fractions to be characteristic of a type III humic acid, based on Stevenson and


1 Goh's classification [32], indicating a relatively young product. In general, it is found that the
2 main IR bands are consistent with other humic acids reported previously, and subsequent
3 assignments follow those given in the literature [18,32-36].

4 All HAs contain broad bands at 3207–3356 cm^{-1} attributable to O-H stretch in both free
5 and dimeric carboxylic groups, as well as relatively small aliphatic C-H stretching bands in
6 the region of 2851 and 2921 cm^{-1} [32]. The spectra in Fig. S1 for HA and NaHA illustrate
7 these features very well, the bands differing only in their relative intensities.

8 However, it is perhaps more pertinent to point out differences seen in the spectra for **I** and
9 **S** fractions in the lower wavenumber region, as these are indicative of differences in chemical
10 structure and composition. Therefore, with reference to the 900–1100 cm^{-1} region of Fig. 3,
11 HA and all the fractions show two distinct absorption bands centered on 1021 cm^{-1} , although
12 these are more distinctive in HA, **I1** and **I2** compared with **S1** and **S2**. These bands have
13 previously been assigned to a C-O stretching mode in polysaccharides or polysaccharide-like
14 substances [35,36], coinciding with Si-O stretching modes (see below), indicating that all
15 these groups may be fewer in number in the soluble fractions, consistent with the above TGA
16 results.

17 The strong band at 1563 cm^{-1} (with a very small shoulder at $\sim 1700 \text{ cm}^{-1}$ in HA and the
18 insoluble fractions), respectively assigned to C=C skeletal vibrations and C=O stretch,
19 indicates a comparably small amount of carbonyl functionality which had been seen
20 previously in humic acid extracted from lignite [18], and also in a commercial humic acid
21 sample from Fluka [25]. The intensity of this band is reduced by complexation with divalent
22 ions (Cu^{2+} , Ca^{2+} , Pb^{2+}) or salt formation with K^+ [36]. The strong 1366 cm^{-1} band is assigned
23 to the symmetric C=O stretch of carboxylic acid groups and C-OH stretch of phenolic groups
24 and is slightly stronger in the soluble fractions **S1** and **S2**. A broad, lower intensity shoulder

1 at $\sim 1250\text{ cm}^{-1}$ was previously assigned to the C-O stretch of esters, ethers and phenols in the
2 humic acid structure [37]. From these comparisons, it appears that some part of the
3 supramolecular structure of HA has partially dissolved to generate the soluble fractions. In
4 the low wavenumber region of the spectra, a band at 531 cm^{-1} , due to Si-O-Si bending
5 vibrations, is present in the original HA, **I1** and **I2**, but absent in **S1** and **S2**. This band is
6 accompanied by Si-O stretching vibrations at 1030 cm^{-1} [38]. The substantial loss of those
7 peaks in **S1** and **S2** indicates that siliceous mineral matter (discussed above as being a known
8 impurity) most likely accumulates in the **I1** and **I2** fractions, or may be present, in part, as a
9 separate colloidal fraction.



10
11 **Fig. 3.** ATR-IR spectra of original HA and its soluble and insoluble fractions (indicated). Arbitrary, but
12 constant, vertical transmittance scale.

13 CPMAS solid-state NMR spectra of HA and its fractions, shown in Fig. S3 (Supporting
14 Material), are in general support of the foregoing. Peaks in solid-state ^{13}C NMR spectra are
15 typically assigned as [39-41]: alkyl C (0–45 ppm), O-alkyl C (45–110 ppm), olefinic or H- or

1 alkyl-substituted aromatic C (110–140 ppm), O- or N-substituted aromatic (mainly phenolic)
 2 C at 140–160 ppm, and carbonyl C (160–220 ppm). All the spectra in Fig. S3 show three
 3 main regions, corresponding to aliphatic, aromatic and carbonyl groups, as seen previously in
 4 solution spectra [42], and agree well with Aldrich HA samples studied by others [31], even
 5 after considerable purification stages to remove insoluble organic and inorganic contaminants
 6 [40,43,44]. In an extreme case, Fuentes et al. [39] obtained a spectrum almost devoid of
 7 aromatic signals which was similar in appearance to the lipid-like HA fraction extracted by
 8 Chilom et al. [13].

9 The present spectra allow relative signal intensities to be determined for the aliphatic,
 10 aromatic and carbonyl group regions, and these are summarized in Table 2, from which it is
 11 evident that the structural components are similar in the soluble and insoluble fractions, but
 12 that their relative intensities are noticeably different. In particular, the aliphatic/aromatic ratio
 13 for the soluble fractions is double that found for the insoluble fractions. The same is true for
 14 the respective carbonyl contents, which is consistent with the higher water solubility of **S1**
 15 and **S2**. The overall aromatic content of the soluble fractions is ~30% higher than in the
 16 insoluble fractions. These features have not been described previously with respect to
 17 fractionation studies of this commercial HA.

18 **Table 2. Compositional data, expressed as intensity ratios, from ¹³C CPMAS spectra for humic acid and**
 19 **its fractions.**

Sample	$I_{\text{Carbonyl/Total C}}$	$I_{\text{Aromatic C/Aliphatic C}}$	$I_{\text{Aromatic C/Total C}}$
HA	0.15	1.14	0.53
S1	0.19	1.99	0.67
S2	0.20	1.92	0.66
I1	0.12	1.00	0.50
I2	0.08	0.99	0.50

3.3. Solution and colloidal properties of the S1 fraction

The aqueous behavior of humic substances is highly dependent on the presence of carboxylic and phenolic groups, which principally define their acid-base characteristics [2,11,16,24,25,27]. In the present study we have separated two water-soluble fractions by dissolution in the pH range 5.7-7. Under these pH conditions, the more acidic carboxylic acid groups will be ionized.


In Table 3 are shown the acid group contributions to **S1** and **S2**, in comparison with a previous analysis of a re-precipitated Aldrich HA [27]. From these data, it is apparent that the two soluble fractions have a similar total number of acid groups, although **S2** has a greater contribution from weaker groups, which is in accord with their respective dissolution behavior, although it is not manifest in the NMR results.

Table 3. Analysis of acid groups of S1 and S2 by precipitation with CTA⁺. The literature data [27] have been included for comparison.

Sample	Acid groups (mmol/g)		
	Strong acid groups (pH 7)	Total acid groups (pH 9.8)	Weak acid groups
S1	2.70 ± 0.16	3.48 ± 0.03	0.78 ± 0.19
S2	2.36 ± 0.07	3.46 ± 0.02	1.10 ± 0.09
Twice re-precipitated HA (Aldrich)* from ref. [27]	2.87	3.61	0.74

Evidence for the anionic colloidal nature of the first soluble fraction comes from the effect of concentration on the zeta potential (ζ) of **S1** in water, shown in Fig. 4(a). A pH change

- 1 from 6.9 to 6.1, within the range originally used to separate **S1**, accompanies the attainment
- 2 of a plateau ζ of ~ 30 mV, which starts at ~ 0.1 g/L.



- 3
- 4 **Fig. 4.** (a) Concentration dependence of the zeta potential for **S1** solutions. (b) Concentration dependence of the
- 5 surface tension for **S1** solutions (open circles), NaHA at their natural pH (9.8) (filled circles), and (c) NaHA at
- 6 pH 11 (open triangles). The inset shows the comparison between surface tensions for **S1** and **S2** solutions. All
- 7 lines have been drawn to guide the eye.

1

2 Corresponding surface tensions of **S1** solutions, together with NaHA (at pH 9.8 and 11),
3 are shown as a function of concentration in Fig. 4(b). **S1** is seen to exhibit only a slight
4 reduction in surface tension at ~3 g/L, compared with the behavior of the NaHA solutions for
5 which the surface tension starts to decrease at ~0.3 g/L (pH 9.8) and ~0.02 g/L (pH 11). This
6 suggests that surface activity may be related to the ionization of the phenolic groups.

7 The surface tension behavior of HA solutions has been determined in several previous
8 studies from which the results were variously interpreted in terms of micelle formation
9 [1,12,45,46], “pseudo-micelle” formation (explained as micelle-like aggregates, but without
10 clear critical micelle concentrations) [2,47], or other “non-micelle-like” humic acid
11 aggregates forming at high concentrations [48]. The surface tension behavior found by
12 Terashima et al. was shown to be more dependent on HA concentration than on pH [46]. The
13 inflection point in the surface tension-pH curve at pH ~5.5 found by these workers did not
14 correspond to the apparent acid dissociation constant which was interpreted as reflecting the
15 heterogeneous nature of HAs [46,49]. In fact, the data also suggested a very weak
16 dependence on pH in the range 6-7 covered in the present study [46].

17 Small differences in surface tension data for the two isolated soluble fractions **S1** and **S2** at
18 their natural pH range of 6-7 are shown in the inset to Fig. 4(b). Over the concentration range
19 studied, **S2** is found to exhibit slightly lower surface tensions than **S1**, as would be expected
20 from the lower solubility of the former, possibly due to the presence of lipid-like groups. In
21 support of this, the partial solubility of humic acid [24,25] containing a lipid-like fraction (as
22 opposed to the humic-like fraction) was previously reported to have a lower surface tension
23 in aqueous solution [13].

24

1


2 *3.3.1. Cation effects on HA dissolution and S1 precipitation behavior*

3 The anionic character of **S1** is supported by the effects of metal chloride salts on HA
4 dissolution and **S1** solubility, as shown in Fig. 5. It is seen that both properties are inhibited
5 as cation concentrations are increased. Previous work by Powell and Town [17] indicated that
6 humic acid solubility decreased with increasing ionic strength, i.e. salting-out upon the
7 addition of KNO₃ (0.1 and 0.6 mol/L), Na₄P₂O₇ (0.1 mol/L) and synthetic seawater. These
8 workers also reported that the effect of KNO₃ was much lower than that of Na₄P₂O₇ when
9 present at the same molar concentrations. The solubility of humic acid in seawater (excluding
10 divalent ions) decreased by ~30%. The addition of Mg²⁺ (0.05 mol/L) and Ca²⁺ (0.1 mol/L)
11 reduced HA solubility by ~60%. From Fig. 5(a), it is seen that the ionic strengths at which
12 each of the salts NaCl, CaCl₂ and LaCl₃ initially inhibit HA dissolution differ by
13 approximately an order of magnitude, the multivalent cations being more effective than
14 monovalent Na⁺. This is consistent with the known preference of HA to complex with
15 multivalent cations [19].

16 A minimum ionic strength can also be identified for each chloride salt, above which HA
17 dissolution is completely inhibited. These values are in the order NaCl (~1 mol/L) >> CaCl₂
18 ≈ LaCl₃ (~0.03 mol/L), and most likely reflect double-layer effects as well as complexation
19 with the HA polyelectrolyte species, since the relative ionic strengths are not fully consistent
20 with the expected $1/z^6$ dependence on cation charge (z) predicted by DLVO theory for critical
21 flocculation or coagulation concentrations of colloids [50]. Therefore, unlike in previous
22 work [17], the present results suggest the importance of cation charge on dissolution through
23 colloidal interactions and complexation, consistent with the anionic character of the HA
24 species.

1 Similar salt effects are seen regarding the stability of **S1** solutions. As the data in Fig. 5(b)
2 illustrate, the presence of the same three salts as above all resulted in the salting-out of **S1**. In
3 these experiments, aliquots of stock **S1** solution (0.62 g/L) were diluted to double their
4 volume with different concentrations of the three aqueous chloride salts. LaCl_3 is the most
5 effective cation in terms of the extent of precipitation and the minimum ionic strength
6 required. CaCl_2 appears to be a considerably less effective, based on the same metrics.
7 Significantly higher ionic strength (>1 mol/L) is required to initiate precipitation in the case
8 of NaCl . It was also qualitatively observed that **S1** precipitation rates appear to vary in the
9 order $\text{NaCl} < \text{CaCl}_2 < \text{LaCl}_3$, suggesting that kinetic effects may also contribute to the
10 measured precipitation. Since it is known that HA hydroxyl, carbonyl and carboxylic acid
11 functional groups can participate in complex formation with divalent ions such as Cu^{2+} , Pb^{2+}
12 and Ca^{2+} [36], it is possible that such reactions could also contribute to the differences shown
13 in Fig. 5(b), rather than solely colloidal, double-layer (charge neutralization) effects [51]. The
14 increased binding of higher valence cations to HA has been identified by other workers
15 [19,52], and utilized in metal ion extraction [53].

16



1

2 **Fig. 5.** (a) The effects of salt concentration on the dissolution of **S1** from HA. **S1** concentrations are relative to
 3 dissolution in deionized water. (b) The effect of salt concentration on the precipitation of **S1** from 0.31 g/L
 4 aqueous solution. For both plots, the lines are to guide the eye.

5


6

7

1 3.3.2. *S1* solution structure

2 On the basis of the foregoing, we were interested to consider further the structure of **S1** as
3 an anionic colloid. Small pieces ($\sim 1 \text{ cm}^2$) of freshly-cleaved muscovite mica were placed in
4 two different **S1** solutions at low ($\sim 0.5 \text{ g/L}$) and high ($\sim 6 \text{ g/L}$) concentration for $\sim 1 \text{ h}$
5 (corresponding to the labeled positions *a* and *b*, respectively, in Fig. 1(a)). The mica pieces
6 were then removed, rinsed with deionized water, and air-dried. Subsequent AFM
7 measurements produced the images shown in Fig. 6, the adsorbed layer derived from the
8 lower concentration solution at position *a* in Fig. 1(a) having a smoother, globular appearance
9 compared to the higher concentration image from position *b*. This is more obvious from the
10 respective section profiles also shown in Fig. 6. The higher concentration solution produced
11 much smaller individual topological features which are similar to those reported elsewhere
12 for the same type of commercial material [54,55], although in the latter studies the solutions
13 were prepared using different procedures, and were of lower concentration (0.1 g/L). Thus,
14 Liu et al. [55] started with commercial NaHA which was dissolved in water and subjected to
15 a precipitation-ultrafiltration process to remove impurities and constrain the molecular weight
16 range to between 5000 and 10000 Da; this produced AFM heights for adsorbed HA of
17 3.1–3.7 and 4.2–5.7 nm depending on the drying regime applied during sample preparation.

18



1

2 **Fig. 6.** Tapping mode 3-D topological AFM images (top) and section profiles through the respective AFM
 3 images (bottom) for mica containing adsorbed **SI** from (a) ~0.5 g/L and (b) ~6 g/L solutions in deionized water
 4 (see also Fig. 1(a) for the solutions used).

5

6 Pläschke et al. [54] purified Aldrich NaHA according to the method of Kim et al. [56] by
 7 dissolution in 0.1 mol/L NaOH in the presence of sodium fluoride (0.2 g/g HA). After several
 8 precipitation/re-dissolution (NaOH)/re-precipitation cycles, the resultant HA was adsorbed on
 9 mica surfaces at different pHs and imaged using AFM. The smallest particle diameters were
 10 found to be in the range 8-13 nm by AFM, compared with spheroids of 9–12 nm diameter

1 observed by transmission electron microscopy (TEM) [57]. Chen et al. [58] also observed
2 similar images to those presented here for HA from an unidentified source.

3 On the other hand, Balnois et al. [59] studied two standard HAs: Suwannee River HA
4 (SRHA; International Humic Substances Society) and a peat-derived HA (PHA; UK
5 Geological Survey). These were sufficiently soluble in deionized water, and were free from
6 non-HA impurities. For SRHA the observed pH-invariant AFM image height was 0.5–2 nm,
7 whereas for the more hydrophobic PHA aggregated structures (~22 nm) dominated, with a
8 smaller fraction of 1–3 nm.

9 With the above studies as background information, it is evident that the present results for
10 **S1** adsorbed on mica show characteristics similar to those seen for other HA samples,
11 including some that had been exhaustively purified. At our lowest concentration (albeit
12 higher than in some previous studies), Fig. 6 (position *a*) is consistent with extended
13 molecules present on the mica surface under the low ionic strength/neutral pH conditions,
14 resulting in the uniform build-up of surface layers [55]. However, the image in Fig. 6
15 (position *b*) produced at higher **S1** concentration indicates the presence of smaller adsorbed
16 units exhibiting more regular, distinct structures. Both images are also consistent with recent
17 adsorption studies as a function of concentration on highly-ordered pyrolytic graphite [60] as
18 well as with modeling [61].

19 However, as was helpfully pointed out by one referee, AFM images of this type refer only
20 to films built up by dried primary units, and are not necessarily representative of the structure
21 in solution. It was also suggested that SAXS measurements, for example, could provide
22 additional information in support of solution structure, particularly if the dispersed units are
23 colloidal, and a brief study of **S1** and **S2** solutions is therefore included here.

1 SAXS allows the probing of different dimensional characteristics of a system as a function
2 of the scattering vector, $q = (4\pi \sin \theta)/\lambda$, where 2θ is the scattering angle and λ is the X-ray
3 wavelength. The approach we have taken here is, admittedly, relatively simplistic, in order to
4 identify differences in scattering behavior of the soluble HA fractions. On a log-log scale, the
5 Porod plots shown in Fig. 7 plots relate scattering intensity, $I(q)$ to the scattering vector, q for
6 **S1** and **S2** solutions in deionized water. In essence, each scattering curve shows two distinct
7 power law regions, in which $I(q) \propto q^{-d}$, where d is the Porod exponent. At high q ($>1 \text{ nm}^{-1}$),
8 the scattering is dominated by the internal molecular arrangements of the structural units. In
9 the lower q range ($0.04 \text{ nm} < q < 0.4 \text{ nm}$ in this case), larger, presumably aggregated,
10 structural units dominate the scattering. Analysis of scattering from the largest length scale
11 units was not possible owing to the limitation of the accessible q range. This also prevented a
12 simple Guinier analysis, according to $I(q) \propto \exp(-R_g^2 q^2/3)$, where R_g is the radius of gyration,
13 as the plots for **S1** and **S2** shown in Fig. S4 are not linear, indicating aggregation or
14 insufficiently low q range.

15 With due consideration for the limitations associated with the analysis of SAXS data
16 alluded to above, we analyzed the present results using the global unified scattering function
17 [62-64] provided in SansView software (v. 2.1, University of Tennessee)². This approach
18 allows for both Guinier and power-law contributions to the scattering curve, and for **S1** and
19 **S2**, two sets of (R_g, d) parameters were found to describe the scattering intensity over the q
20 range studied. The global unified model fitting parameters obtained from SansView are given
21 in Table 4.

22

23

² Later versions of this software are known as SasView.


Table 4. Fitting parameters from SAXS data based on the global unified model.

	S1 concentration		S2 concentration
	2.03 g/L	11.9 g/L	1.59 g/L
$R_g(1)$ (nm)	9.07	11.4	8.95
$R_g(2)$ (nm)	46.6 *	51.5 *	47.5 *
$d(1)$	1.22	1.25	1.72
$d(2)$	2.82	2.98	2.84

*These should be regarded as minimum sizes, as they will be limited by the lowest accessible q value.

On this basis, the fitting data indicate two scattering regions. At high q , the R_g value of ~ 10 nm and low Porod exponents for **S1**, suggest that the scattering units responsible are semi-flexible chains (with $1 < d < 5/3$) [62,65]. For the **S2** solution, the higher d is characteristic of a mass fractal (for which $5/3 < d < 3$) with a swollen chain structure [65] and the Porod exponent as defined above corresponds to the fractal dimension. Additionally, at low q , the larger scattering units of at least 50 nm for both **S1** and **S2** are also consistent with mass fractals, but here they consist of clustered network structures, for which $d \approx 3$ [65]. Fig. 7 contains a simple schematic representation of the solution structure of the two soluble fractions in solution, based on these SAXS characteristics. Thus, at the molecular level (high q), scattering arises from **S1** and **S2** chains (low d) with radii of gyration ~ 10 nm. This is a value similar to the characterization length of 7.8 nm obtained by Shang and Rice for HA dissolved in alkaline solution [66]. Probing the solution structure on a greater length scale at low q identifies larger mass fractal scattering units with $R_g \sim 50$ nm, although here, this value will be governed by the lowest q value used in the analysis since a true Guinier region has not been attained. It can be seen from Fig. 6, that these dimensions are not reflected in the AFM image height (z -axis), mainly as a result of the relatively concentrated solutions saturating the mica surfaces, thereby obscuring the $z = 0$ position. On the other hand, the section taken in

1 the xy plane, also shown in Fig. 6, indicates the spatial dimensions of assumed close-packed
 2 adsorbed units to be ~ 100 nm and ~ 50 nm for 0.5 and 6 g/L, respectively. This is consistent
 3 with results from other AFM [54] and TEM [57] studies. Based on Porod log-log fits to
 4 scattering data, other workers have obtained d values between 2.2-2.4 for different humic
 5 substances in alkaline solution [66-68]. Osterberg and Morton also estimated from a SANS
 6 analysis an average cluster size of ~ 50 nm for two different HAs, and an R_g value of 32 nm
 7 for one of them [68].



8
 9 **Fig. 7.** SAXS Porod plots for S1 (upper) and S2 (lower) solutions at the indicated concentrations in deionized
 10 water. The solid lines are unified (power law and Guinier) fits to the data (see text for details). The lower plot
 11 also contains schematic representations of the length scales probed by the SAXS measurements.

1

2 **3.4. Implications for the colloidal structure of HA**

3 Characterization of the solubility fractions in the present work has revealed very significant
4 structural differences between the soluble and insoluble fractions on the basis of TGA,
5 elemental analysis, and infrared and solid-state ^{13}C NMR spectroscopy. The latter
6 spectroscopic data indicate that **S1** and **S2** are structurally similar, with molecular weight and
7 acid/base properties [69] likely to account for their respective solubility behavior. However,
8 differences between **S1** and **S2** and their insoluble counterparts **I1** and **I2** are more
9 significant. CPMAS ^{13}C NMR spectroscopy reveals higher aromaticity in the soluble species.
10 This is reinforced by TGA results which indicate relatively higher aromatic content in the
11 soluble fractions, and infrared spectroscopy which identifies lower levels of polysaccharide in
12 **S1** and **S2**.

13 As described earlier, there have been numerous fractionation studies on HAs, and the
14 specific species obtained from a particular HA would be expected to be dependent on the
15 fractionation method used. Herein, HA fractionation on the basis of water solubility has led
16 us to some considerations regarding HA structure, but it also provides further evidence for
17 their colloidal nature. Although the present findings will apply particularly to the commercial
18 HA sample studied, the wider application to other humic substances should also be
19 considered. Soluble HA fractions at near-neutral pH conditions are likely to have important
20 functions in soils, for example, including the regulation of ionic composition and determining
21 the surface chemistry of minerals.

22 As already considered above, the structure of humic substances in general, and HAs in
23 particular, has been the subject of debate over the years. The earliest models regarded humic
24 materials as random coil ionic polymers, the conformation of which responded to their

1 environmental conditions of pH and electrolyte concentration (e.g. [70]). Subsequently,
2 Wershaw [71,72] suggested a “membrane-micelle” model in which mixed micelles are
3 formed from an ordered aggregation of plant-derived lipids and charged amphiphilic
4 molecules held together by hydrogen bonding and π - π interactions. More recently, a
5 consensus is developing that humic substances are supramolecular systems [22,23]. The
6 building blocks involved in such suprastructures originate from smaller heterogeneous
7 species derived from aromatic lignins, polysaccharides, proteins and lipids [22,73], evidence
8 for which is present in the fractions derived in the present work. Each of these components
9 exhibits different polarity/solubility characteristics, but their structures allow for mutual
10 interactions, as proposed, for example, in the lignin-carbohydrate complex (LCC) model [74]
11 and through interactions between lipids and “humic-like fractions”, creating what the authors
12 term a “structure-within-a-structure architecture” [13,75].

13 Through its behavior in water, the HA sample of the present study has been shown, both
14 here and elsewhere [69], to be a multi-component mixture of species classes which are not
15 covalently bonded, also supporting separation on the basis of adsorption [21] and
16 ultrafiltration [19,20].

17 The colloidal character of HAs has been identified by AFM by a number of workers [1,54-
18 59] although to the best of our knowledge this is the first time that such an HA sub-fraction
19 has been imaged by AFM. The topological features of the **S1** fraction indicates a relatively
20 small size for adsorbed units on mica, which is seen to be concentration dependent (Fig. 6),
21 with the higher concentration producing the smaller-sized units. In part, the SAXS analyses
22 agree with the AFM findings based on data at low q , from which the scattering units in **S1**
23 and **S2** are suggested to have radii of gyration of at least 50 nm. In the high q region, the
24 scattering from $R_g \sim 10$ nm units is consistent with the characteristic length determined by

1 Shang and Rice for HA molecules in alkaline solution [66]. Here, we propose that this latter
2 dimension is characteristic of the constituents of the larger aggregated structures.

3 We have also focused mainly on the colloidal properties of **S1**, which has the highest water
4 solubility. In respect of its sensitivity towards flocculation in the presence of indifferent
5 cations, **S1** exhibits the classical properties of an anionic colloid, as also found for a purified
6 HA sample in the presence of Mg^{2+} by Wang et al. [76].

7

8 **4. CONCLUSIONS**

9

10 The nature of humic substances continues to be the subject of ongoing investigations, but it
11 is only relatively recently that the supramolecular association of heterogeneous species has
12 been generally accepted as defining their structural organization [22,23]. The result is that
13 humic substances are multicomponent colloidal systems, the dispersed phase comprising a
14 diverse range of self-assembled components, building on the membrane-micelle model of
15 Wershaw [71,72]. In the present work we have fractionated a commercial HA sample by
16 sequential dissolution in deionized water and shown that one of the sub-fractions, in
17 particular, behaves as a hydrophilic anionic colloid. AFM and SAXS measurements suggest
18 an approximate size for the colloidal dimensions. The results lend some support for Lehmann
19 and Kleber's "emergent view" of the nature of humic substances inasmuch as we have
20 identified water-soluble "forms that are actually soluble in water" [11]. The results also
21 provide a link with the earlier "polymer model" in which humic substances were considered
22 as ionic polymers [77].

23 We have used a multi-technique approach to the study of the composition and colloidal
24 properties of a commercial sample of HA. As reported previously by others [24,25], we also

1 observed unusual solubility behavior of this HA sample, which enabled us to isolate four
2 fractions by sequential dissolution under near-neutral pH conditions – two water-soluble
3 products (**S1** and **S2**) and two water-insoluble residues (**I1** and **I2**). Although not carried out
4 here, it is possible that a more exhaustive separation scheme, perhaps including pH and ionic
5 strength variation, would lead to a greater number of narrower class fractions as are being
6 studied by Klučáková and her coworkers [78-81].

7 The analytical results highlight significant differences between the soluble and insoluble
8 HA fractions. The former contain proportionally more oxygen (revealed by a combination of
9 elemental analysis and TGA), with increased carbonyl functionality ($>C=O$ evident from ^{13}C
10 CPMAS NMR and ATR-IR) and phenol groups (aromatic C-OH from ATR-IR).
11 Additionally, the soluble fractions contain a greater aromatic/aliphatic carbon ratio (from ^{13}C
12 CPMAS NMR and TGA) and a lower polysaccharide C-O content (ATR-IR). Thus, the
13 components identified here are consistent with known structural features in all HAs.

14 As pointed out by Tarasevich et al. [1], to date there have been relatively few systematic
15 investigations of the effects of concentration on the form taken by HA in solution. However,
16 using a simple dissolution approach, we have shown that the original commercial humic acid
17 used in the present study contains non-covalently bonded species with different polarity and
18 water solubility. In this way, the present findings concur with the description of HA as a
19 supramolecular system, with macromolecules as well as molecular aggregates being present,
20 as concluded by Baigorri et al. [82].

21

22 **ACKNOWLEDGEMENTS**

23

24 We are grateful to BP America for supporting WMS under the EngD CDT program in
25 Materials Technology, as well as for establishing the CPSC at the University of Surrey. The

1 EPSRC Solid-State NMR Service facility at Durham University is thanked for the ¹³C
2 CPMAS analysis. We acknowledge Dr Adam Squires, University of Bath, for facilitating the
3 SAXS measurements at the Diamond Light Source and we are also indebted to Violeta
4 Doukova, Rifat Shaikh and Judy Peters for their expert assistance with TGA, AFM and
5 microanalysis, respectively. Finally, we are indebted to several referees for constructive
6 comments.

7

8 REFERENCES

- 9 [1] Y.I. Tarasevich, S.A. Dolenko, M.Y. Trifonova, E.Y. Alekseenko, *Colloid J.* 75
10 (2013) 207-213.
- 11 [2] R. von Wandruszka, *Geochem. Trans.*, 1 (2000) 10-15.
- 12 [3] L. Gutierrez, M. Pawlik, *Int. J. Min. Proc.*, 126 (2014) 117–125.
- 13 [4] L. Gutierrez, M. Pawlik, *Int. J. Min. Proc.*, 126 (2014) 126–135.
- 14 [5] J.L. Margeson, V. Hornof, G.H. Neale, *J. Can. Pet. Tech.*, 28 (1989) 57-62.
- 15 [6] L.S. Kotlyar, B.D. Sparks, H. Kodama, P.E. Grattan-Bellew, *Energy Fuels*, 2 (1988)
16 589-593.
- 17 [7] M. Ishiguro, W. Tan, L.K. Koopal, *Colloids Surfaces, A* 306 (2007) 29.
- 18 [8] I. Christl, R. Kretzschmar, *Environ. Sci. Technol.*, 41 (2007) 1915-1920.
- 19 [9] L.K. Koopal, T.P. Goloub, T.A. Davis, *J. Colloid Interface Sci.* 275 (2004) 360-367.
- 20 [10] P. Conte, A. Agretto, R. Spaccini, A. Piccolo, *Environ. Poll.*, 135 (2005) 515–522.
- 21 [11] J. Lehmann, M. Kleber, *Nature*, 528 (2015) 60-68.

- 1 [12] T.F. Guetzloff, J.A. Rice, *Science Total Environ.*, 152 (1994) 31–35.
- 2 [13] G. Chilom, A.S. Bruns, J.A. Rice, *Org. Geochem.*, 40 (2009) 455–460.
- 3 [14] C. Young, R. von Wandruszka, *Geochem. Trans.*, 2 (2001) 16-20.
- 4 [15] L.M. Yates, R.R. Engebretson, T.J. Haakenson, R. von Wandruszka, *Anal. Chim.*
5 *Acta*, 356 (1997) 295-300.
- 6 [16] A. Nebbioso, A. Piccolo, *Anal. Bioanal. Chem.*, 405 (2013) 109–124.
- 7 [17] H.K.J. Powell, R.M. Town, *Anal. Chim. Acta*, 267 (1992) 47–54.
- 8 [18] P. Conte, R. Spaccini, D.Smejkalova, A. Nebbioso, A. Piccolo, *Chemosphere*, 69
9 (2007) 1032–1039.
- 10 [19] L. Carlsen, P. Lassen, P. Warwick, A. Randall, *Chemosphere*, 33 (1996) 659-670.
- 11 [20] O. Francioso, S. Sánchez-Cortés, D. Casarini, J.V. Garcia-Ramos, C. Giavatta, C.
12 Gessa, *J. Mol. Struct.* 609 (2002) 137-147.
- 13 [21] A. Pitois, L.G. Abrahamsen, P.I. Ivanov, N.D. Bryan, *J. Colloid Interface Sci.*, 325
14 (2008) 93–100.
- 15 [22] A. Piccolo, *Soil Sci.*, 166 (2001) 810-832.
- 16 [23] R. Sutton, G. Sposito, *Environ. Sci. Technol.*, 39 (2005) 9009-9015.
- 17 [24] M. Klučáková, M. Pekař, *Colloids Surfaces, A* 252 (2005) 157–164.
- 18 [25] M. Klučáková, M. Pekař, *Colloids Surfaces, A* 318 (2008) 106–110.
- 19 [26] C. Gamboa, A.F. Olea, *Colloids Surfaces, A* 278 (2006) 241–245.
- 20 [27] M. De Nobili, M. Contin, L. Leila, *Can. J. Soil Sci.*, 70 (1990) 531-536.
- 21 [28] R.L. Johnson, K. Schmidt-Rohr, *J. Magn. Reson.*, 239 (2014) 44–49.

- 1 [29] R.M.B.O. Duarte, A. C. Duarte, *Atmospheric Environ.*, 42 (2008) 6670–6678.
- 2 [30] V.I. Esteves, A.C. Duarte, *Marine Chem.*, 63 (1999) 225–233.
- 3 [31] R.L. Malcolm, P. MacCarthy, *Environ. Sci. Technol.*, 20 (1986) 904–911.
- 4 [32] F.J. Stevenson, K.M. Goh, *Geochim. Cosmochim. Acta*, 35 (1971) 471–483.
- 5 [33] A.L. Mafra, N. Senesi, G. Brunetti, A.A.W. Miklós, A.J. Melfi, *Geoderma*, 138
6 (2007) 170–176.
- 7 [34] J. Polak, M. Bartoszek, M. Ządło, A. Kos, W.W. Sułkowski, *Chemosphere*, 84 (2011)
8 1548–1555.
- 9 [35] M. Tatzber, M. Stemmer, H. Spiegel, C. Katzlberger, G. Haberhauer, A. Mentler,
10 M.H. Gerzabek, *J. Plant Nutr. Soil Sci.*, 170 (2007) 522–529.
- 11 [36] A. Piccolo, F.J. Stevenson, *Geoderma*, 27 (1982) 195–208.
- 12 [37] D.C.W. Tsang, N.J.D. Graham, I.M.C. Lo, *Chemosphere*, 75(2009) 1338–43.
- 13 [38] P. Innocenzi, *J. Non-Cryst. Solids*, 316 (2003) 309–319.
- 14 [39] M. Fuentes, R. Baigorri, G. González-Gaitano, J.M. García-Mina, *Org. Geochem.*, 38
15 (2007) 2012–2023.
- 16 [40] F. Monteil-Rivera, E.B. Brouwer, S. Masset, Y. Deslandes, J. Dumonceau, *Anal. Chim.*
17 *Acta*, 424 (2000) 243–255.
- 18 [41] R.L. Johnson, K. Schmidt-Rohr, *J. Magn. Res.*, 239 (2014) 44–49.
- 19 [42] A. Majid, J.A. Ripmeester, *Fuel*, 69 (1990) 1527–1536.
- 20 [43] H.-S. Shin, J.M. Monsallier, G.R. Choppin, *Talanta*, 50 (1999) 641–647.
- 21 [44] N.A. Wall, G.R. Choppin, *Appl. Geochem.*, 18 (2003) 1573–1582.

- 1 [45] K. Hayase, H. Tsubota, *Geochim. Cosmochim. Acta*, 47 (1983) 947–952.
- 2 [46] M. Terashima, M. Fukushima, S. Tanaka, *Colloids Surfaces, A* 247 (2004) 77–83.
- 3 [47] R.R. Engebretson, R. Von Wandrusrka, *Environ. Sci. Technol.*, 28 (1994) 1934–1941.
- 4 [48] A. Piccolo, D. Smejkalova, *Environ. Sci. Technol.*, 42 (2008) 699–706.
- 5 [49] M. Fukushima, S. Tanaka, K. Hasebe, M. Taga, H. Nakamura, *Anal. Chim. Acta*, 302
6 (1995) 365-373.
- 7 [50] J. Lyklema, *Colloids Surfaces, A* 460 (2014) 468–472.
- 8 [51] J. Lyklema, *J. Colloid Interface Sci.*, 392 (2013) 102–104.
- 9 [52] K. Furukawa, Y. Takahashi, *Chemosphere*, 73 (2008) 1272–1278.
- 10 [53] T.A. Sabirjanovna, S. Pasa, D.U. Zhumasilovich, H. Temel, N.G. Oryntayevna, *Desal.*
11 *Water Treatment*, 57 (2016) 776–790.
- 12 [54] M. Pläschke, J. Römer, R. Klenze, J.I. Kim, *Colloid Surfaces, A* 160 (1999) 269–279.
- 13 [55] A. Liu, R.C. Wu, E. Eschenazi, K. Papadopoulos, *Colloids Surfaces, A* 174 (2000)
14 245–252.
- 15 [56] J. I. Kim, G. Buckau, G. H. Li , H. Duschner, N. Psarros, *Fresenius J. Anal. Chem.*,
16 338 (1990) 245-252.
- 17 [57] S.C.B. Myneni, J.T. Brown, G.A. Martinez, W. Meyer-Ilse, *Science*, 286 (1999)
18 1335-1337.
- 19 [58] C. Chen, X. Wang, H. Jiang, W. Hu, *Colloids Surfaces, A* 302 (2007) 121-125.
- 20 [59] E. Balnois, K.J. Wilkinson, J.R. Lead, J. Buffle, *Environ. Sci. Technol.*, 33 (1999)
21 3911-3917.
- 22 [60] Z. Liu, Y. Zu, R. Meng, Z. Xing, S. Tan, L. Zhao, T. Sun, Z. Zhou, *Microscopy*
23 *Microanal.*, 17 (2011) 1015-1021.
- 24 [61] L.T. Sein, J.M. Varnum, S.A. Jansen, *Environ. Sci. Technol.*, 33 (1999) 546-552.

- 1 [62] G. Beaucage, *J. Appl. Cryst.* 28 (1995) 717-728.
- 2 [63] G. Beaucage, *J. Appl. Cryst.* 29 (1996) 134-146.
- 3 [64] G. Beaucage, H.K. Kammler, S.E. Pratsinis, *J. Appl. Cryst.* 37 (2004) 523-535.
- 4 [65] B. Hammouda, *J. Appl. Cryst.* 43 (2010) 1474-1478.
- 5 [66] C. Shang, J.A. Rice, *J. Colloid Interface Sci.*, 305 (2007) 57–61.
- 6 [67] J.A. Rice, E. Tombàcz, K. Malekani, *Geoderma*, 88 (1999) 251-264.
- 7 [68] R. Osterberg, K. Mortensen, *Eur. Biophys. J.*, 21 (1992) 163-167.
- 8 [69] M. Klučáková, *React. Functional Polymers*, 109 (2016) 9–14.
- 9 [70] M.M. Kononova, *Soil Organic Matter. Its Nature, Its Role in Soil Formation and in*
10 *Soil Fertility*. Pergamon Press, New York, 1961.
- 11 [71] R.L. Wershaw, *Env. Health Perspect.*, 83 (1989) 191-203.
- 12 [72] R.L. Wershaw, *US Geological Survey Water-Supply Paper*, 1994, Vol.2410.
- 13 [73] K.H. Tan, *Humic Matter in Soil and the Environment. Principles and Controversies*,
14 *CRC Press, Boca Raton, 2nd Edition, 2014, Chaps. 2 and 5.*
- 15 [74] C. Colombo, G. Palumbo, R. Angelico, H.G. Cho, O. Francioso, A. Ertani, S. Nardi,
16 *Chemosphere*, 138 (2015) 821–828.
- 17 [75] G. Chilom, A. Baglieri, C.A. Johnson-Edler, J.A. Rice, *Org. Geochem.*, 57 (2013)
18 119–126.
- 19 [76] L.-F. Wang, L.-L. Wang, X.-D. Ye, W.-W. Li, X.-M. Ren, G.-P. Sheng, H.-Q. Yu,
20 X.-K. Wang, *Environ. Sci. Technol.*, 47 (2013) 5042–5049.
- 21 [77] F.J. Stevenson, *Humus Chemistry: Genesis, Composition, Reactions*, 2nd Edition,
22 *John Wiley & Sons, Ltd, New York, 1994.*
- 23 [78] M. Klučáková, R. Kolajová, *React. Funct. Polymers*, 78 (2014) 1–6.
- 24 [79] M. Klučáková, M. Kalina, *J. Soils Sediments*, 15 (2015) 1900-1908.
- 25 [80] M. Klučáková, *Environ. Sci. Pollut. Res*, 23 (2016) 7722–7731.

- 1 [81] M. Klučáková, *React. Funct. Polymers*, 109 (2016) 9–14.
- 2 [82] R. Baigorri, M. Fuentes, G. González-Gaitano, J.M. García-Mina, *Colloids Surfaces,*
- 3 *A* 302 (2007) 301–306.
- 4