

RESEARCH ARTICLE

Open Access

Characterization and gene expression analysis of the *cir* multi-gene family of *plasmodium chabaudi chabaudi* (AS)

Jennifer Lawton¹, Thibaut Brugat¹, Yam Xue Yan², Adam James Reid³, Ulrike Böhme³, Thomas Dan Otto³, Arnab Pain^{3,4}, Andrew Jackson³, Matthew Berriman³, Deirdre Cunningham¹, Peter Preiser² and Jean Langhorne^{1*}

Abstract

Background: The *pir* genes comprise the largest multi-gene family in *Plasmodium*, with members found in *P. vivax*, *P. knowlesi* and the rodent malaria species. Despite comprising up to 5% of the genome, little is known about the functions of the proteins encoded by *pir* genes. *P. chabaudi* causes chronic infection in mice, which may be due to antigenic variation. In this model, *pir* genes are called *cirs* and may be involved in this mechanism, allowing evasion of host immune responses. In order to fully understand the role(s) of CIR proteins during *P. chabaudi* infection, a detailed characterization of the *cir* gene family was required.

Results: The *cir* repertoire was annotated and a detailed bioinformatic characterization of the encoded CIR proteins was performed. Two major sub-families were identified, which have been named A and B. Members of each sub-family displayed different amino acid motifs, and were thus predicted to have undergone functional divergence. In addition, the expression of the entire *cir* repertoire was analyzed via RNA sequencing and microarray. Up to 40% of the *cir* gene repertoire was expressed in the parasite population during infection, and dominant *cir* transcripts could be identified. In addition, some differences were observed in the pattern of expression between the *cir* subgroups at the peak of *P. chabaudi* infection. Finally, specific *cir* genes were expressed at different time points during asexual blood stages.

Conclusions: In conclusion, the large number of *cir* genes and their expression throughout the intraerythrocytic cycle of development indicates that CIR proteins are likely to be important for parasite survival. In particular, the detection of dominant *cir* transcripts at the peak of *P. chabaudi* infection supports the idea that CIR proteins are expressed, and could perform important functions in the biology of this parasite. Further application of the methodologies described here may allow the elucidation of CIR sub-family A and B protein functions, including their contribution to antigenic variation and immune evasion.

Background

An important feature of the malaria parasite is the export of proteins to the surface of infected red blood cells (iRBCs). Surface proteins or variant surface antigens (VSA) have been identified so far in three species infecting humans: *Plasmodium falciparum*, *P. vivax* and *P. knowlesi*, as well as in the rodent malaria parasites *P. chabaudi* and *P. yoelii* [1-5]. These proteins are implicated in antigenic variation and immune evasion, as well

as parasite accumulation or sequestration in host tissues; features which may be critical in determining the outcome of malaria reviewed by [6,7]. Since VSAs are recognized by antibodies, they are also likely to be important targets for a protective immune response eg. [8,9].

In most *Plasmodium* species VSAs are encoded by multi-gene families, usually located in sub-telomeric chromosome regions [2,10-12]. In *P. falciparum*, several VSA gene families have been identified such as *var*, *rif*, *stevor* and *surf* reviewed by [7]. Among them, the best characterized is the *var* gene family encoding PfEMP1 proteins, which is implicated in both antigenic variation

* Correspondence: jlangho@nimr.mrc.ac.uk

¹Division of Parasitology, MRC National Institute for Medical Research, London, UK

Full list of author information is available at the end of the article

and sequestration [1,13]. This family consists of 60 genes that can be grouped into several families (A-E) according to chromosomal location, coding and non-intergenic sequences, direction of transcription, and domain arrangements [14,15]. Similarly, the *rif* genes, the largest multi-copy gene family in *P. falciparum* (circa 150-200 genes), has been divided into subgroups A and B on the basis of sequence similarity [16,17]. A- and B-type RIFINs have different sub-cellular localizations, in that only the A-type RIFINs appear to be exported towards the surface of iRBCs [16]. Thus, A-type RIFINs may be more likely to play a role in the host/parasite relationship during the blood stages of *P. falciparum*.

Plasmodium interspersed repeat (*pir*) genes have been identified in *Plasmodium vivax* (*vir*), *P. knowlesi* (*kir*), *P. berghei* (*bir*), *P. chabaudi* (*cir*), and *P. yoelii* (*yir*) [2,18,19]. Together, *pirs* form the largest multi-gene family identified to date in *Plasmodium* species, and occupy up to 5% of the parasite genome. On completion of the *P. vivax* Salvador I genome sequence, a total of 12 *vir* sub-families were identified [20-22]. Similarly, 5 sub-families have been identified within the *yir* repertoire [23]. However, so far no function has been ascribed to the subgroups of *yir* and *vir* families.

Microarray studies carried out on *P. vivax* and *P. yoelii* suggest that approximately 50% of the *yir* and *vir* repertoires are expressed in a population of iRBCs during infection, with no evidence of preferential transcription according to their chromosomal location or phylogenetic sub-groups [2,4,20,24]. Surprisingly, considering the large proportion of *yir* genes transcribed in the parasite population, only 1-3 *yirs* are transcribed in a single iRBC [4]. This suggests that transcription of these genes is under tight control. The function(s) of PIR proteins, however, remain unknown.

Rodent malaras offer the only feasible system in which the contribution of PIR proteins to immune evasion and to sequestration and pathology can be examined. Of these, *P. chabaudi* is the only rodent species that naturally produces a chronic infection in mice. These chronic infections are likely to be perpetuated by antigenic variation [25-27]; however, the antigens involved have not yet been determined. *P. chabaudi* also exhibits other important features observed in human *Plasmodium* infections, including rosetting and adhesion to host endothelial cells [5,28], and thus is an ideal model in which to investigate the role of PIR proteins.

The *cir* genes comprise the largest gene family in *P. chabaudi*, and are located in sub-telomeric regions of chromosomes [10,29]. Recently, restriction fragment length polymorphism (RFLP) analysis indicated that *cir* genes may have tissue specific patterns of expression

[30]. However, as such techniques are not gene specific, they are only able to provide clues about general *cir* expression. Furthermore, very little is known about the function of CIRs in stimulating or evading host immunity during *P. chabaudi* infection.

Here we describe two major sub-families within the *cir* repertoire of the recently completed *P. chabaudi* AS genome. Using an approach similar to that applied to the RIFIN repertoire [17], we found both conserved and sub-family specific amino acid motifs, and predict functional divergence between the proteins from different CIR sub-families.

We have analyzed in detail the expression of the entire *cir* repertoire via RNA sequencing (Illumina RNA-seq) and microarray. Up to 40% of the *cir* gene repertoire was expressed in the parasite population during infection, and dominant *cir* transcripts could be identified, with some differences in the pattern of expression between the *cir* subgroups. Finally, we found specific *cir* genes were expressed at different time points during asexual blood stages. Together these data will allow future investigation of the CIR family to elucidate their roles in the host/pathogen relationship.

Methods

Ethics statement

This study was carried out in strict accordance with the UK Animals (Scientific Procedures) Act 1986 and was approved by the Ethical Committee of the MRC National Institute for Medical Research, and the British Home Office (PPL: 80/2538).

Annotation of *cir* genes

cir genes were annotated using Artemis release 11 [31,32] onto the eight-fold coverage assembly of the *Plasmodium chabaudi chabaudi* AS genome [33]. Putative coding sequences containing conserved features of previously published *cir* genes identified from the three-fold coverage genome assembly [34]. These were then searched for similarity to the *pir* superfamily genes via a combination of basic local alignment search tool (BLAST) and Hidden Markov Model (HMM) on PIR super-family proteins [CIR_BIR_YIR (PF06022)], available in the PFAM database [35], following previously described methods [35,36].

Detection of conserved motifs

Motif Elicitation analysis (MEME, [37]) was used to identify up to 20 conserved amino acid motifs within the CIR repertoire. The average motif locations were identified and plotted onto each gene. WebLogos were generated with the MEME program, using an adaptation of the WebLogo software [38].

Analysis of sequence similarity

Amino acid sequences of 183 *cir* genes were aligned using the Multiple Sequence Comparison by Log-Expectation algorithm (Muscle, [39]). Sequences aligning poorly with the other CIRs were excluded from the alignment and regions containing large insertions were deleted (Additional file 1). The sequence similarity along the alignment is attached in Additional file 2.

Since phylogenetic trees only represent a bifurcating lineage and imply phylogenetic relationships, reticulate networks were created within the program Splitstree 4.0 [40]. All networks used the algorithms NeighborNet [41] for calculation of distances and Equal angle [42] for calculation of splits. 500 bootstrap replicates were generated (Additional file 3).

To support the network, a phylogenetic tree was also constructed using the Maximum Likelihood method from the PhyML server [43]. Here, the evolutionary model applied was Le Gascuel (LG) [44], and the branch support was calculated by approximate likelihood ratio test (aLRT) [45]. Three YIR [46] and three BIR sequences (Ulrike Böhme, WTSI, personal communication) were added to the CIR alignment to allow a root to be placed within this tree (Additional file 4).

Clades identified by both methods with high branch support values contained highly similar CIR sequences. Small clades within each major sub-family were denoted numerically: A1-A5 and B1-B4. Members of each clade are tabulated in Additional file 5.

Detection of phylogenetic incompatibilities between *cir* genes

Phylogenetic incompatibilities within the alignment of 183 CIRs, and each identified sub-family, were analyzed using the pairwise homoplasy index (PHI) in Splitstree v4.0 [40].

Phylogenetic profiling was used to detect phylogenetic inconsistencies between four *cir* DNA sequences, selected at random, using a hidden Markov model method within the TOPALi platform v2.5 [47,48]. The probability of generating each of the three possible tree topologies for the four sequences was modelled in a given 100 nucleotide window. Possible recombination breakpoints were identified where the most probable topology altered at different positions along the alignment.

Five *cir* quartets, chosen at random, were analyzed per clade by each method described above (Additional file 6).

Function shift analysis

The alignment of 183 CIRs was split into two files containing only A- and B-type CIRs. The two alignments were then used to apply the 'FunShift' methodology [49], and predict whether these 2 groups of proteins

may perform different functions. Positions containing only gaps in a subfamily were not counted.

The method used was the same as described previously by Abhiman and Sonnhammer [49] with a few modifications. Rate-Shifting Sites (RSS) were defined as positions conserved in one sub-family but variable in the other, and were identified using the likelihood ratio test (LRT) program [50]. The U-values generated by this program indicate the likelihood of rate change for each position in the alignment between the two sub-families. U-values above 4.0 were considered significant at the 5% significance level, as previously described [50].

Conservation-Shifting Sites (CSS) were defined as positions that were conserved in both groups, but containing different residues in each. CSS were detected using the method developed by Abhiman and Sonnhammer [49]. This calculates a Z-score based on the normalized cumulative relative entropy at each position of the alignment, between the two sub-families. Z-scores exceeding 0.5 per alignment position were considered significant [49].

CSS and RSS are plotted within a sub-section of the alignment in Additional file 7.

Mice and parasites

Female BALB/c and C57BL/6 mice aged 6-8 weeks were obtained from the specific pathogen-free unit at the MRC National Institute for Medical Research (NIMR), London. For experimental purposes, mice were housed conventionally with sterile bedding, food and irradiated water on a 12 hour light-dark cycle.

A cloned line of *Plasmodium chabaudi chabaudi* (AS) was used in this study [51]. Stabilates were cryo-preserved in blood from BALB/c mice. To obtain parasites for experimental infection, an aliquot of the stabilate was injected intraperitoneally (i.p.) into immunodeficient BALB/c *RAG2*^{-/-} mice [52]. Blood was taken from the donor mice 7 days after infection and experimental mice were infected by injecting 10⁵ infected erythrocytes i.p. Parasitaemia was monitored by examination of Giemsa-stained blood films as previously described [53].

Blood was collected from each mouse by cardiac puncture under terminal anaesthesia into Krebs saline (114 mM NaCl, 4.57 mM KCl, 1.15 mM MgSO₄) containing 0.2% glucose and 25 U/ml heparin (Leo Pharmaceuticals) 7 days after infection. Leukocytes were removed via Plasmodipur filtration (Euro-Diagnostica) according to manufacturer's instructions. Blood was then stored at -80°C in TRIZOL reagent (Invitrogen) for subsequent RNA extraction.

RNA extraction

RNA was extracted from *P. chabaudi* infected blood samples by guanidinium thiocyanate-phenol-chloroform

extraction according to standard methods [54] and DNase digested using Turbo DNase (Ambion) according to the manufacturer's instructions.

Microarray hybridization and analysis

P. chabaudi AS is a highly synchronous parasite for which development in the blood follows its host's circadian rhythm. Twelve time-points were then collected; one every two hours, to cover the entire 24 h cycle of blood stage development. At the peak of parasitaemia, one mouse was sacrificed at each time point and thin blood films were made and stained with Giemsa for optical microscopy. The pan-rodent microarray was designed using the OligoRankPick program as previously described [55]. The RNA preparation, Cy-dye coupling to cDNA, hybridization and slide scanning were performed as described by Bozdech and colleagues [56].

Data processing and analysis (including the Fast Fourier Transform) were carried out as described by Bozdech and colleagues [24]. The phaseogram (Figure 7) contains genes with $> 1.7 \log_2$ ratio of change in mRNA abundance across the IDC. A list of identified genes can be found in Additional file 8.

RNA sequencing

10 μg of *P. chabaudi* total RNA obtained from BALB/c and C57BL/6 mice was used for this analysis. RNA sequencing was performed using an Illumina GAIIx following the methodology described by Otto and colleagues [57] and was used to create 76 bp paired-end reads. TopHat [58] was used to map reads against the *Plasmodium chabaudi chabaudi* AS reference genome [33], with maximum intron size set to 10000 and inner-mate distance set to 100. Gene expression levels (RPKM) were calculated as defined by Mortazavi and colleagues [59] with minor modifications. Non-uniquely mapping reads were excluded and read-length windows of protein-coding regions that were non-unique were excluded from the gene length term used in calculating RPKM.

Each sequencing run contained different amounts of noise (eg. reads mapping to generally unexpressed parts of the genome: introns and intergenic regions). Therefore we calculated RPKMs over 500 bp windows of exonic and intronic sequences on chromosome 14. For each sequencing run, we took an RPKM cutoff above which only 10% of intron sequences were expressed. A list of identified genes can be found in Additional file 9, with the threshold calculation in Additional file 10. Legends for Additional files 1-10 can be found in Additional file 11.

A Kolmogorov-Smirnov test [60] was used to compare the distribution of *cir* sub-groups according to their level of expression.

Results

I) Bioinformatic analysis of the CIR multi-gene family

a) Identification of *cir* genes

Completion of the *P. chabaudi* AS genome sequencing and revised assembly allowed the initial *cir* annotation to be manually revisited. Conserved features from the initially identified *cir* genes [18,29], such as the relative exon lengths, splice sites and amino acid sequences were used to identify putative *cir* coding sequences in the assembled contigs. 117 *cir* genes were first identified and manually annotated. A hidden Markov model (HMM) was then constructed on the basis of the identified *cir* repertoire and was used to detect more divergent *cir* genes, bringing the total of identified *cirs* to 196 [33], including 3 additional *cir*-like genes, containing some but not all expected features of *cir* family members.

Only three partial *cir* genes were identified, arising from the few unresolved contig assemblies for the *P. chabaudi* AS genome. Three long *cir* genes were also identified, containing an extended first exon. The majority of *cir* genes contained one predicted TM domain including some of the divergent *cirs*.

b) Sequence similarity of CIR proteins

In order to investigate similarity between CIRs, the amino acid sequences identified during *cir* gene annotation were aligned using Muscle [39], and refined manually (Additional file 1a). Upon addition of more divergent *cir* genes to the repertoire, 16 sequences aligned poorly with the majority of CIRs and were excluded from the analysis (tabulated in Additional file 1b). These included the three partial CIRs, eight CIR sequences with low C- or N-terminal similarity to other members of the repertoire, two CIRs encoded by genes with an atypical structure and three sequences which had been identified as *cir*-like genes, containing some but not all features of *cir* genes.

The sequence similarity of the alignment was determined using Plotcon [61]. The most conservation was found between amino acids 75-120 and 350-385 (Additional file 2).

To determine the relationships between CIR sequences, a network was created (Figure 1, Additional file 3), using the NeighborNet and Equal Angle algorithms [41,42] in the Splitstree program [40]. This methodology prevented bias from inferred evolutionary relationships, common to phylogenetic analyses [62]. Assumptions of linear evolution are not truly appropriate in the analysis of multi-gene family members which are likely to undergo frequent recombination, as the *var* and *sicavar* genes are known to do [19,63,64]. Instead, production of a network enables visualization of box-like structures (reticulations), where recombination may




Figure 1 Relationships between CIR amino acid sequences. Similarities between CIR sequences were visualized using a network created in SplitsTree4 [40], using the NeighborNet and Equal Angle algorithms [41,42]. This network is shown without branch support values, for ease of view. Clades within each major sub-family are indicated by red (sub-family A) or blue (sub-family B) brackets.

have occurred between *cirs* and linear evolution cannot be assumed (reviewed by [62]).

Figure 1 shows two distinct sub-families, each of which comprises multiple smaller clades, the members of which are shown in Additional file 5. The large sub-family A contained more divergent CIR sequences, many of which were identified by the CIR HMM. Five clades containing more than five CIR sequences were present within this sub-family: A1-A5 (highlighted in red, Figure 1), of which clade A1 was the most distinct. Sub-family B was comprised of the most conserved CIR sequences, and could also be further sub-divided into 4 clades: B1-B4 (highlighted in blue, Figure 1). The definition of the A1 group as a part of sub-family A was further reinforced by the creation of a phylogenetic tree to support the clades identified by the network in Figure 1. The Maximum Likelihood tree contained the same sub-families and smaller clades as observed in Figure 1, even after three YIR and three BIR sequences ([46], Böhme *et al.*, unpublished) were added to the CIR alignment to enable a root to be placed (Additional file 4 TreeBase study accession URL <http://purl.org/phylo/treebase/phyloids/study/TB2:S12458> [65-67]). The clade A1 continued to cluster within CIR sub-family A, indicating that these CIR sequences were more similar to each other than to members of CIR sub-family B.

In addition, the visualization of CIR sequence relationships using network methodology indicated phylogenetic inconsistencies within the *cir* repertoire, as many box-like structures (reticulations) were present (Figure 1). Across the whole *cir* repertoire, different regions of *cir* genes were found to display different phylogenetic relationships with each other (Additional file 6). Such phylogenetic incompatibilities may arise from either data ambiguity, convergent evolution or recombination events [62], suggesting that recombination may have played a strong role in shaping the *cir* repertoire.

c) Identification of conserved amino acid motifs

The identification of conserved amino acid sequences may provide indications of possible protein function(s). Multiple Expectation maximization for Motif Elicitation analysis (MEME, [37]) identified 14 motifs in the CIR amino acid sequences. Each motif is represented in Figure 2a as a WebLogo image [38], where the height of each letter indicates the proportion of CIR sequences containing that residue.

Motifs 1 and 3 were the most conserved, being present in 98.66% of CIR sequences. Motif 1 contained the almost unanimous YK residues, corresponding to the start of the third *cir* exon and part of the predicted CIR transmembrane domain. Motif 3 was found within the second *cir* exon and contained the highly conserved sequence YAILWLSY. Motif 10, present in 91.28% of CIRs, contained some degeneracy, but a clear conserved methionine indicated the N-terminus of all CIR proteins. All CIRs possessing motif 10 also contained a cysteine six residues into the sequence. Conserved cysteine residues were also located within motifs 5, 7 and 14. Motif 10 was found at the N-terminus of almost all CIR proteins (91%). Motifs 9, 3, 11, 5, 4 and 1 were also found in members of all sub-families.

The remaining motifs appeared to have sub-family specific distributions, with motifs 2, 6 and 7 only found within sub-family B sequences. By contrast, motifs 8, 12, 13 and 14 were specific to members of CIR sub-family A. The arrangement of motifs within the amino acid sequence is indicated for a typical member of each major sub-family in Figure 2b.

Part of motif 8 (unique to sub-family A) lies within the predicted TM domain and the TM domains of A-type CIRs contain more proline residues than B-type CIRs. In addition the majority of CIRs containing more than one TM domain are found within sub-family A. Together, these differences indicate that the TM domain of CIR sub-type A proteins may have altered properties compared to the rest of the CIRs.

d) Similarities between CIRs and RIFINS and function shift analysis

The identification of two major CIR sub-families in Figure 1 resembled the *P. falciparum* RIFIN repertoire




Figure 2 Conserved amino acid motifs within CIR sequences. Amino acid motifs present within the CIR repertoire were identified by MEME analysis (Bailey and Elkan, 1994). These are shown as Weblogo images [38], A). Hydrophobic residues are shown in blue, polar, non-charged residues in green, acidic residues in pink and positively charged residues in red. E-values refer to the significance of each motif found within the CIR sequences. The frequency (percentage) of CIR sequences within the whole repertoire, clade A1 or whole sub-families A and B containing each motif is shown. Examples of motif arrangement in CIR proteins are shown, B), using PCHAS_040110 and PCHAS_070130, which belong to the sub-families A (clade A1) and B, respectively. Exon: exon boundaries in the encoding *cir* genes are indicated by dotted lines.

organization [16,17]. As the *rif* and *stevor* multi-gene families have been suggested to be distantly related to the *pir* genes [18], we compared the CIR and RIFIN repertoires.

The RIFIN sub-family A is defined by the presence of a 25 amino acid sequence, which is absent from RIFIN-B types [16,17]. An insertion sequence could be detected only in members of CIR sub-family A, which was most conserved in the clade A1. This was located between position 253 and 316 of the alignment of 183 CIRs (Additional file 7a). Furthermore, in this insertion

(displayed as a weblogo, [38]), showed some similarities with the A-type RIFIN insertion sequence (Additional file 6b and c, [16,17]). Approximately a third of the residues in each insertion sequence were hydrophobic and very few basic residues were present. Notably, both sequences included two conserved cysteine residues.

The two RIFIN sub-families have been shown to display different sub-cellular localizations [16], and thus A- and B-type RIFINs may have different functions. This hypothesis has been supported by bio-informatic analysis of the whole RIFIN repertoire in the *P. falciparum*

clones 3D7, DD2 and HB3 [17]. To investigate whether the members of each major CIR sub-type could also have functionally diverged, the alignment of 183 CIRs was split into the two major sub-families, between which Rate- and Conservation- Shifting Sites were compared (RSS and CSS, respectively), as previously described [17,49]. Briefly, RSS measures the probability that each position in the alignment has a different mutation rate in the two sub-families, whilst CSS compares the amino acid distribution between members of each sub-family. Significant CSS and RSS sites are plotted in Figure 3a. 77 RSS (15.4% of all positions) and 158 CSS (31.8% of all positions) were identified along the alignment, strongly suggesting that functional divergence between the CIR sub-families may have occurred according to the criteria generated with protein families of known function [36].

The most significant CSS and RSS sites above the threshold indicated in Figure 3a are highlighted in a section of the CIR amino acid alignment, Figure 3b. Notably, several of the sites had altered conservation of cysteine residues (for example: at site 210 a conserved cysteine residue was present in sub-family A CIRs, but the residues N, K, S, G or D could be found at this position in members of sub-family B, Table 1) or changes in residue conservation within the predicted TM domain, between residues 330-352 (at sites 330, 349 and 350, Figure 3b and Table 1). Similar shifts have also been observed between the RIFIN A and B sub-families, with four highly significant shifts in cysteine conservation,

and seven shifted sites detected within the RIFIN TM domain [17].

II) Analysis of *cir* gene expression during *plasmodium chabaudi* infection

a) RNA sequencing analysis

To establish whether the different subgroups of CIR might encode for proteins with different biological functions, we determined the patterns of *cir* gene expression during an infection. For this, mRNA was purified from *Plasmodium chabaudi* AS iRBCs at the late trophozoite stage from four BALB/c mice and sequenced using Illumina/Solexa technology [Array Express accession number: E-ERAD-25 and ENA Study Accession Number: ERP000983]. Results obtained from intergenic regions allowed the determination of a threshold of detection, above which a gene was considered significantly expressed in each sample (Additional file 10). Accordingly, the expression of up to 40% of *cir* genes could be detected in each sample analyzed (Figure 4a).

In addition, mRNA was also sequenced from parasites passaged in two C57BL/6 mice. Both the BALB/c and C57BL/6 inbred strains of mice are commonly used hosts for *P. chabaudi* infection (for example: [68-71]), thus it was important to determine whether either host genetic background could influence *cir* expression. While in the C57BL/6 samples only 17% and 12% of the *cir* repertoire were expressed above background, these lower values were at least partly the result of a higher RPKM cutoff due to relatively high expression of




Table 1 Highly significant rate and conservation shifting sites identified between the two major CIR sub-families

Position in alignment	Residues in CIR sub-family:		Z score
	A	B	
Significant conservation shifting sites (CSS):			
145	GWSRHYINLQD	K	2.24
172	I	L	2.09
176	IVLFS	Y	2.95
196	CSWKLFYIQ	FLIN	2.60
210	C	NKSGD	2.49
221	K	SLF	2.62
233	I AVTSMNYKHL	CFRYL	2.09
330	YERGVKAIR	PSALT	2.50
349	L	S	2.74
350	STKALVPYFI	L	2.21
353	WRQSIGC	F	2.79
354	RTKGAVN	DGH	2.09
356	KEMNTY	R	2.15
358	KEFTNI	QHQRN	2.44
360	K	Q	2.21
361	K	Y	2.36
363	M	R	2.93
367	I	K	2.90
368	N	K	2.42
369	L	TAIKV	2.01
Significant rate shifting sites (RSS):			
8	M	MDKTNIL	19.66
25	VNKG A	TYDVSEKNQPGI	16.86
315	EKTASQPISDRLYGN	EPFLT	19.99
316	IQDENPASGHLKVTR	SEDGQR	29.63
322	IPSVTNAKEQLCYHG	SFNTPL	45.64
366	IVLD	LRVIP	18.26

The most significant CSS ($Z > 2$) and RSS ($U > 15$) are listed, with the possible amino acids for each of the two sub-families at each position of the alignment. x refers to the IUPAC code, representing any amino acid.


annotated introns and thus probably do not reflect differences in *cir* expression between BALB/c and C57BL/6 mice.

A list of the ten most highly expressed *cir* genes was established for each sample (Figure 4b). As their expression was most highly detected in a population of parasites (either due to higher levels of expression in selected iRBCs or expression in a higher number of iRBCs), these will henceforth be referred to as “dominant *cirs*”. When we compared the lists of dominant *cirs* thus established, it was evident that most of these genes (> 60%) were different between the samples analyzed (Figure 4c). This indicates that the relative level of *cir* transcripts may vary from one mouse to another; which could reflect antigenic variation, immune selection of iRBCs or sampling differences upon infection of the mice. Whilst the overall level of *cir* expression was significantly lower than for other parasite genes (Kolmogorov-Smirnov test; $D = 0.6742$, P -value < $2.2e-16$,

Figure 4d), the dominant *cir* transcripts were found amongst the most highly expressed parasite genes.

Surprisingly, the same gene (PCHAS_110030) was expressed more than ten fold higher than any other *cir* genes in all the BALB/c samples analyzed. Similarly, three other *cirs* (PCHAS_140140, PCHAS_070130, and PCHAS_000430) were also dominant in these four samples. Furthermore, all of these genes, except PCHAS_140140, were dominant in at least one of the two C57BL/6 mice studied (Additional file 9). This indicates that some *cir* genes are consistently highly represented in a population of *Plasmodium chabaudi* AS parasites at the late trophozoite stage during the peak of parasitemia (8 days post infection).

To evaluate whether there was a relationship between CIR sub-families and functions, a comparison of their expression patterns was undertaken. In our analysis, more of the expressed *cirs* belonged to sub-family B (on average, $57.2\% \pm 4.7\%$ were B-type *cirs* and $42.3\% \pm$





4.7% were A-types; Figure 5a). A similar result was obtained in both analyzed C57BL/6 mice (58.33% \pm 8.41% of *cirs* belonged to sub-family B and 41.66% \pm 8.41% were A-type *cirs*, data not shown). When the expression levels of *cir* genes from each sub-family were compared, no statistically significant differences could be detected (K-S test, $P = 0.46$). It thus seems that the overall level of expression between *cir* genes belonging to sub-families A and B was similar.

However, the most highly expressed *cirs* were all members of sub-family A (Figure 5b). Indeed, PCHAS_110030, the most highly expressed *cir* gene in the six samples analyzed, belonged to this sub-family. We investigated whether this was also the case for the other dominant genes identified previously, Figure 4b. However, in each sample analyzed, about half of the dominant *cir* genes identified belonged to each sub-family (on average, 52.5% \pm 18.9% of the dominant genes were A-type *cirs* and 47.5% \pm 18.9% were members of sub-family B, Figure 5c, indicated by black dots in Additional file 3). This observation was confirmed in the C57BL/6 mice (on average, 50% \pm 14.14% of the dominant *cirs* belonged to sub-family A and 50% \pm 14.14% were B-type *cirs*).

b) Microarray analysis

Timing of transcription could be an alternative mechanism by which CIRs of different subgroups mediate different functions, therefore we investigated the transcriptional pattern of these genes throughout the intra-erythrocytic developmental cycle (IDC) in *P. chabaudi*. For this purpose parasites were collected at 2-hour intervals from mice with a synchronous infection of *P. chabaudi* for a total of 24 hrs, representing one complete IDC of the parasite. The RNA was extracted and *cir* transcription was determined using the pan-rodent *Plasmodium* spp microarray, which has been described previously [55].

Figure 6 shows the detected *cir* expression patterns throughout the IDC, where each time-point represents parasites collected from a single mouse [accession number in GEA: GSE33333]. Of the total 96 unique *cir* genes represented on the array, 49 showed clear transcriptional activation throughout the IDC. Each *cir* showed a single peak of transcription in line with previous reports that in *Plasmodium*, genes are only activated once during the IDC [24,56]. Importantly, not all *cir* genes were activated at the same time but rather showed progressive transcriptional activation throughout



the IDC with approximately one third of the analyzed *cirs* respectively showing peak transcriptional activity at the ring, trophozoite and schizont stages. Similar observations have also been made in *P. vivax*, where members of the *vir* gene family showed a comparable transcription pattern [24].

Differences in absolute levels of transcription of the two *cir* sub-families could not be quantified using this approach as two-channel hybridization only allows relative abundance comparison for each gene and not between different genes. However, many members of each sub-family could be detected at every time point. These data also strongly suggest that different *cirs* are indeed transcribed at different stages of parasite development in the erythrocyte in line with their products having different functions.

Interestingly, several of the dominant *cir* transcripts from the RNA sequencing experiments were also detected in this microarray timecourse, despite the fact that this material was obtained from separate *P. chabaudi* infections. These *cirs* were: PCHAS_110030, PCHAS_140140, PCHAS_110020, PCHAS_001090, PCHAS_070130, PCHAS_000110 and PCHAS_000120, which comprised approximately a third of the dominant *cirs* identified by RNA sequencing analysis. In addition, as each time point represents the material from a single mouse, the progression of *cir* transcription appeared to be conserved in the mice that were infected with a single starting parasite population and would suggest that limited switching occurs within a single cycle of asexual development. Importantly, the peak timing of expression of these *cirs* detected by microarray is consistent with the parasite stage used for the RNA sequencing analysis, further supporting the notion that the timing of expression of different CIR is conserved in different independent infections.

Discussion

In total, 196 *cir* genes have been identified and annotated in the *P. chabaudi* genome. The *cir* genes distribute into two major groups, according to sequence similarity. Many *cirs* from both sub-families are transcribed during *P. chabaudi* infection, although dominant *cir* transcripts are present. Differences between the sub-families, such as sub-family specific amino acid motifs and altered residue conservation, particularly within the predicted TM domain, indicate that the two sub-families could be functionally divergent. Recombination of *cir* genes may have occurred within the *cir* repertoire, consistent with that observed in other *Plasmodium* multi-gene families including *var* and *sicavar* (for example: [63,64]).

Similar to the VIR and YIR gene families [2,20,22,23], we confirm that there are also sub-groups within the CIR repertoire in agreement with Ebbinghaus and Krucken [30]. As each of the major groups defined here represented almost half of the CIRs, we defined these as the sub-families: A, comprising the more divergent CIRs; and B, containing more conserved CIRs. These CIR sub-families could each be further sub-divided into smaller clades. Despite the different analysis of CIRs recently described [30], the same groups could be identified within our network (as clade A1 and sub-family B), indicating that the sequences in each of these groups are indeed highly similar to each other. While the overall conservation of PIRs across species is relatively low, some evidence exists for sub-families being shared between *P. yoelii* and *P. berghei* [23], as well as *P. chabaudi*, *P. yoelii* and *P. berghei* [30] suggesting common functions.

The extensive sequence variation often seen in multi-gene families like *var*, *rif* and *stevor* in *P. falciparum*, or the *pirs* in *P. vivax* and rodent malaria parasites, is thought to reflect that these genes are under immune pressure (eg. [2,29,72-74]). In our analysis however, we have identified 14 conserved amino acid motifs, suggesting that some parts of the CIR sequences are under constraints important for the biological functions of these proteins. While some motifs were found in the majority of CIRs, motifs unique to each of the two sub-families were also detected. This divergence of conserved amino acid sequence motifs in the different sub-groups presents strong evidence that the different CIR sub-types may indeed carry out distinct but related functions in the parasite, as suggested also for PIR and RIFIN sequences [18]. Since the CIR A sub-family contained the most divergent CIRs, these proteins may be exposed to stronger selective pressure, and it is possible that their diverse nature enables immune evasion *in vivo*.

In line with the functional diversification of CIR sub-types A and B, some CIRs belonging to sub-type A possessed a unique amino acid motif (motif 8) at the beginning of the predicted TM domain. This, in addition to different conservation of proline residues and other amino acids within the predicted TM domains of each sub-family, suggests that CIR sub-family A proteins may have a functionally different TM domain than the B-type CIRs, perhaps altering which membrane the proteins are able to span.

Janssen and colleagues have predicted that *rif* and *stevor* could be ancestral to the *pir* genes due to similarities in terms of size, amino acid motifs and conservation of the first intron [18]. The observations that the CIR repertoire could be separated into two major sub-types namely thanks to an insertion within CIR sub-type A, a section of which contained two highly conserved cysteine residues [similar to RIFIN sub-type A, [16,17]], is strong evidence that the CIR family displays further parallels to the RIFINs. The variation we observed in the number and position of cysteine residues in the two CIR sub-families could significantly impact on the overall protein tertiary structure, and thereby function. As predicted for the A and B type RIFINs [17,49], we found high proportions of rate and conservation shifting sites between the two major CIR sub-families, providing further support for the idea of functional divergence.

Transcriptional differences in both in amount as well as timing can indicate functional differences between related genes. Until now, only Northern blot, degenerate PCR and restriction fragment length polymorphism (RFLP) analyses have been used to give indications of *cir* gene expression during *P. chabaudi* infection [29,30]. Here, we have used both mRNA sequencing as well as microarray studies to obtain a detailed picture of *cir*

transcription *in vivo*. The expression of up to 40% of the *cir* genes could be detected during an infection. Different *cirs* were transcribed at different times during the IDC of the parasite, with distinct *cirs* being transcribed in ring, trophozoite and schizont stages. The timing of transcription for a specific *cir* appeared to be conserved from one animal to another (as seen in the microarray experiments) as well as across different experiments (as seen with the good overlap between the genes transcribed in the trophozoite stage as detected by both RNA sequencing and microarray), suggesting that CIRs may carry out distinct ring-, trophozoite- or schizont-specific functions. This is further supported by the observation that *pir* transcription follows a similar pattern in both *P. vivax* and *P. yoelii*, where 59% of the *vars* and 42% of the *yirs* were expressed respectively [4,24].

Whilst most *cir* genes were transcribed at lower levels than other parasite genes, this most likely reflects differences in the proportion of parasites that actually express a particular *cir*, resulting in populations of iRBCs each expressing different *cir* genes. This would be consistent with observations made for *vars*, *yirs* and *vars* [1,4,75], that individual iRBCs transcribe only one or a small number of these genes. It is therefore of particular interest that three *cir* genes were represented at relatively high levels in all six samples analyzed by RNA sequencing (PCHAS_110030, PCHAS_070130, and PCHAS_000430), with PCHAS_110030 being detected at a level more than 10 fold higher than any other *cir* in all the samples analyzed. In a classic model of antigenic variation, it would be expected that a single or relatively small number of a gene family would be transcribed in any population, with other members of the gene family being transcribed at much lower levels. This is exactly what we have detected during *P. chabaudi* infection.

While our data suggest that *cirs* belonging to sub-family A are more highly transcribed than those belonging to sub-family B, these differences are relatively small. At this stage there it is not clear whether there is a preference for the transcription of A or B sub-families. What is evident though, is that both A and B sub-family members were transcribed at the peak of a blood-stage *P. chabaudi* infection. Considering that the sequence analysis carried out here strongly suggests functional differences between the two sub-types, the expression of both sub-families may indicate that A and B-type CIRs provide complementary but non-overlapping roles at different stages during infection.

Conclusions

We have here presented a thorough analysis of the *cir* repertoire, highlighting similarities between these genes and other multi-gene families. The two CIR sub-families

have been predicted to have different functions, as shown for the RIFINs of *P. falciparum*. Whilst these function(s) remain unknown, the large number of *cir* genes and their expression throughout the IDC indicates that CIR proteins are likely to play key roles in the biology of the parasite. These may include antigenic variation and immune evasion. Our application of RNA sequencing during *P. chabaudi* infection has enabled the detection of dominant *cir* transcripts for the first time, supporting such roles. Further application of these methodologies may elucidate the functions of CIR proteins and help to clarify the roles that members of the CIR sub-families play during infection.

Additional material

- Additional file 1: Alignment of 183 CIR sequences.
- Additional file 2: CIR alignment conservation.
- Additional file 3: CIR network showing bootstrap values.
- Additional file 4: Maximum likelihood tree of CIR sequences.
- Additional file 5: Sub-families identified from the alignment of 183 CIRs.
- Additional file 6: Detection of phylogenetic incompatibilities between *cir* genes.
- Additional file 7: Identification of similarities between the CIR and RIFIN repertoires.
- Additional file 8: Raw microarray data.
- Additional file 9: Raw RNA sequencing data.
- Additional file 10: *cir* gene expression threshold of detection determination.
- Additional file 11: Legends to Additional Files 1-10.

Acknowledgements

We are grateful to the PFAM group at the Wellcome Trust Sanger Institute for creating a HMM based on the *cir* genes, and to Saraswathi Abhiman and Erik L. Sonnhammer for allowing us to use the Funshift program in our analysis [49].

Funding

This work was supported by the Medical Research Council, UK (file reference U117584248); Singapore A*Star-UK MRC Joint Grant (A*Star reference 10/1/22/24/630); Singapore Nanyang Technological University (WBS N° M47080081); and the European Community's Seventh Framework Programme (FP7/2007-2013) under grant agreement N° 242095. Jennifer Lawton was in receipt of an MRC PhD scholarship.

Author details

¹Division of Parasitology, MRC National Institute for Medical Research, London, UK. ²Division of Genomics and Genetics, Nanyang Technological University Singapore, Singapore. ³Parasite Genomics, Wellcome Trust Sanger Institute, Hinxton, UK. ⁴Pathogen Genomics Group, Computational Bioscience Research Center, Chemical Life Sciences and Engineering Division, 4700 King Abdullah University of Science and Technology, Thuwal 23955-6900, Kingdom of Saudi Arabia.

Authors' contributions

JL^(1st author) carried out the *cir* gene annotation in collaboration with UB, AP and MB; carried out the bioinformatic analyses, supervised by AJ; prepared RNA samples for RNAseq and drafted the manuscript in collaboration with TB. RNAseq data were analyzed by AJR, TDO and TB. Microarray analyses were performed by YXY and PP. JL^(last author) conceived of and supervised

the study, and edited the manuscript with DC, AJ, AJR, AP, MB and PP. All authors read and approved the final manuscript.

Competing interests

The authors declare that they have no competing interests.

Received: 4 November 2011 Accepted: 29 March 2012

Published: 29 March 2012

References

- Joergensen L, Bengtsson DC, Bengtsson A, Ronander E, Berger SS, Turner L, Dalgaard MB, Cham GK, Victor ME, Lavstsen T, et al: **Surface co-expression of two different P/EEMP1 antigens on single Plasmodium falciparum-infected erythrocytes facilitates binding to ICAM1 and PECAM1.** *PLoS Pathogens* 2010, **6(9)**:e1001083.
- del Portillo HA, Fernandez-Becerra C, Bowman S, Oliver K, Preuss M, Sanchez CP, Schneider NK, Villalobos JM, Rajandream MA, Harris D, et al: **A superfamily of variant genes encoded in the subtelomeric region of Plasmodium vivax.** *Nature* 2001, **410(6830)**:839-842.
- Howard RJ, Barnwell JW, Kao V: **Antigenic variation of Plasmodium knowlesi malaria: identification of the variant antigen on infected erythrocytes.** *Proc Natl Acad Sci USA* 1983, **80(13)**:4129-4133.
- Cunningham D, Fonager J, Jarra W, Carret C, Preiser P, Langhorne J: **Rapid changes in transcription profiles of the Plasmodium yoelii yir multigene family in clonal populations: lack of epigenetic memory?** *PLoS One* 2009, **4(1)**:e4285.
- Gilks CF, Walliker D, Newbold CI: **Relationships between sequestration, antigenic variation and chronic parasitism in Plasmodium chabaudi chabaudi-a rodent malaria model.** *Parasite Immunology* 1990, **12(1)**:45-64.
- Rowe JA, Claessens A, Corrigan RA, Arman M: **Adhesion of Plasmodium falciparum-infected erythrocytes to human cells: molecular mechanisms and therapeutic implications.** *Expert Reviews in Molecular Medicine* 2009, **11**:e16.
- Scherf A, Lopez-Rubio JJ, Riviere L: **Antigenic variation in Plasmodium falciparum.** *Annu Rev Microbiol* 2008, **62**:445-470.
- Marsh K, Howard RJ: **Antigens induced on erythrocytes by P. falciparum: Expression of diverse and conserved determinants.** *Science* 1986, **231(4734)**:150-153.
- Marsh K, Otoo L, Hayes RJ, Carson DC, Greenwood BM: **Antibodies to blood stage antigens of Plasmodium falciparum in rural Gambians and their relation to protection against infection.** *Transactions of the Royal Society of Tropical Medicine and Hygiene* 1989, **83(3)**:293-303.
- Fischer K, Chavchich M, Huestis R, Wilson DW, Kemp DJ, Saul A: **Ten families of variant genes encoded in subtelomeric regions of multiple chromosomes of Plasmodium chabaudi, a malaria species that undergoes antigenic variation in the laboratory mouse.** *Mol Microbiol* 2003, **48(5)**:1209-1223.
- Rubio JP, Thompson JK, Cowman AF: **The var genes of Plasmodium falciparum are located in the subtelomeric region of most chromosomes.** *EMBO J* 1996, **15(15)**:4069-4077.
- Cunningham D, Lawton J, Jarra W, Preiser P, Langhorne J: **The pir multigene family of Plasmodium: antigenic variation and beyond.** *Mol Biochem Parasitol* 2010, **170(2)**:65-73.
- Bachmann A, Predehl S, May J, Harder S, Burchard GD, Gilberger TW, Tannich E, Bruchhaus I: **Highly co-ordinated var gene expression and switching in clinical Plasmodium falciparum isolates from non-immune malaria patients.** *Cellular Microbiology* 2011, **13(9)**:1397-409.
- Kraemer SM, Smith JD: **Evidence for the importance of genetic structuring to the structural and functional specialization of the Plasmodium falciparum var gene family.** *Mol Microbiol* 2003, **50(5)**:1527-1538.
- Lavstsen T, Salanti A, Jensen AT, Arnot DE, Theander TG: **Sub-grouping of Plasmodium falciparum 3D7 var genes based on sequence analysis of coding and non-coding regions.** *Malar J* 2003, **2**:27.
- Petter M, Haeggstrom M, Khattab A, Fernandez V, Klinkert MQ, Wahlgren M: **Variant proteins of the Plasmodium falciparum RIFIN family show distinct subcellular localization and developmental expression patterns.** *Mol Biochem Parasitol* 2007, **156(1)**:51-61.
- Joannin N, Abhiman S, Sonnhammer EL, Wahlgren M: **Sub-grouping and sub-functionalization of the RIFIN multi-copy protein family.** *BMC Genomics* 2008, **9**:19.
- Janssen CS, Phillips RS, Turner CM, Barrett MP: **Plasmodium interspersed repeats: the major multigene superfamily of malaria parasites.** *Nucleic Acids Res* 2004, **32(19)**:5712-5720.
- Pain A, Bohme U, Berry AE, Mungall K, Finn RD, Jackson AP, Mourier T, Mistry J, Pasini EM, Aslett MA, et al: **The genome of the simian and human malaria parasite Plasmodium knowlesi.** *Nature* 2008, **455(7214)**:799-803.
- Carlton JM, Adams JH, Silva JC, Bidwell SL, Lorenzi H, Caler E, Crabtree J, Angiuoli SV, Merino EF, Amedeo P, et al: **Comparative genomics of the neglected human malaria parasite Plasmodium vivax.** *Nature* 2008, **455(7214)**:757-763.
- del Portillo HA, Lanzer M, Rodriguez-Malaga S, Zavala F, Fernandez-Becerra C: **Variant genes and the spleen in Plasmodium vivax malaria.** *Int J Parasitol* 2004, **34(13-14)**:1547-1554.
- Merino EF, Fernandez-Becerra C, Durham AM, Ferreira JE, Tumilasci VF, d'Arc-Neves J, da Silva-Nunes M, Ferreira MU, Wickramarachchi T, Udagama-Randeniya P, et al: **Multi-character population study of the vir subtelomeric multigene superfamily of Plasmodium vivax, a major human malaria parasite.** *Mol Biochem Parasitol* 2006, **149(1)**:10-16.
- Fonager J, Cunningham D, Jarra W, Koernig S, Henneman AA, Langhorne J, Preiser P: **Transcription and alternative splicing in the yir multigene family of the malaria parasite Plasmodium y. yoelii: identification of motifs suggesting epigenetic and post-transcriptional control of RNA expression.** *Molecular and Biochemical Parasitology* 2007, **156(1)**:1-11.
- Bozdech Z, Mok S, Hu G, Imwong M, Jaidee A, Russell B, Ginsburg H, Nosten F, Day NP, White NJ, et al: **The transcriptome of Plasmodium vivax reveals divergence and diversity of transcriptional regulation in malaria parasites.** *Proc Natl Acad Sci USA* 2008, **105(42)**:16290-16295.
- McLean SA, Pearson CD, Phillips RS: **Plasmodium chabaudi: antigenic variation during recrudescence parasitaemias in mice.** *Exp Parasitol* 1982, **54(3)**:296-302.
- McLean SA, Pearson CD, Phillips RS: **Plasmodium chabaudi: relationship between the occurrence of recrudescence parasitaemias in mice and the effective levels of acquired immunity.** *Exp Parasitol* 1982, **54(2)**:213-221.
- McLean SA, Pearson CD, Phillips RS: **Antigenic variation in Plasmodium chabaudi: analysis of parent and variant populations by cloning.** *Parasite Immunology* 1986, **8(5)**:415-424.
- Mota MM, Jarra W, Hirst E, Patnaik PK, Holder AA: **Plasmodium chabaudi-infected erythrocytes adhere to CD36 and bind to microvascular endothelial cells in an organ-specific way.** *Infect Immun* 2000, **68(7)**:4135-4144.
- Janssen CS, Barrett MP, Turner CM, Phillips RS: **A large gene family for putative variant antigens shared by human and rodent malaria parasites.** *Proceedings of the Royal Society B: Biological Sciences* 2002, **269(1489)**:431-436.
- Ebbinghaus P, Krucken J: **Characterization and tissue-specific expression patterns of the Plasmodium chabaudi cir multigene family.** *Malaria Journal* 2011, **10**:272.
- Rutherford K, Parkhill J, Crook J, Horsnell T, Rice P, Rajandream MA, Barrell B: **Artemis: sequence visualization and annotation.** *Bioinformatics* 2000, **16(10)**:944-945.
- Carver T, Berriman M, Tivey A, Patel C, Bohme U, Barrell BG, Parkhill J, Rajandream MA: **Artemis and ACT: viewing, annotating and comparing sequences stored in a relational database.** *Bioinformatics* 2008, **24(23)**:2672-2676.
- The Plasmodium chabaudi homepage in GeneDB. [http://www.genedb.org/Homepage/Pchabaudi].
- Hall N, Karras M, Raine JD, Carlton JM, Kooij TWA, Berriman M, Florens L, Janssen CS, Pain A, Christophides GK, et al: **A comprehensive survey of the Plasmodium life cycle by genomic, transcriptomic, and proteomic analyses.** *Science* 2005, **307(5706)**:82-86.
- Finn RD, Mistry J, Tate J, Coggill P, Heger A, Pollington JE, Gavin OL, Gunasekaran P, Ceric G, Forslund K, et al: **The Pfam protein families database.** *Nucleic Acids Res* 2010, **38 Database**: D211-D222.
- Sonnhammer EL, Eddy SR, Birney E, Bateman A, Durbin R: **Pfam: multiple sequence alignments and HMM-profiles of protein domains.** *Nucleic Acids Res* 1998, **26(1)**:320-322.
- Bailey TL, Elkan C: **Fitting a mixture model by expectation maximization to discover motifs in biopolymers.** *Proceedings of the Second International Conference on Intelligent Systems for Molecular Biology* 1994, **2**:28-36.

38. Crooks GE, Hon G, Chandonia JM, Brenner SE: **WebLogo: a sequence logo generator.** *Genome Res* 2004, **14**(6):1188-1190.
39. Edgar RC: **MUSCLE: multiple sequence alignment with high accuracy and high throughput.** *Nucleic Acids Res* 2004, **32**(5):1792-1797.
40. Huson DH: **SplitsTree: analyzing and visualizing evolutionary data.** *Bioinformatics* 1998, **14**(1):68-73.
41. Bryant D, Moulton V: **Neighbor-net: an agglomerative method for the construction of phylogenetic networks.** *Mol Biol Evol* 2004, **21**(2):255-265.
42. Gambette P, Huson DH: **Improved layout of phylogenetic networks.** *IEEE/ACM Transactions on Computational Biology and Bioinformatics* 2008, **5**(3):472-479.
43. Guindon S, Lethiec F, Duroux P, Gascuel O: **PHYML Online-a web server for fast maximum likelihood-based phylogenetic inference.** *Nucleic Acids Res* 2005, **33** Web Server: W557-W559.
44. Le SQ, Gascuel O: **An improved general amino acid replacement matrix.** *Mol Biol Evol* 2008, **25**(7):1307-1320.
45. Anisimova M, Gascuel O: **Approximate likelihood-ratio test for branches: A fast, accurate, and powerful alternative.** *Syst Biol* 2006, **55**(4):539-552.
46. Carlton JM, Angiuoli SV, Suh BB, Kooij TW, Perteu M, Silva JC, Ermolaeva MD, Allen JE, Selengut JD, Koo HL, et al: **Genome sequence and comparative analysis of the model rodent malaria parasite *Plasmodium yoelii yoelii*.** *Nature* 2002, **419**(3):512-519.
47. Milne I, Wright F, Rowe G, Marshall DF, Husmeier D, McGuire G: **TOPALI: software for automatic identification of recombinant sequences within DNA multiple alignments.** *Bioinformatics* 2004, **20**(11):1806-1807.
48. Milne I, Lindner D, Bayer M, Husmeier D, McGuire G, Marshall DF, Wright F: **TOPALI v2: a rich graphical interface for evolutionary analyses of multiple alignments on HPC clusters and multi-core desktops.** *Bioinformatics* 2009, **25**(1):126-127.
49. Abhiman S, Sonnhammer EL: **FunShift: a database of function shift analysis on protein subfamilies.** *Nucleic Acids Res* 2005, **33** Database: D197-D200.
50. Knudsen B, Miyamoto MM: **A likelihood ratio test for evolutionary rate shifts and functional divergence among proteins.** *Proc Natl Acad Sci USA* 2001, **98**(25):14512-14517.
51. Slade SJ, Langhorne J: **Production of interferon-gamma during infection of mice with *Plasmodium chabaudi chabaudi*.** *Immunobiology* 1989, **179**(4-5):353-365.
52. Shinkaia Y, Rathbun G, Lama KP, Oltza EM, Stewart V, Mendelsohn M, Charona J, Datta M, Younga F, Stalla AM, et al: **RAG-2-deficient mice lack mature lymphocytes owing to inability to initiate V(D)J rearrangement.** *Cell* 1992, **68**(5):855-867.
53. von der Weid T, Honarvar N, Langhorne J: **Gene-targeted mice lacking B cells are unable to eliminate a blood stage malaria infection.** *J Immunol* 1996, **156**(7):2510-2516.
54. Kyes S: **Reliable RNA preparation for *Plasmodium falciparum*.** In *Methods in Malaria Research*. Fourth edition. Edited by: Ljungström I, Perlmann H, Schlichtherle M, Artur Scherf A, Wahlgren M. Manassas, Virginia; 2004:175-178.
55. Liew KJL, Hu G, Bozdech Z, Peter PR: **Defining species specific genome differences in malaria parasites.** *BMC Genomics* 2010, **11**(1):128.
56. Bozdech Z, Llinas M, Pulliam BL, Wong ED, Zhu J, DeRisi JL: **The transcriptome of the intraerythrocytic developmental cycle of *Plasmodium falciparum*.** *PLoS Biology* 2003, **1**(1):E5.
57. Otto TD, Wilinski D, Assefa S, Keane TM, Sarry LR, Bohme U, Lemieux J, Barrell B, Pain A, Berriman M, et al: **New insights into the blood-stage transcriptome of *Plasmodium falciparum* using RNA-Seq.** *Mol Microbiol* 2010, **76**(1):12-24.
58. Trapnell C, Pachter L, Salzberg SL: **TopHat: Discovering splice junctions with RNA-Seq.** *Bioinformatics* 2009, **25**(9):1105-1111.
59. Mortazavi A, Williams BA, McCue K, Schaeffer L, Wold B: **Mapping and quantifying mammalian transcriptomes by RNA-Seq.** *Nature Methods* 2008, **5**:621-628.
60. Kolmogorov A: **Sulla determinazione empirica di una legge di distribuzione.** *Giornale dell'Istituto Italiano degli Attuari* 1933, **4**:83-91.
61. **Plotcon.** [http://emboss.bioinformatics.nl/cgi-bin/emboss/plotcon].
62. Morrison DA: **Networks in phylogenetic analysis: new tools for population biology.** *Int J Parasitol* 2005, **35**(5):567-582.
63. Corredor V, Meyer EV, Lapp S, Corredor-Medina C, Huber CS, Evans AG, Barnwell JW, Galinski MR: **A *SICA*var switching event in *Plasmodium knowlesi* is associated with the DNA rearrangement of conserved 3' non-coding sequences.** *Mol Biochem Parasitol* 2004, **138**(1):37-49.
64. Frank M, Kirkman L, Costantini D, Sanyal S, Lavazec C, Templeton TJ, Deitsch KW: **Frequent recombination events generate diversity within the multi-copy variant antigen gene families of *Plasmodium falciparum*.** *Int J Parasitol* 2008, **38**(10):1099-1109.
65. Morell V: **TreeBASE: the roots of phylogeny.** *Science* 1996, **273**:569-600.
66. Piel W, Donoghue MJ, Sanderson MJ: **TreeBASE: a database of phylogenetic knowledge.** In *To the interoperable "Catalog of Life" with partners Species 2000 Asia Oceania Research Report from the National Institute for Environmental Studies No 171*. Edited by: Shimura J, Wilson KL, Gordon D. Tsukuba, Japan; 2002:41-47.
67. Sanderson MJ, Donoghue MJ, Piel W, Eriksson T: **TreeBASE: a prototype database of phylogenetic analyses and an interactive tool for browsing the phylogeny of life.** *American Journal of Botany* 1994, **81**(6):183.
68. Belyaev NN, Brown DE, Diaz ALG, Rae A, Jarra W, Thompson J, Langhorne J, Potocnik AJ: **Induction of an IL7-R + c-Kit hi myelolymphoid progenitor critically dependent on IFN-γ signaling during acute malaria.** *Nat Immunol* 2010, **11**(6):477-485.
69. Spence PJ, Cunningham D, Jarra W, Lawton J, Langhorne J, Thompson J: **Transformation of the rodent malaria parasite *Plasmodium chabaudi*.** *Nat Protoc* 2011, **6**(4):553-561.
70. Sponaas AM, Do Rosario APF, Voisine C, Mastelic B, Thompson J, Koernig S, Jarra W, Renia L, Mauduit M, Potocnik AJ, et al: **Migrating monocytes recruited to the spleen play an important role in control of blood stage malaria.** *Blood* 2009, **114**(27):5522-5531.
71. Stephens R, Langhorne J: **Effector memory Th1 CD4 T cells are maintained in a mouse model of chronic malaria.** *PLoS Pathogens* 2010, **6**(11):e1001208.
72. Su XZ, Heatwole VM, Wertheimer SP, Guinet F, Herrfeldt JA, Peterson DS, Ravetch JA, Wellems TE: **The large diverse gene family *var* encodes proteins involved in cytoadherence and antigenic variation of *Plasmodium falciparum*-infected erythrocytes.** *Cell* 1995, **82**(1):89-100.
73. Kyes S, Horrocks P, Newbold C: **Antigenic variation at the infected red cell surface in malaria.** *Annual Review of Microbiology* 2001, **55**:673-707.
74. Niang M, Xue YY, Preiser PR: **The *Plasmodium falciparum* STEVOR multigene family mediates antigenic variation of the infected erythrocyte.** *PLoS Pathogens* 2009, **5**(2):e1000307.
75. Fernandez-Becerra C, Pein O, de Oliveira TR, Yamamoto MM, Cassola AC, Rocha C, Soares IS, de Braganca Pereira CA, del Portillo HA: **Variante proteins of *Plasmodium vivax* are not clonally expressed in natural infections.** *Mol Microbiol* 2005, **58**(3):648-658.

doi:10.1186/1471-2164-13-125

Cite this article as: Lawton et al: Characterization and gene expression analysis of the *cir* multi-gene family of *plasmodium chabaudi chabaudi* (AS). *BMC Genomics* 2012 **13**:125.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

