

## Antarctic Science

<http://journals.cambridge.org/ANS>

Additional services for **Antarctic Science**:

Email alerts: [Click here](#)

Subscriptions: [Click here](#)

Commercial reprints: [Click here](#)

Terms of use : [Click here](#)



---

## Magnetic properties of Upper Quaternary sediments from the Scotia Sea, Antarctica

Norman Hamilton and Carol J. Pudsey

Antarctic Science / Volume 13 / Issue 01 / March 2001, pp 61 - 66  
DOI: 10.1017/S0954102001000098, Published online: 27 April 2004

**Link to this article:** [http://journals.cambridge.org/abstract\\_S0954102001000098](http://journals.cambridge.org/abstract_S0954102001000098)

### How to cite this article:

Norman Hamilton and Carol J. Pudsey (2001). Magnetic properties of Upper Quaternary sediments from the Scotia Sea, Antarctica. *Antarctic Science*, 13, pp 61-66 doi:10.1017/S0954102001000098

**Request Permissions :** [Click here](#)

# Magnetic properties of Upper Quaternary sediments from the Scotia Sea, Antarctica

NORMAN HAMILTON<sup>1</sup> and CAROL J. PUDSEY<sup>2</sup>

<sup>1</sup>*School of Ocean and Earth Science, University of Southampton, Southampton Oceanography Centre, Southampton SO14 3ZH, UK*

<sup>2</sup>*British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge CB3 0ET, UK*

**Abstract:** Magnetic properties of bulk sediment samples taken from three cores from the Scotia Sea, Antarctica were determined using a fully-automated variable field translation balance. Fine-grained detrital magnetite is identified as the principal carrier of remanence in these Upper Quaternary sediments which were deposited under the influence of the Antarctic Circumpolar Current. Inferred magnetite grain-size is consistent with published bulk grain-size data for these cores. Pseudo-single domain grains characterize Holocene samples, and larger, multi-domain grains occur in glacial samples from two of the cores, whereas samples from the northernmost core site show dominantly multi-domain behaviour.

Received 17 August 2000, accepted 22 November 2000

**Key words:** environmental magnetic parameters, Quaternary, Scotia Sea, sediments

## Introduction

Southern Ocean sediments are sensitive recorders of the state of glaciation of Antarctica. They preserve evidence of changes in sea ice cover, iceberg rafting, biological productivity and deep ocean circulation. Deep circulation has been studied using a variety of proxy measurements including stable isotopes as water mass tracers (Charles & Fairbanks 1992), radiogenic isotopes and reworked microfossils as indicators of sediment redistribution (Burckle 1981, Frank *et al.* 1996), and grain size as an indicator of palaeocurrent strength (Pudsey 1992). In other areas of the ocean, environmental magnetic parameters are being developed as climate proxies (e.g. Stoner *et al.* 1995). Sagnotti *et al.* (1998) utilised the environmental magnetic record of glaciomarine sediments of the CIROS-1 core, McMurdo Sound, Antarctica to infer palaeoclimate during the Eocene/Oligocene boundary interval.

Composition and grain size of magnetic minerals in areas distant from continental margin sources provide a useful adjunct to bulk sediment texture and composition. This present study is the first to describe the environmental magnetic parameters of Scotia Sea sediments other than their bulk magnetic susceptibility which was previously reported by Pudsey & Howe (1998).

The sediment samples used in this investigation were taken from two piston cores (PC29 and PC31) together with a Kasten core (KC64) obtained during cruises on the RRS *Discovery* (1987/88) and RRS *James Clark Ross* (1992/93), respectively. Core PC29 is located in the north-central part of the Scotia Sea and PC31 is situated some 260 km to the south-west; the site of KC64 lies south of the North Scotia Ridge (Fig. 1 & Table I).


Lithologically, the cores are composed of an upper biogenic unit which is dominantly a diatom ooze or foraminifer-bearing diatom mud underlain by a terrigenous diatom mud

unit (Pudsey & Howe 1998). Some thin vitric ash layers, often preserved as smectitic alteration products, are also present (Moreton & Smellie 1998). The unaltered ash layers can be identified by a magnetic susceptibility peak in downcore susceptibility records (Pudsey & Howe 1998). Cores PC29 and PC31 also contain a lower biogenic unit of muddy diatom ooze and diatom mud. Within the dominantly terrigenous component of the sediments deposited during the Last Glacial Maximum at site KC64, Pudsey & Howe (1998) reported magnetite as a prominent constituent of the fine-grained fraction.

Jordan & Pudsey (1992) defined a local diatom stratigraphy for the Scotia Sea, extending back to approximately 60 000 years. However, for PC29 and PC31, studies of radiolarian and palaeomagnetic stratigraphy (Jordan & Pudsey 1992) indicate that sediment recovery in the base of these piston cores is no older than *c.* 30 ka. Both cores contain a dominant peak in the relative abundance of the radiolarian species *Cycladophora davisiana* which has been correlated with the Last Glacial Maximum at 18 ka. Evidence for the high sedimentation rates in these cores and their interglacial/glacial sedimentology is described by Pudsey & Howe (1998). They demonstrated that the core locations are in depositional environments characterized by contourite sedimentation. The sediments record the fluctuating history of palaeoflow of the Antarctic Circumpolar Current during the late Quaternary. Hence, the remanence carriers may be strongly influenced by the nature of terrigenous and biogenic sediment supplied under the control of the Antarctic Circumpolar Current.

## Methods

Magnetic parameters including high-temperature thermomagnetic properties, saturation isothermal remanence



**Fig. 1.** Map of the Scotia Sea showing the location of cores 29, 31 and 64 in relation to the axis of the Antarctic Circumpolar Current (heavy dashed line, from Orsi *et al.* 1995) and the northern limit of sea ice (light dashed line, from Gloersen *et al.* 1992). Areas shallower than 2000 m shaded. Inset shows simplified core logs and magnetic susceptibility data (SI units  $\times 10^{-5}$  from Pudsey & Howe 1998). Upper and lower biosiliceous units shown in dotted ornament, with the foraminifer-bearing mud at the top of core 64 distinguished by irregular dots; dashed ornament indicates the terrigenous unit. Small black arrows are sample depths. Isotope stages (circled numbers) from data in Jordan & Pudsey (1992), supplemented by AMS  $^{14}\text{C}$  dates from Howe & Pudsey (1999) and Moreton (1999).

( $M_{rs}$ ), saturation magnetization ( $M_s$ ), coercive force ( $B_c$ ), coercivity of remanence ( $B_{cr}$ ) and hysteresis loops, were measured using an automated variable field translation balance (VFTB) installed in the palaeomagnetic laboratory of the Southampton Oceanography Centre, Southampton, UK. All measurements were made on dried bulk sediment samples of a few hundred milligrams. The sediment was taken from a suite of discrete samples used for an earlier palaeomagnetic study (O'Brien 1989). The VFTB system offers the advantage of direct determination of magnetic properties on bulk samples rather than requiring magnetic separates. Hysteresis loops, acquisition of saturation remanence ( $M_{rs}$ ) and back-field demagnetization of  $M_{rs}$  were completed up to maximum fields of 0.8 T.

## Results and discussion

The magnetic properties of the samples are summarized in

**Table I.** Core locations, water depths and lengths.

Core	Latitude	Longitude	Water depth (m)	Length (m)
PC29	56°25'S	41°10'W	3369	4.39
PC31	58°03'S	44°20'W	2745	4.24
KC64	53°52'S	48°20'W	4304	3.20

Table II. The measured values of the coercivity of remanence ( $B_{cr}$ ) are in the range 28–64 mT. Representative isothermal remanence acquisition and back-field demagnetization results from the three cores are shown in Fig. 2. All samples have broadly consistent isothermal remanence acquisition curves, with saturation being achieved at field strengths between 150 and 250 mT. This is strongly indicative of the presence of a ferrimagnetic mineral, such as titanomagnetite, as the dominant magnetic mineral, although Roberts (1995) cautions that the iron sulphide, greigite, produces similar behaviour.

The change in magnetization during heating to 700°C and cooling in air is illustrated by the representative thermomagnetic curves shown in Fig. 3. The majority of the heating curves have an inflexion point between 310–330°C. This may be due to the presence of some pyrrhotite (Dekkers 1989), greigite (Roberts 1995), or titanomagnetite. This is particularly apparent in the samples from core KC64. However, the majority of the samples show a very well-defined and abrupt loss of magnetization as the Curie point of magnetite is reached near 580°C. The decreased magnetization observed during cooling indicates that magnetite has oxidized to hematite for most of the samples. Few curves show reversible thermomagnetic behaviour. On the basis of the Curie point determinations, it seems probable that pure magnetite, rather than low-Ti titanomagnetite, is the major magnetic mineral




Fig.2. Representative isothermal remanence acquisition and back-field demagnetization curves.

Table II. Magnetic properties of samples used in this study.

Sample	Lithology	Depth (mbsf)	Bc (mT)	Bcr (mT)	Mrs ( $10^{-3} \text{Am}^2 \text{kg}^{-1}$ )	Ms ( $10^{-3} \text{Am}^2 \text{kg}^{-1}$ )	Mrs/Ms	Bcr/Bc
K641X05	fbm	0.05	6.8	52.4	13.8	210	0.07	7.72
K641X29	fbm	0.29	6.8	53.7	11.5	200	0.06	7.91
K641X59	fbm	0.59	5.1	38.4	8.2	165	0.05	7.53
K641X89	do	0.89	4.8	31.7	10.7	260	0.04	6.60
K641X119	dbm	1.19	4.7	27.6	11.8	300	0.04	5.87
KC642D01	dbm	1.51	6.9	34.2	16.3	280	0.06	4.99
KC642D11	dbm	1.61	6.7	34.3	16.6	330	0.05	5.09
KC642D24	dbm	1.74	6.8	34.2	15.8	295	0.05	5.00
KC642D41	dbm	1.91	6.9	34.2	16.1	310	0.05	4.99
KC642D61	dbm	2.11	6.8	34.2	19.6	330	0.06	5.01
KC642D74	dbm	2.24	6.8	31.0	20.6	340	0.06	4.54
KC642D88	dbm	2.38	6.8	34.2	18.4	370	0.05	5.04
TC29006	do	0.06	19.8	57.2	10.0	35	0.29	2.90
PC291036	do	0.36	19.8	63.9	7.6	27	0.28	3.23
PC291056	do	0.56	19.7	57.8	10.9	38	0.29	2.94
PC292046	do	2.00	11.8	46.0	7.6	65	0.12	3.90
PC292106	dbm	2.60	10.1	40.1	14.4	140	0.10	3.97
PC293006	mdo/dm	3.11	9.6	40.1	11.8	115	0.10	4.18
PC293066	mdo/dm	3.71	13.3	49.0	8.6	55	0.16	3.68
PC293126	dm	4.31	9.6	39.8	13.2	130	0.10	4.15
PC311026	do	0.26	19.6	48.0	5.9	27	0.22	2.45
PC311106	dm	1.06	12.0	40.0	15.3	130	0.12	3.33
PC312066	dm	2.00	11.1	40.0	17.5	160	0.11	3.60
PC312106	mdo/dm	2.40	12.0	38.0	11.1	90	0.12	3.17
PC313105	mdo/dm	3.89	15.0	57.9	13.7	75	0.18	3.86

Lithology codes: dbm = diatom-bearing mud, do = diatom ooze, fbm = foraminifer-bearing mud, mdo = muddy diatom ooze, dm = diatom mud




Fig.3. Typical thermomagnetic curves for heating and cooling of bulk sediment in air.

present in these sediments. This is in agreement with the visual mineralogical identification of magnetite, as described by Pudsey & Howe (1998).

Typical hysteresis loops are shown in Fig. 4. Samples from KC64 are characterized by narrow hysteresis loops, whereas loops from PC29 and PC31 are generally broader and more varied. The hysteresis ratio parameters recorded in Table II are confined to a range of values between 0.04 and 0.29 for the remanence ratio parameter  $M_{rs}/M_s$  together with an average value of 4.6 for the coercivity ratio  $B_{cr}/B_c$ . These values are consistent with the view that magnetite and/or titanomagnetite grains are present (Wasilewski 1973, Day *et al.* 1976, Dunlop 1986, Dunlop & Ozdemir 1997).

Magnetic grains from cores PC29 and PC31 lie dominantly in the pseudo-single-domain (PSD) grain size field in a hysteresis ratio plot (Day *et al.* 1977), whereas predominantly multidomain (MD) type behaviour is observed for core KC64 (Fig. 5). This is in agreement with the northward-coarsening trend in bulk sediment texture of Scotia Sea cores (Pudsey & Howe 1998). We have compared the magnetite grain diameter inferred from  $B_c$  values (Day *et al.* 1977, Dunlop 1986) with median diameters from Sedigraph analyses of bulk sediment. The Sedigraph measures equivalent settling diameter assuming a grain density of  $2.65 \text{ g cc}^{-1}$  (quartz). Allowing for the greater

density of magnetite (a quartz sphere is 1.26 x the diameter of a magnetite sphere of the same mass), the agreement is good.

Holocene samples from cores PC29 and PC31 have median diameters of 1.8–2 microns while glacial samples from these cores have median diameters of 2.5–4 microns, consistent with  $B_c$  values of 19–20 and 10–15 mT respectively (Table II). Low magnetic susceptibility (Fig. 1) and the predominance of quartz in the sand fraction of these cores were explained by derivation from sedimentary and metasedimentary rocks on the east side of the Antarctic Peninsula and the islands of the South Scotia Ridge (Pudsey & Howe 1998). In general, sediments deposited from the ACC are coarser in glacial intervals than interglacials, reflecting more vigorous ACC flow (Pudsey & Howe 1998, fig. 12).

Core KC64 is coarser throughout, reflecting its proximity to the ACC axis (Fig. 1). Holocene median diameters range from 8–16 microns and glacial values are 7–10 microns, again consistent with  $B_c$  values of 1.5–2.4 and less than 1.2 mT, respectively; the grain size of Holocene samples shows considerable scatter by either technique. Entrainment of terrigenous detritus from the Antarctic Peninsula and southernmost South America was invoked by Pudsey & Howe (1998) to explain high magnetite susceptibility and common volcanic lithic grains in this core. The coarse values of





Fig. 4. Representative hysteresis loop measurements demonstrating the narrow loops that characterize the northern Scotia Sea sediments of core 64 compared to the slightly broader loops of cores 29 and 31 (Note: sample TC29006 is from the trigger core of PC29).

Holocene grain size at this site (see also Howe & Pudsey 1999, fig. 8) are not fully understood; they cannot be attributed to localized input of ice-rafted material, since the ice-rafted debris content of core KC64 is very low (O'Coiffaigh *et al.* unpublished data).

### Conclusions

Magnetic properties investigated in this study of pelagic sediments from three Scotia Sea cores reveal the dominance of fine-grained magnetite as the remanence carrier. It is likely

that this is detrital in origin, as biogenic magnetite, if present, would yield magnetic parameter data closer to SD hysteresis values than those reported in this study. The Upper Quaternary sediments from the northern and central Scotia Sea have similar magnetic parameter values which reflect current-controlled contourite deposition of particles with uniform size and absence of significant post-depositional diagenetic alteration. These conclusions concerning the magnetic mineralogy are compatible with earlier sedimentological inferences that a fine fraction terrigenous component is transported by the Antarctic Circumpolar Current to sites in



**Fig. 5.** Hysteresis ratio plot after Day *et al.* (1977). SD = single domain, PSD = pseudo-single-domain, MD = multidomain, and SP = superparamagnetic particles. Holocene refers to stage 1 on core logs on Fig. 1, glacial refers to stages 2 and 3.

the Scotia Sea. This work demonstrates the potential utility of magnetic parameters as proxies for particle size and hence the strength of palaeocurrent flow; unlike bulk sediment measurements, they are unaffected by the presence of biogenic grains in the sample.

### Acknowledgements

Dr John Foster provided initial guidance and advice on the operation of the VFTB system. We thank Dr Andrew Roberts for reviewing an early version of this manuscript. M.E. Evans and an anonymous reviewer are thanked for their helpful and constructive comments.

### References

- BURCKLE, L.H. 1981. Displaced Antarctic diatoms in the Amirante Passage. *Marine Geology*, **39**, 39–43.
- CHARLES, C.D. & FAIRBANKS, R.G. 1992. Evidence from Southern Ocean sediments for the effect of North Atlantic deep-water flux on climate. *Nature*, **335**, 416–419.
- DAY, R., FULLER, M.D. & SCHMIDT, V.A. 1976. Magnetic hysteresis properties of synthetic titanomagnetites. *Journal of Geophysical Research*, **81**, 873–880.
- DAY, R., FULLER, M.D. & SCHMIDT, V.A. 1977. Hysteresis properties of titanomagnetites: grain size and compositional dependence. *Physics of the Earth and Planetary Interiors*, **13**, 260–267.
- DEKKERS, M.J. 1989. Magnetic properties of natural pyrrhotite. II. High- and low-temperature behaviour of  $J_{re}$  and TRM as a function of grain size. *Physics of the Earth and Planetary Interiors*, **57**, 266–283.
- DUNLOP, D.J. 1986. Hysteresis properties of magnetite and their dependence on particle size: a test of pseudo-single-domain remanence models. *Journal of Geophysical Research*, **91**, 9569–9584.
- DUNLOP, D.J. & OZDEMIR, O. 1997. *Rock magnetism: fundamentals and frontiers*. Cambridge: Cambridge University Press, 573 pp.
- FRANK, M., GERSONDE, R., RUTGERS VAN DER LOEFF, M., KUHN, G. & MANGINI, A. 1996. Late Quaternary sediment dating and quantification of lateral sediment redistribution applying  $^{230}\text{Th}_{ex}$ : a study from the eastern Atlantic sector of the Southern Ocean. *Geologische Rundschau*, **85**, 554–566.
- GLOERSEN, P., CAMPBELL, W.J., CAVALIERI, D.J., COMISO, J.C., PARKINSON, C.L. & ZWALLY, H.J. 1992. Arctic and Antarctic sea ice, 1978–1987: satellite passive microwave observations and analysis. NASA SP-511. Washington, DC: National Aeronautics & Space Administration, 290 pp.
- HOWE, J.A. & PUDSEY, C.J. 1999. Antarctic Circumpolar Deepwater: a Quaternary palaeoflow record from the northern Scotia Sea, South Atlantic Ocean. *Journal of Sedimentary Research*, **69**, 847–861.
- JORDAN, R.W. & PUDSEY, C.J. 1992. High-resolution diatom stratigraphy of Quaternary sediments from the Scotia Sea. *Marine Micropaleontology*, **19**, 201–237.
- MORETON, S.G. 1999. *Quaternary tephrochronology of the Scotia Sea and Bellingshausen Sea, Antarctica*. PhD thesis, Cheltenham & Gloucester CHE, 164 pp. [Unpublished.]
- MORETON, S.G. & SMELLIE, J.L. 1998. Identification and correlation of distal tephra layers in deep sea cores, Scotia Sea, Antarctica. *Annals of Glaciology*, **27**, 285–289.
- O'BRIEN, P.D. 1989. *The magnetostratigraphy of marine sediments from Jane Basin, southeast of the South Orkney Microcontinent, Antarctica*. PhD thesis, University of Southampton, 300 pp. [Unpublished].
- ORSI, A.H., WHITWORTH III, T. & NOWLIN JR, W.D. 1995. On the meridional extent and fronts of the Antarctic Circumpolar Current. *Deep-Sea Research*, **42**, 641–673.
- PUDSEY, C.J. 1992. Late Quaternary changes in Antarctic Bottom Water velocity inferred from sediment grain size in the northern Weddell Sea. *Marine Geology*, **107**, 9–33.
- PUDSEY, C.J. & HOWE, J.A. 1998. Quaternary history of the Antarctic Circumpolar Current: evidence from the Scotia Sea. *Marine Geology*, **148**, 83–112.
- ROBERTS, A.P. 1995. Magnetic properties of sedimentary greigite ( $\text{Fe}_3\text{S}_4$ ). *Earth and Planetary Science Letters*, **134**, 227–236.
- SAGNOTTI, L., FLORINDO, F., VEROSUB, K.L., WILSON, G.S. & ROBERTS, A.P. 1998. Environmental magnetic record of Antarctic palaeoclimate from Eocene/Oligocene glaciomarine sediments, Victoria Land Basin. *Geophysical Journal International*, **134**, 653–662.
- STONER, J.S., CHANNELL, J.E.T. & HILLAIRE-MARCEL, C. 1995. Magnetic properties of deep-sea sediments off southwest Greenland: Evidence for major differences between the last two deglaciations. *Geology*, **23**, 241–244.
- WASILEWSKI, P.J. 1973. Magnetic hysteresis in natural materials. *Earth and Planetary Science Letters*, **20**, 67–72.