

SLEEP HABITS IN CHILDREN AFFECTED BY AUTISM SPECTRUM DISORDERS: A PRELIMINARY CASE-CONTROL STUDY

FRANCESCO PRECENZANO¹, MARIA RUBERTO², LUCIA PARISI³, MARGHERITA SALERNO⁴, AGATA MALTESE³, DOMENICO VERDE¹, GABRIELE TRIPPI^{5,6}, PALMIRA ROMANO¹, ANNABELLA DI FOLCO³, TERESA DI FILIPPO³, GIOVANNI MESSINA⁷, MICHELE ROCCELLA³

¹Clinic of Child and Adolescent Neuropsychiatry; Headache Center for children and adolescents, Department of Mental Health and Physical and Preventive Medicine, Second University of Naples, Italy - ²Department of Medical-Surgical and Dental Specialties, Second University of Naples, Italy - ³Department of Psychological, Pedagogical and Educational Sciences, University of Palermo, Italy - ⁴Sciences for Mother and Child Health Promotion, University of Palermo, Italy - ⁵Department PROSAMI, University of Palermo, Italy - ⁶Childhood Psychiatric Service for Neurodevelopmental Disorders, CH Chinon, France - ⁷Department of Experimental Medicine, Section of Human Physiology and Unit of Dietetics and Sports Medicine, Department of Clinical and Experimental Medicine, University of Foggia, Foggia, Italy

**Francesco Precenzano and Maria Ruberto equally contributed to the manuscript*

ABSTRACT

Introduction: The core of autism spectrum disorders (ASD) consists of alterations of neurological functions that affect the typical developmental trajectory leading to deficits in social interaction and non-verbal behaviors. In general, ASD is diagnosed at age three and many other frequent neurological signs may be present, such as sleep disorders.

The aim of study is evaluating sleep habits in a sample of ASD children.

Material and methods: The study population consists of 65 ASD children (43 males and 22 females), aged between 2 and 11 years (mean 5.73 ± 2.39 years). The control group consists of 114 children with typical development (68 males and 46 females), aged between 3 and 10 years (mean 5.91 ± 2.23 years).

The diagnosis of ASD was confirmed by the evaluation ADOS -2, while the evaluation of sleep habits with the SDSC tests.

Results: The two groups were matched for age ($p = 0.633$) and sex distribution ($p = 0.483$).

The ASD subjects show a significantly higher rate of sleep disorders in all categories explored by SDSC tests (Table 1).

Discussion: The sleep of people with ASD is compromised at any age and regardless of the severity of social impairment and communication, but not the level of development.

Still, ASD is a very important risk factor for the development of sleep disorders, although the report could also be interpreted in the opposite manner.

Keywords: Autism Spectrum Disorders, sleep problems, DIMS, SDSC.

DOI: 10.19193/0393-6384_2017_3_059

Received October 30, 2016; Accepted January 02, 2017

Introduction

The core of autism spectrum disorders (ASD) consist in alterations among neurological functions impacting typical developmental trajectory with consequent deficits in social interaction and non-verbal behaviors. Generally, ASD is diagnosed at the age of three years and many frequent other neu-

rological signs may be present such as sleep troubles⁽¹⁻⁶⁾.

The more common sleep problems identified in ASD children include delayed sleep onset, night wakings after sleep onset, early morning wakings, decreased total sleep time, bedtime resistance, and sleep-onset association problems (eg, sleeping with a parent, sleeping in places other than the child's

own bed, and requiring idiosyncratic objects to initiate sleep onset). In general, sleep disturbances can interfere with cognition, attention, memory consolidation, and daytime behavioral adjustment in general. Therefore, in ASD children, sleep troubles may amplify already delayed social interactions, repetitive behaviors, affective problems, and inattention/hyperactivity⁽⁷⁻¹²⁾.

The aim of the present study is evaluating the sleep habits in a sample of ASD children.

Material and methods

Population

65 children (43 males and 22 females) affected by ASD according to the DSM-5 criteria, aged 2 to 11 years (mean age 5.73 ± 2.39) were recruited.

The diagnosis of ASD according to the Diagnostic and Statistical Manual of Mental Disorders 5th ed. (DSM-5) criteria was confirmed by an Autism Diagnostic Observation Schedule-2 (ADOS-2) assessment considered the gold-standard assessments. The average score at the ADOS scale in ASD children was 12.24 ($SD \pm 3.29$).

Exclusion criteria were the following: overweight ($z\text{-BMI} > 85$ pc) and obesity ($z\text{-BMI} > 95$ pc), cognitive disability ($IQ < 70$), neurological disorders (ie headaches, epilepsy), chromosomal syndromes (eg. Down, Prader-Willi, Crouzon, Pierre-Robin, trisomy 18), psychiatric illness (ie. mood disorders, anxiety disorders, psychosis) and specific neuropsychological disorders.

The control group consists of 114 typically developing children (68 males and 46 females), aged between 3 and 10 years (mean age 5.91 ± 2.23). Children of both groups are all Caucasians, and were recruited in the same urban area and similar for socio-economic status.

All parents have signed a specific informed consent prior psychological assessment and dissemination of results to scientific purposes.

The clinical study was conducted according to the principles of the Declaration of Helsinki.

Autism Diagnostic Observation Schedule (ADOS -2)

The ADOS is a semi-structured standardized observational assessment in which a clinician rates a participant's communication, reciprocal social behavior, and repetitive behaviors and stereotyped interest patterns. The ADOS consists of five modules and selection of the appropriate module is

based on an individual's chronological age and expressive language level. Module 4 emphasizes interview questions and conversations and is appropriate for use with verbally fluent older adolescents and adults who possess at least a minor level of independence in terms of relationships and goals. Inter-rater reliability for the ADOS algorithm ranged from 83 to 100 % for exact agreement, with j ranging from .41 to .84⁽¹⁻⁶⁾.

Sleep Disturbance Scale for Children (SDSC)

To evaluate sleep habits and disturbances, all of the subjects' mothers filled out the Sleep Disturbances Scale for Children (SDSC), 30 a standardized questionnaire for the assessment of sleep problems during development, consisting of 26 items grouped into six subscales: Disorders in Initiating and Maintaining Sleep (DIMS), Sleep Breathing Disorders (SBD), Disorders of Arousal (DA), Sleep-Wake Transition Disorders (SWTD), Disorders Of Excessive Somnolence (DOES), and Nocturnal Hyperhidrosis (SHY). Both the original or modified versions of this measure are widely used in school-aged children. According to the SDSC validation criteria,³⁰ scores ≥ 71 for the SDSC total score, ≥ 17 for DIMS, ≥ 7 for SBD, ≥ 6 for DA, ≥ 14 for SWTD, ≥ 13 for DOES, and ≥ 7 for SHY⁽³⁹⁾.

Statistic analysis

Chi-square and t-test were performed when appropriated, in order to compare the two population (ASD and typical developing children) for age, gender, sleep habits assessment.

Moreover, the logistic regression analysis (OR) was performed to evaluate the putative inter-relationship between ASD and sleep troubles. P values ≤ 0.05 were considered as statistical significant.

Results

The two groups are comparable for age ($p = 0.633$) and sex distribution ($p = 0.483$).

The ASD subjects show a significantly higher rate of sleep disorders in all categories explored by SDSC tests (Table 1).

Discussion

ASD is a neurodevelopmental disorder and can present with varying severity. During developmental age, sleep problems in ASD subjects may

ranging to about 80% versus the 25% in general pediatric population. Particularly, insomnia tends to range between 40-80% of ASD subjects.

	ASD (n = 65) (%)	Controlli (n = 114) (%)	OR	95% IC	Z	p
DIMS	30.77	2.63	16.44	4.655 - 58.088	4.349	0.0001
DRS	53.85	10.53	9.92	4.583 - 21.455	5.827	0.0001
DA	36.92	8.77	6.09	2.677 - 13.843	4.31	0.0001
DTVS	49.23	10.53	8.24	3.813- 17.817	5.363	0.0001
DES	29.23	7.89	4.82	2.027- 11.45	3.561	0.0004
IPN	23.08	4.39	6.54	2.252 - 18.991	3.453	0.0006
TOT	18.46	3.51	4.82	1.916- 20.224	3.043	0.0023

Table 1: shows comparison between children affected by autism spectrum disorders (ASD) and typical developing ones among Sleep Disturbance Scale for Children (SDSC) scales: Disorders in Initiating and Maintaining Sleep (DIMS); Sleep Breathing Disorders (SBD); Disorders of Arousal (DA); Sleep–Wake Transition Disorders (SWTD); Disorders Of Excessive Somnolence (DOES); Nocturnal Hyperhidrosis (SHY).

For comparison between the two groups (ASD and controls) t- Test was applied.

Logistic regression analysis was performed in order to calculate odd ratio (OR) and interval of confidence (IC). P values < 0.05 were considered statistically significant.

Among ASD children sleep problems are multifactorial, with genetic, environmental, immunological, and neurological factors such as melatonin rhythms impairment with alterations in this synchronization of the melatonin rhythm causing sleep problems⁽⁴⁰⁻⁴⁵⁾. Moreover, other neurotransmitters such as serotonin, and GABA are required for establishing a regular sleep wake cycle.

In autism, GABAergic interneurons migration and maturation may be affected. A region of genetic susceptibility has been identified on chromosome 15q that contains GABA-related genes. ASD represents a complex disease and articulated in several respects. Besides the core symptoms, in fact, there are many other complications that the clinician, therapist and parents are forced to face. The neurodegenerative disorders are, among these, the key for feeding difficulties notes, to pain and sensory perception. In this plight, an impairment in role of the orexin system regulation on the hypothalamus-pituitary-thyroid axis in ASD subjects may not be excluded⁽⁴⁰⁻⁴⁵⁾.

The sleep of people with ASD is compromised at any age and independently of the severity of impaired social and communicative sphere, but not the level of development (excluded in our study)⁽⁴⁶⁻⁵⁰⁾.

What emerges from the results of our survey shows that ASD is a very significant risk factor for developing sleep disorders, although the report could also be interpreted in the opposite manner: that have a disturbed sleep can be an aggravating factor in the nuclear symptoms ASD subjects, although further studies are needed in this respect

References

- 1) Parisi, L Di Filippo T, Roccella M. The child with autism spectrum disorders (ASD): behavioral and neurobiological aspects. *Acta Medica Mediterranea*, 2015, 31: 1187.
- 2) Parisi L, Di Filippo T, Roccella M. Behavioral Phenotype and Autism Spectrum Disorders in Cornelia de Lange Syndrome. *Ment Illn.* 2015 Sep 30;7(2):5988. doi: 10.4081/mi.2015.5988. eCollection 2015 Sep 30.
- 3) Parisi L, Di Filippo T, Roccella M. Autism spectrum disorder in Kabuki syndrome: clinical, diagnostic and rehabilitative aspects assessed through the presentation of three cases. *Minerva Pediatr.* 2015 Aug;67(4):369-75.
- 4) Vecchio D, Salzano E, Vecchio A, Di Filippo T, Roccella M. A case of femoral-facial syndrome in a patient with autism spectrum disorders. *Minerva Pediatr.* 2011 Aug; 63(4): 341-4.
- 5) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Gallai B, Marotta R, Lavano SM, Lavano F, Roccella M. Visual-spatial training efficacy in children affected by migraine without aura: a multicenter study. *Neuropsychiatr Dis Treat.* 2017 Jan 27; 13: 253-258. doi: 10.2147/NDT.S119648.
- 6) Miano S, Bruni O, Elia M, Trovato A, Smerieri A, Verrillo E, Roccella M, Terzano MG, Ferri R. Sleep in children with autistic spectrum disorder: a questionnaire and polysomnographic study. *Sleep Med.* 2007 Dec; 9(1): 64-70.
- 7) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Vagliano C, Messina G, Di Folco A, Di Filippo T, Michele Roccella. Executive functioning in preschool children affected by autism spectrum disorder: a pilot study. *Acta Medica Mediterranea*, 2017, 33: 35-39; DOI: 10.19193/0393-6384_2017_1_005;
- 8) Precenzano F, Lombardi P, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Magliulo RM, Messina G, Roccella M. Internalizing symptoms in children affected by childhood absence epilepsy: a preliminary study. *Acta Medica Mediterranea*, 2016, 32: 1749-1753; DOI: 10.19193/0393-6384_2016_6_158;
- 9) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Della Valle I, Visco G, Magliulo RM, Messina G, Roccella M. ADHD-like symptoms in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea*, 2016,

- 32: 1755-1759; DOI: 10.19193/0393-6384_2016_6_159;
- 10) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, D'alessandro I, Grappa MF, Magliulo RM, Messina G, Roccella M. Borderline intellectual functioning and parental stress: an italian case-control study. *Acta Medica Mediterranea*, 2016, 32: 1761-1765; DOI: 10.19193/0393-6384_2016_6_160;
 - 11) Ruberto M, Precenzano F, Parisi L, Salerno M, Maltese A, Messina G, Roccella M. Visuomotor integration skills in children affected by obstructive sleep apnea syndrome: a case-control study. *Acta Medica Mediterranea*, 2016, 32: 1659; DOI: 10.19193/0393-6384_2016_5_146;
 - 12) Parisi L, Ruberto M, Precenzano F, Di Filippo T, Russotto C, Maltese A, Salerno M, Roccella M. The quality of life in children with cerebral palsy. *Acta Medica Mediterranea*, 2016, 32: 1665; DOI: 10.19193/0393-6384_2016_5_147;
 - 13) Esposito M, Gallai B, Roccella M, Marotta R, Lavano F, Lavano SM, Mazzotta G, Bove D, Sorrentino M, Precenzano F, Carotenuto M. Anxiety and depression levels in prepubertal obese children: a case-control study. *Neuropsychiatr Dis Treat*. 2014 Oct 3; 10: 1897-902. doi: 10.2147/NDT.S69795
 - 14) Carotenuto M, Esposito M, Parisi L, Gallai B, Marotta R, Pascotto A, Roccella M. Depressive symptoms and childhood sleep apnea syndrome. *Neuropsychiatr Dis Treat*. 2012; 8: 369-73. doi: 10.2147/NDT.S35974
 - 15) Perillo L, Esposito M, Caprioglio A, Attanasio S, Santini AC, Carotenuto M. Orthodontic treatment need for adolescents in the Campania region: the malocclusion impact on self-concept. *Patient Prefer Adherence*. 2014 Mar 19; 8: 353-9. doi: 10.2147/PPA.S58971
 - 16) Carotenuto M, Esposito M, Di Pasquale F, De Stefano S, Santamaria F. Psychological, cognitive and maternal stress assessment in children with primary ciliary dyskinesia. *World J Pediatr*. 2013 Nov; 9(4): 312-7. doi:10.1007/s12519-013-0441-1
 - 17) Esposito M, Parisi L, Gallai B, Marotta R, Di Dona A, Lavano SM, Roccella M, Carotenuto M. Attachment styles in children affected by migraine without aura. *Neuropsychiatr Dis Treat*. 2013; 9: 1513-9. doi: 10.2147/NDT.S52716
 - 18) Esposito M, Roccella M, Gallai B, Parisi L, Lavano SM, Marotta R, Carotenuto M. Maternal personality profile of children affected by migraine. *Neuropsychiatr Dis Treat*. 2013; 9: 1351-8. doi: 10.2147/NDT.S51554.
 - 19) Perillo L, Esposito M, Contiello M, Lucchese A, Santini AC, Carotenuto M. Occlusal traits in developmental dyslexia: a preliminary study. *Neuropsychiatr Dis Treat*. 2013; 9: 1231-7. doi: 10.2147/NDT.S49985.
 - 20) Esposito M, Marotta R, Gallai B, Parisi L, Patriciello G, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Temperamental characteristics in childhood migraine without aura: a multicenter study. *Neuropsychiatr Dis Treat*. 2013; 9: 1187-92. doi:10.2147/NDT.S50458.
 - 21) Esposito M, Gallai B, Parisi L, Castaldo L, Marotta R, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. Self-concept evaluation and migraine without aura in childhood. *Neuropsychiatr Dis Treat*. 2013; 9: 1061-6. doi: 10.2147/NDT.S49364.
 - 22) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Maternal stress and childhood migraine: a new perspective on management. *Neuropsychiatr Dis Treat*. 2013; 9: 351-5. doi:10.2147/NDT.S42818.
 - 23) Esposito M, Ruberto M, Pascotto A, Carotenuto M. Nutraceutical preparations in childhood migraine prophylaxis: effects on headache outcomes including disability and behaviour. *Neurol Sci*. 2012 Dec; 33(6): 1365-8. doi: 10.1007/s10072-012-1019-8.
 - 24) Carotenuto M, Esposito M, Precenzano F, Castaldo L, Roccella M. Cosleeping in childhood migraine. *Minerva Pediatr*. 2011 Apr; 63(2): 105-9.
 - 25) Guzzetta A, Pizzardi A, Belmonti V, Boldrini A, Carotenuto M, D'Acunto G, Ferrari F, Fiori S, Gallo C, Ghirri P, Mercuri E, Romeo D, Roversi MF, Cioni G. Hand movements at 3 months predict later hemiplegia in term infants with neonatal cerebral infarction. *Dev Med Child Neurol*. 2010 Aug; 52(8): 767-72. doi:10.1111/j.1469-8749.2009.03497.x.
 - 26) Carotenuto M, Santoro N, Grandone A, Santoro E, Pascotto C, Pascotto A, Perrone L, del Giudice EM. The insulin gene variable number of tandem repeats (INS-VNTR) genotype and sleep disordered breathing in childhood obesity. *J Endocrinol Invest*. 2009 Oct; 32(9): 752-5. doi: 10.3275/6398.
 - 27) Esposito M, Carotenuto M, Roccella M. Primary nocturnal enuresis and learning disability. *Minerva Pediatr*. 2011 Apr; 63(2): 99-104.
 - 28) Esposito M, Carotenuto M. Borderline intellectual functioning and sleep: the role of cyclic alternating pattern. *Neurosci Lett*. 2010 Nov 19; 485(2): 89-93. doi:10.1016/j.neulet.2010.08.062.
 - 29) Carotenuto M, Esposito M, D'Aniello A, Ripa CD, Precenzano F, Pascotto A, Bravaccio C, Elia M. Polysomnographic findings in Rett syndrome: a case-control study. *Sleep Breath*. 2013 Mar; 17(1): 93-8. doi: 10.1007/s11325-012-0654-x.
 - 30) Esposito M, Antinolfi L, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Precenzano F, Carotenuto M. Executive dysfunction in children affected by obstructive sleep apnea syndrome: an observational study. *Neuropsychiatr Dis Treat*. 2013; 9: 1087-94. doi: 10.2147/NDT.S47287.
 - 31) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Patriciello G, Precenzano F, Carotenuto M. Visuomotor competencies and primary monosymptomatic nocturnal enuresis in prepubertal aged children. *Neuropsychiatr Dis Treat*. 2013; 9: 921-6. doi: 10.2147/NDT.S46772.
 - 32) Esposito M, Parisi P, Miano S, Carotenuto M. Migraine and periodic limb movement disorders in sleep in children: a preliminary case-control study. *J Headache Pain*. 2013 Jul 1; 14:57. doi: 10.1186/1129-2377-14-57.
 - 33) Carotenuto M, Gimigliano F, Fiordelisi G, Ruberto M, Esposito M. Positional abnormalities during sleep in children affected by obstructive sleep apnea syndrome: the putative role of kinetic muscular chains. *Med Hypotheses*. 2013 Aug; 81(2): 306-8. doi: 10.1016/j.mehy.2013.04.023.
 - 34) Carotenuto M, Esposito M. Nutraceuticals safety and efficacy in migraine without aura in a population of children affected by neurofibromatosis type I. *Neurol Sci*. 2013 Nov; 34(11): 1905-9. doi: 10.1007/s10072-

- 013-1403-z.
- 35) Esposito M, Carotenuto M. Intellectual disabilities and power spectra analysis during sleep: a new perspective on borderline intellectual functioning. *J Intellect Disabil Res.* 2014 May; 58(5): 421-9. doi: 10.1111/jir.12036.
- 36) Carotenuto M, Gallai B, Parisi L, Roccella M, Esposito M. Acupuncture therapy for insomnia in adolescents: a polysomnographic study. *Neuropsychiatr Dis Treat.* 2013; 9: 157-62. doi: 10.2147/NDT.S41892.
- 37) Esposito M, Pascotto A, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Gritti A, Mazzotta G, Carotenuto M. Can headache impair intellectual abilities in children? An observational study. *Neuropsychiatr Dis Treat.* 2012; 8: 509-13. doi:10.2147/NDT.S36863.
- 38) Esposito M, Verrotti A, Gimigliano F, Ruberto M, Agostinelli S, Scuccimarra G, Pascotto A, Carotenuto M. Motor coordination impairment and migraine in children: a new comorbidity? *Eur J Pediatr.* 2012 Nov;171(11):1599-604. doi:10.1007/s00431-012-1759-8.
- 39) Carotenuto M, Esposito M, Cortese S, Laino D, Verrotti A. Children with developmental dyslexia showed greater sleep disturbances than controls, including problems initiating and maintaining sleep. *Acta Paediatr.* 2016 Sep; 105(9): 1079-82. doi: 10.1111/apa.13472.
- 40) Pasquali D, Carotenuto M, Leporati P, Esposito M, Antinolfi L, Esposito D, Accardo G, Carella C, Chiovato L, Rotondi M. Maternal hypothyroidism and subsequent neuropsychological outcome of the progeny: a family portrait. *Endocrine.* 2015 Dec; 50(3): 797-801. doi: 10.1007/s12020-015-0564-3.
- 41) Santamaria F, Esposito M, Montella S, Cantone E, Mollica C, De Stefano S, Mirra V, Carotenuto M. Sleep disordered breathing and airway disease in primary ciliary dyskinesia. *Respirology.* 2014 May; 19(4): 570-5. doi: 10.1111/resp.12273.
- 42) Esposito M, Ruberto M, Gimigliano F, Marotta R, Gallai B, Parisi L, Lavano SM, Roccella M, Carotenuto M. Effectiveness and safety of Nintendo Wii Fit Plus™ training in children with migraine without aura: a preliminary study. *Neuropsychiatr Dis Treat.* 2013; 9: 1803-10. doi: 10.2147/NDT.S53853.
- 43) Di Filippo T, Orlando MF, Concialdi G, La Grutta S, Lo Baido R, Epifanio MS, Esposito M, Carotenuto M, Parisi L, Roccella M. The quality of life in developing age children with celiac disease. *Minerva Pediatr.* 2013 Dec; 65(6): 599-608.
- 44) Garon NM, Piccinin C, Smith IM. Does the BRIEF-P Predict Specific Executive Function Components in Preschoolers? *Appl Neuropsychol Child.* 2016; 5(2): 110-8. doi: 10.1080/21622965.2014.1002923;
- 45) Messina A, De Fusco C, Monda V, Esposito M, Moscatelli F, Valenzano A, Carotenuto M, Viggiano E, Chieffi S, De Luca V, Cibelli G, Monda M, Messina G. Role of the Orexin System on the Hypothalamus-Pituitary-Thyroid Axis. *Front Neural Circuits.* 2016 Aug 25; 10: 66. doi: 10.3389/fncir.2016.00066.
- 46) Messina G, Di Bernardo G, Viggiano A, De Luca V, Monda V, Messina A, Chieffi S, Galderisi U, Monda M. Exercise increases the level of plasma orexin A in humans. *J Basic Clin Physiol Pharmacol.* 2016 Sep 26. pii: /j/jbcpp.ahead-of-print/jbcpp-2015-0133/jbcpp-2015-0133.xml. doi: 10.1515/jbcpp-2015-0133.
- 47) Parisi L, Faraldo MA, Ruberto M, Salerno M, Maltese A, Di Folco A, Messina G, Di Filippo T, Roccella M. Life events and primary monosymptomatic nocturnal enuresis: a pediatric pilot study. *Acta Medica Mediterranea,* 2017, 33: 23-27; DOI: 10.19193/0393-6384_2017_1_003.
- 48) Precenzano F, Ruberto M, Parisi L, Salerno M, Maltese A, Vagliano C, Messina G, Di Folco A, Di Filippo T, Roccella M. Executive functioning in preschool children affected by autism spectrum disorder: a pilot study. *Acta Medica Mediterranea,* 2017, 33: 35-39; DOI: 10.19193/0393-6384_2017_1_005.
- 49) Parisi L, Fortunato MR, Salerno M, Maltese A, Di Folco A, Di Filippo T, Roccella M. Sensory perception in preschool children affected by autism spectrum disorder: a pilot study. *Acta Medica Mediterranea,* 2017, 33: 49-53; DOI: 10.19193/0393-6384_2017_1_007.
- 50) Toldo I, Rattin M, Perissinotto E, De Carlo D, Bolzonella B, Nosadini M, Rossi LN, Vecchio A, Simonati A, Carotenuto M, Scalas C, Sciruicchio V, Raieli V, Mazzotta G, Tozzi E, Valeriani M, Cianchetti C, Balottin U, Guidetti V, Sartori S, Battistella PA. Survey on treatments for primary headaches in 13 specialized juvenile Headache Centers: The first multicenter Italian study. *Eur J Paediatr Neurol.* 2016 Dec 31. pii: S1090-3798(16)30299-9. doi: 10.1016/j.ejpn.2016.12.009

Corresponding author

MICHELE ROCCELLA; MD; PhD
 Department of Psychological,
 Pedagogical and Educational Sciences
 University of Palermo
 (Italy)