

Experimental extractable work-based multipartite separability criteria

Mario A. Ciampini,^{1,*} Luca Mancino,^{1,†} Adeline Orioux,^{1,2} Caterina
Vigliar,¹ Paolo Mataloni,¹ Mauro Paternostro,³ and Marco Barbieri⁴

¹*Dipartimento di Fisica, Sapienza Università di Roma, P.le Aldo Moro 5, 00185, Rome, Italy*

²*LTCI, CNRS, Télécom ParisTech, Université Paris-Saclay, 75013, Paris, France*

³*Centre for Theoretical Atomic, Molecular and Optical Physics,*

School of Mathematics and Physics, Queen's University Belfast, Belfast BT7 1NN, United Kingdom

⁴*Dipartimento di Scienze, Università degli Studi Roma Tre, Via della Vasca Navale 84, 00146, Rome, Italy*

A thermodynamic theory of quantum entanglement, as well as the establishment of rigorous formal connections between the laws of thermodynamics and the phenomenology of entanglement are currently open areas of investigation. In this quest, a core problem is embodied by the understanding of the role that entanglement plays in processes of work extraction. Here, by considering information thermodynamics-based protocols, we answer the question “*Is it possible to infer, quantitatively, quantum correlations by considering work-extraction schemes?*”. Our experimental settings consist of suitably designed multi-photon optical interferometers able to address the case of both bipartite and multipartite entangled states. We compare the performance of such criteria to that of witnesses of entanglement based on the violation of Bell-like tests, showing their inherently different nature. Our work contributes strongly to the ongoing efforts in establishing photonic systems as a platform for experiments in information thermodynamics.

* marioarnolfo.ciampini@uniroma1.it

† Present address: Dipartimento di Scienze, Università degli Studi Roma Tre, Via della Vasca Navale 84, 00146, Rome, Italy

A. Introduction

Thermodynamics is one of the pillars of physical, chemical and biological sciences [1]. It predicts and explains the occurrence and efficiency of complex chemical reactions and biological processes. In physics and engineering, the conduction of heat across a medium, the concept of the arrow of time [2, 3], and the efficiency of motors are formulated in thermodynamic terms [4–7]. In information theory, the *mantra* that “information is physical” simply entails that, by being encoded in physical supports, information must obey the laws of thermodynamics [8, 9]. Even more, the tightness of the link between information and thermodynamics can be appreciated from the thermodynamic interpretation of the landmark embodied by Landauer’s erasure principle [10], Jaynes principle of maximum entropy [11, 12], or the information theoretical exorcism of Maxwell’s demon operated by Charlie Bennett [13–15]. More recently, it has been realized that thermodynamics can be used to provide a new way of assessing and exploiting quantum dynamics, build super-efficient nano- and micro-engines that take advantage of quantum coherence, and assess information theory from the perspective of thermodynamic costs [16–19].

The identification of the specific features of quantum systems that might influence their thermodynamic performance is still an open challenge [20, 21]: quite controversially, perfectly reversible work extraction from an ensemble of non-interacting quantum information carriers appears to be possible without the need for creating any entanglement [22]. Moreover, while entangling quantum operations allow for the retrieval of the same amount of work as in the classical case but with more economic protocols, such advantage appear to fade away in the case of a large number of systems [23].

Within such an interesting quest for the origin of “quantumness” in quantum thermodynamics, a core achievement is represented by thermodynamics-inspired witness of (multipartite) entanglement in a working medium [24, 25]. Such criteria can be cast as Bell-like games, in which the payoffs do correspond univocally to thermodynamics-inspired quantities. Having in mind the Szilard engine-version of Maxwell’s paradox, one can build one such game, corresponding to an enhanced extractable work from an hypothetical heat bath when using an entangled working medium. This all provides thermodynamic-inspired inequalities that are satisfied by any separable state of the working medium, but are violated by entangled states.

Such approaches to inseparability in the case of physical working media prepared in quantum laboratories has not yet been addressed due to the scarcity of experiments addressing the quantum thermodynamic framework, of which we only have a handful to date [26–30].

B. Results

Here, we bridge this gap by investigating experimentally thermodynamics-based separability criteria using non-ideal entangled two- and three-qubit quantum working media. We demonstrate the attainability of a maximal violation of the thermodynamic separability criterion, which would be thus equivalent to the achievement of the maximal extractable work allowed by the availability of non-classical correlations. The similarity between the realisation of the game at the basis of the thermodynamic inseparability criterion studied here and Bell-like tests allows for the comparison between such approaches to the characterisation of quantum correlations. We experimentally show that non-classical correlations empowering the extractable work bear differences to those responsible for the violation of Bell-type inequalities. We believe that, through so far unexplored theoretical tools, our results pioneer novel experimental methods towards the understanding of how non-classical correlations can effect the efficiency of quantum thermodynamic processes, at equal footing with thermodynamic potentials. This is a topical problem of both foundational and pragmatic relevance for the development of future engines operating at the nanoscale and in the fully quantum regime.

Let us set the thermodynamic context that serves the basis for the interpretation of the thermodynamic criterion for separability. Consider a single-particle gas in a Szilard machine-like device, i.e. a split chamber that be put in contact with a thermal reservoir at temperature T . By determining whether the particle occupies the left or right half-chamber, we can extract $k_B T \ln 2$ of work out of the reservoir simply by letting the gas equilibrate with it (isothermally). Here, k_B is Boltzmann’s constant. Suppose now that only partial information about the position of the particle in the chamber is available. In this case, the work that could in principle be extracted is $(k_B T \ln 2)[1 - H(X)]$, where $H(X)$ is the Shannon entropy of the binary position X of the particle [24]. Here, $k_B T \ln 2$ is a single scaling factor whose inclusion little adds to the physics of the problem. From here on, we thus rescale work in units of $k_B T \ln 2$, and focus solely on the “information-theoretical” gain $1 - H(X)$, which encompasses the core of the extractable-work scheme illustrated here. In this sense, while it should now be clear that we are addressing explicitly an information-thermodynamic problem, we can elevate the reservoir at the role of a virtual entity with limited physical relevance in the context here illustrated, besides the establishment of a “gauge” for the quantification of extractable work.

Let us now extend the scenario to the quantum domain and use projective operations to acquire information on the state of a two-level system that embodies the counterpart of the one-particle gas in the split chamber. In this case, we can store the measurement results in classical bits, so that the same process as above can be applied. Clearly, the entropy $H(X)$ is associated to the outcomes of a measurement on a two-level state. The work that could be extracted is then associated purely to the measurement and not to the dynamics of the state. This allows us to focus on the information brought about by the state of the system, rather than the implementation of an actual protocol for the extraction.

Moving from the single-particle to the bipartite case, we take inspiration from the fact that Maxwell had in mind Greek daimones, i.e. powerful entities bridging the human and the divine. We shall thus formulate our approach in the language of a general game involving two daemons, Aletheia ($A\lambda\theta\eta\epsilon\iota\alpha$) and Bia ($B\iota\alpha$), each attempting at extracting work from a two-qubit system.

They share an ensemble of qubit pairs, each prepared in the same bipartite state ρ . Moreover, they agree on the choice of two sets of measurement operators (one per daemon), that densely cover a great circle on the Bloch sphere of a single qubit. For half of the ensemble, Aletheia should choose from the set $\{\hat{A}_\theta\}$ the projection operator to use in order to measure her qubit. For the remaining half of the ensemble, Bia shall pick her measurement operator from the set $\{\hat{B}_{\theta'}\}$. In these sets, θ and θ' stand for the angular position, over a great circle of their respective single-qubit Bloch sphere, of the measurements performed by the daemons and $\theta, \theta'=0$ correspond to the z axis. In a given run of the game, Aletheia measures operator \hat{A}_θ and communicates the corresponding outcome A_θ to Bia, who attempts at evaluating the extractable work owing to the information gathered, locally, by Aletheia's measurement. In fact, the latter allows Bia to reduce the entropy associated to her measurement result from the value $H(B_{\theta'})$ to $H(B_{\theta'}|A_\theta)$ [24]. With such information, using a Szilard-like machine, Bia would be able to extract an amount of work $1-H(B_{\theta'}|A_\theta)$ from a hypothetical heat reservoir. An analogous reasoning can be carried out when exchanging the role of the daemons, concluding that, for a given setting, the extractable work per run of the game is

$$W_\rho(A_\theta, B_{\theta'}) = 1 - H(A_\theta, B_{\theta'}) + \frac{1}{2}[H(A_\theta) + H(B_{\theta'})], \quad (1)$$

where $H(A_\theta, B_{\theta'})$ is the joint entropy of variables A_θ and $B_{\theta'}$. For a sufficiently dense covering of the great circle, and for $\theta' = \theta$, the average extractable work achieved through the game, maximised over all the possible great circles, is

$\mathcal{W}(\rho) = \max_\phi \int_0^{2\pi} W_\rho(A_\theta, B_\theta) d\theta / 2\pi$, where ϕ is the azimuthal angle for the great circles being considered. In Ref. [25] it was shown that any separable state ρ is such that

$$\mathcal{W}(\rho) \leq \mathcal{W}_f = 0.443 \text{ bits}. \quad (2)$$

where \mathcal{W}_f is the maximum average work evaluated for any pure factorized state. Therefore, any surplus of extractable work above \mathcal{W}_f arises in virtue of entanglement shared by the daemons. Eq. (2) can then be used as a thermodynamically rooted entropic witness for inseparability.

We employ the thermodynamics-based separability criterion in Eq. (2) for the characterisation of photonic entanglement. As stated, our aim is to use information thermodynamics to assess entanglement, rather than replicating the behaviour of a working medium in a photonic setup [26]. Polarization-entangled pairs of qubits, ideally prepared in $|\Phi\rangle = \cos\varphi|HH\rangle + \sin\varphi|VV\rangle$, are produced by using the double-pass source in Ref. [31] [see Fig. 1(a)]. Here, $|H\rangle$ ($|V\rangle$) stands for a photon in the horizontal (vertical) polarization state. Further, we explore how the thermodynamic bound evolves through a depolarising channel (of strength $1 - \mu$) degrading the initial state to the mixture $\rho = \mu|\Phi\rangle\langle\Phi| + (1 - \mu)\mathbb{I}/4$. This has been implemented by using two liquid crystals, with their axes at 0° and 45° , onto the path of one photon. The two crystals are tuned in such a way that they perform a rapid succession of Pauli operators $\hat{\sigma}_x, \hat{\sigma}_y, \hat{\sigma}_z$, thus generating white noise on the state [32, 33].

In the inset of Fig. 1 (a), we show typical experimental curves for the work $W_\rho(\theta, \theta)$ calculated according to Eq. (1), as we inspect the set of operator pairs used by Aletheia and Bia, labelled by the angle θ on the great circle of the Bloch sphere corresponding to the linear polarizations (i.e. the equator of the single-qubit Bloch sphere). This is the set of directions allowing for optimal work extraction (cf. Methods). The observed oscillations are due to the polarization unbalance, while the average level is affected by the purity of the state. For each choice of θ , we have calculated the Shannon entropy associated to single-particle operators, as well as the joint entropy associated to the joint operator. We have verified by numerical simulations that our covering with 19 different directions is sufficiently dense for the continuous approximation to hold. In addition, we have fully characterised our states with quantum state tomography, from which the maximal value of the Bell operator S has been extracted for a comparison with a standard entanglement witness.

We measured the extractable thermodynamic work for several sets of parameters φ and μ in the resource state, which were estimated by reconstructing the density matrix of the experimentally engineered state ρ_{exp} via complete quantum state tomography for each position of the QWP in the source (setting φ) and every configuration of the LCs (fixing μ). Extractable work has been calculated by measuring Shannon entropy spanned over the whole big circle of the Bloch Sphere.

In order to highlight the inherently different nature of the test implemented here and a standard Bell-CHSH test, we now consider the Bell function $S = \text{Tr}[\rho(\hat{O}_1 - \hat{O}_2 + \hat{O}_3 + \hat{O}_4)]$ with $\hat{O}_1 = \sigma_x \otimes \sigma_x(\alpha)$, $\hat{O}_2 = \sigma_x \otimes \sigma_z(\alpha)$, $\hat{O}_3 = \sigma_z \otimes \sigma_x(\alpha)$, and $\hat{O}_4 = \sigma_z \otimes \sigma_z(\alpha)$. Here, $\sigma_k(\alpha) = \hat{R}(\alpha)\sigma_k\hat{R}^\dagger(\alpha)$ with $\hat{R}(\alpha) = \cos\alpha\mathbb{1} + i\sin\alpha\sigma_y$ and σ_k is the $k = x, y, z$ Pauli matrix. Local realistic theories bound such function as $|S| \leq 2$, while quantum mechanically $|S| \leq 2\sqrt{2}$. Using state ρ , we find $S = \mu[1 + \sin(2\varphi)][\cos(2\alpha) + \sin(2\alpha)]$, which we have used to estimate the value of the Bell function using ρ_{exp} for each set value of μ and φ .

The results of our experiment are summarised in Fig. 2, where we show the measured extractable work and the Bell function for two resource states with different values of φ . The comparison with Bell's test reveals how both quantities capture the degradation resulting from the depolarisation channel, and that they are similarly robust against the bias between the $|HH\rangle$

and $|VV\rangle$ contributions. A complete analysis shows that the relation between the thermodynamics-inspired criterion and Bell's inequalities is not trivial. In fact, we were able to find values of μ and φ for which it is possible to violate just one of the two criteria, and vice versa. We have considered an experimental state with large bias ($\cos \varphi = 0.25 \pm 0.03$) and purity ($\mu = 0.97$) that offers no violation of the local realistic bound ($S = 1.977 \pm 0.009$). However, the corresponding extractable work is $\mathcal{W} = 0.491 \pm 0.021$, which exceeds the separability threshold \mathcal{W}_f by 2 standard deviations. This implies that the use of entanglement for powering a thermal machine is qualitatively different from that necessary for communication tasks, such as measurement-independent key distribution. These considerations are extended in the analysis reported in the Supplementary Information.

We now extend our investigation to the case of a tripartite system. It is well known that, for pure states, only two non-equivalent classes of genuine tripartite entanglement exist [35], either the GHZ or W class. Here, we focus on the ability of distinguishing GHZ-type entanglement from W -type one using a criterion inspired on thermodynamics rather than state fidelity [36]. The protocol now involves a further daemon, Charis ($X\acute{\alpha}\rho\iota\zeta$), which is set to act on her subsystem by acting in a way conditioned on information provided by Aletheia and Bia. The strategy they agreed on consists of the following steps: **a)** Aletheia and Bia perform a projective measurement along a common axis u in the single-qubit Bloch sphere chosen among the three Pauli settings; **b)** Charis receives information on the outcomes of such measurements. In light of such information, she can extract work from her system by performing a projective measurement along a suitably chosen direction v on the Bloch sphere; **c)** In order to exclude bipartite entanglement and thus link the excess extractable work to the presence of genuine multipartite entanglement in our resource state, the protocol must be repeated by permuting the role of the three daemons.

Clearly, the amount of work that can be obtained in a single run of such a tripartite game is $W_\rho(A_u, B_u, C_v) = 1 - H(C_v|A_u, B_u)$ with $H(C_v|A_u, B_u)$ the Shannon entropy of variable C_v conditioned on the outcomes A_u and B_u performed by Aletheia and Bia. A bound for the average extractable work, obtained by maximizing over the choice of v , can be established for separable states as

$$\mathcal{W}(\rho) = \max_v \frac{1}{3} \sum_u W_\rho(A_u, B_u, C_v) \leq \mathcal{W}_f = \frac{1}{3}. \quad (3)$$

A violation of this bound signals the presence of entanglement in the state. Furthermore, it can be shown that W -type entanglement can only provide an extractable work of at most $7/9 = 0.778$ [25]. Therefore, this thermodynamic criterion is well suited for identifying the entanglement class to which the state belongs: any extractable work in excess of 0.778 would signal GHZ character.

In order to achieve useful benchmarks for the performance of our experiments, we have performed a thorough theoretical analysis of the protocol for both pure and mixed-state resources. In particular, we have considered the achievable value of the extractable work \mathcal{W} when using the resource state $\rho_3 = \mu|\Psi\rangle\langle\Psi| + (1 - \mu)\mathbb{I}/8$ with $|\Psi\rangle$ a tripartite pure state. We have first focused on the case of $\mu = 1$ with $|\Psi\rangle$ embodied by either a tripartite GHZ-Cluster state of the form

$$|GHZc\rangle = \frac{1}{2}(|000\rangle - |110\rangle + |011\rangle + |101\rangle)_{ABC} \quad (4)$$

where $\{|0\rangle, |1\rangle\}$ are the logical states of the three qubits, labelled as A , B , and C and owned by Aletheia, Bia, and Charis respectively. This state can be shown to give rise to the same performance as a standard GHZ resource. The W -type resource is instead

$$|W\rangle = \frac{1}{\sqrt{3}}(|001\rangle + |010\rangle + |100\rangle)_{ABC}. \quad (5)$$

In Supplementary Information we have looked at the expected behaviour of \mathcal{W} against the angular direction of the great circle on the single-qubit Bloch sphere identified by the angles (θ, ϕ) , both for pure and mixed states.

Such predictions have then been tested in a two-photon implementation of both a three-qubit GHZ-Cluster type state and a W type one. Such states are generated by adding a path-encoded qubit to the two-qubit setup, as shown in Fig. 1 **(b)** and **(c)** (cf. Methods). This arrangement allowed us to engineer both a GHZ-Cluster state of the form $(|HH0\rangle - |VV0\rangle + |HV1\rangle + |VH1\rangle)_{ABC}/2$ and a W -type resource state of the form $|W\rangle = (|HH1\rangle + |HV0\rangle + |VH0\rangle)/\sqrt{3}$. Path-qubit analysis was performed by either selecting one of the two paths, which implements a projection in the computational basis, or selecting the proper phase shift ϕ between the two modes, thus enabling the projection on the diagonal basis ($\phi = 0, \pi$) and the circular basis ($\phi = \pi/2, -\pi/2$).

In all cases, there is a significant violation of the separability bound. Moreover, the symmetry among the results that we have achieved capture unambiguously the GHZ-Cluster character of the experimental resource. In order to show this feature more clearly, we have repeated the experiment when complete decoherence is introduced in one of the qubits of the system, in an attempt to render our resource state bi-separable and wash out its genuine tripartite entanglement. When full decoherence is introduced in the path-encoded qubit by inserting a thick glass plate on one of the paths, allowing to disrupt coherence between the two modes (independent of polarization), we get $\mathcal{W}_1 = 0.799 \pm 0.006$. However, decoherence lessens the values of the other

173 two witnesses, to values that are very close to the separability threshold in Eq. (3), demonstrating the disappearance of tripartite
174 entanglement.

175 A similar analysis has been conducted using the experimental W -type resource, whose corresponding experimental results are
176 reported in Table II. As done for the GHZ-Cluster state, we have assessed both the case of a close-to-ideal resource and that of
177 a strongly decohered one close to the separability threshold. The diversity (in terms of entanglement-sharing structure) of such
178 resources is fully captured by the extractable work.

179 A comparison with non-locality-based criteria similar to the one reported for the study of bipartite entanglement can be
180 made in this tripartite context. Here, rather than concentrating on two-qubit correlations, we shall investigate their genuinely
181 multipartite nature, whose characterisation is clearly a much more demanding problem to tackle. A powerful tool in this respect
182 is embodied by genuinely tripartite versions of a Bell inequality, such as the inequality proposed by Svetlichny [37], whose
183 violation witnesses the occurrence of multipartite non-local correlations. Such inequality has been successfully experimentally
184 violated using a photonic GHZ state [38] and used to ascertain the properties of the ground state of a many-body system in a
185 photonics quantum simulator [39].

186 In the Svetlichny game, the daemons locally rotate their respective qubit by an angle α_j ($j = A, B, C$) through the operator
187 $\hat{\mathcal{R}}(\alpha_j) = \cos \alpha_j \sigma_z + \sin \alpha_j \sigma_x$ and project it over the basis of σ_z . If state $|0\rangle$ ($|1\rangle$) is found, they attach a dichotomic variable
188 the value $+1$ (-1). This allows them to build the the statistical correlation function for local spin measurements $E(\alpha_A, \alpha_B, \alpha_C)$
189 and, in turn, construct the Mermin-Ardehali-Belinskii-Klyshko function [40, 41]

$$M_3 = E(\alpha_A, \alpha_B, \alpha'_C) + E(\alpha_A, \alpha'_B, \alpha_C) + E(\alpha'_A, \alpha_B, \alpha_C) - E(\alpha'_A, \alpha'_B, \alpha'_C). \quad (6)$$

190 The Svetlichny function is thus $\mathcal{S}_3 = |M_3 + M'_3|$, where M'_3 is the same as M_3 with $\alpha_j \leftrightarrow \alpha'_j$. Any biseparable state satisfies
191 the inequality $|\mathcal{S}_3| \leq 4$. Tripartite entangled states violate the inequality up to the maximum value of $4\sqrt{2} = 5.65685$, which
192 is obtained using GHZ-like states. In fact, state $|GHZc\rangle$ allows to achieve such a maximum violation for $(\alpha_A, \theta_B, \theta_C) =$
193 $(3\pi/8, -\pi/4, 0)$ and $(\alpha'_A, \theta'_B, \theta'_C) = (\pi/8, 0, \pi/4)$. We have used the experimental quantum state tomographies of the GHZ-
194 Cluster and W resource states to evaluate the Svetlichny function. After a global optimisation over all the angles involved, we
195 have found $\mathcal{S}_3^{GHZc} = 4.83 \pm 0.07$ and $\mathcal{S}_3^W = 3.39 \pm 0.06$. This shows that the experimental GHZ-Cluster state is consistently
196 found to be entangled in a genuinely tripartite sense by both the extractable work-based criterion and the Svetlichny one. On the
197 other hand, the non locality-based entanglement criterion fails to detect the tripartite entangled nature of the experimental W
198 resource, which is instead well captured by the sensitivity exhibited by the extractable work.

199 C. Discussion

200 We have demonstrated experimentally a fundamental result of information thermodynamics, showing that communication-
201 assisted games based on Maxwell's daemon are able to provide core information on the quantum correlated nature of the state
202 of a multipartite system. Such protocols are both practical and fundamentally interesting in light of their distinct nature from,
203 say, non locality-based entanglement witnesses. This work has significant implications for both technological and conceptual
204 aspects. Concerning the technological advances that it entails, the photonic platform we described has the capability of preparing,
205 controlling and measuring, with high fidelity and a reduced experimental complexity, states of a multipartite working medium,
206 addressing questions that are genuinely inspired by thermodynamic problems. General evolutions can be easily implemented or
207 simulated, thus paving the way to the implementation of a fully fledged experimental assessment of non-equilibrium dynamics.
208 The latter has so far been limited to either classical systems or nuclear magnetic resonance setups, where any quantum feature is
209 weakened by the typically strong environmental actions. As for the fundamental aspects, we have highlighted how non-classical
210 correlations within the working medium are strongly linked to the performance of thermodynamic processes, not differently
211 from quantities with a counterpart in classical thermodynamics. This points to an interesting direction for understanding the
212 emergence of ordinary world from its quantum microscopic constituents.

213 D. Methods

Two-qubit protocol The bound on the extractable work is found by averaging over different projectctions by the daemons; these are performed by removing the quarter waveplates (QWPs) from the polarization analysis-module shown in Fig. 1, and using only the half waveplates (HWPs) to determine the direction of the projection. As we used two avalanche photo-diodes (APDs) after the polarising beam splitters (PBSs), we had to implement four different types of measurements for each of the 19 choices of angular direction on the equator of the single-qubit Bloch sphere. In particular, we had to consider the four sets of directions $(\theta_A, \theta'_B) = (\theta, \theta), (\theta, \theta + 45^\circ), (\theta + 45^\circ, \theta), (\theta + 45^\circ, \theta + 45^\circ)$, where the first (second) angular direction is for Aletheia's (Bia's) measurements choice. The corresponding detected coincidences at the APDs are labelled as N_{θ_A, θ'_B} . A full

circulation of the Bloch-sphere equator is then achieved by rotating the HWPs from 0° to 45° in 19 steps. This implies that a total of 76 measurements were needed in order to acquire the necessary information to reliably evaluate the performance of the scheme. Upon suitable normalization of the detected coincidence counts, we have the following set of probabilities

$$p_{AB}(\theta) = N_{\theta,\theta+45^\circ}/D(\theta), \quad (7)$$

$$p_A(\theta) = (N_{\theta,\theta} + N_{\theta+45^\circ,\theta})/D(\theta), \quad (8)$$

$$p_B(\theta) = (N_{\theta,\theta} + N_{\theta,\theta+45^\circ})/D(\theta) \quad (9)$$

with $D(\theta) = N_{\theta,\theta} + N_{\theta,\theta+45^\circ} + N_{\theta+45^\circ,\theta} + N_{\theta+45^\circ,\theta+45^\circ}$. Example of measured coincidence counts are $N_{0,0} = 3578$, $N_{0,45^\circ} = 58$, $N_{45^\circ,0} = 173$, $N_{45^\circ,45^\circ} = 4328$, achieved in 30s of measurement. These are instrumental to the evaluation of the Shannon entropies needed to calculate $W_\rho(\theta, \theta)$ and, in turn, \mathcal{W} .

Three-qubit protocol Our two-photon implementation of three-qubit states relies on the addition of a third qubit, encoded in the two paths within a Sagnac interferometer (consisting of a 50:50 BS and three mirrors) on one of the two photons; the logical state $|0\rangle$ correspond to the clockwise circulation of the photon inside the interferometer, and $|1\rangle$ to the anticlockwise circulation. The phase ϕ between such logical state can be tuned by tilting a thin glass plate, placed in one of the paths. This arrangement allowed us to engineer a GHZ-Cluster state of the form $(|HH0\rangle - |VV0\rangle + |HV1\rangle + |VH1\rangle)_{ABC}/2$ by introducing a HWP at 0° on the clockwise path $|0\rangle$ and a HWP at 45° on the anticlockwise one [cf. Fig. 1 (b)]. On the other hand, a W -type resource state of the form $|W\rangle = (|HH1\rangle + |HV0\rangle + |VH0\rangle)/\sqrt{3}$ can be easily engineered by modifying the configuration for the GHZ-Cluster state. After changing the label of the two paths, a PBS can be added in the setup to perform a polarization-to-path mapping, and thus eliminate the contribution from state $|VV1\rangle$ [cf. Fig. 1 (c)].

E. Acknowledgements

We are grateful to M. Vidrighin and V. Vedral for insightful feedback on this manuscript.

F. Contributions

MP MB and PM conceived the project. The experiment has been designed by AO and MAC, and performed by MAC with the assistance of LM and CV. Data was analysed and discussed by all the authors. MP wrote the manuscript, with input from all the authors.

G. Competing interests

The authors declare no conflict of interest.

H. Fundings

This work was partially supported by the project FP7-ICT-2011-9-600838 (QWAD Quantum Waveguides Application and Development; www.qwad-project.eu). MB is supported by a Rita Levi-Montalcini fellowship of MIUR. MP acknowledges financial support from the EU FP7 Collaborative Project TherMiQ, the John Templeton Foundation (grant number 43467), the Julian Schwinger Foundation (grant number JSF-14-7-0000), and the UK EPSRC (grant number EP/M003019/1).

I. References

-
- [1] Zemansky, M. W. & Dittman, R. H. *Heat and Thermodynamics* (McGraw-Hill, New York, NY, 1981).
 - [2] Eddington, A. S., *The Nature of the Physical World*, (McMillan, New York, NY, 1928).
 - [3] Campisi, M. & Hänggi, P. Fluctuation, Dissipation and the Arrow of Time, *Entropy* **13**, 2024 (2011).
 - [4] Van Wylen, G. J. & Sonntag, R. E. *Fundamentals of Classical Thermodynamics* (John Wiley & Sons, Hoboken, NJ, 1985).
 - [5] Callen, H. B. *Thermodynamics and an Introduction to thermostatistics* (John Wiley & Sons, Hoboken, NJ, 185).

- 245 [6] Sinitsyn, N. A. Fluctuation Relation for heat Engines, *J. Phys. A: Math. Theor.* **44**, 405001 (2011).
- 246 [7] Curzon, F. L. & Ahlborn, B. Efficiency of a Carnot Engine at Maximum Power, *Am. J. Phys.* **43**, 24 (2975).
- 247 [8] Landauer, R. Information is physical, *Phys. Today* **44**, 23 (1991).
- 248 [9] Lutz, E. & Ciliberto, S. Information: From Maxwell's demon to Landauer's eraser, *Phys. Today* **68**, 30 (2015).
- 249 [10] Landauer, R. Irreversibility and heat generation in the computing process, *IBM J. Res. Develop.* **5**, 183 (1961).
- 250 [11] Jaynes, E. T. Information Theory and Statistical Mechanics. II, *Phys. Rev.* **108**, 171 (1957).
- 251 [12] Cox, R. T. The Algebra of Probable Inference, *Am. J. Phys.* **31**, 66 (1963).
- 252 [13] Bennett, C. H. Notes on Landauer's principle, Reversible Computation and Maxwell's Demon, *Stud. Hist. Philos. Modern Phys.* **34**, 501
- 253 (2003).
- 254 [14] Plenio, M. B. & Vitelli, V. The physics of forgetting: Landauer's erasure principle and information theory, *Contemp. Phys.* **42**, 25 (2001).
- 255 [15] Maruyama, K., Nori, F. & Vedral, V. The Physics of Maxwell's demon and information, *Rev. Mod. Phys.* **81**, 1 (2009).
- 256 [16] Goold, J., Huber, M., Riera, A., del Rio, L. & Skrzypczyk, P. The role of quantum information in thermodynamics: a topical review, *J. Phys. A: Math. Theor* **49** 143001 (2016).
- 257 [17] Kosloff, R. Quantum Thermodynamics, *Entropy* **15**, 2100 (2013).
- 258 [18] Vinjanampathy, S. & Anders, J. Quantum Thermodynamics, *Contemporary Physics* **57** 545-579 (2016).
- 259 [19] Gelbwaser-Klimovsky, D., Niedenzu, W. & Kurizki, G. Thermodynamics of quantum systems under dynamical control, *Adv. Atom. Mol. Opt. Phys.* **64**, 329 (2015).
- 260 [20] Scully, M., Zubairy, M. S., Agarwal, G. S. & Walther, H. Extracting Work from a Single Heat Bath via Vanishing Quantum Coherence, *Science* **299**, 862 (2003).
- 261 [21] Uzdin, R., Levy, A. & Kosloff, R. Quantum equivalence and quantum signatures in heat engines, *Phys. Rev. X* **5**, 031044 (2015).
- 262 [22] Hovhannisyan, K. V., Perarnau-Llobet, M., Huber, M. & Acín, A. Entanglement Generation is Not Necessary for Optimal Work Extrac-
263 tion, *Phys. Rev. Lett.* **111**, 240401 (2013).
- 264 [23] Perarnau-Llobet, M., Hovhannisyan, K. V., Huber, M., Skrzypczyk, P., Brunner, N. & Acín, A. Extractable Work from Correlations,
265 *Phys. Rev. X* **5**, 041011 (2015).
- 266 [24] Maruyama, K., Morikoshi, F. & Vedral, V. Thermodynamical detection of entanglement by Maxwell's demons, *Phys. Rev. A* **71**, 012108
267 (2005).
- 268 [25] Vigiú, V., Maruyama, K. & Vedral, V. Work extraction from tripartite entanglement, *New J. Phys.* **7**, 195 (2005).
- 269 [26] Batalhão, T. B. *et al.*, Experimental reconstruction of work distribution and verification of fluctuation relations at the full quantum level,
270 *Phys. Rev. Lett.* **113**, 140601 (2014).
- 271 [27] Silva, J. P. P. *et al.* Experimental demonstration of information to energy conversion in a quantum system at the Landauer Limit, *Proc. R. Soc. A* **472**, 20150813 (2016).
- 272 [28] Raitz, C., Souza, A. M., Auccaise, R., Sarthour, R. S. & Oliveira, I. S. Experimental implementation of a nonthermalizing quantum
273 thermometer, *Quant. Inf. Process.* **14**, 37 (2015).
- 274 [29] Batalhão, T. B. *et al.* Irreversibility and the arrow of time in a quenched quantum system, *Phys. Rev. Lett.* **115**, 190601 (2015).
- 275 [30] Vidrighin, M. D. *et al.* *Phys. Rev. Lett.* **116** 050401 (2016).
- 276 [31] Cinelli, C., Di Nepi, G., De Martini, F., Barbieri, M. & Mataloni, P. Parametric source of two-photon states with a tunable degree of
277 entanglement and mixing: Experimental preparation of Werner states and maximally entangled mixed states, *Phys. Rev. A* **70**, 022321
278 (2004).
- 279 [32] Chiuri, A., Giacomini, S., Macchiavello, C. & Mataloni, P. Experimental achievement of the entanglement-assisted capacity for the
280 depolarizing channel, *Phys. Rev. A* **87**, 022333 (2013).
- 281 [33] Chiuri, A. *et al.* Experimental Realization of Optimal Noise Estimation for a General Pauli Channel, *Phys. Rev. Lett.* **107**, 253602 (2011).
- 282 [34] Horodecki, R., Horodecki, P., Horodecki, M. & Horodecki, K. Quantum entanglement, *Rev. Mod. Phys.* **81**, 865 (2009).
- 283 [35] Dür, W., Vidal, G. & Cirac, J. I. Three qubits can be entangled in two inequivalent ways, *Phys. Rev. A* **62**, 062314 (2000).
- 284 [36] Bourennane, M. *et al.* Witnessing multipartite entanglement, *Phys. Rev. Lett.* **92**, 087902 (2004).
- 285 [37] Svetlichny, G. Distinguishing three-body from two-body non-separability by a Bell-type inequality, *Phys. Rev. D* **35**, 3066 (1987).
- 286 [38] Lavoie, J., Kaltenbaek, R. & Resch, K. J. Experimental violation of Svetlichny's inequality, *New J. Phys.* **11**, 073051 (2009).
- 287 [39] Orioux, A., Boutari, J., Barbieri, M., Paternostro, M. & Mataloni, P. Experimental linear-optics simulation of the ground state of an
288 interacting quantum spin-ring, *Sci. Rep.* **4**, 7184 (2014).
- 289 [40] Mermin, N. D. Extreme quantum entanglement in a superposition of macroscopically distinct states, *Phys. Rev. Lett.* **65**, 1838 (1990).
- 290 [41] Klyshko, D. N. The Bell and GHZ theorems: a possible three-photon interference experiment and the question of nonlocality, *Phys. Lett. A* **172**, 399 (1993).
- 291
- 292
- 293
- 294
- 295

J. Figure Legends

(a) (b) (c)

Figure 1. **(a)** Experimental setup for the extractable work-based inseparability of bipartite states: The entangled photon source uses a 1.5mm thick β Barium-Borate (BBO) crystal pumped with 100 mW of laser at 355 nm, in conjunction with a spherical mirror (M), and delivers approximately 200 coincidences/s through 5 nm (FWHM) interference filters (IF). Details on the source can be found in Ref. [31]. We encode the logical states of each qubit in the horizontal and vertical polarization states $|H\rangle$ and $|V\rangle$ of each photon. Using quantum state tomography, we estimate a fidelity $F = 0.961 \pm 0.007$ of the entangled resource with the maximally entangled state $\frac{1}{\sqrt{2}}(|HH\rangle + |VV\rangle)$. The corresponding value of tangle is $T = 0.911 \pm 0.008$ [34]. The relative weight ($\cos \varphi$) of the two polarization contributions in such state can be tuned by the quarter waveplate (QWP_m) in the source. One of the photons passes through a depolarizing channel, consisting of two liquid crystals (LCs) and the associated control electronics, which selects the value of μ . Finally, polarization measurements are performed at a polarisation-analysis module consisting of a QWP, a half waveplate (HWP), a polarising beam splitter (PBS), and an avalanche photodiode (APD) per mode. We also show the single-mode fibers (SMF) used to convey the photonics signal to the polarization analysis module. Inset: measured $W_\rho(\theta, \theta)$ as a function of the measurement angle θ for state $|\rho\rangle$ with $\cos \varphi = 0.85$, and $\mu = 0.98$ (blue); $\cos \varphi = 0.62$, and $\mu = 0.98$ (purple); $\cos \varphi = 0.62$, and $\mu = 0.51$ (red). **(b)** Experimental setup for a GHZ-Cluster state. **(c)** Analogous setup for a W -type resource.

(a) (b)

Figure 2. Experimental results for two different resource states corresponding to $\cos \varphi = 0.62$ [panel **(a)**], and $\cos \varphi = 0.85$ [panel **(b)**] for different strengths of the depolarising channels μ . Red dots: experimental points for \mathcal{W} calculated from the experimental count rates measured in 30s. Magenta dots: experimental points corresponding to the values taken by the Bell parameter, evaluated from the fully reconstructed state tomographies of the resource ρ_{exp} . The value of $\cos \varphi$ is evaluated by direct inspection of the coincidence rates, averaged over all the experimental states. The error bars take into account the Poissonian statistics of the measured rates and derived from direct error propagation for \mathcal{W} and Monte Carlo simulation for S . The green and blue dashed curves represent theoretical predictions based on state ρ , while the solid straight lines denote the entanglement thresholds and the Bell function. The dashed vertical lines identify the values of μ for which \mathcal{W} and S cross their respective bounds.

Figure 3. Diagrammatic representation of the scheme extractable work-based tripartite inseparability. Each panel illustrates one of the steps of the protocol described in the main text.

K. Tables

\mathcal{W}_1^{GHZc}	\mathcal{W}_2^{GHZc}	\mathcal{W}_3^{GHZc}
0.798 ± 0.006	0.779 ± 0.005	0.795 ± 0.004
0.799 ± 0.006	0.304 ± 0.003	0.305 ± 0.003

Table I. Extractable work in bits for the three permutations of roles in the protocol for extractable work-based inseparability evaluated using a GHZ -Cluster resource. The three columns are for the value of the extractable work achieved from the path-encoded qubit, the first polarization-encoded qubit, and the second polarization-encoded one, respectively. The first row refers to a tripartite entangled GHZ -Cluster state whose fidelity with the ideal resource $(|HH0\rangle - |VV0\rangle + |HV1\rangle + |VV1\rangle)/2$ is found to be $F = 0.851 \pm 0.008$. The second row refers to the same resource where any coherence between path and polarization qubit is removed. Correspondingly, \mathcal{W}_2^{GHZ} and \mathcal{W}_3^{GHZ} are below the threshold for separability, showing that in the corresponding configuration we only have bipartite entanglement.

\mathcal{W}_1^W	\mathcal{W}_2^W	\mathcal{W}_3^W
0.4715 ± 0.0036	0.6625 ± 0.0050	0.5966 ± 0.0050
0.4715 ± 0.0036	0.3205 ± 0.0023	0.3282 ± 0.0027

Table II. Extractable work in bits for the three permutations of roles in the protocol for extractable work-based inseparability evaluated using a W resource. The three columns are for the value of the extractable work achieved from the path-encoded qubit, the first polarization-encoded qubit, and the second polarization-encoded one, respectively. The first row refers to a tripartite entangled W state very close to an ideal $|W\rangle$ state: The fidelity between the experimental resource and the ideal state $(|HH1\rangle + |HV0\rangle + |VH0\rangle)/\sqrt{3}$ is $F = 0.823 \pm 0.006$. The second row refers to the same W configuration where, however, any coherence between path-encoded and polarization-encoded qubits is removed. Correspondingly, \mathcal{W}_2^W and \mathcal{W}_3^W are below the threshold for separability, showing that in the the corresponding configuration we only have bipartite entanglement.