



Oyeyiola, A. O. and Davidson, C. M. and Olayinka, K. O. and Oluseyi, T. O. and Alo, B. I. (2013) Multivariate analysis of potentially toxic metals in sediments of a tropical coastal lagoon. Environmental Monitoring and Assessment, 185 (3). pp. 2167-2177. ISSN 0167-6369 , <http://dx.doi.org/10.1007/s10661-012-2697-7>

This version is available at <https://strathprints.strath.ac.uk/43019/>

Strathprints is designed to allow users to access the research output of the University of Strathclyde. Unless otherwise explicitly stated on the manuscript, Copyright © and Moral Rights for the papers on this site are retained by the individual authors and/or other copyright owners. Please check the manuscript for details of any other licences that may have been applied. You may not engage in further distribution of the material for any profitmaking activities or any commercial gain. You may freely distribute both the url (<https://strathprints.strath.ac.uk/>) and the content of this paper for research or private study, educational, or not-for-profit purposes without prior permission or charge.

Any correspondence concerning this service should be sent to the Strathprints administrator: strathprints@strath.ac.uk

1 **Multivariate Analysis of Potentially Toxic Metals in Sediments of a Tropical**
2 **Coastal Lagoon**

3 A.O Oyeyiola,^a C.M Davidson,^b K.O Olayinka,^a T.O Oluseyi^a and B.I. Alo^a

4 ^aDepartment of Chemistry, University of Lagos Akoka- Yaba, Lagos, Nigeria

5 ^bWestCHEM, Department of Pure and Applied Chemistry, University of Strathclyde, 295
6 Cathedral Street, Glasgow G1 1XL, UK

7
8
9 **Abstract**

10 Surface sediments collected from the Lagos Lagoon, Nigeria, and three adjoining rivers were
11 analysed for their physicochemical properties and pseudototal concentration of the potentially
12 toxic metals (PTM) Cd, Cr, Cu, Pb and Zn. The concentration of the PTM varied seasonally and
13 spatially. Odo-Iyaaloro was observed to be the most polluted river, with highest concentrations of
14 42.1 mg kg⁻¹, 102 mg kg⁻¹, 185 mg kg⁻¹, 154 mg kg⁻¹ and 1040 mg kg⁻¹ of Cd, Cr, Cu, Pb and Zn
15 respectively, while Ibeshe River was the least contaminated, apart from a site affected by Cu from
16 the textile industry. Some of the sediments were found to be above the consensus-based probable
17 effect concentrations and Dutch sediment guideline for metals. Overall metal concentrations were
18 similar to those reported for other tropical lagoon and estuarine systems affected by anthropogenic
19 inputs as a result of rapid urbanisation. Due to the large number of samples, principal component
20 analysis was used to examine relationships within the data set. Generally, sediments collected
21 during the dry season were observed to have higher concentration of PTM than those collected
22 during the rainy season. This means that PTM could accumulate over a prolonged period and then
23 be released relatively rapidly, on an annual basis, into tropical lagoon systems.

24
25 *Keywords:* Lagos Lagoon, Potentially toxic metals, principal component analysis, sediments
26
27
28
29
30
31
32

1 **1. Introduction**

2 Rapid urbanisation and industrialisation can lead to the release of a wide variety of pollutants,
3 from both point and diffuse sources, and thus place considerable pressure on aquatic resource such
4 as coastal and riverine systems. This is of particular concern in developing countries where
5 expansion of urban area may be relatively unregulated and where environmental protection may
6 be, or may historically have been, inadequate (Fonesca *et al.* 2011; Li *et al.* 2007; Manning 2011;
7 Nriagu 1992; Ramessur *et al.* 2010).

8
9 Sediments in tropical lagoon and estuarine systems of developing countries are increasingly
10 studied because they can act as both sinks for potentially toxic metals (PTM), allowing records of
11 past inputs to be reconstructed where no contemporary monitoring data exists, and as sources, with
12 potential impacts on ecosystems and human health (Acevedo-Figueroa *et al.* 2006; Ahmad *et al.*
13 2010; Alagarsamy 2006; Alaoui *et al.* 2010; El Ati Hellal *et al.* 2011; Fonesca *et al.* 2011; Jara-
14 Marini *et al.* 2008; Li *et al.* 2007; Pereira *et al.* 2010; Vázquez-Sauceda *et al.* 2011; Xu *et al.*
15 2009; Zourarah *et al.* 2007). Pollutant metals may affect the population directly through contact
16 with contaminated sediment or water and *via* the food chain. A further exposure route exists where
17 contaminated sediment is dredged and used for land reclamation or construction.

18
19 Lagos State is Africa’s biggest city and the fastest growing metropolis in the world. It is the most
20 heavily industrialised city in Nigeria, with much of the nation’s wealth and economic activities
21 located there. The City of Lagos is currently undergoing an extensive program of expansion and
22 development with the aim of becoming a ‘megacity’ and major international focus for trade and
23 industry (Howden 2010; Thisday’s Special Intenational Project 2007). The state is reported
24 already to be home to over 70% of the country’s medium and large-scale manufacturing industries
25 (Oketola and Osibanjo 2007). Some of the manufactured goods produced in the city include
26 machinery, motor vehicles, electronic equipment, chemicals, beer, processed food, and textiles.
27 The urban area is built on a number of flood plains and encompasses a network of marshes,
28 swamps, streams, creeks and rivers which receive large quantities of storm water run-off, together
29 with domestic, municipal and industrial waste effluents, and these receptors empty into the Lagos
30 Lagoon. Odo-Iyaaloro, Shasha and Ibeshe Rivers are three main trans-urban rivers in the state,
31 receiving domestic and industrial wastes from numerous sources.

1

2 The Lagos Lagoon is a brackish coastal lagoon – the largest in the West African coast – located
3 between longitude 3° 23' and 3° 40'E and between latitude 6° 27' and 6° 48'N. It is a fairly
4 shallow expanse of water (0.3-3 m deep) which is about 50 km long and 3 to 13 km wide and
5 separated from the Atlantic Ocean by a narrow strip of barrier bar complex. The lagoon is used for
6 fishing, irrigation and recreation. Its pollution by PTM has potential adverse effects on both
7 aquatic life and on human beings whose life and livelihood depend on the Lagos Lagoon (Abayomi
8 *et al.* 2011; Isebor *et al.* 2006; WES 1997). Dredging of sediment from the lagoon, although
9 illegal, is commonplace to satisfy local demand for building materials.

10

11 Relatively few researchers have worked previously on the pollutant levels in Lagos Lagoon and its
12 environs. Ihenyen (1991) measured high concentrations of PTM in sediments from areas within
13 the Lagos Harbour, which he attributed to the impact of industrial effluent. Otitolaju and
14 colleagues conducted experiments on the uptake of Cd, Cu, Hg, Pb and Zn by benthic organisms
15 from the lagoon (Otitolaju 2003; Otitolaju and Don-Pedro 2004; Otitolaju *et al.* 2007). In their
16 study, it was observed that sediment samples from the western part of the lagoon generally had
17 higher concentrations of PTM compared to the eastern region. Most recently, Abayomi *et al.*
18 (2011) studied the contribution of roadside soil to phosphorus loading in the Lagos Lagoon, but
19 this was not related to the concentration of PTM in the water or sediments. Thus, despite rapid
20 industrialization in this region, there has been a dearth of recent robust information on sediment-
21 metal concentrations and relationship between sediment properties and metal levels. Further, there
22 is no information on seasonal-variations in metal-pollutant loads, which may be of importance in
23 tropical regions affected by distinct dry and rainy seasons.

24

25 Many studies focusing on the aquatic environment now use multivariate analysis techniques to aid
26 in interpretation of results. For example, Altm *et al.* (2009) in their study on the assessment of
27 seasonal variations of surface water quality characteristics of Porsuk stream used a variety of
28 methods to display the information concealed in the variables observed in a water quality
29 monitoring network. Some of the multivariate statistical techniques which have been used in
30 literature include cluster analysis, principal component analysis, discriminant analysis and
31 multiscaling analysis. Kikuchi *et al.* (2009) used both cluster analysis and principal component

1 analysis to evaluated spatial and seasonal variations in the concentrations of PTM in Nhue River
2 and one of its tributaries, the To Lich River. The two multivariate analyses showed that the level
3 of PTM in water of To Lich River was distinctly different from that of Nhue.

4
5 The aims of the current study are to (i) quantify the levels of Cd, Cr, Cu, Pb and Zn in sediments
6 of the Lagos Lagoon systems. (ii) examine relationships between pH, % organic matter, CEC,
7 grain size, Cd, Cr, Cu, Pb and Zn in the sediments using principal component analysis (iii)
8 investigate seasonal variations in the PTM contents in sediments of these waterbodies, and (iv)
9 compare concentrations with other aquatic systems and with sediment quality standards to inform
10 the development of appropriate management strategies for the lagoon.

11 12 13 **2. Experimental**

14 15 **2.1 Study areas**

16 The study areas include the Lagos Lagoon and three rivers: Odo-Iyaalaro, Shasha and Ibeshe
17 Rivers. Sediments samples were collected from each of the rivers six times during the year at
18 different seasons (rainy and dry). Samples were collected at three points along each river, at the
19 points of discharge of each of the rivers, and at points approximately 1 km offshore and to the east
20 and west of the discharge of the rivers into the Lagos Lagoon (Figure 1 shows the sampling
21 stations). The Lagos Lagoon was sampled three times during the year. Samples were only
22 collected once from points 1 and 2 due to lack of access thereafter. Thus, 26 sediment samples
23 were collected from Odo-Iyaalaro River, 36 samples each were collected from Shasha and Ibeshe
24 Rivers, and 15 samples were collected from the Lagos Lagoon (a total of 103 sediment samples).

25 26 **2.2 Sediment samples**

27 The samples were collected using a stainless steel grab sampler deployed from the riverbank, a
28 bridge, or a boat. Samples were placed in polyethylene containers and taken to the laboratory.
29 Once in the laboratory, the sediment samples were air dried, homogenized in a mortar, sieved to
30 pass through a 2 mm stainless steel sieve and stored in polyethylene containers prior to analysis.

1 The pH of the sediments was determined according to BS ISO 10390 (1995), using a Philips PW
2 9420 pH meter and the organic matter content (%OM) was estimated by Walkley-Black method
3 (1934). The cation exchange capacity (CEC) was determined by first saturating the sediment with
4 sodium ion, the sodium ion was then exchanged for ammonium ion. The amount of sodium
5 exchanged was determined by flame atomic absorption spectrometry (FAAS), using caesium as an
6 ionisation suppressant (Ryan *et al.* 2001). Grain size analysis was carried out on the dry sediment
7 samples by laser granulometry using the method of Webster *et al.* (2000). The concentrations of
8 Cd, Cr, Cu, Pb and Zn were determined by digesting 1 g of each of the sediment samples with 20
9 ml *aqua-regia* in a microwave (CEM MDS-2000) (British Standard, ISO 11466 1995) with
10 microwave power from 0 to 630 W depending on the number of samples digested. The total period
11 of digestion was 50 mins. The digests were filtered using Fisherbrand QL 100 filter paper (11 cm)
12 into 100 ml standard flasks, made up to the mark with distilled water and stored in plastic bottles
13 at a temperature of 4 °C prior to analyses. The sediment samples were digested in triplicate. The
14 quantification of the analytes was performed by FAAS using a Perkin Elmer AAnalyst 200
15 instrument with air-acetylene flame under optimal conditions. The detection limit of the
16 instrument ($DL_{inst.}$) for each of the metal of interest was determined using the equation below.
17 Procedural detection limits ($DL_{pro.}$) were also calculated taking into consideration the digestion
18 methods used (Vandecasteele and Block 1997).

19
20
$$DL_{inst.} = \frac{3s}{b}$$

21 where

22 s = standard deviation of the ten replicate measurements of the calibrant solution

23 b = slope of the calibration curve

24

25 **2.3 Quality control**

26 For the determination of potentially toxic metals, the precision and accuracy of the method used
27 was determined by using reference material BCR CRM 143R. This is a sewage sludge amended
28 soil which is a certified reference material for *aqua-regia* soluble metals. The digestion of the
29 sediments was carried out in triplicates for each sample, and for each batch of digestion there was
30 a procedural blank.

1 **2.4 Multivariate analysis**

2 In order to reduce the relatively large number of variables to a smaller number of orthogonal
3 factors, the original data obtained was processed by multivariate statistical methods. The
4 associations of parameters in sediments were determined by principal components analysis (PCA)
5 and applying varimax with Kaiser Normalization rotation method to facilitate easier interpretation.
6 Varimax rotation maximizes the sum over components of the variances of the squared loadings,
7 thereby emphasizing cluster recognition. The eigenvalues were used to determine the number of
8 PCs to be retained in order to comprehend the trends in the data. The PCA was carried out with
9 factors that have eigenvalues greater than 1. The PCA was carried out using the SPSS for windows
10 software version 13.0

11
12
13 **3. Results and discussion**

14
15 **3.1 Quality control**

16 Table 1 shows the results of PTM concentration in the certified reference material BCR CRM
17 143R. Found values were within two standard deviations of the certified values for all metals in
18 the *aqua regia* digests, and a recovery of 95 to 105% was obtained. This indicated that the analysis
19 was under control. Further, the relative standard deviation obtained for analysis of triplicate test
20 portions of the lagoon sediment samples was < 10% in most cases.

21
22 **3.2 Sediment properties and pseudototal metal concentration**

23 Sediment properties and pseudototal metal concentration during the rainy and the dry seasons are
24 summarized in Table 2, while the actual values for the different sampling points are given in
25 online resources 1 and 2.

26
27 **3.2.1 Odo-Iyaalaro River**

28 For Odo-Iyaalaro River, the sediment pH ranged from 2.8 to 5.6 during the dry season, and from
29 3.6 to 5.1 during the rainy season, indicating that the sediments were acidic throughout the year.
30 The % organic matter ranged from 0.8 to 21.0 and 1.4 to 15.7, while the CEC of the sediments
31 ranged from 2.0 to 14.6 meq/100g and 4.2 to 14.2 meq/100g during the dry and rainy seasons

1 respectively. Sediment samples collected from points 5 and 6 were on most occasions observed to
2 have higher organic matter content than those from other points in the river. This could arise from
3 decaying plant material as there are large quantities of plants growing at the bank of the river at
4 these sites. Samples from points 5 and 6 in Odo-Iyaaloro River were also observed to generally
5 have higher concentrations of metals compared to other points along the river (apart from site 1,
6 discussed below) for both dry and rainy seasons. This could be due to the high organic matter
7 content of the sediment since organic matter is known to influence PTM adsorption in sediments
8 (Loska and Wiechu 2003). Furthermore, the high concentration of metals at points 5 and 6 could
9 be because that they are close to other tributaries emptying into the Lagos Lagoon, and the water
10 from these tributaries may contain high concentrations of PTM.

11
12 Samples collected from points 3 to 6 showed distinct seasonal trends, with generally higher metal
13 concentrations found in sediments collected during the dry season (up to double the amount found
14 in the rainy season for Cd and Zn at sites 5 and 6). The month of March, which is the peak of dry
15 season, had particularly higher concentration of metals when compared with samples collected
16 from the same point in other months (21.3, 139, 119 and 777 mg kg⁻¹ for point 5 and 34.4, 185,
17 154 and 1040 mg kg⁻¹ for point 6, for Cd, Cu, Pb and Zn respectively). This may be as a result of
18 accumulation of these PTM in sediment during the dry season, and remobilisation during the rainy
19 season. The current study did not investigate whether the metals are mobilised in solution or
20 suspension, but the importance of particulate transport for PTM in aquatic systems is well
21 established (Viers *et al.* 2009) and the change observed in average sediment grain size
22 distributions at sites 3-6 from 52% sand + 18% silt + 30 % clay in the dry season to 64% sand +
23 14% silt + 22% clay in the rainy season shows that, as would be expected, fine particles are lost
24 when river flow rate increases (results of grain size analysis for all sediment samples are provided
25 in online resources 3 and 4).

26
27 Previous workers have reported seasonal influence on sediment metal concentrations in tropical
28 systems, but findings are contradictory. Similar to the current study, Bahena-Manjarrez *et al.*
29 (2002) found highest metal concentrations in the dry season in the Coatzacoalcos River, Mexico,
30 and this was supported by Alagarsamy (2006) who reported lowest metal concentrations during
31 the monsoon in the Mandovi estuary, India. However, whereas a significant difference between

1 pre-monsoon and monsoon concentrations of Cd in sediments of Buriganga River (Dhaka,
2 Bangladesh) was noted recently (Ahmad *et al.* 2010) no seasonal differences were found in the
3 same study for levels of Cu, Cr, Ni or Pb. Indeed, Jara-Marini (2008) reported that sediment metal
4 concentrations in the Mazatlan Harbour increased during the rainy season, although that was as a
5 site where deposition and accumulation of material washed down from the upper reaches of the
6 river system might have been important.

7
8 The sample collected from point 1 had amongst the highest concentrations of Cd, Pb and Zn found
9 in the Odo-Iyaaloro River (42, 108 and 805 mg kg⁻¹ respectively). This point is under a road bridge
10 and so the high concentrations could be due to anthropogenic input from vehicle wear and
11 exhausts. Descriptive statistics (Table 2) showed Odo-Iyaaloro to be the most polluted of the three
12 trans-urban rivers in terms of the average concentrations of Cd (9.6 and 4.8 mg kg⁻¹), Pb (56.7 and
13 63.5 mg kg⁻¹) and Zn (292 and 147 mg kg⁻¹) during the dry and rainy seasons respectively.

14 15 3.2.2 Shasha River

16 For Shasha River, the pH ranged from 4.3 to 7.3 during the dry season, and from 4.2 to 7.7 during
17 the rainy season. This indicates acidic to neutral sediments. For this river, the highest
18 concentrations of PTM were observed at point 7 during both the rainy and dry seasons. This may
19 be because the sampling point is very close to a bridge which normally has a high traffic density
20 and is also close to industries. Lead levels as high as 202 mg kg⁻¹ were found which, as in the case
21 with samples from point 1 from Odo-Iyaaloro, could be caused by the exhausts of vehicles since
22 tetraethyl lead is still used in petrol in Nigeria. The concentration of Cd was found to be generally
23 less than the procedural detection limit of 0.9 mg kg⁻¹, except in a few cases. This showed that Cd
24 is not a pollutant in this river. The PTM concentrations in sediments from Shasha River were
25 observed to decrease with decreasing distance from the Lagos Lagoon. This variation may be
26 correlated with the location of industries and their waste disposal system: there are more industries
27 closer to point 7 than other points along this river and their effluents are reflect in the higher PTM
28 levels upstream, whereas the sediments far away from these source will be diluted with
29 uncontaminated material. The average concentration of PTE in Shasha River were lower than in
30 Odo-Iyaaloro River, except for Cr in the rainy season, and no clear seasonal trends were evident.

1 3.2.3 Ibeshe River

2 In Ibeshe River, Cu was observed to be the major metal pollutant. The low concentration of other
3 PTM could be because the water body is located in a relatively rural area where there is very little
4 emission from either industrial effluents or vehicular traffic. There was an increase in Cu
5 concentration at site 14 relative to site 13, which is probably due to the discharge of effluents by a
6 textile company that uses Cu as one of the components of dye. The highest concentrations of Cu at
7 point 14 were observed in the months of February (332 mg kg⁻¹) and March (246 mg kg⁻¹) which is
8 the dry season while the concentration became lower (average < 50 mg kg⁻¹) during the rainy
9 season. The concentration of Cu was also found to decrease downstream from point 14 to point 18
10 which is in the Lagos Lagoon.

11

12 3.2.4 Lagos Lagoon

13 In the Lagos Lagoon, points 19, 20 and 21 were observed to have higher concentration of Cd, Cu,
14 Pb and Zn than points 22 and 23 for all the months sampled. This is in agreement with Otitolaju *et*
15 *al.* (2007) who observed higher concentration of metals in the western part of the lagoon. The
16 higher concentrations in this area could be because these points are close to a bridge or other site
17 where human activities take place, and because the sediments differ in composition from those in
18 the centre of the lagoon. Sediment from sites 19-21 have approximately 70% of particles in the
19 sand size fraction and contain 6.3% organic matter on average, whereas samples from sites 22 and
20 23 are dominated (> 95%) by sand and contain ≤ 0.5% organic matter. Compared to the rivers
21 under investigation, Cr concentration in the Lagos Lagoon is quite high and the range is small.
22 This may indicate that the source of Cr in this water body is geogenic. No systematic differences
23 were observed between concentrations of metals in rainy and dry seasons in the lagoon, nor were
24 there marked seasonal differences in grain size distribution, as had been observed in some of the
25 tributary rivers. This is to be expected because according to Lund-Hansen *et al.* (1999) re-
26 suspension rate was observed to be 10 times higher than sedimentation rate in shallow coastal
27 lagoons and the depth of Lagos Lagoon water body means that deposition, as well as re-
28 suspension, is likely to occur throughout the year.

29

30 3.2.5 Comparison with literature and regulatory concentrations

1 Table 3 gives historical data on the Lagos Lagoon and some recent literature values for PTM
2 concentration in other tropical and semi-tropical lagoons and river estuaries. Levels of PTM found
3 in lagoon sediments in the present study are broadly similar to those reported previously (Otitolaju
4 *et al.* 2007) allowing for the fact that the current study sampled from both the western and eastern
5 parts of the lagoon. The highest concentrations of Cd, Cu, Pb and Zn found overall i.e. including
6 points from the three trans-urban rivers, are similar to maxima reported in aquatic systems
7 impacted by rapid urbanisation in other developing countries, such as the Pearl River in China (Li
8 *et al.* 2011), the Buriganga River in Bangladesh (Ahmad *et al.* 2010) and the Rodrigo de Freitas
9 Lagoon in Brazil (Fonseca *et al.* 2011), but the concentration of Cd at point 1 (42 mg/kg) is
10 remarkably high.

11
12 There is currently no legislation in Nigeria governing acceptable levels of PTM in sediments and
13 so it is not possible to assess the current findings within a local regulatory framework. Included in
14 Table 3 are consensus-based probable effect concentrations (PEC) recommended by MacDonald
15 (2000) and Dutch sediment guideline (Grimwood and Dixon 1997). Of the 103 sediments studied,
16 11 exceed the PEC for Cd (all in the Odo-Iyaaloro); two exceed the PEC for Cr (sites 7 and 8 in
17 the Shasha River); four exceed the PEC for Cu (one from Odo-Iyaaloro and three from Ibeshe
18 River); three exceed the PEC for Pb (including site 7 in Shasha River in both dry and rainy
19 seasons); and four exceed the PEC for Zn (three in Odo-Iyaaloro and, again, site 7 in Shasha
20 River). Forty of the sediments were above the Dutch sediment guideline for Cu and four (three
21 from Odo-Iyaaloro during the dry season) exceed the guideline for Zn.

22

23 **3.3 Principal component analysis**

24 The data were subjected to PCA to elucidate geochemical factors influencing PTM distribution in
25 the Lagos Lagoon system. Eigenvalues were used to determine the number of principal
26 components that should be retained for further study. For both seasons, the first three principal
27 components had eigenvalues greater than 1, and explained 71.19 % of the variance during the dry
28 season, and 74.95% of the variance during the rainy season.

29

30 For the dry season, the first factor (PC1) explaining 40.89% of the total variance was observed to
31 be strongly and positively correlated to % OM, CEC, Pb, Cd, and Zn. The second factor (PC2)

1 explained 20.00% of the variance, and was observed to be positively correlated to % silt and clay
2 and negatively correlated to % sand, while the third factor (PC3), explaining 10.30% of the
3 variance, was positively correlated to Cr, and negatively correlated to pH (Fig. 2a).

4
5 According to several studies, (Bloemen *et al.* 1995; Davies 1997; Ujevic *et al* 2000) an association
6 of Cd, Pb and Zn with each other suggests anthropogenic influence. These metals often associate
7 in urban systems, and so their relationship in the Lagos Lagoon system indicates that it is being
8 affected by the rapid urbanization in Lagos State. The plot of scores shows that samples 1D, 9D,
9 10D, 15D and 21D (samples collected from points 1, 5 and 6 in Odo-Iyaaloro and points 7 in
10 Shasha River) are highly correlated to PC1, and this reflects the presence of high concentration of
11 Cd, Zn Pb and % organic matter and CEC. This indicates that these two rivers pollute the Lagos
12 Lagoon in terms of those anthropogenic metals as compared to Ibeshe River. It is also consistent
13 with, for example, the findings of Loska and Wiechu, (2003) who applied PCA to surface
14 sediments from a reservoir Poland and found that all the metals analysed were positively
15 correlated to organic matter.

16
17 In the rainy season, the first PC which explained 42.60 % of the total variance was observed to be
18 correlated to Cd, CEC and %OM, and negatively correlated to pH, while the second PC, which
19 accounts for 18.32 % of the total variance, is correlated to Pb, Zn, Cu, and Cr (Fig.3a). The plot of
20 scores (Fig. 3b) shows that samples 4R, 8R, 2R, 3R, 12R, 7R, and 11R – all in the Odo-Iyaaloro
21 River – have the highest association with PC1. All these samples have relatively high %OM
22 and/or CEC and they represent the only points where Cd was detected at appreciable
23 concentrations in the rainy season. Samples 51R and 56R collected from the Lagos Lagoon were
24 also observed to have a high correlation with PC1 due to their high %OM and CEC. Sample 13R
25 collected from point 7 in Shasha River (immediately below a road bridge) in May was observed to
26 have the highest association with PC2 due to the presence of high concentrations of Cr, Cu, Pb and
27 Zn. May is just at the onset of the rainy season, and the volume of rainfall at this time may not
28 have been enough to flush out the pollutants from the sediments.

31 **4. Conclusion**

1 The physicochemical parameters and PTM concentration of sediments from three rivers and the
2 Lagos Lagoon has been determined and the data subjected to statistical analysis. Odo-Iyaaloro was
3 the most contaminated of the rivers in terms of Cd, Pb and Zn, and this can be attributed to the
4 various industries in the environs which empty their untreated effluents into the Lagos Lagoon via
5 this river. Samples collected from points either under a bridge or close to a bridge with high traffic
6 density was observed to have high concentrations of, in particular, Pb. Sediments from Ibeshe
7 River were generally less contaminated than other tributaries, but specific points were observed to
8 be highly enriched in Cu from the effluent of the textile industry. PTM concentrations in lagoon
9 sediment were generally low at locations towards the centre of the lagoon, but higher along the
10 western margin where the major urban and industrialised areas are located. The concentration of
11 Cr in the Lagos Lagoon was observed to be fairly constant which may indicate that Cr is a natural
12 part of the sediments from this waterbody. Maximum analyte concentrations were similar to the
13 highest values reported in other rapidly expanding urban areas in developing countries, with
14 potentially harmful levels present at some sites. Sediments collected during the dry season were
15 often observed to have higher concentration of PTM compared to those collected during the rainy
16 season. This supports the hypothesis that PTM accumulate in catchment areas during the dry
17 season and are then remobilised during the rainy season into tropical lagoon systems. Further work
18 is required to determine the forms of metals transported and retained in the sediments and the
19 timescale over which they are released at the onset of the rains. In the meantime, steps should be
20 taken to minimise human exposure to PTM in the Lagos Lagoon system. These could include
21 prevention of access to the most highly contaminated areas, temporal restrictions on activities such
22 as dredging so that material is collected when pollutant concentrations are minimal, or flushing
23 away of fine-grained material from dredgings to retain only the sand-sized fraction.

24

25

26 **Acknowledgements**

27 AOO wishes to acknowledge the financial support of the University of Lagos, Nigeria, and the
28 hospitality of the Department of Pure and Applied Chemistry, University of Strathclyde, during a
29 period as a visiting researcher in the UK.

30

1 **References**

- 2 Abayomi, A., Nimmo, M., Williams, C., Olayinka, K. O., Osuntogun, B., Alo, B., & Worsfold, P.
3 (2011). The contribution of roadside soil to phosphorus loading in the eutropic Lagos
4 Lagoon, Nigeria. *Journal of Environmental Monitoring*, 13, 1884-1889.
- 5 Acevedo-Figueroa, D., Jiménez, B. D., & Rodríguez-Sierra, C. J. (2006). Trace metals in
6 sediments of two estuarine lagoons from Puerto Rico. *Environmental Pollution*, 141,
7 336-342
- 8 Ahmad, M. K., Islam, S., Rahman, S., Haque, M. R., & Islam, M. M. (2010). Metals in water,
9 sediment and some fishes of Buriganga River, Bangladesh. *International Journal of*
10 *Environmental Research*, 4(2), 321-332
- 11 Ahmed, F., Bibi, M. H., Fukushima, T. M., Seto, K., & Ishiga, H. (2011). Recent sedimentary
12 environment of coastal lagoon in southwestern Japan: evidence from major and trace
13 elements. *Environmental Monitoring and Assessment*, 173, 167-180
- 14 Alagarsamy, R. (2006). Distribution and seasonal variation of trace metals in surface sediments of
15 the Mandovi Estuary, west coast of India. *Estuarine, Coastal and Shelf Science*, 67, 333-
16 339
- 17 Alaoui A. M., Choura, M., Maanan, M., Zourarah, B., Robin, M., Conceicao, M. F., Andrade, C.,
18 Khalid, M., & Carruesco, C. (2010). Metal fluxes to the sediments of the Moulay
19 Bouselham Lagoon, Morocco. *Environmental Earth Sciences*, 61, 275-286
- 20 Altm, A., Filiz, Z., & Iscen, C. F. (2009). Assessment of seasonal variations of surface water
21 quality characteristics of Porsuk Stream. *Environmental Monitoring and Assessment*, 158,
22 51-65
- 23 Bahena-Manjarrez, J. L., Rosales-Hoz, L., & Carranza-Edwards, A. (2002). Spatial and temporal
24 variation of heavy metals in a tropical estuary. *Environmental Geology*, 42, 575-582
- 25 Bloemen, M. L., Markert, B., & Leith, H. (1995). The distribution of Cd, Cu, Pb, and Zn in
26 topsoils of Osnabruck in relation to land use. *Science of the Total Environment*, 166, 137-
27 148.
- 28 British Standard. Chemical methods- Extraction of trace elements soluble in *aqua-regia*. BS
29 7755-3.9: 1995 ISO 11466 1995
- 30 British Standard. Determination of pH. BS 7755-3.2 1995, ISO 10390 1995

- 1 Davies, B.E. (1997). PTM contaminated soils in an old industrial area of Wales, Great Britain:
2 source identification through statistical data interpretation. *Water, Air and Soil Pollution*,
3 94, 85-98
- 4 Denton, G. R. W., Morrison, R. J., Bearden, B. G., Houk, P., Starmer, J. A., & Wood., H. R.
5 (2009). The impact of the coastal dump in tropical lagoon on trace metal concentrations
6 in surrounding marine biota: A case study from Saipan, Commonwealth of the
7 Northern Mariana Islands (CNMI). *Marine Pollution Bulletin*, 58, 424-455
- 8 El Ati Hellal, M., Hellal, F., Khemissi, Z., Jebali, R., & Dachraoui, M. (2011). Trace metals in
9 algae and sediments from the north-eastern Tunisian Lagoons. *Bulletin of*
10 *Environmental Contamination and Toxicology*, 86, 194-198
- 11 Fonseca, E. M., Baptista Neto, J. A., Fernandez, M. A., McAlister, J., & Smith, B. (2011).
12 Geochemical behaviour of heavy metals in different environments in Rodrigo de Freitas
13 Lagoon Brazil-RJ/Brazil. *Anais da Academia Brasileira de ciencias*, 83, 457-469
- 14 Grimwood, M. J., & Dixon, E. (1997). Assessment of risks posed by List II metals to Sensitive
15 Marine Areas (SMAs) and adequacy of existing environmental quality standards (EQSs)
16 for SMA protection. *Report to English Nature*.
- 17 Howden, D. (2010). Lagos: Inside the ultimate mega-city. *The independent Africa*.
18 [http://www.independent.co.uk/news/world/africa/lagos-inside-the-ultimate-megacity-](http://www.independent.co.uk/news/world/africa/lagos-inside-the-ultimate-megacity-1945246.html)
19 [1945246.html](http://www.independent.co.uk/news/world/africa/lagos-inside-the-ultimate-megacity-1945246.html)
- 20 Ihenyen, A. E. (1991). Trace metals in the bottom sediments of the harbour of Lagos, Nigeria.
21 *Environmental Monitoring and Assessment*, 16, 259-264.
- 22 Isebor, C. E., Awosika, A., & Smith, S. V. (2006). Preliminary water, salt, and nutrient budgets
23 for Lagos Lagoon, Nigeria. <http://nest.su.se/MNODE/Africa/Lagos/lagosbud.htm>.
- 24 Jara-Marini, M. E., Soto-Jimenez, M.F., & Paez-Osuna, F. (2008). Bulk and bioavailable heavy
25 metals (Cd, Cu, Pb and Zn) in surface sediments from Mazatlan Harbor (SE Gulf of
26 California). *Bulletin of Environmental Contamination and Toxicology*, 80, 150-153
- 27 Kikuchi, T., Furuichi, T. M., Trung Hai, H., & Tanaka, S. (2009). Assessment of PTM pollution
28 in river water of Hanoi, Vietnam Using Multivariate Analyses. *Bulletin of Environmental*
29 *Contamination and Toxicology*, 83, 575-582
- 30 Li, Q., Wu, Z., Chu, B., Zhang, N., Cai, S., & Fang, J. (2007). Heavy metals in coastal wetland
31 sediments of the Pearl River Estuary, China. *Environmental Pollution*, 149, 158-164

- 1 Loska, K; & Wiechu, D. (2003) Application of principal component analysis for the estimation of
2 source of heavy metal contamination in surface sediments from the Rybnik Reservoir.
3 *Chemosphere*, 51, 723-733
- 4 Lund-Hansen, L. C., Petersson, M., & Nurjaya, W. (1999). Vertical sediment fluxes and wave-
5 induced sediment resuspension in a shallow-water coastal lagoon. *Estuaries*, 22, 39-46
- 6 Maanan, M., Zourarah, B., Carruesco, C., Aajjare, A., & Naud, J. (2004). The distribution of
7 PTM in the Sidi Moussa Lagoon sediments (Atlantic Moroccan Coast). *Journal of*
8 *African Earth Sciences*, 39, 473-483
- 9 MacDonald, D. D., Ingersoll, C. G., & Berger, T. A. (2000). Development and evaluation of
10 consensus-based sediment quality guidelines for fresh water ecosystems. *Archives of*
11 *Environmental Contamination and Toxicology*, 39, 20-31
- 12 Manning, W. J. (2011). Urban environment: Defining its nature and problems and developing
13 strategies to overcome obstacles to sustainability and quality of life. *Environmental*
14 *Pollution* 159, 1963-1964
- 15 Morrison, R. J., & Brown, P. L. (2003). Trace metals in Fanga'uta Lagoon, Kingdom of Tonga.
16 *Baseline / Marine Pollution Bulletin*, 46, 139-152
- 17 Nriagu, J. O. (1992). Toxic metal pollution in Africa. *Science of the Total Environment*, 121, 1-37
- 18 Oketola, A. A., & Osibanjo, O. (2007). Estimating sectoral pollution load in Lagos by Industrial
19 Pollution Projection System (IPPS). *Science of the Total Environment*, 377, 125-141
- 20 Otitoloju, A. A. (2003). Relevance of joint action toxicity evaluations in setting realistic
21 environmental safe limits of PTM. *Journal of Environmental Management*, 67, 121-128.
- 22 Otitoloju, A. A., & Don-Pedro K. N. (2004). Integrated laboratory and field assessment of PTM
23 accumulation in edible periwinkle *Tympanotonus fuscatus* var *radula* (L.) *Ecotoxicology*
24 *and Environmental Safety*, 57, 354-362.
- 25 Otitoloju, A. A., Don-Pedro, K. N., & Oyewo, E. O. (2007). Assessment of potential ecological
26 disruption based on PTM toxicity, accumulation and distribution in media of Lagos
27 Lagoon. *African Journal of Ecology*, 45, 454-463A.
- 28 Pereira, A. A., Hattum, B., Boer, J., Bodegom, P. M., Rezende, C. E., & Salomons, W. (2010).
29 Trace elements and carbon and nitrogen stable isotopes in organisms from a tropical
30 coastal lagoon. *Archives of Environmental Contamination and Toxicology*, 59, 464-477

- 1 Ramessur, R. T. & Ramjeawon, T. (2002). Determination of Pb, Cr and Zn in sediments from an
2 urbanized river in Mauritius. *Environment International*, 28, 315- 324
- 3 Ramessur, R.T., Boodhoo, K., Balgobin, J., & Aurello, A. (2010). Statistical comparison
4 between consecutive winter and summer concentrations in zinc and lead from sediments
5 in a tropical urban estuary in Mauritius. *Environmental Monitoring and Assessment*,
6 168, 345- 351
- 7 Rosales-Hoz, L., Cundy, A. B., & Bahena-Manjarrez, J. L. (2003). Heavy metals in sediment
8 cores from a tropical estuary affected by anthropogenic discharges: Coatzacoalcos
9 estuary, Mexico. *Estuarine, Coastal and Shelf Science*, 58, 117-126.
- 10 Ryan, J., Estefan, G., & Rashid, A. (2001). Soil and plant analysis laboratory manual. 2nd edition.
11 International Centre for Agricultural Research in the Dry Areas (ICARDA) and National
12 Agricultural Research Centre (NARC) pp 49
- 13 Soto-Jimenez, M. F., & Paez-Osuna, F. (2001). Distribution and normalization of PTM
14 concentration in mangrove and lagoonal sediments from Mazatlan Harbour (SE gulf of
15 California). *Estuarine, Coastal and Shelf Science*, 53, 259-274
- 16 Thisday's Special International Summit, Ney York, USA (2007); Lagos city project.-New city on
17 the Atlantic” [http://www.africanloft.com/lagos-mega-city-project-new-city-on-the-](http://www.africanloft.com/lagos-mega-city-project-new-city-on-the-atlantic/)
18 [atlantic/](http://www.africanloft.com/lagos-mega-city-project-new-city-on-the-atlantic/)
- 19 Ujevicâ, I., Odziak, N., & Baric, A. (2000). Trace metal accumulation in different grain size
20 fractions of the sediments from a semi-enclosed bay heavily contaminated by urban and
21 industrial wastewaters. *Water Research*, 34, 3055-3061
- 22 Vandecasteele C., & Block C. B. (1997). Modern methods for trace element determination. John
23 Wiley and Sons, Chichester, p.68
- 24 Vazquez-Sauceda, M., Aguirre-Guzman, G., Sanchez-Martinez, J.G., & Perez-Castaneda, R.
25 (2011). Cadmium, lead and zinc concentrations in water, sediment and oyster of San
26 Andres Lagoon, Mexico. *Bulletin of Environmental Contamination and Toxicology*, 86,
27 410-414
- 28 Viers J., Dupre, B., & Gaillardet, J. (2009). Chemical composition of suspended sediments in
29 World Rivers: New insights from a new database. *Science of the Total Environment*, 407,
30 853-868

- 1 Walkley, A., & Black, I. A. (1934). An examination of Degtjareff method for determining soil
2 organic matter and a proposed modification of the chromic acid titration method. *Soil*
3 *Science*, 37, 29-37
- 4 Webster, L., McIntosh, A. D., Moffat, C. F., Dalgarno, E. J., Brown, N. A & Fryer, R. J. (2000).
5 Analysis of sediments from Shetland Island voes for polycyclic aromatic hydrocarbons,
6 steranes and triterpanes. *Journal of Environmental Monitoring*, 2, 29-38.
- 7 World Environmental Systems (WES) (1997). Special report for the Lagos state Ministry of
8 Environment and Physical Planning, Alausa – Ikeja. 85pp.
- 9 Xu, B., Yang, X., Gu, Z., Zhang, Y., Chen, Y., & Lv, Y. (2009). The trend and extent of heavy
10 metal accumulation over the last hundred years in Liadong Bay China. *Chemosphere*, 75,
11 442- 446
- 12 Zourarah, B., Maanan, M., Carruesco, C., Aajjane, A., Mehdi, K., & Freitas, M. C. (2007). Fifty-
13 year sedimentary record of heavy metal pollution in the lagoon of Oualidia (Moroccan
14 Atlantic coast). *Estuarine, Coastal and Shelf Science*, 72, 359-369.
- 15
- 16

1 Table 1 Results of analysis of certified reference material BCR CRM 143R (mg kg⁻¹)

	<i>Cd</i>	<i>Cr</i>	<i>Cu</i>	<i>Pb</i>	<i>Zn</i>
Certified	72 ± 1.8	426±12	128±7*	174±5	1063±16
Found (n = 5)	71.5 ± 3.5	412±8	122±6	172±8	1074±12

2 Found results are mean values ± standard deviation for n = 5 (pseudototal digestion)

3 * = indicative (non-certified) value

4

1 Table 2 Descriptive statistics of physicochemical properties and PTM concentration in
 2 sediments of dry and rainy seasons

Parameter	Dry			Rainy		
	Minimum	Maximum	Mean	Minimum	Maximum	Mean
<i>Odo-Iyaaloro</i>						
pH	2.8	5.6	4.0	3.6	5.1	4.2
OM (%)	0.8	21	7.8	1.4	15.7	8.5
CEC	2.0	14.6	7.6	4.2	14.2	9.6
Cd ($mg\ kg^{-1}$)	< 0.9	42.1	9.6	< 0.9	12.8	4.8
Cr ($mg\ kg^{-1}$)	< 17.5	102	31.0	<17.5	28.9	22.9
Cu ($mg\ kg^{-1}$)	15.6	185	60.9	15.8	105	55
Pb ($mg\ kg^{-1}$)	< 15.8	154	56.7	21.6	108	63.5
Zn ($mg\ kg^{-1}$)	53.1	1040	292	18.7	377	147
<i>Shasha River</i>						
pH	4.3	7.3	5.6	4.2	7.7	6.2
OM (%)	0.1	8.2	2.4	0.2	7.4	2.1
CEC	1.6	9.8	3.9	1.6	10.2	3.8
Cd ($mg\ kg^{-1}$)	< 0.9	1.2	0.9	<0.9	1.7	1.0
Cr ($mg\ kg^{-1}$)	<17.5	56.7	27.8	<17.5	140	40.6
Cu ($mg\ kg^{-1}$)	< 2.8	78.5	31.3	<2.8	106	32.0
Pb ($mg\ kg^{-1}$)	< 15.8	189	38.6	<15.8	202	45.5
Zn ($mg\ kg^{-1}$)	1.7	467	101	13	641	104
<i>Ibeshe River</i>						
pH	3.6	7.4	5.1	3.2	7.3	4.5
OM (%)	0.3	7.2	2.3	0.2	7.6	2.4
CEC	1.6	4.2	2.5	1.8	5.2	3.0
Cd ($mg\ kg^{-1}$)	< 0.9	<0.9	0.9	<0.9	<0.9	0.9
Cr ($mg\ kg^{-1}$)	<17.5	48.3	27.1	<17.5	54.5	31.3
Cu ($mg\ kg^{-1}$)	< 2.8	332	62.0	<2.8	115	23.3
Pb ($mg\ kg^{-1}$)	<15.8	26.3	16.2	< 15.8	21.8	15.8
Zn ($mg\ kg^{-1}$)	4.6	158	35.9	6.0	57.5	21.3
<i>Lagos Lagoon</i>						
pH	4.3	6.6	5.4	4.1	7.0	5.6
OM (%)	0.1	12.6	4.2	0.1	10.8	3.7
CEC	1.8	11.2	5.0	2	11.0	4.9
Cd ($mg\ kg^{-1}$)	<0.9	2.1	1.2	<0.9	<0.9	<0.9
Cr ($mg\ kg^{-1}$)	34.4	51.7	44.7	23.8	51.7	35.8
Cu ($mg\ kg^{-1}$)	< 2.8	33.7	20.6	< 2.8	43.0	18.8
Pb ($mg\ kg^{-1}$)	< 15.8	39	28.4	<15.8	39.2	25.6

	Zn ($mg\ kg^{-1}$)	1.3	190	103	1.3	246	118
1	<hr/>						
2							

1 Table 3 Literature values for pseudototal metal concentration of PTM in sediments of
 2 Lagos Lagoon and other areas (mg kg⁻¹)

Site	Cd	Cr	Cu	Pb	Zn
<i>Lagos Lagoon Harbour (Ihenyen 1991)</i>	*	97-120	16-41	16-52	10-43
<i>Lagos Lagoon West (Otitoloju et al. 2007)</i>	0.12-0.53	*	11-26	300-420	120-170
<i>Lagos Lagoon East (Otitoloju et al. 2007)</i>	0.10 0.26	*	1.9-12	93-140	8.4-34
<i>Mazatlan Harbour, Mexico (Soto-Jimnez and Paez-Osuna 2001)</i>	*	7.6-42	7.7-91	*	46-350
<i>Coatzacoalcos River, Mexico (Bahena-Manjarrez et al. 2002)</i>	0.97-2.8	40-60	22-42	11-58	88-109
<i>St Louis River, Mauritius (Ramessur and Ramjeawon 2002)</i>	*	100	*	14	170
<i>Coatzacoalcos estuary, Mexico (Rosales-Hoz et al. 2003)</i>	*	37-73	12-120	21-40	69-240
<i>Fanga'uta Lagoon, Kingdom of Tonga (Morrison and Brown 2003)</i>	≤ 0.1	*	7-15	3-8	13-38
<i>Sidi Moussa Lagoon, Morocco (Maanan et al. 2004)</i>	*	97	30	*	50
<i>Joyuda Lagoon Puerto Rico (Acevedo-Figueroa et al. 2006)</i>	0.10	*	22	7.6	52
<i>Mandovi estuary, India (Alagarsamy 2006)</i>	*	*	12-78	4.5-46	20-84
<i>San Jose Lagoon Puerto Rico (Acevedo-Figueroa et al. 2006)</i>	1.8	*	100	220	530
<i>Oualidia Lagoon, Morocco (Zourarah et al. 2007)</i>	*	46-62	20-90	31-88	200-260
<i>Pearl River estuary, China (Li et al. 2007)</i>	1.8-6.4	14-190	≤ 350	12-86	120-480
<i>Mazatlan Harbour, Mexico (Jara-Marini et al. 2008)</i>	3.1-3.3	*	32-45	50-54	220-320
<i>Tanapag Lagoon, Commonwealth of the Northern Mariana Islands (Denton et al. 2009)</i>	<0.17-1.7	1.4-18	0.50-100	0.65-160	2.4-360
<i>Buriganga River, Bangladesh (Ahmad et al. 2010)</i>	2.4-4.2	120-220	22-32	65-77	*
<i>Mãe-Bá Lagoon, Brazil (Pereira et al. 2010)</i>	0.03-0.92	2.4-29	0.47-12	1.5-21	1.6-32
<i>Moulay Bousselham Lagoon, Morocco (Alaoui et al. 2010)</i>	0.02 - 0.84	19-110	22-310	6.2-32	170-760
<i>Lake Shinji, Japan (Ahmed et al. 2011)</i>	*	18-240	3-43	11-39	16-200
<i>North-Eastern Tunisian Lagoons (El Ati Hellal et al. 2011)</i>	*	8.7-62	11-36	13-60	60-230
<i>Rodrigo de Freitas Lagoon, Brazil (Fonseca et al. 2011)</i>	*	10-150	5-120	5-220	≤ 390
<i>Sable Noir estuary, Mauritius (Ramessur et al. 2011)</i>	*	*	*	44	200
<i>San Andres Lagoon, Mexico (Vázquez-Sauceda et al. 2011)</i>	1.1	*	*	0.89-1.0	8.4-9.3
<i>Probable effect concentration (MacDonald et al. 2000)</i>	4.98	111	149	128	459
<i>Dutch sediment guideline (Grimwood and Dixon 1997)</i>	*	380	36	530	480

1 Concentrations are in $\mu\text{g/g}$. * Not reported.

1 **Fig. 1** Map of the Lagos Lagoon and tributary rivers

2

3 **Fig. 2** (a) Plot of loadings during the dry season

4 (b) Plot of scores during the dry season


5 (Odo-Iyaalaaro – blue; Shasha – red; Ibeshe – green; Lagos Lagoon – black)

6


7 **Fig. 3** (a) Plot of loadings during the rainy season

8 (b) Plot of scores during the rainy season


9 (Odo-Iyaalaaro – blue; Shasha – red; Ibeshe – green; Lagos Lagoon – black)



1
2




(a)




(b)

- 1
- 2
- 3



(a)



(b)

- 1
- 2
- 3
- 4
- 5

Supplementary Information

Online Resource 1

Physicochemical properties and PTM concentration in sediments during the dry season.

Sample no	Sampling Month	Sampling Points	pH	OM %	CEC (meq/100g)	Cd (mg/kg)	Cr (mg/kg)	Cu (mg/kg)	Pb (mg/kg)	Zn (mg/kg)
<i>Odo-Iyaalaro</i>										
1D	January	1	3.8	6.2	4.0	42.1±1.8	31±2.4	94.5±5.4	108±3	805±25
2D	January	2	4.2	1.4	2.0	2.1±0.3	24.8±1.9	15.6±2.2	41.3±1.9	100±6
3D	January	3	3.8	4.3	5.8	3.6±0.2	31.5±2.1	32.9±1.3	43.5±3.2	110±6
4D	January	4	4.6	8.2	10.4	2.9±0.2	30.8±2.0	25.2±1.2	36±2.4	84.5±4.8
5D	January	5	4.2	11.6	7.4	1.9±0.1	<17.5	15.9±2.6	25.5±1.7	64.5±3.2
6D	January	6	3.9	16.4	7.5	8±0.5	29±1.5	62.2±2.3	54.8±3.7	266±8.4
7D	March	3	3.1	5.1	4.2	4.8±0.4	102±3.3	54.4±2.3	45.3±3.7	157±3
8D	March	4	2.8	7.8	11.2	1.9±0.3	<17.5	38.8±0.8	<15.8	75.2±3.4
9D	March	5	3.9	10.2	14.6	21.3±0.6	<17.5	139±2	119±8	777±16
10D	March	6	4.8	21	12.7	34.4±0.9	30.2±1.9	185±2	154±3	1040±5
11D	November	3	5.6	0.8	3.8	<0.9	<17.5	29.3±2.2	19.7±0.5	53.1±2.7
12D	November	4	3.3	5.9	6.8	<0.9	<17.5	44.1±0.8	28±3.7	82.6±3.7
13D	November	5	5.0	6.1	7.7	6.9±0.4	<17.5	67.4±3.4	60.6±3.6	298±4.2
14D	November	6	3.4	4.4	8.9	2.2±0.1	<17.5	48.6±1.1	43.1±3.7	165±3
<i>Shasha River</i>										
15D	January	7	5.3	5.8	9.8	<0.9	43.1±2.6	48.6±3.7	189 ±7	433±7
16D	January	8	4.9	1.4	2.2	<0.9	24.2±3.2	42.7±1.9	24.5±2.4	87.6±3.2
17D	January	9	4.3	2.1	3.2	<0.9	<22	33.8±1.6	32.6±2.1	43.6±1.9
18D	January	10	5.2	0.8	2.2	<0.9	26.1±1.3	26.6±1.6	36.8±4.2	34.2±1.8
19D	January	11	4.3	8.2	6.4	<0.9	<17.5	44.8±4.7	19.4±1.8	65.9±2.8
20D	January	12	7.2	0.6	4.8	<0.9	<17.5	62.8±3.5	<15.8	40.2±2.5
21D	March	7	4.9	4.8	7.6	1.2±0.2	56.7±2.8	78.5±4.2	93.4±5.5	467±17

22D	March	8	5.2	0.9	2.4	<0.9	53.2±3.8	32.4±2.1	54.2±3.8	93.4±1.3
23D	March	9	4.3	3.2	3.4	<0.9	26.7±2.6	28.6±1.9	56.8±5.6	74.6±2.9
24D	March	10	4.6	0.9	1.8	<0.9	<17.5	15.9±1.9	<15.8	43.2±3.5
25D	March	11	6.7	7.6	3.2	<0.9	<17.5	78.2±1.7	34.2±2.2	129±4
26D	March	12	7.2	0.6	1.6	<0.9	<17.5	16.3±1.4	<15.8	20.1±1.2
27D	December	7	5.7	1.4	7.8	<0.9	25.1±1.5	11.9±0.8	30.4±2.5	53±2.2
28D	December	8	6.1	0.3	2.4	<0.9	<17.5	<2.8	<15.8	1.7±0.3
29D	December	9	6.4	0.4	3.6	<0.9	<17.5	3.4±0.5	<15.8	9.7±0.8
30D	December	10	5	1.6	2.4	<0.9	<17.5	16±1.6	<15.8	82.8±3.3
31D	December	11	5.4	2.2	4.2	<0.9	25.1±1.5	17.7±0.9	<15.8	139±8
32D	December	12	7.3	0.1	2.0	<0.9	<17.5	<2.8	<15.8	3.6±0.4
<i>Ibeshe River</i>										
33D	February	13	6.2	4.1	2.0	<0.9	<17.5	4.2±0.3	<15.8	20.7±2.2
34D	February	14	5.8	6.1	3.2	<0.9	<17.5	332±11	<15.8	96.9±8.1
35D	February	15	5.2	1.8	2.2	<0.9	<17.5	264±10	<15.8	98.3±2.1
36D	February	16	3.7	0.4	2.0	<0.9	<17.5	4.2±0.2	<15.8	4.6±0.1
37D	February	17	4.8	0.4	1.8	<0.9	<17.5	3.4±0.9	<15.8	6±0.5
38D	February	18	6.5	0.3	1.8	<0.9	<17.5	3.6±0.2	<15.8	8.6±1.1
39D	March	13	6.3	3.8	2.6	<0.9	<17.5	<2.8	<15.8	11.3±1.7
40D	March	14	4.2	7.2	3.3	<0.9	32.1±1.3	246±4	<15.8	53.7±0.6
41D	March	15	4.6	2.1	3.2	<0.9	<17.5	84.6±3.1	<15.8	45.9±2.1
42D	March	16	3.9	1.0	2.2	<0.9	42.1±1.8	24.5±2.5	<15.8	10.3±1.2
43D	March	17	4.6	0.6	2.0	<0.9	48.3±2.4	7.2±1.3	<15.8	7.6±0.4
44D	March	18	4.8	0.4	1.6	<0.9	41.2±1.6	9.3±0.9	<15.8	7.6±0.2
45D	December	13	7.4	3.6	3.4	<0.9	<17.5	15.7±1.5	26.3±2.2	158±6
46D	December	14	5.7	4.1	4.2	<0.9	<17.5	13.7±1.1	18±3.3	43.3±1.1
47D	December	15	4.6	1.8	3.2	<0.9	<17.5	13.8±1.3	<15.8	19.7±1.4
48D	December	16	4.8	0.8	2.0	<0.9	34.4±1.2	14.2±0.7	<15.8	13.5±0.7
49D	December	17	3.6	0.8	2.0	<0.9	25.7±1.5	68.2±2.4	<15.8	27.5±1.5
50D	December	18	4.5	1.2	2.2	<0.9	<17.5	5.2±0.7	<15.8	13.1±1.2

<i>Lagos Lagoon</i>										
51D	December	19	4.5	3.5	5.1	<0.9	34.4±2.1	33.7±2.1	33.2±0.7	164±8
52D	December	20	5.4	4.1	4.8	1.3±0.2	51.7±3.2	30±0.8	39±2.3	146±5
53D	December	21	4.3	12.6	11.2	2.1±0.2	51.7±4.8	33.7±1.5	39±3.4	190±4
54D	December	22	6.4	0.1	2.0	<0.9	42.5±1.3	<2.8	<15.8	1.3±0.1
55D	December	23	6.6	0.5	1.8	<0.9	43.0±3.2	<2.8	<15.8	14.8±1.9

Results are mean values ± standard deviation for n = 3; < indicates a value below the detection limit

Online Resource 2

Physicochemical properties and PTM concentration in sediments during the rainy season.

Sample no	Sampling Month	Sampling Points	pH	OM %	CEC (meq/100g)	Cd (mg/kg)	Cr (mg/kg)	Cu (mg/kg)	Pb (mg/kg)	Zn (mg/kg)
<i>Odo-Iyaaloro</i>										
1R	May	3	4.5	1.4	5.2	<0.9	<17.5	27.5±2.0	21.6±3.7	18.7±2.8
2R	May	4	4.2	11.3	8.6	12.8±1.9	<17.5	40.2±1.9	52.3±4.2	68.3±4.5
3R	May	5	3.9	12.3	12.8	6.9±0.3	<17.5	67.8±1.6	43.8±3.6	105±10
4R	May	6	4.3	15.7	12.6	8.6±0.6	<17.5	105±4	46.9±2.1	224±22
5R	July	3	4.9	5.1	4.2	<0.9	<17.5	15.8±2.2	98.2±5.8	134±11
6R	July	4	3.9	7.2	10.2	<0.9	<17.5	38.2±2.1	48.5±3.6	64.3±3.6
7R	July	5	4.5	10.9	11.2	5.5±0.7	25.6±1.7	58.9±3.7	108± 3	264±8
8R	July	6	4.2	15.3	12.4	10.2±1.1	28.9±1.6	92.4±2.4	89.7±4.8	197±13
9R	September	3	5.1	2.9	4.6	<0.9	<17.5	41.1±2.0	59.5±2.7	64.8±5.9
10R	September	4	3.6	4.5	7.8	<0.9	<17.5	43.8±1.9	48.5±3.6	59.9±2.9
11R	September	5	4.2	8.1	14.2	2.6±0.4	<17.5	51.1±2.4	58.2±3.2	190±7
12R	September	6	3.6	7.5	11.8	6.1±0.2	<17.5	77.9±1.3	86.7±4.5	377±5
<i>Shasha River</i>										
13R	May	7	5.9	6.6	8.2	<0.9	140±3	106±2	202±5	641±25
14R	May	8	4.4	1.6	3.4	<0.9	112±8	33.8±2.6	44.2±1.9	112±12
15R	May	9	4.5	2.6	4.4	<0.9	34.2±3.3	15.4±2.1	67.8±3.5	40.1±2.1
16R	May	10	4.4	1.0	2.3	<0.9	<17.5	28±0.4	39.2±1.1	42.1±1.9
17R	May	11	6.8	1.2	3.8	<0.9	<17.5	15.9±1.1	<15.8	23.8±1.5
18R	May	12	7.7	0.4	2.2	<0.9	26.1±3.8	23.8±1.3	<15.8	21.4±1.2
19R	July	7	4.8	7.4	10.2	1.4±0.2	47.2±3.8	98.6±4.3	75.2±3.8	301±4
20R	July	8	5.6	0.9	2.4	<0.9	35.5±3.2	<2.8	18.6±1.4	102±10
21R	July	9	4.2	3.8	4.8	<0.9	29.8±2.8	<2.8	43.2±1.5	32.1±1.9
22R	July	10	7.2	0.8	2.2	<0.9	45.5±3.3	26.8±2.0	19.9±0.8	34.7±2.8
23R	July	11	7.7	3.1	2.8	<0.9	25.9±2.2	54.4±4.1	35.6±3.2	124±6
24R	July	12	7.6	0.5	2.0	<0.9	<17.5	44.2±4.1	<15.8	34.2±2.9
25R	October	7	5.6	1.4	6.4	<0.9	<17.5	27.9±1.9	39±1.2	34.6±3.1
26R	October	8	6.3	0.2	3.2	<0.9	40.7±2.4	<2.8	<15.8	13±1.3
27R	October	9	5.4	0.3	3.6	<0.9	25.6±2.1	<2.8	<15.8	18.6±1.9

28R	October	10	7.5	3.5	2.2	1.7±0.4	35.5±2.1	64±3.1	104±7	190±4
29R	October	11	7.7	1.9	3.2	<0.9	<17.5	13.6±1.1	22.4±1.2	70.8±4.1
30R	October	12	7.7	0.4	1.6	<0.9	<17.5	11.5±1.0	30.4±2.9	38±3.8
<i>Ibeshe River</i>										
31R	May	13	6.0	0.7	2.4	<0.9	<17.5	<2.8	<15.8	17.6±2.0
32R	May	14	4.0	7.6	4.8	<0.9	<17.5	115±2	<15.8	24.8±1.0
33R	May	15	3.4	1.4	3.2	<0.9	<17.5	42.2±1.6	<15.8	10.5±1.3
34R	May	16	4.8	5.3	2.0	<0.9	52.1±2.6	10.2±0.4	<15.8	33.1±2.7
35R	May	17	4.7	0.2	2.2	<0.9	54.5±4.2	5.7±1.1	<15.8	6±0.9
36R	May	18	3.8	0.5	2.4	<0.9	48.8±4.7	8±0.4	<15.8	11.2±1.4
37R	July	13	7.3	3.0	4.6	<0.9	45.6±3.9	15.2±0.4	21.8±2.4	57.5±0.9
38R	July	14	4.1	5.2	4.8	<0.9	34.5±2.1	13.6±0.4	<15.8	54.2±1.2
39R	July	15	4.3	1.5	3.2	<0.9	<17.5	6.8±1.2	<15.8	21.8±0.5
40R	July	16	3.8	1.0	2.0	<0.9	37.2±4.2	13.5±1.3	<15.8	18.5±1.2
41R	July	17	4.3	4.7	2.2	<0.9	35.9±2.1	74.7±3.7	<15.8	45.6±2.7
42R	July	18	4.5	0.3	1.8	<0.9	33.3±1.6	56.7±2.7	<15.8	16.8±0.9
43R	October	13	6.9	2.8	4.0	<0.9	<17.5	3.6±0.5	<15.8	7.8±0.4
44R	October	14	3.6	6.2	5.2	<0.9	24.3±1.9	12.4±1.7	<15.8	18.2±1.2
45R	October	15	3.2	1.3	2.2	<0.9	<17.5	12.2±1.7	<15.8	12.2±0.7
46R	October	16	3.6	1.0	2.6	<0.9	<17.5	12.7±0.8	<15.8	11.1±0.8
45R	October	17	4.8	0.4	2.4	<0.9	<17.5	6.2±0.9	<15.8	9.7±0.4
48R	October	18	4.5	0.3	2.8	<0.9	<17.5	7.8±0.4	<15.8	7.3±0.3
<i>Lagos Lagoon</i>										
49R	July	19	4.6	3.7	5.4	<0.9	23.8±0.9	31±1.3	29.8±1.3	146±6
50R	July	20	5.9	4.3	6.2	<0.9	28.1±2.4	16.5±1.6	39.2±0.7	112±5
51R	July	21	4.4	10.8	11.0	<0.9	26.5±1.1	29.2±1.9	34.1±2.8	223±2
52R	July	22	6.7	0.2	2.2	<0.9	45.6±0.8	4.2±0.6	19.2±1.9	16.7±0.9
53R	July	23	6.5	0.1	2.0	<0.9	51.7±2.7	<2.8	<15.8	43.3±1.8
54R	October	19	4.1	3.3	4.2	<0.9	34.4±1.8	35.2±2.8	33.2±1.8	238±14
55R	October	20	7	4.4	4.4	<0.9	34.4±1.4	19.5±1.6	<15.8	142±11
56R	October	21	4.1	9.8	8.8	<0.9	25.7±0.9	43±3.4	39±3.1	246±11
57R	October	22	6.6	0.1	2.2	<0.9	51.7±1.9	<2.8	<15.8	1.3±0.3
58R	October	23	6.4	0.2	2.2	<0.9	36.2±1.3	4.2±0.6	<15.8	13.2±0.9

Results are mean values ± standard deviation for n = 3; < indicates a value below the detection limit

Online Resource 3

Grain size analysis of sediments collected during the dry season.

Sample no	% sand	% silt	% clay	Sample no	% sand	% silt	% clay
<i>Odo-Iyaaloro</i>							
1D	31.48	29.89	38.63	8D	61.61	13.37	25.02
2D	40.64	24.52	34.84	9D	59.7	18.41	21.89
3D	54.44	12.85	32.71	10D	67.21	13.44	19.35
4D	49.96	16.06	33.98	11D	47.32	16.46	36.22
5D	42.26	27.37	30.37	12D	48.23	14.96	36.81
6D	43.25	28.81	27.94	13D	45.82	22.93	31.25
7D	73.31	10.68	16.01	14D	38.14	18.21	43.65
<i>Shasha River</i>							
15D	64.57	10.67	24.76	24D	77.67	13.47	8.86
16D	95.81	1.92	2.27	25D	32.17	17.79	50.04
17D	44.88	16.09	39.03	26D	74.08	11.94	13.98
18D	77.67	13.47	8.86	27D	58.83	17.84	23.33
19D	32.17	17.79	50.04	28D	95.20	2.40	2.40
20D	77.08	11.94	13.98	29D	91.02	2.78	6.20
21D	54.58	17.77	27.65	30D	67.15	17.94	14.91
22D	79.42	13.73	6.85	31D	79.29	13.37	7.34
23D	41.64	17.1	41.26	32D	97.66	0.35	1.99
<i>Ibeshe River</i>							
33D	35.27	19.76	44.97	42D	93.35	6.47	0.18
34D	46.44	20.17	33.39	43D	82.81	12.36	4.83
35D	64.78	15.28	19.94	44D	95.08	4.92	0.00
36D	75.67	14.21	10.12	45D	29.7	17.53	52.77
37D	87.28	9.28	3.44	46D	19.78	13.92	66.30
38D	96.71	3.12	0.17	47D	93.88	3.32	2.80
39D	42.28	18.4	39.32	48D	54.03	34.71	11.26
40D	54.20	18.23	27.61	49D	68.21	19.92	11.87
41D	64.28	19.27	16.45	50D	46.98	32.17	20.85
<i>Lagos Lagoon</i>							
51D	61.08	18.95	19.97	54D	100.00	0.00	0.00
52D	79.76	8.87	11.37	55D	91.12	8.83	0.05
53D	68.84	13.53	17.63				

Online Resource 4

Grain size analysis of sediments collected during the rainy season.

Sample no	% sand	% silt	% clay	Sample no	% sand	% silt	% clay
<i>Odo-Iyaaloro</i>							
1R	86.43	5.18	8.39	7R	55.23	25.15	19.62
2R	57.85	14.95	27.20	8R	57.18	27.21	15.61
3R	60.18	15.31	24.51	9R	80.36	7.88	11.76
4R	63.62	11.37	25.01	10R	59.14	11.58	29.28
5R	74.86	7.28	17.86	11R	54.03	20.35	25.62
6R	59.76	18.21	22.03	12R	57.90	13.13	28.97
<i>Shasha River</i>							
13R	55.92	17.05	27.02	22R	79.03	11.28	9.69
14R	50.79	17.61	31.6	23R	90.24	3.71	6.04
15R	82.76	8.56	8.68	24R	91.28	2.73	5.99
16R	76.68	13.61	9.71	25R	52.18	20.17	27.65
17R	89.99	8.87	1.14	26R	81.19	10.12	8.69
18R	96.04	0.70	3.26	27R	79.92	10.71	9.37
19R	56.42	19.20	24.38	28R	92.37	3.76	3.87
20R	77.72	16.54	5.74	29R	94.28	3.10	2.62
21R	90.81	7.26	1.93	30R	91.76	2.78	5.46
<i>Ibeshe River</i>							
31R	1.09	6.71	92.20	40R	80.51	18.21	1.28
32R	16.07	20.70	63.23	41R	76.47	11.22	12.31
33R	98.95	1.05	0.00	42R	68.27	16.15	15.58
34R	62.25	16.82	20.93	43R	33.29	12.69	54.02
35R	98.59	1.34	0.07	44R	84.96	7.20	7.84
36R	85.52	11.02	3.46	45R	80.49	7.23	12.28
37R	30.21	18.24	51.55	46R	54.36	29.23	16.41
38R	31.27	46.01	22.72	45R	60.21	16.74	23.05
39R	82.61	7.52	9.87	48R	47.23	21.00	31.77
<i>Lagos Lagoon</i>							
49R	71.25	18.21	10.54	54R	68.01	17.26	14.73
50R	75.32	19.27	5.41	55R	77.21	9.86	12.93
51R	61.62	20.12	18.26	56R	62.67	17.21	20.12
52R	97.64	1.12	1.24	57R	95.27	4.61	0.12
53R	95.76	4.18	0.06	58R	96.72	3.28	0.00