

Renshaw, A. E. & Haberman, S. (1999). An empirical study of claim and sickness inception transition intensities (aspects of the UK permanent health insurance experience) (Report No. Actuarial Research Paper No. 121). London, UK: Faculty of Actuarial Science & Insurance, City University London.



**CITY UNIVERSITY  
LONDON**

[City Research Online](#)

**Original citation:** Renshaw, A. E. & Haberman, S. (1999). An empirical study of claim and sickness inception transition intensities (aspects of the UK permanent health insurance experience) (Report No. Actuarial Research Paper No. 121). London, UK: Faculty of Actuarial Science & Insurance, City University London.

**Permanent City Research Online URL:** <http://openaccess.city.ac.uk/2257/>

#### **Copyright & reuse**

City University London has developed City Research Online so that its users may access the research outputs of City University London's staff. Copyright © and Moral Rights for this paper are retained by the individual author(s) and/ or other copyright holders. All material in City Research Online is checked for eligibility for copyright before being made available in the live archive. URLs from City Research Online may be freely distributed and linked to from other web pages.

#### **Versions of research**

The version in City Research Online may differ from the final published version. Users are advised to check the Permanent City Research Online URL above for the status of the paper.

#### **Enquiries**

If you have any enquiries about any aspect of City Research Online, or if you wish to make contact with the author(s) of this paper, please email the team at [publications@city.ac.uk](mailto:publications@city.ac.uk).

**An Empirical Study of Claim and  
Sickness Inception Transition  
Intensities (Aspects of the UK  
Permanent Health Insurance  
Experience)**

by

**A E RENSHAW and S HABERMAN**

**Actuarial Research Paper No. 121**

**Department of Actuarial Science and Statistics  
City University  
London**

**November 1999**

**ISBN 1 901615 41 3**

**“Any opinions expressed in this paper are my/our own and not necessarily those of my/our employer or anyone else I/we have discussed them with. You must not copy this paper or quote it without my/our permission”.**

AN EMPIRICAL STUDY OF CLAIM AND SICKNESS INCEPTION TRANSITION INTENSITIES  
(ASPECTS OF THE UK PERMANENT HEALTH INSURANCE EXPERIENCE)

By A.E. RENSHAW, Ph.D AND S. HABERMAN, Ph.D, F.I.A.

(of City University, London)

ABSTRACT

We consider the relationship between the claim inception transition intensity and the sickness inception transition intensity. Using data from the CMI Bureau for Permanent Health Insurance (PHI) policies, we propose models for representing the underlying trends in the sickness inception intensities over an extended recent period, *viz* 1975-94. As part of this study, we note the dependence of this relationship on the underlying recovery intensities and we consider, in detail, the extent to which the recovery intensities for the policies with different deferred periods differ significantly (when matched by sickness duration and age).

KEYWORDS

Sickness inception intensities; Claim inception intensities

1. INTRODUCTION

We focus on the relationship between claim and sickness inception transition intensities from the healthy state, and explore its possible implications for modelling. This work may be seen as the natural extension to Renshaw & Haberman (1999) in which time trends, over the period 1975-94, are identified and modelled in respect of claim inception transition intensities (as well as for sickness recovery transition intensities and for the force of mortality when sick) for UK permanent health insurance (PHI) data. We also acknowledge aspects of the pioneering work of the CMI Bureau which, using the 1975-78 data base, established their PHI multiple state model (CMI Committee (1991)) and which are relevant to this work. Note, however, that we choose to operate wholly within a generalised linear modelling setting: as in Renshaw & Haberman (1995,1999).

In this study, it should be noted that we focus only on the male PHI UK experience, modelling the data either in 'dynamic' mode represented by individual calendar years, or in 'static' mode with the data grouped into one of five consecutive quadrennia defined sequentially within the twenty year period 1975-94 prior to analysis and with each quadrennium modelled separately. It should be noted that it is a straight forward matter to move from the first mode to the second mode by suppressing the relevant suffix in  $t$  (actual or implied) in any expressions and by pre-setting any time related parameters in the linear predictors to zero.

Given our stated interest in the 'dynamic' mode of analysis, it is a relatively simple matter to generalise the definition of the underpinning semi-Markovian process (CMI Committee (1991)), and to

develop the associated Kolmogorov forward equations, so as to accommodate this feature. Summary details are presented in Appendix I.

Issues concerning the implementation of the basic identity connecting claim and sickness transition intensities are discussed in Section 2. The chief of these, concerning the extent to which the recovery intensities, for different deferred periods based on a specific quadrennium, differ significantly, when matched by sickness duration and age is investigated in Section 3. The practical determination of the identity is described in Section 4. The sickness transition intensity is modelled as a function of age and calendar period effects in Section 5, while the implications of a viable alternative way of achieving this under the proposed modelling assumptions, are investigated in Section 6. The paper is summarised in Section 7.

## 2. A BASIC RELATIONSHIP

Let  $\tau_{tx}$  and  $\sigma_{tx}$  denote the respective claim and sickness inception transition intensities, for a healthy male, aged  $x$  in calendar year  $t$ , at sickness inception. They are related by the formula

$$\tau_{tx} = \pi_{txd} \sigma_{tx}$$

in which  $\pi_{txd}$  denotes the probability that the time spent by an individual in the sick state exceeds the deferred period  $d$  weeks, before which benefits become payable under the terms of the sickness insurance policy. This relationship plays a role in the calculation of expected present values of disability annuities in Sweden: see Mattsson (1956), Dillner (1969), Haberman & Pitacco (1999). It is also implicit in the approach used to graduate sickness inception intensities for the 1975-78 UK experience based on a normal approximation to an overdispersed Poisson distribution: see CMI Committee (1991). Typically, such UK experiences involve deferred periods  $d = 1, 4, 13$ , or 26 weeks, also referred to as DP1, DP4, DP13 and DP26 respectively.

The computation of  $\pi_{txd}$  is crucial to the targeting of  $\sigma_{tx}$ . Within the context of a suggested 'dynamic' version of the multiple state PHI model, it is defined by

$$\pi_{txd} = \exp - \int_0^{d/52} (\rho_{t+u, x+u, u} + \nu_{t+u, x+u, u}) du \quad (2.1)$$

where  $\rho_{txz}$  and  $\nu_{txz}$  denote the respective recovery transition intensity when sick and the force of mortality when sick, for an individual with attained age  $x$  years, for calendar year  $t$  and current sickness duration  $z$  weeks. When we consider a specific quadrennium, we shall suppress the first of the suffices. Note also that the expression for  $\pi_{txd}$  partitions into two multiplicative components, with the recovery component dominant over the mortality component, given the parameter estimates in CMI Committee (1991), Renshaw & Haberman (1995,1999).

As recognised within the context of the data set for the 1975-78 quadrennium (CMI Committee 1991), the evaluation of the integral in (2.1) poses intrinsic practical difficulties because data associated with the deferred period concerned are not available for the estimation of the functional forms of  $\rho$  and  $\sigma$  within the range of the integral. That is, observations are only available for sickness durations in excess of the respective deferred periods. Consequently, a plausible way forward is to resort to the use of functional forms for  $\rho$  and  $\nu$  based on the data associated with those deferred periods which are shorter than the one in question, where this is feasible. In so far as the contribution to  $\pi_{txd}$  from  $\nu$  is concerned, the difficulty is resolved by the practical necessity of having to group the data over all deferred periods prior to the modelling of  $\nu$ , matched by calendar period, age and sickness duration, because of the paucity of reported deaths.

With respect to the more dominant contribution from  $\rho$ , the CMI Committee (1991) study, based on the 1975-78 experience, has identified two characteristics in the data which are subsequently exploited in the determination of the contribution to  $\pi_{txd}$  from  $\rho$ . The first characteristic is the presence of tapering convergent sickness duration 'run-in' periods in which the recovery rates for DP4, DP13 and DP26 policies, in the first four weeks when claims are allowed under the terms of these policies, lag significantly behind the corresponding recovery rates for DP1 policies, matched for sickness duration and age. The second characteristic is that, once beyond these tapering 'run-in' thresholds, the recovery rates for DP1, DP4, DP13 and DP26 policies do not differ significantly when matched by sickness duration and age. Hence, the data in these cells effectively converge and may be grouped, prior to the modelling of  $\rho$  as a function of  $x$  and  $z$ . It is of interest to ascertain the extent to which these two characteristics extend from the 1975-78 experience to subsequent quadrennia. This problem is considered in Section 3.

### 3. A COMPARISON OF MATCHING RECOVERY INTENSITIES

In this section, our objective is designed to ascertain the extent to which the recovery intensities, for different deferred periods based on a specific quadrennium (so that the suffix  $t$  is suppressed throughout this section), differ significantly, when matched by sickness duration and age. In this respect we are mindful of the pattern of significant differences established for the 1975-78 quadrennium *viz* the 'run-in' thresholds described above. The approach used is based on the analysis of paired differences in log crude recovery intensities. It is preferred as an alternative to the approach adopted by the CMI Committee (1991) since it acknowledges the interactive nature of the dependence of  $\rho$  on  $x$  and  $z$  prior to testing. By way of cross-reference, we also reinvestigate the 1975-78 experience in addition to the other four quadrennial experiences.

For the purpose of the analysis we focus on the following cross-classification of data cells

gender male  
quadrennia 75-78, 79-82, 83-86, 87-90, 91-94  
deferred periods (dp) 1, 4, 13, 26 wks  
ages (x) 18-, 25-, 30-, 35-, 40-, 45-, 50-, 55-, 60-64 yrs (coded 1,2, ..., 9)  
sickness durations(z) 4,5,6,7,8,9,10,11-12; 13-14,15-17,18-21,22-25; 26-29,30-38,39-51 wks; 1yr,2-11yrs

This selection of categories agrees with that used by the CMI Committee (1996). In particular, we note that the four duration categories labelled 4, 5, 6 & 7 weeks constitute the so-called four week 'run-in' period for DP4, that the two duration categories labelled 13-14 & 15-17 effectively constitute the four week 'run-in' period for DP13, and that the single duration category labelled 26-29 constitutes the four week 'run-in' period for DP26.

For a specific quadrennium and for deferred period  $d$  ( $d = 1, 4, 13, 26$ ), let

$$\begin{aligned} r_{xz}^d &= \text{number of recoveries in cell } (d, x, z) \\ e_{xz}^d &= \text{exposure (in years) to the possibility of recovery in cell } (d, x, z) \end{aligned}$$

and compute the differences in log crude recovery intensities

$$y_{xz}^d = \log \left\{ \frac{r_{xz}^d}{e_{xz}^d} \right\} - \log \left\{ \frac{r_{xz}^d}{e_{xz}^d} \right\}$$

where  $d = 4, 13$  or  $26$ , as appropriate. By this means, we can compare log crude recovery intensities, matched by age and sickness duration, for each deferred period DP4, DP13, DP13, relative to DP1.

Let

$$E(Y_{xz}^d) = \eta_{xz}^d$$

denote the expected value of the difference in the log crude recovery intensities,  $Y_{xz}^d$ . Then, in the light of the CMI Committee (1991) findings described above, we formulate the null hypothesis:

$$H_0: \eta_{xz}^d \begin{cases} \neq 0 \quad \forall (z, x), \text{ within the 4 week 'run-in' threshold} \\ = 0 \quad \forall (z, x), \text{ otherwise} \end{cases} \quad \forall d = 4, 13, 26$$

Given the focus on *differences*, matched age for age as well as by sickness duration, we can test this hypothesis by fitting the one-way main effects parameterised structure

$$E(Y_{xz}^d) = \eta_{xz}^d = \alpha_z \tag{3.1}$$

for  $d = 4, 13, 26$  respectively. Under  $H_0$ , it is expected that the resulting predictor estimates  $\hat{\eta}_{xz}^d = \hat{\alpha}_z$  will differ significantly from zero inside the sickness duration 'run-in' threshold *only*. Further, the

establishment of (statistically significant) decreasing predictor estimates with increasing duration within the 'run-in' period (observable for DP4 and DP13) would be consistent with the tapering of recovery rates within the 'run-in' period. Clearly, non-significant predictor estimates beyond the 'run-in' threshold would be consistent with the convergence of the respective matched recovery rates.

As an added precaution, we also test for main age effects using the two-way additive structure *viz*

$$E(Y_{zz}^d) = \eta_{zz}^d = \alpha_z + \beta_x. \quad (3.2)$$

Given the emphasis here on matched differences in log recovery intensities, we do not anticipate this to be a statistically significant factor on the basis of the CMI Committee (1991) findings in relation to the 1975-78 experience.

The structures are fitted by the method of weighted least squares, with the  $y_{zz}^d$  statistics as the responses, and with weights

$$\frac{r_{zz}^1 r_{zz}^d}{r_{zz}^1 + r_{zz}^d}.$$

One aspect of this choice of weights is to ensure that in the event of a reported zero recovery cell count, for which the response variable goes undefined, such cells are weighted out of the analysis.

The parameter estimates  $\hat{\alpha}_z$ , coupled with their t-statistics, for the main sickness duration effects model structure (3.1) are presented systematically in Tables 3.1(a,b,c). The F-statistics associated with the addition of main age effects into the model structure are systematically reported in Table 3.2.

Scrutiny of the first column in each of Tables 3.1(a,b,c), representing the 1975-78 quadrennium, reveals that with one exception, all of the  $\alpha_z$ s differ significantly from zero within the durational 'run-in' periods and that again with one exception, all of the  $\alpha_z$ s do not differ significantly from zero outside the 'run-in' period. In addition, the values of the predictor estimates imply a pronounced tapering effect within the 'run-in' for the DP4 and DP13 experiences. We note, however, that there is no statistically significant support for a four week 'run-in' period for the DP26 experience from this analysis. Scrutiny of the first column in Table 3.2, again representing the 1975-78 quadrennium, provides strong evidence that, as anticipated, age effects are not statistically significant. Such results provide strong confirmation of the convergent 'run-in' periods for the DP4 and DP13 experiences identified by the CMI Committee (1991) for their PHI model and also lend strong support for the modelling of a common set of recovery intensities across deferred periods outside of the four week durational 'run-in' periods.

The situation is not quite so clear cut, however, in the more recent quadrennia under investigation, the results for which are recorded in the remaining columns of Tables 3.1(a,b,c) and Table 3.2. Tables 3.1(a,b,c) confirm the presence of statistically significant non-zero  $\alpha_z$ s, with one



exception (DP13 for 1991-94), within the durational 'run-in' periods for each of the four quadrennia. These results indicate the persistence of the initial four week differential durational effect for recovery intensities when comparing DP4, DP13, and DP26 policies on the one hand and DP1 policies on the other hand, matched by sickness duration and by age. However, the distinctive tapering effect for the DP4 experiences across the four individual weeks concerned is not quite so pronounced for the 1983-86 and 1991-94 quadrennia (Table 3.1(a)). More importantly, there is a noteworthy occurrence of statistically significant non-zero  $\alpha_{zs}$  outside the durational 'run-in' period coupled with the added complication of one significant main age effects F-statistic (and a further two marginally significant main age effects F-statistics: see Table 3.2). This runs somewhat counter to the rationale of grouping matching data cells across different deferred periods, outside the durational 'run-in' periods, prior to the modelling of recovery rates. We summarise these results by tabulating the tally of statistically significant  $\alpha_{zs}$  outside the 'run-in' period thus:

	<i>Quadrennium</i>					<i>Total possible</i>
	<i>75-78</i>	<i>79-82</i>	<i>83-86</i>	<i>87-90</i>	<i>91-94</i>	
<i>DP4</i>	0	1*	0	4	3	13
<i>DP13</i>	0	2	0†	3†	4	7
<i>DP26</i>	1	3	1	2	0	4

(\* denotes statistical significant age effects which should be taken into account)

(† denotes marginally significant age effects which are also investigated)

Tally of statistically significant non-zero  $\alpha_{zs}$  outside the durational 'run-in' period

This summary table provides support for grouping in the case of the 1983-86 quadrennium (in addition to the 1975-78 quadrennium). The case for grouping is not well supported by these results when it comes to the other three quadrennia, although there are no obviously discernible, systematic patterns in the non-zero  $\alpha_{zs}$  (outside of the four week 'run-in' periods) with one exception. This exception concerns DP4 policies in the 1987-90 quadrennium for which there is evidence of an extended 'run-in' of up to 12 weeks duration.

Next, we investigate the implications of the statistically significant F-statistic for DP4 policies in the 1979-82 quadrennium, established on fitting the two-way main effects structure, equation (3.2). On contrasting the resulting age effects, it is possible to reduce the number of effective age parameters  $\beta_x$ , without a significant loss of information, to just two, by setting

$\beta_1 = 0$  representing jointly the youngest age group [18,30) and the combined oldest ages [50,65]

$\beta_2 \neq 0$  representing the single age group [30,35)

$\beta_3 \neq 0$  representing the combined age groups [35,50)

(Note that the first of these parameters is set equal to zero out of necessity in order to avoid over-parameterisation in a technical sense). The resulting estimated parameter values

$$\hat{\eta}_{x'z}^4 = \hat{\alpha}_z + \hat{\beta}_{x'}$$

together with the t-statistics are reported, in the appropriate format, in Table 3.3. While these results are supportive of a tapering four week 'run-in' period, the results relating in particular to the 30-34 age group, and to a slightly lesser degree, the 35-49 age group, also establish significant differences in recovery intensities beyond the four week 'run-in' threshold.

Finally, for the record, it is also of interest to record the implications of the marginally significant F-statistic (p-value 5.5%) for DP13 policies in the 1983-86 quadrennium. This time, following an analysis of the contrasts in age effects, we are prompted to use five effective parameters  $\beta_{x'}$ , achieved by using the same parameter for the youngest and the oldest age groups,  $[18,30) \cup [60,65]$ , and by combining the adjacent age groups  $[45,50) \cup [50,55]$  into a single grouping. The resulting estimated parameter values  $\hat{\eta}_{x'z}^{13}$ , together with the associated t-statistics are recorded in Table 3.4.

We have not pursued further the marginally significant F-statistic (p-value 8.3%) for DP13 policies in the 1987-90 quadrennium.

#### 4. COMPUTATION OF $\pi_{txd}$

In this section, we need to consider the underlying transition intensities necessary for the calculation of  $\pi_{txd}$ . For the purpose of this study, we make use of equation (2.1) in combination with the formulae

$$\begin{aligned} \log \rho_{t+z, x+z, z} = \\ \mu + \alpha\sqrt{z} + \beta z + \gamma_1 t + \gamma_2 t\sqrt{z} + \gamma_3 tz + \theta x + \theta_2 x^2 + \theta_3 x^3 + \phi x\sqrt{z} + \psi xz \end{aligned} \quad (4.1)$$

$$\log \nu_{t+z, x+z, z} = \mu + \gamma t + \theta x + \beta z + \sum_{j=1}^n \beta_j (z - z_j)_+ \quad (4.2)$$

to compute values of  $\pi_{txd}$ . These formulae are established in Renshaw & Haberman (1999) in an attempt to model the respective trends in the recovery intensities for the male 1975-94 DP1 experience, (4.1), and the mortality from sickness intensities for the male 1975-94 experience for all DPs combined, (4.2). Given the somewhat inconclusive nature of the test results of Section 3 for sickness durations in excess of the 'run-in' thresholds, we have opted not to group the data over deferred period in any matching cells, prior to fitting the first of these formulae. In addition, with the 'static' mode of analysis in mind, the data have been grouped by quadrennia and modelled separately. It transpires that each of these cases is satisfactorily represented by a member of the same class of formulae, with obviously  $\gamma_1, \gamma_2, \gamma_3, \gamma$  pre-set to zero together with certain of the other parameters, as appropriate. The results are summarised in Table 4.1 and Table 4.2 in which details of the parameters estimates are presented, and from which details of the formulae structures for each case may be deduced. Details of

the integration of (2.1) are outlined in Appendix II. It is a relatively simple matter to program the integration under the 'static' mode as a special case of the integration under the 'dynamic' mode.

#### 5. TARGETING $\sigma_{tx}$

The same data sets are used to target both the sickness inception intensity  $\sigma_{tx}$  and the claim inception intensity  $\tau_{tx}$ . For a specific gender and specific deferred period, they comprise claim inception counts  $i_{tx}$  representing the number of sicknesses starting in cell  $(t, x)$  and which last beyond the deferred period of the class of policies concerned, together with matching exposures  $e_{tx}$  measured in years.

It should be noted that the exposures do not include any allowance for the exposure time spent as sick but not claiming, the effect of which is reportedly (CMI Committee 1996) "to overstate the exposure by about 0.5%". In addition, as with the targeting of  $\tau_{tx}$  (Renshaw & Haberman 1999), we have not inflated the reported claim inception counts by a hypothetical figure for "the number of recoveries within four weeks after the end of the deferred period which were not reported as claims or recoveries", as described for the 1975-78 quadrennial (DP4, DP13, DP26) experiences in Part C Section 3 CMI Committee (1991). In particular, it should be noted that this adjustment is not insubstantial, being of the order of at least 25% over the whole of the reported age range in the case of the DP4 experience (Table C1 p56, CMI Committee 1991), and consequently has a material effect on the resulting 'standard' sickness intensities based on the 1975-78 experience (Tables C7, C8, C14, CMI Committee 1991). We take as justification for our approach, the subsequent need to re-adjust any *expected* claim inception counts based on the 'standard' sickness inception intensities, in order to counteract the effects of the hypothetical unreported claims built into the 1975-78 'standard' tables, prior to comparison with the actual claim counts in quadrennia subsequent to 1975-78 (CMI Committee 1996).

The data have been cross-classified in the first instance by

gender	male
deferred periods (dp)	1, 4, 13, 26 wks
age at inception (x)	18-, 25-, 30-, 35-, 40-, 45-, 50-, 55-, 60-64 yrs
calendar time at inception (t)	1975, 76, 77, ..., 94

with the further option of pooling the data into the five separate successive quadrennia 75-78, 79-82, ..., while at the same time relaxing the grouping with respect to  $x$ , prior to analysis. [Inset to follow]

For a specific deferred period  $d$ , we follow Renshaw & Haberman (1995) in targeting  $\sigma_{tx}$  within the generalised linear modelling framework by declaring independent overdispersed Poisson responses  $i_{tx}$  such that

$$m_{tx} = E(i_{tx}) = e_{tx} \pi_{txd} \sigma_{tx}, \text{Var}(i_{tx}) = \phi m_{tx},$$

where the scale parameter  $\phi$  is included in recognition of duplicates among the policies contributing to the data base. The associated deterministic structure is based on the log-link predictor relationship

$$\eta_{tx} = \log m_{tx} = \log(e_{tx} \pi_{txd}) + \log \sigma_{tx}$$

where  $\eta_{tx}$  denotes the linear predictor and  $\log(e_{tx} \pi_{txd})$  the offset term.

Under this modelling formulation, when  $\pi_{txd} = 1 \forall t, x$  and specific  $d$ ,  $\sigma_{tx}$  becomes  $\tau_{tx}$  and we revert to the modelling of claim inception transition intensities. Here we are interested in modelling the trends in the sickness inception transitions and in comparing such trends with the trends established in the claim inception transitions (Renshaw & Haberman (1999)).

The role played by the (constant) scale parameter  $\phi$  is of potential interest. Given that it impacts on the second moment properties and not on the first, it has no bearing on the values of the parameter estimates in the linear predictor, and hence on the fitted values  $\hat{m}_{tx}$  (and  $\hat{\sigma}_{tx}$  or  $\hat{\tau}_{tx}$ ). Instead it forms a multiplicative factor  $\sqrt{\phi}$  in the construction of the standard errors for the parameter estimates.

In the generalised linear modelling setting, it is customary to estimate  $\phi$  as the model deviance  $D$ , divided by the model degrees-of-freedom. For the current choice of modelling distribution

$$D = \sum_{tx} d_{tx} = 2 \sum_{tx} \left\{ i_{tx} \log \frac{i_{tx}}{\hat{m}_{tx}} - (i_{tx} - \hat{m}_{tx}) \right\}.$$

Also, given the presence of the free standing scale parameter  $\phi$  in this analysis, we make use of Studentised deviance residuals

$$\text{sign}(i_{tx} - \hat{m}_{tx}) \sqrt{\frac{d_{tx}}{\phi(1 - h_{tx})}}$$

throughout, where  $h_{tx}$  are the diagonal entries in the so called hat-matrix of the fit (eg McCullagh & Nelder (1989), Francis, Green & Payne (1993)). Typically here, values of  $h_{tx}$  are of the order of 0.05 and consequently have little impact on the values of such residuals in general, while the contribution from the scale parameter  $\phi$  is analogous to the scaling of residuals under the more familiar normal  $N(\mu, \sigma^2)$  modelling assumption (scale parameter  $\sigma^2$ ).

As with the targeting of  $\tau$  (Renshaw & Haberman (1999)), bivariate polynomial structures

$$\log \sigma_{t'x} = \alpha_0 + \sum_{i=1}^r \alpha_i t'^i + \sum_{j=1}^s \beta_j x^j + \sum_{i=1}^r \sum_{j=1}^s \gamma_{ij} t'^i x^j$$

are fitted for separate DPs, where  $x$  is allocated values at the centre of the relevant age cell (as defined above) and  $t' = (t - 1984.5)/9.5$  taking equally spaced values in the interval  $[-1, 1]$  corresponding to the calendar years  $t = 1975, 76, \dots, 94$ . Details of the fitted structures may be deduced from Table 5.1 in which the parameter estimates, their standard errors and 't-statistics' are tabulated. For completeness, we also tabulate details of the parameter estimates associated with the modelling of  $\tau$  as reported in Renshaw & Haberman (1999). In the case of  $\sigma$ , the fitted structures comprise

<i>DP1</i>	quintic in $t$ effects + cubic in $x$ effects + 2nd & 3rd order product effects
<i>DP4</i>	quintic in $t$ effects + cubic in $x$ effects + 2nd order product effects
<i>DP13</i>	cubic in $t$ effects + cubic in $x$ effects
<i>DP26</i>	quintic in $t$ effects + quadratic in $x$ effects

which are identical in form to the case of  $\tau$  with the exception of DP26 (cubic  $t$  effects plus cubic  $x$  effects). Each structure has been selected on the basis of exploratory data analysis involving an examination of deviance profiles, residual plots, and the statistical significance or otherwise of the associated parameter estimates. In particular, residuals plotted against both age and calendar year of sickness inception are especially informative. For economy of space, we tabulate the residuals for each DP in Table 5.2 instead of reproducing the plots. By scanning both the columns and rows of these rectangular tabulations (age by period), it is possible to make an assessment of the goodness-of-fit. With the possible exception of the years 1981, 1992 (DP1), the year 1982 (DP13 & DP26), and ages 18-24 (DP26), where negative residuals are over represented, the tabulations (and underlying plots) are supportive of the fitted model structures. Any such clusters of negative residuals imply that there is a tendency for the predicted sickness inception intensities to be overstated at these locations. Further, with the obvious exception of the case of DP26 at ages 18-24, it appears (from Figure 5.1) that these negative residuals occur at turning points in the predicted trends, which is a possible reflection of the rigidity of low order bivariate polynomial structures in accommodating sharp turning points. The apparent lack of fit at ages 18-24 for DP26 is attributable to the paucity of data in these cells, comprising relatively low exposures coupled mainly with zero reported transitions. As with this approach to modelling in general, any structures selected by these methods are demonstrably plausible, but are of necessity, not unique.

The role played here by the scale parameter  $\phi$  is that of a token variance ratio, assumed to be *constant* over all the data cells  $(t, x)$ . (The information necessary to compute the variance ratios for individual cells is not available). On the evidence of the typical set of variance ratios computed at each age for the 1979-82 male assured lives experience (Table 17.6, Forfar *et al.* 1988), estimated values,  $\hat{\phi}$ , in the range 1 to 2 are consistent with these tabulated values. While this is the case for three of the four deferred periods considered (Table 5.1), the estimated value,  $\hat{\phi}$ , for the DP1 experience is excessive. An examination of the individual deviance components reveals that this is due in no small part to relative large contributions from relatively few of the data cells. It is possible to identify these

cells by reference to Table 5.2. These are the cells associated with the largest residuals in absolute value. Note that these occur in roughly equal proportions of positives and negatives and are seemingly randomly scattered. In addition, we note that the value of the deviance and hence the estimated value,  $\hat{\phi}$ , can generally be reduced through the introduction of more complex predictor structures. Having considered this possibility, we have concluded that we do not feel justified in doing this within the context of bivariate low degree polynomial predictor structures. We also remark, that as a guide to parameter retention when referring to Table 5.1, that if a *smaller* value  $\tilde{\phi}$  of  $\phi$  is imposed while at the same time retaining the same predictor giving rise to the estimated value  $\hat{\phi}$  of  $\phi$ , the resulting values of the t-statistics are *increased* by a factor of  $\sqrt{(\hat{\phi}/\tilde{\phi})}$ . That said, it is important to keep this issue in perspective, namely that it relates only to the second moment properties of the modelling process.

The resulting smoothed log sickness inception intensities generated by these models for each DP, plotted against calendar year for each age, are illustrated in the upper frames of Figures 5.1(a-d). Also by way of comparison, the matching smoothed log claim inception intensities reported in Renshaw & Haberman (1999) are reproduced in the respective lower frame of each of these figures.

The families of parallel curves in the case of the DP13 and DP26 experiences reflect the additive nature of the linear predictor in age and calendar period effects. For the DP1 and DP4 experiences the interactive nature of these curves is more pronounced in the DP1 experience compared with the DP4 experience. For the DP13 and DP26 experiences, with the exception of the extreme ages (below age 30 for claim inceptions and above age 55 for sickness inceptions), a noteworthy feature of these graphs concerns the reversal of the ordering of members of the same family of curves (age effects) when comparing log sickness with matching log claim inception intensities. This feature also has a strong presence in the DP4 experience, and, to a lesser extent, in the DP1 experience, although these cases are complicated by the interactive nature of the family of curves. In essence, claim inception intensities initially decrease before increasing with increasing age, approximately above the age of 30. On the other hand, sickness inception intensities decrease with increasing age, approximately below the age of 55 before increasing, subject to possible interaction effects for the DP1 experience in particular and, to a lesser extent, for the DP4 experience. We note that these patterns are largely consistent with the patterns reported in Table C16 (CMI Committee (1991)) for graduated values of  $\sigma_x$  based on the 1975-78 quadrennial male experience but are only partially consistent (for ages over 30) with the patterns in Table C17 for  $\tau_x$  which increase with age over the whole of the age range for the 1975-78 quadrennial experience.

The cyclical appearance of these curves leads to the suggestion of an association with indices of the economic cycle, and a plausible causal link. In particular, there is a pronounced local minimum in the years 1981-82 (the depth of which is somewhat more extreme than that predicted by the model) coupled with a localised maximum in the years 1990-91. We note that Haberman & Walsh (1998) found difficulties in establishing a plausible causal link with a range of economic measures, albeit for a more restricted data set. This remains an area for future investigation.

## 6. ALTERNATIVE WAYS OF MODELLING $\sigma_{tx}$

The modelling process described in Section 5 gives rise to a choice of methods when targeting  $\sigma_{tx}$  (or  $\tau_{tx}$ ). Thus in Section 5, for a given deferred period  $d$ , by incorporating the probabilities  $\pi_{txd}$  into the offset, we have targeted  $\sigma_{tx}$  directly (so that  $\tau_{tx}$  can then be targeted indirectly using the identity  $\tau_{tx} = \pi_{txd}\sigma_{tx}$ ). Alternatively, by suppressing  $\pi_{txd}$  in the offset, we can target  $\tau_{tx}$  directly, as in Renshaw & Haberman (1999), and subsequently use the inverse identity  $\sigma_{tx} = \pi_{txd}^{-1}\tau_{tx}$  to target  $\sigma_{tx}$ , indirectly. For a given deferred period, let  $\hat{\sigma}_{tx}$  and  $\hat{\tau}_{tx}$  denote the respective fitted sickness and claim inception intensities generated by these two modelling approaches respectively: see Table 5.1 for details of the respective fits. It then follows from the very nature of the separate fitting processes, that  $\hat{\sigma}_{tx}$  and  $\hat{\tau}_{tx}$  will depart from the basic relationship  $\hat{\tau}_{tx} = \pi_{txd}\hat{\sigma}_{tx}$ . It is of interest to investigate the magnitude of this effect in order to gain a better understanding of the implications of these choices. It is apparent that such an investigation can only be conducted on an empirical level.

Given  $\hat{\tau}_{tx}$  or  $\hat{\sigma}_{tx}$  define  $\hat{\tau}_{tx}^*$  or  $\hat{\sigma}_{tx}^*$  respectively, such that

$$\begin{aligned} \hat{\tau}_{tx}^* &= \pi_{txd}\hat{\sigma}_{tx} && \text{when } \sigma_{tx} \text{ is the target of the model fitting exercise, and} \\ \hat{\sigma}_{tx}^* &= \pi_{txd}^{-1}\hat{\tau}_{tx} && \text{when } \tau_{tx} \text{ is the target of the model fitting exercise.} \end{aligned}$$

Then the values

$$\frac{\hat{\sigma}_{tx}^* - \hat{\sigma}_{tx}}{\hat{\sigma}_{tx}} \times 100\% \quad \left( = \frac{\hat{\tau}_{tx} - \hat{\tau}_{tx}^*}{\hat{\tau}_{tx}} \times 100\% \right) \quad (6.1)$$

provide a measure of the extent by which  $\hat{\sigma}^*$  and  $\hat{\sigma}$  differ, (and by implication, the extent by which  $\hat{\tau}$  and  $\hat{\tau}^*$  differ). Alternatively, the roles of the symbols  $\hat{\cdot}$  and  $\hat{\cdot}^*$  can be reversed in this ratio, leading to slightly different but equivalent measures of these effects. For practical purposes, it is sufficient to quote only one of these, and we present the former. The values based on the model structures reported in Table 5.1, matched for DP, are presented in Table 6.1, for alternate calendar years. On the basis of these figures it is possible to draw the broad conclusions that (a) there is a tendency for the results to diverge with increasing deferred period  $d$ : recall that the domain of the integral which determines  $\pi_{txd}$  in (2.1), and which lies outside the domain of the data set, increases with  $d$ , and that (b) the divergence of results is especially great in the age range 18-25 where the inception transition counts are generally low.

The same effects can also be investigated in more detail for specific quadrennia with the data classified by individual years of age  $x = 18, 19, \dots, 64$  rather than in grouped format (and the suffix  $t$  suppressed): the same format as that used in the construction of the standard PHI model (CMI Committee (1991)) based on the 1975-78 quadrennium. By way of illustration, we concentrate on the 1991-94 quadrennium and fit the following polynomial age effects structures

$$\log \sigma_{x'} \text{ or } \log \tau_{x'} = \sum_{j=0}^s \beta_j x'^j$$

with  $x' = (x - 41)/23$ , ranging from -1 to 1. The values of  $\pi_{x,d}$  are determined through the numerical evaluation of the integral expression (2.1) using the versions of formulae (4.1) and (4.2) for  $\rho_{xz}$  and  $\nu_{xz}$  implied by the respective final columns of Tables 4.1 and 4.2. Details of the fitted structures may be summarised thus:

	<i>DP1</i>	<i>DP4</i>	<i>DP13</i>	<i>DP26</i>
<i>target <math>\sigma</math></i>	cubic	quintic	quartic	quintic
<i>target <math>\tau</math></i>	quadratic	quintic	quartic	quintic

1991-94 quadrennial experience: fitted age effects

with the parameter details given in Table 6.2. The values of the ratio (6.1) based on these results are presented in Table 6.3 which are in agreement with the broad conclusions stated above.

## 7. SUMMARY

The same raw data are used to target both the claim inception intensity  $\tau_{tx}$  and the sickness inception intensity  $\sigma_{tx}$ . For a given deferred period  $d$ , these intensities are related by the identity

$$\tau_{tx} = \pi_{txd} \sigma_{tx}$$

where  $\pi_{txd}$  is the probability that the time spent by an individual in the sick state exceeds the deferred period  $d$  in question. A knowledge of  $\pi_{txd}$  is essential when targeting  $\sigma_{tx}$  but is not essential when targeting  $\tau_{tx}$ .

The evaluation of expression (2.1), viz

$$\pi_{txd} = \exp - \int_0^{d/52} (\rho_{t+u, x+t+u, u} + \nu_{t+u, x+t+u, u}) du$$

is problematic in the sense that formulae for  $\rho$  and  $\nu$  have to be assumed, within the limits of the integral, for the specific deferred  $d$  in question. The recovery transition intensity  $\rho$  is the dominant term in the integral. For the 1975-78 quadrennial experience (CMI Committee (1991)), four week 'run-in' periods in the DP4, DP13 and DP26 experiences have been identified, outside of which, a common formula for  $\rho$  is established across all deferred periods. This formula is then used to determine the contribution to the integral from  $\rho$  by extrapolating backwards to week zero. We have devised a method (Section 3) for testing these features and applied them to each of the five quadrennia 75-78, 79-82, 83-86, 87-90 and 91-94 with somewhat mixed results. While there is strong supportive evidence of persistent four week 'run-in' periods across all of the five quadrennia examined, the evidence for grouping the data outside these regions across all deferred periods is less compelling, to varying degrees, in the quadrennia subsequent to the initial 1975-78 quadrennium.

Bivariate polynomial structures have been used to model  $\sigma_{tx}$  as a function of calendar year  $t$  and



age  $x$ , for each deferred period  $d$  (Section 5), using values of  $\pi_{txd}$  based on the DP1 study of  $\rho_{txz}$  and the combined DP study of  $\nu_{txz}$  (Renshaw & Haberman (1999)). Additionally, the patterns in the sickness inception intensity  $\sigma_{tx}$  have been compared and contrasted with those in the claim inception intensity  $\tau_{tx}$ , matched for deferred period. Two noteworthy features, common to both inception intensities and all deferred periods, are a local minimum in the years 1981-82 and a local maximum in the years 1990-91.

An empirical investigation has been conducted (Section 6) into the differences which can arise in the fitted values of  $\sigma_{tx}$  under the two alternative modes of implementing the proposed model formulation. While these differences are shown to be very small over the bulk of the rectangular domain of the function  $\sigma_{tx}$ , there is a general tendency for the differences to increase with increasing deferred period. In addition, large differences are recorded for the youngest age grouping 18-24, where there is a paucity of data. The ultimate choice as to which method of targeting  $\sigma_{tx}$  to employ, may well be dictated by practical necessity. Thus, not only is the so-called 'direct' method, in which  $\log \pi_{txd}$  features in the offset, intuitively appealing given the central role played by  $\sigma$  in the formulation of the PHI multi-state model, it also has the advantage of leading to a closed formula for  $\sigma_{tx}$ . This in turn has distinct computational advantages when it comes to the numerical computation of the associated probabilities and sickness benefits.

Finally it is noted that the Kolmogorov forward equations, underpinning the 'static' multiple state model based on the 1975-78 quadrennial experience (CMI Committee (1991)), generalise to allow explicitly for possible variation with calendar time over a longer time scale.

#### ACKNOWLEDGEMENTS


The authors wish to thank the CMI Bureau for providing both financial support and the data.

#### REFERENCES

- CMI Committee (1991). The analysis of Permanent Health Insurance data. *CMIR*, 12, 1-263.
- CMI Committee (1996). Recovery rates under PHI policies, individually 1975-90 and grouped 1975-86. *CMIR*, 15, 1-50.
- DILLNER, C.G. (1969). New bases for non-cancellable sickness insurance in Sweden. *Skandinavisk Aktuarietidskrift*, 52, 113-124.
- FRANCIS, B., GREEN, M. & PAYNE C. (1993). The Glim system. Release 4 Manual. *Clarendon Press, Oxford*.
- FORFAR, B.A., MCCUTCHEON, J.J. & WILKIE A.D. (1988). On graduation by mathematical formula. *Journal of the Institute of Actuaries*, 115, 1-135.
- HABERMAN, S. & PITACCO, E. (1999). Actuarial Models for Disability Insurance. Chapman and Hall/CRC, Boca Raton. (Chapter 3.3).
- HABERMAN, S. & WALSH, D.J.P. (1998). Analysis of trends in PHI claim inceptions data. *Trans 26th International Congress of Actuaries*, 6, 159-186.
- MATTSSON, P. (1956). New bases for non-cancellable sickness insurance in Sweden. *Skandinavisk Aktuarietidskrift*, 39, 198.
- MCCULLAGH P. & NELDER J.A. (1989). Generalised linear models. *Chapman & Hall*.
- RENSHAW, A.E. & HABERMAN, S. (1995). On the graduations associated with a multiple state model for permanent health insurance. *Insurance: Mathematics and Economics*, 18, 1-17.
- RENSHAW, A.E. & HABERMAN, S. (1999). Modelling the recent time trends in UK Permanent Health Insurance recovery, mortality and claim inception transition intensities. *Submitted*.

APPENDIX I

Consider the three state Markovian process depicted by



The process is defined by

$$\{Y(t, x), Z(t, x): x \geq 0, t \geq 0\}$$

where  $Y(t, x) = H$  (for healthy),  $S$  (for sick) or  $D$  (for death), is a *state variable*, and  $Z(t, x) \in \mathbb{R}_{>0}$  is the duration of the sojourn so far in the current state  $Y(t, x)$ , for a life now aged  $x$  and time  $t$ . For example:

$$\{Y(t, x) = S, Z(t, x) = z\}$$

denotes that an individual is sick at age  $x$ , time  $t$ , and that the duration of the current sickness is  $z$ .

Define

$${}_sP_{txz}^{jk} = P(Y(t+s, x+s) = k \mid Y(t, x) = j \text{ and } Z(t, x) = z)$$

subject to the following implied rationalisation, summarised in tabular format below

$j \backslash k$	${}_sP_{txz}^{jk}$		
	$H$	$S$	$D$
$H$	${}_sP_{tx}^{HH}$	${}_sP_{tx}^{HS}$	${}_sP_{tx}^{HD}$
$S$	${}_sP_{tx}^{SH}$	${}_sP_{tx}^{SS}$	${}_sP_{tx}^{SD}$
$D$	0	0	1

Also define

1)  ${}_sP_{tx}^{\bar{H}\bar{H}} = P(Y(t+s, x+s) = H \text{ and } Z(t+s, x+s) \geq s \mid Y(t, x) = H)$

the probability that an individual remains healthy from  $(t, x)$  to  $(t+s, x+s)$  given the individual was healthy at  $(t, x)$

2)  ${}_sP_{txz}^{\bar{S}\bar{S}} = P(Y(t+s, x+s) = S \text{ and } Z(t+s, x+s) = z + s \mid Y(t, x) = S \text{ and } Z(t, x) = z)$

the probability that an individual remains sick from  $(t, x)$  to  $(t+s, x+s)$  given the individual was sick at  $(t, x)$  with duration of sickness  $z$ ; with the special case when  $z = 0$  denoted by  ${}_sP_{tx0}^{\bar{S}\bar{S}} = {}_sP_{tx}^{\bar{S}\bar{S}}$

3)  ${}_{ws}P_{tx}^{HS} = P(Y(t+s, x+s) = S \text{ and } Z(t+s, x+s) \leq w \mid Y(t, x) = H)$

For this process, the transitions in a small interval of time  $(s, s+\delta s]$  behave as follows:

$$P(Y(t+s+\delta s, x+s+\delta s) = S | Y(t+s, x+s) = H) = \sigma_{t+s, x+s} \delta s + o(\delta s)$$

$$P(Y(t+s+\delta s, x+s+\delta s) = D | Y(t+s, x+s) = H) = \mu_{t+s, x+s} \delta s + o(\delta s)$$

$$P(Y(t+s+\delta s, x+s+\delta s) = H | Y(t+s, x+s) = S \text{ and } Z(t+s, x+s) = z) = \rho_{t+s, x+s, z} \delta s + o(\delta s)$$

$$P(Y(t+s+\delta s, x+s+\delta s) = D | Y(t+s, x+s) = S \text{ and } Z(t+s, x+s) = z) = \nu_{t+s, x+s, z} \delta s + o(\delta s)$$

$$P(2 \text{ or more transitions}) = o(\delta s)$$

where

$$\lim_{\delta s \rightarrow 0} \frac{o(\delta s)}{\delta s} = 0.$$

1. Focus on

$${}_s P_{tx}^{\bar{H}\bar{H}} = P(Y(t+s, x+s) = H \text{ and } Z(t+s, x+s) \geq s | Y(t, x) = H).$$

Get

$${}_{s+\delta s} P_{tx}^{\bar{H}\bar{H}} = {}_s P_{tx}^{\bar{H}\bar{H}} (1 - \mu_{t+s, x+s} \delta s - \sigma_{t+s, x+s} \delta s) + o(\delta s)$$

which implies

$$\frac{\partial {}_s P_{tx}^{\bar{H}\bar{H}}}{\partial s} = -{}_s P_{tx}^{\bar{H}\bar{H}} (\mu_{t+s, x+s} + \sigma_{t+s, x+s}).$$

Integration gives

$${}_s P_{tx}^{\bar{H}\bar{H}} = \exp\left(-\int_0^s (\mu_{t+u, x+u} + \sigma_{t+u, x+u}) du\right).$$

2. Focus on

$${}_s P_{tz}^{\bar{S}\bar{S}} = P(Y(t+s, x+s) = S \text{ and } Z(t+s, x+s) = z + s | Y(t, x) = S \text{ and } Z(t, x) = z).$$

Get

$${}_{s+\delta s} P_{tz}^{\bar{S}\bar{S}} = {}_s P_{tz}^{\bar{S}\bar{S}} (1 - \rho_{t+s, x+s, z+s} \delta s - \nu_{t+s, x+s, z+s} \delta s) + o(\delta s)$$

which implies

$$\frac{\partial {}_s P_{tz}^{\bar{S}\bar{S}}}{\partial s} = -{}_s P_{tz}^{\bar{S}\bar{S}} (\rho_{t+s, x+s, z+s} + \nu_{t+s, x+s, z+s}).$$

Integration gives

$${}_s P_{tz}^{\bar{S}\bar{S}} = \exp\left(-\int_0^s (\rho_{t+u, x+u, z+u} + \nu_{t+u, x+u, z+u}) du\right).$$


Note in particular

$$\pi_{txd} = {}_{d/52} P_{tx0}^{\bar{S}\bar{S}} = {}_{d/52} P_{tx}^{\bar{S}\bar{S}}.$$

3. Focus on

$${}_sP_{tx}^{HH} = P(Y(t+s, x+s) = H | Y(t, x) = H).$$

Consider



Get

$${}_{s+\delta s}P_{tx}^{HH} = {}_sP_{tx}^{HH} (1 - \mu_{t+s, x+s} \delta s - \sigma_{t+s, x+s} \delta s) \quad (\text{contribution, route 1})$$

$$+ \int_{u=0}^s {}_uP_{tx}^{HH} \sigma_{t+u, x+u} {}_{s-u}P_{t+u, x+u}^{\bar{S}} \rho_{t+s, x+s, s-u} du \delta s \quad (\text{contribution, route 2})$$


which implies:

$$\frac{\partial {}_sP_{tx}^{HH}}{\partial s} = -{}_sP_{tx}^{HH} (\mu_{t+s, x+s} + \sigma_{t+s, x+s}) + \int_0^s {}_uP_{tx}^{HH} \sigma_{t+u, x+u} {}_{s-u}P_{t+u, x+u}^{\bar{S}} \rho_{t+s, x+s, s-u} du.$$

4. Focus on

$${}_wP_{tx}^{HS} = P(Y(t+s, x+s) = S \text{ and } Z(t+s, x+s) \leq w | Y(t, x) = H).$$

Consider



Get

$$\frac{\partial {}_w s P_{tx}^{HH}}{\partial w} = {}_{s-w} P_{tx}^{HH} \sigma_{t+s-w, x+s-w} \bar{u} P_{t+s-w, x+s-w}^{\bar{S}\bar{S}} \quad 0 \leq w < t$$

$$= 0 \quad w > t.$$


Integration implies

$${}_w s P_{tx}^{HS} = \int_{\max(0, s-w)}^s {}_u P_{tx}^{HH} \sigma_{t+u, x+u} {}_{s-u} P_{t+u, x+u}^{\bar{S}\bar{S}} du.$$

5. Focus on

$${}_s P_{tx}^{HD} = P(Y(t+s, x+s) = D | Y(t, x) = H).$$

Consider



Get

$$\frac{\partial {}_s P_{tx}^{HD}}{\partial s} = {}_s P_{tx}^{HH} \mu_{t+s, x+s} \quad (\text{contribution, route 1})$$

$$+ \int_0^s {}_u P_{tx}^{HH} \sigma_{t+u, x+u} {}_{s-u} P_{t+u, x+u}^{\bar{S}\bar{S}} \nu_{t+s, x+s, s-u} du. \quad (\text{contribution, route 2})$$

Integration implies

$${}_s P_{tx}^{HD} = \int_0^s {}_u P_{tx}^{HH} \mu_{t+u, x+u} du + \int_0^s {}_u P_{tx}^{HH} \sigma_{t+u, x+u} \left\{ \int_0^{s-u} {}_w P_{t+u, x+u}^{\bar{S}\bar{S}} \nu_{t+u+w, x+u+w, w} dw \right\} du.$$

APPENDIX II

The integral expression

$$\pi_{txd} = \exp - \int_0^{d/52} (\rho_{t+v,x+v,v} + \nu_{t+v,x+v,v}) dv$$

when

$$\log \rho_{t+z,x+z,z} = \mu + \alpha\sqrt{z} + \beta z + \gamma_1 t + \gamma_2 t\sqrt{z} + \gamma_3 tz + \theta x + \theta_2 x^2 + \theta_3 x^3 + \phi x\sqrt{z} + \psi xz$$

$$\log \nu_{t+z,x+z,z} = \mu + \gamma t + \theta x + \beta z + \sum_{j=1}^n \beta_j (z - z_j) +$$

and all three of  $t, x, z$  are measured in years, may be evaluated as follows.

1) To determine the contribution from  $\rho$ , write

$$\log \rho_{t+z,x+z,z} = a_{tx} + b_{tx}\sqrt{z} + c_{tx}z$$

where

$$a_{tx} = \mu + \gamma_1 t + \theta x + \theta_2 x^2 + \theta_3 x^3, \quad b_{tx} = \alpha + \gamma_2 t + \phi x, \quad c_{tx} = \beta + \gamma_3 t + \psi xz.$$

Then

$$J(t,x; z) = \int_0^z \rho_{t+v,x+v,v} dv = k_{tx} \int_0^z \exp \left\{ c_{tx} \left( \sqrt{v} + \frac{b_{tx}}{2c_{tx}} \right)^2 \right\} dv$$

reducing to

$$J(t,x; z) = [\exp(a_{tx}) \{ \exp(b_{tx}\sqrt{z} + c_{tx}z) - 1 \} - k_{tx} b_{tx} \int_{\frac{b_{tx}}{2c_{tx}}}^{\frac{b_{tx}}{2c_{tx}} + \sqrt{z}} \exp(c_{tx} s^2) ds] / c_{tx}$$

where

$$k_{tx} = \exp \left( a_{tx} - \frac{b_{tx}^2}{4c_{tx}} \right).$$

The definite integral is evaluated numerically by Simpson's rule. Since  $c_{tx} > 0 \forall (t, x)$ , it cannot be equated with the area under a standard normal variate (which in any event is evaluated numerically).

2) To determine the contribution from  $\nu$ : for

$$z_j < z \leq z_{j+1}, \quad j = 0, 1, \dots, n; \quad 0 = z_0 < z_1 < z_2 < \dots < z_{n+1} = \infty$$

let

$$\log \nu_{t+z,x+z,z} = A_{tx}^{(j)} + B^{(j)} z$$

where

$$A_{tx}^{(0)} = \mu + \gamma t + \theta x, \quad B^{(0)} = \beta; \quad A_{tx}^{(j)} = A_{tx}^{(j-1)} - \beta_j z_j, \quad B^{(j)} = B^{(j-1)} + \beta_j; \quad j = 1, 2, \dots, n.$$

Then for  $z_j < z \leq z_{j+1}$ , define

$$I_j(t, x; z) = \int_{z_j}^z \exp(A_{tx}^{(j)} + B^{(j)} v) dv = \exp A_{tx}^{(j)} \{ \exp(B^{(j)} z) - \exp(B^{(j)} z_j) \} / B^{(j)}$$

so that, for  $z_g < z \leq z_{g+1}$

$$I(t, x; z) = \int_0^z \nu_{t+v, x+v, v} dv = \sum_{j=0}^{g-1} I_j(t, x; z_j) + I_g(t, x; z).$$

The summation term does not come into play when  $g = 0$ .

3) Hence the required result

$$\pi_{txd} = \{ \exp -J(t, x; \frac{d}{52}) \} \{ \exp -I(t, x; \frac{d}{52}) \}.$$

\*\*\*\*\*

a) In the event that the expression for  $\log \rho$  takes the form

$$\begin{aligned} \log \rho_{t+z', x+z', z'} &= \mu + \alpha' \sqrt{z'} + \beta' z' + \gamma_1 t + \gamma_2 t \sqrt{z'} + \gamma_3 t z' + \theta x + \theta_2 x^2 + \theta_3 x^3 + \phi' x \sqrt{z'} + \psi' x z' \\ &= a_{tx} + b'_{tx} \sqrt{z'} + c'_{tx} z' \end{aligned}$$

in which  $z'$  is expressed in weeks, the change of scale

$$z = z'/w \text{ with inverse } z' = wz \text{ where } w = 52$$

implies that

$$\begin{aligned} \alpha &= \sqrt{w} \alpha', \quad \gamma_2 = \sqrt{w} \gamma_2', \quad \phi = \sqrt{w} \phi'; \quad b_{tx} = \sqrt{w} b'_{tx} \\ \beta &= w \beta', \quad \gamma_3 = w \gamma_3', \quad \psi = w \psi'; \quad c_{tx} = w c'_{tx} \end{aligned}$$

and

$$b_{tx} z = b'_{tx} z', \quad c_{tx} z = c'_{tx} z', \quad k_{tx} = \exp \left( a_{tx} - \frac{b'^2_{tx}}{4 c'_{tx}} \right).$$

b) In the event that the expression for  $\log \nu$  takes the form

$$\log \nu_{t+z, x+z, z} = \mu' + \gamma' t' + \theta x + \beta' z' + \sum_{j=1}^n \beta'_j (z' - z'_j)_+$$

where  $t' = 1, 2, 3, 4, 5$  denote consecutive quadrennia, and  $z'$  is in weeks, the transformation

$$t = 4t' - 3 \Leftrightarrow t' = (t + 3)/4$$

mapping consecutive quadrennia into a run of consecutive matching years 1, 2, ..., 20; coupled with the change of scale

$$z = z'/w \text{ with inverse } z' = wz \text{ where } w = 52,$$

implies that

$$\mu = \mu' + 3\gamma'/4; \quad \gamma = \gamma'/4; \quad \beta = w\beta', \quad \beta_j = w\beta'_j.$$



Table 3.1(a) Estimates  $\hat{\eta}_{zz}^A = \hat{\alpha}_z$  and t-statistics by duration  $z$ , for quadrennia (all ages  $x$ )

duration (weeks)	quadrennium				
	75-78	79-82	83-86	87-90	91-94
4	1.1943 (11.94)*	1.2711 (11.82)*	1.4127 (12.1)*	1.3104 (12.8)*	1.2169 (11.3)*
5	0.7065 (7.27)*	0.5973 (5.74)*	0.6622 (6.21)*	0.7521 (7.46)*	0.7424 (6.76)*
6	0.3986 (3.97)*	0.4759 (4.65)*	0.5438 (4.87)*	0.5057 (5.01)*	0.7454 (6.11)*
7	0.2198 (2.12)*	0.3750 (3.09)*	0.6446 (5.21)*	0.4512 (3.92)*	0.6766 (5.51)*
8	0.0650 (0.54)	0.1864 (1.34)	-0.0756 (-0.59)	0.2797 (2.26)*	0.2695 (2.04)*
9	0.0495 (0.36)	0.4593 (2.87)*	0.1515 (1.06)	0.3431 (2.62)*	0.2859 (1.68)
10	0.1416 (0.90)	-0.0675 (-0.37)	0.2883 (1.81)	0.2968 (1.90)†	0.2361 (1.25)
11-12	-0.0293 (-0.24)	0.0004 (0.00)	0.2002 (1.48)	0.2824 (2.21)*	-0.0794 (-0.62)
13-14	-0.1597 (-1.00)	0.2809 (1.78)	0.2979 (1.97)†	-0.0573 (-0.34)	0.1211 (0.72)
15-17	0.0919 (0.64)	0.1579 (1.03)	0.0175 (1.11)	-0.1857 (-1.07)	-0.4223 (-2.60)*
18-21	-0.0521 (-0.31)	-0.0954 (-0.52)	-0.2090 (-1.14)	-0.0991 (-0.60)	0.0187 (0.09)
22-25	0.0515 (0.23)	0.2199 (0.89)	0.0458 (0.23)	-0.4042 (-1.66)	-0.9493 (-3.60)*
26-29	-0.2615 (-1.03)	-0.2316 (-0.80)	0.1316 (0.46)	0.3326 (1.18)	-0.1442 (-0.60)
30-38	-0.4035 (-1.66)	0.4652 (1.64)	-0.3558 (-1.35)	-0.2823 (-1.23)	-0.1758 (-0.71)
39-51	-0.3819 (-1.11)	0.1488 (0.43)	-0.2254 (-0.93)	-0.0386 (-0.15)	-0.2736 (-1.02)
1-yr	-0.1617 (-0.75)	-0.1058 (-0.45)	0.0263 (0.12)	-0.9981 (-3.63)*	-0.1330 (-0.66)
2-11yrs	-0.3571 (-0.58)	0.0875 (0.26)	0.1441 (0.56)	-0.1483 (-0.50)	-0.3549 (-1.75)

(\* statistically significant difference from zero, † of marginal statistical significance)

Table 3.1(b) Estimates  $\hat{\eta}_{zz}^{13} = \hat{\alpha}_z$  and t-statistics by duration  $z$ , for quadrennia (all ages  $x$ )

duration (weeks)	quadrennium				
	75-78	79-82	83-86	87-90	91-94
13-14	0.8598 (4.20)*	1.1436 (5.83)*	1.2852 (7.91)*	0.8478 (4.39)*	1.1560 (6.35)*
15-17	0.3303 (2.34)*	0.5860 (3.63)*	0.5665 (3.82)*	0.4063 (2.38)*	0.2665 (1.71)
18-21	0.0358 (0.22)	0.1956 (1.02)	0.0248 (0.15)	-0.0142 (-0.09)	0.1199 (0.72)
22-25	0.1150 (0.59)	0.6281 (2.45)*	0.2030 (1.15)	-0.5101 (-2.34)*	-0.5193 (-2.20)*
26-29	-0.1894 (-0.75)	0.1009 (0.34)	0.0617 (0.27)	-0.0042 (-0.02)	0.0570 (0.29)
30-38	-0.3341 (-1.56)	0.8687 (3.16)*	-0.1088 (-0.49)	-0.3289 (-1.54)	-0.2086 (-1.11)
39-51	0.2814 (0.94)	0.2361 (0.78)	0.1934 (0.90)	-0.1729 (-0.77)	-0.5156 (-2.51)*
1- yr	0.2775 (1.41)	0.0015 (0.01)	0.0375 (0.21)	-0.7759 (-3.01)*	-0.4309 (-2.79)*
2-11 yrs	-0.4474 (-1.07)	-0.0408 (-0.14)	0.3591 (1.65)	-0.7601 (-3.62)*	-0.4355 (-2.77)*

(\* denotes statistically significant difference from zero)

Table 3.1(c) Estimates  $\hat{\eta}_{zz}^{26} = \hat{\alpha}_z$  and t-statistics by duration  $z$ , for quadrennia (all ages  $x$ )

duration (weeks)	quadrennium				
	75-78	79-82	83-86	87-90	91-94
26-29	0.3861 (1.62)	0.8504 (2.08)*	0.9084 (2.17)*	0.8706 (2.84)*	0.8936 (3.14)*
30-38	-0.0322 (-0.15)	0.7345 (2.61)*	0.2116 (0.64)	-0.0694 (-0.34)	0.3404 (1.52)
39-51	0.0418 (0.15)	0.4530 (1.17)	0.6085 (2.06)*	0.1514 (0.66)	-0.0424 (-0.18)
1- yr	0.5234 (2.55)*	0.7050 (2.64)*	0.4076 (1.65)	-0.5860 (-2.42)*	0.0775 (0.43)
2-11 yrs	0.2077 (0.58)	0.6767 (2.22)*	0.2834 (1.14)	-0.5710 (-2.89)*	-0.1099 (-0.60)

(\* denotes statistically significant difference from zero)

Table 3.3 Estimates  $\hat{\eta}_{z'}^4 = \hat{\alpha}_z + \hat{\beta}_{z'}$  and t-statistics, by duration  $z$  and age  $z'$ , 1979-82

duration (weeks)	age			
	18-29	30-34	35-49	50-65
4	1.1088 (10.40)*	1.5890 (10.64)*	1.3610 (12.69)*	1.1088 (10.40)*
5	0.4474 (4.35)*	0.9276 (6.20)*	0.6996 (6.71)*	0.4474 (4.35)*
6	0.3298 (3.28)*	0.8100 (5.51)*	0.5820 (5.60)*	0.3298 (3.28)*
7	0.2384 (2.05)*	0.7186 (4.55)*	0.4906 (4.02)*	0.2384 (2.05)*
8	0.0283 (0.21)	0.5085 (3.00)*	0.2805 (2.08)*	0.0283 (0.21)
9	0.3215 (2.12)*	0.8017 (4.26)*	0.5737 (3.73)*	0.3215 (2.12)*
10	-0.2127 (-1.22)	0.2675 (1.33)	0.0395 (0.22)	-0.2127 (-1.22)
11-12	-0.1351 (-1.04)	0.3452 (2.04)*	0.1171 (0.87)	-0.1351 (-1.04)
13-14	0.1640 (1.10)	0.6442 (3.50)*	0.4162 (2.68)*	0.1640 (1.10)
15-17	0.0329 (0.23)	0.5131 (2.75)*	0.2851 (1.93)	0.0329 (0.23)
18-21	-0.2103 (-1.21)	0.2700 (1.29)	0.0420 (0.24)	-0.2103 (-1.21)
22-25	0.1194 (0.52)	0.5997 (2.31)*	0.3716 (1.60)	0.1194 (0.52)
26-29	-0.3520 (-1.30)	0.1283 (0.44)	-0.0998 (-0.36)	-0.3520 (-1.30)
30-38	0.3870 (1.47)	0.8672 (2.99)*	0.6392 (2.39)*	0.2870 (1.47)
39-51	0.0303 (0.09)	0.5106 (1.51)	0.2825 (0.88)	0.0303 (0.09)
1-yr	-0.2501 (-1.14)	0.2302 (0.94)	0.0021 (0.01)	-0.2501 (-1.14)
2-11yrs	-0.0288 (-0.09)	0.4515 (1.38)	0.2234 (0.72)	-0.0288 (-0.09)

(\* denotes statistically significant difference from zero)

Table 3.4 Estimates  $\hat{\eta}_{x^j z}^{I3} = \hat{\alpha}_z + \hat{\beta}_j$ , and t-statistics, by duration  $z$  & age  $x^j$ , 1983-86

duration (weeks)	age						
	18-29	30-34	35-39	40-44	45-54	55-59	60-65
13-14	1.6756 (8.63)*	1.5195 (5.93)*	1.2743 (6.23)*	1.4164 (7.03)*	0.9913 (5.87)*	1.3539 (7.08)*	1.6756 (8.63)*
15-17	0.9475 (5.11)*	0.7915 (3.30)*	0.5463 (2.76)*	0.6884 (3.55)*	0.2633 (1.66)	0.6259 (3.47)*	0.9475 (5.11)*
18-21	0.3603 (1.79)	0.2042 (0.85)	-0.0410 (-0.19)	0.1011 (0.50)	-0.3240 (-1.77)	0.0386 (0.20)	0.3603 (1.79)
22-25	0.5966 (2.90)*	0.4406 (1.67)	0.1954 (0.93)	0.3375 (1.55)	-0.0876 (-0.48)	0.2750 (1.39)	0.5966 (2.90)*
26-29	0.3479 (1.44)	0.1919 (0.68)	-0.0533 (-0.21)	0.0888 (0.35)	-0.3363 (-1.43)	0.0263 (0.11)	0.3479 (1.44)
30-38	0.2767 (1.18)	0.1206 (0.42)	-0.1246 (-0.53)	0.0175 (0.07)	-0.4076 (-1.86)	-0.0450 (-0.18)	0.2767 (1.18)
39-51	0.5699 (2.51)*	0.4138 (1.45)	0.1686 (0.69)	0.3107 (1.27)	-0.1144 (-0.54)	0.2482 (1.07)	0.5699 (2.51)*
1- yr	0.3659 (1.86)	0.2099 (0.84)	-0.0353 (-0.16)	0.1068 (0.48)	-0.3184 (-1.72)	0.0443 (0.22)	0.3659 (1.86)
2-11 yrs	0.6883 (2.93)*	0.5323 (1.79)	0.2871 (1.15)	0.4292 (1.63)	0.0041 (0.02)	0.3667 (1.74)	0.6883 (2.93)*

(\* denotes statistically significant difference from zero)

Table 4.1 Estimates and t-statistics, modelling  $\rho$  for specified quadrennia

	75-94	75-78	79-82	83-86	87-90	91-94
$\mu$	2.7618 6.58	5.7524 30.45	3.7310 4.74	3.0168 4.10	2.8511 3.31	*
$\alpha$	-9.9601E-1 -19.43	-1.2064 -10.88	-1.1325 -10.22	-1.2422 -24.19	-1.4525 -13.56	-1.6170 -12.76
$\beta$	3.2393E-2 7.15	6.3075E-2 5.48	4.7333E-2 5.00	3.3063E-2 27.41	5.9901E-2 7.69	6.6951E-2 8.83
$\theta$	1.6973E-1 5.64	-3.0086E-2 -5.60	1.2275E-1 2.17	1.7735E-1 3.30	2.0786E-1 3.39	4.1711E-1 36.30
$\phi$	5.7027E-3 5.79	8.9599E-3 3.74	7.0503E-3 3.11	4.8709E-3 4.77	7.3811E-3 3.34	9.5578E-3 3.73
$\psi$	-3.4675E-4 -4.05	-9.6510E-4 -3.70	-5.3929E-4 -2.70	* *	-5.3417E-4 -3.17	-6.2393E-4 -3.87
$\theta_2$	-4.4399E-3 -6.33	* *	-3.5295E-3 -2.66	-4.4978E-3 -3.55	-5.2586E-3 -3.69	-9.8314E-3 -23.48
$\theta_3$	2.9982E-5 5.66	-1.9764E-6 -2.52	2.4205E-5 2.39	2.9881E-5 3.09	3.4947E-5 3.25	6.6818E-5 17.22
$\gamma_1$	4.2984E-2 15.52	* *	* *	* *	* *	* *
$\gamma_2$	-2.6384E-2 -16.04	* *	* *	* *	* *	* *
$\gamma_3$	1.3679E-3 20.21	* *	* *	* *	* *	* *

Table 4.2 Parameter estimates and t-statistics, modelling  $\nu$  for specified quadrennia

	75-94	75-78	79-82	83-86	87-90	91-94
$\mu$	-4.1676 -18.59	-3.3758 -8.35	-4.8298 -8.74	-4.0984 -11.44	-5.4552 -9.37	-5.9343 -7.81
$\theta$	2.3393E-2 8.78	1.8262E-2 2.62	2.6243E-2 3.76	2.6055E-2 4.37	2.6239E-2 4.69	2.1733E-2 3.41
$\beta$	1.0295E-1 5.00	3.4399E-2 1.94	1.4406E-1 2.91	2.6306E-2 2.64	1.5081E-1 2.69	2.2238E-1 3.30
$\beta_j: z_j=10.5$	-8.8884E-1 -3.56	* *	-1.2731E-1 -2.12	* *	-1.2186E-1 -1.90	-2.3039E-1 -3.41
$\beta_j: z_j=15$	* *	-4.4799E-2 -2.43	* *	* *	* *	* *
$\beta_j: z_j=28$	-4.3287E-2 -3.98	* *	-6.6398E-2 2.43	6.8885E-2 -3.71	-7.1999E-2 -3.26	* *
$\beta_j: z_j=45.5$	2.1700E-2 4.03	* *	4.2750E-2 2.96	3.7041E-2 3.22	3.5881E-2 3.39	* *
$\beta_j: z_j=182$	7.8231E-3 9.62	1.2460E-2 5.03	8.3114E-3 3.67	5.2665E-3 2.88	7.6188E-3 4.82	8.0105E-3 5.72
$\gamma$	-1.4287E-1 -8.16	* *	* *	* *	* *	* *

Table 5.1 Inceptions: parameter estimates, standard errors, t-statistics

	<i>targeting <math>\sigma</math></i>				<i>targeting <math>\tau</math></i>			
	<i>DP1</i>	<i>DP4</i>	<i>DP13</i>	<i>DP26</i>	<i>DP1</i>	<i>DP4</i>	<i>DP13</i>	<i>DP26</i>
$\alpha_0$	-5.840	-4.604	-2.303	3.315	-5.428	-2.672	*	3.617
(s.e.)	6.165E-1	7.781E-1	1.181	4.782E-1	6.045E-1	7.915E-1		1.774
t-stat.	-9.47	-5.92	-1.95	6.93	-8.98	-3.38		2.04
$\alpha_1$	2.648E-1	1.448	3.323E-1	1.078	-5.173E-1	5.530E-1	2.280E-1	7.005E-1
(s.e.)	3.062E-1	1.474E-1	7.414E-2	1.469E-1	2.973E-1	1.461E-1	7.110E-2	8.667E-2
t-stat.	0.86	9.82	4.48	7.34	-1.74	3.79	3.21	8.08
$\alpha_2$	*	7.773E-1	*	*	*	7.389E-1	*	*
(s.e.)		1.514E-1				1.503E-1		
t-stat.		5.13				4.92		
$\alpha_3$	-2.293	-1.245	-3.120E-1	-2.826	-2.301	-1.262	-3.261E-1	-5.962E-1
(s.e.)	2.520E-1	3.731E-1	1.059E-1	5.448E-1	2.432E-1	3.703E-1	1.013E-1	1.202E-1
t-stat.	-9.10	-3.34	-2.95	-5.19	-9.46	-3.41	-3.22	-4.96
$\alpha_4$	-3.019E-1	-5.766E-1	*	*	-3.384E-1	-5.777E-1	*	*
(s.e.)	8.985E-2	1.583E-1			8.659E-2	1.571E-1		
t-stat.	-3.36	-3.64			-3.91	-3.68		
$\alpha_5$	1.564	6.860E-1	*	1.916	1.570	6.947E-1	*	*
(s.e.)	2.064E-1	3.061E-1		4.478E-1	1.992E-1	3.038E-1		
t-stat.	7.58	2.24		4.28	7.88	2.29		
$\beta_1$	4.346E-1	3.231E-1	2.276E-1	-2.121E-1	2.378E-1	-1.713E-1	-4.582E-1	-8.180E-1
(s.e.)	4.418E-2	5.680E-2	8.353E-2	2.004E-2	4.325E-2	5.757E-2	9.750E-3	1.207E-1
t-stat.	9.84	5.69	2.72	-10.58	5.50	-2.98	-46.99	-6.78
$\beta_2$	-1.110E-2	-9.334E-3	-8.063E-3	1.947E-3	-5.512E-3	4.062E-3	1.038E-2	1.879E-2
(s.e.)	1.025E-3	1.333E-3	1.910E-3	2.062E-4	1.001E-3	1.347E-3	4.016E-4	2.663E-3
t-stat.	-10.83	-7.00	-4.22	9.44	-5.51	3.02	25.85	7.06
$\beta_3$	8.441E-5	7.707E-5	7.124E-5	*	4.170E-5	-2.352E-5	-6.573E-5	-1.250E-4
(s.e.)	7.686E-6	1.008E-5	1.414E-5		7.505E-6	1.017E-5	4.062E-6	1.910E-5
t-stat.	10.98	7.60	5.04		5.56	-2.31	-16.18	-6.54
$\gamma_{11}$	4.697E-2	-1.632E-2	*	*	6.051E-2	-4.199E-3	*	*
(s.e.)	1.362E-2	2.286E-3			1.324E-2	2.263E-3		
t-stat.	3.45	-7.14			4.57	-1.86		
$\gamma_{12}$	-6.286E-4	*	*	*	-6.737E-4	*	*	*
(s.e.)	1.493E-4				1.452E-4			
t-stat.	-4.21				-4.64			
$\gamma_{21}$	8.077E-3	*	*	*	8.091E-3	*	*	*
(s.e.)	1.844E-3				1.778E-3			
t-stat.	4.38				4.55			
$\tilde{\phi}$	5.439	1.906	1.678	1.480	5.068	1.878	1.547	1.495

Table 5.2 Scaled deviance residuals by age and period

	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
18-24	0.98	1.46	0.18	-0.08	-0.31	-0.15	-1.12	-0.38	0.40	-0.09
25-29	0.54	0.56	-0.61	1.60	-0.36	0.51	-0.94	-0.12	-0.19	0.28
30-34	-0.56	-0.79	-1.53	-0.32	-0.72	0.50	-1.03	0.63	0.96	-0.01
35-39	-0.03	1.08	-0.56	-0.31	0.02	0.54	-1.57	-0.64	0.87	-0.23
40-44	0.53	-0.69	-1.17	0.29	0.58	-0.71	-0.36	-1.19	1.05	-0.07
45-49	0.82	0.53	-0.92	0.37	1.63	-0.09	-0.86	-1.11	-0.16	-1.20
50-54	-0.47	0.36	1.23	0.61	1.16	1.12	-2.25	-2.54	0.66	-1.03
55-59	-1.00	0.04	-1.28	0.83	-0.08	2.27	0.06	0.13	1.72	0.13
60-64	0.13	-0.78	-0.81	0.80	0.80	0.19	-0.29	-0.72	-0.15	-0.18

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
18-24	-0.31	-1.60	-1.81	-1.00	-0.29	-0.53	-0.55	-1.55	0.51	-0.39
25-29	0.76	-0.39	-1.29	0.70	-0.03	-2.45	-0.65	-1.15	-0.14	-0.34
30-34	1.46	0.92	2.33	1.59	1.41	0.48	0.54	-0.23	2.45	0.11
35-39	0.40	0.04	-1.74	-0.28	-0.15	0.87	0.11	-0.53	2.96	-0.56
40-44	0.70	0.24	-0.57	-0.13	1.04	-3.38	-1.50	-1.53	1.78	-0.94
45-49	1.20	0.97	0.47	-1.33	0.14	-1.90	-0.66	-0.22	0.49	-2.21
50-54	1.01	0.35	-0.66	0.97	0.97	-0.47	1.63	-0.22	1.36	-0.46
55-59	-1.06	-0.75	0.50	0.81	1.10	1.31	1.12	-0.37	0.44	0.71
60-64	-0.20	-0.98	-1.09	-0.97	0.44	-1.02	-0.81	-0.30	0.70	0.07

Deferred period 1 week

	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
18-24	-0.66	-0.64	0.18	1.28	1.37	-0.76	-0.28	0.13	0.63	0.70
25-29	-0.38	-1.03	-0.95	0.60	-0.13	0.18	-0.09	-0.33	0.38	-0.12
30-34	-0.15	1.30	-0.26	-0.50	-0.08	-0.06	-1.71	-1.76	1.59	1.08
35-39	0.72	0.67	2.03	0.27	1.36	-0.75	0.38	-0.23	0.22	1.35
40-44	1.20	-1.64	-0.60	0.98	0.01	0.56	-0.98	0.61	-2.27	1.08
45-49	-0.48	-1.24	1.55	-1.02	-0.36	-0.60	-0.19	1.57	-0.22	0.65
50-54	-0.18	0.05	0.96	-1.43	-0.50	0.93	-0.78	-1.43	0.77	-0.30
55-59	-0.78	-0.03	-0.65	0.72	0.05	0.25	0.51	0.06	-0.28	0.16
60-64	0.72	-0.66	-0.02	0.32	0.62	-0.43	-1.67	-1.25	-1.09	2.73

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
18-24	0.15	-2.15	-1.08	0.09	0.54	-0.73	0.92	2.13	-0.31	-1.58
25-29	0.50	-0.52	-0.52	-1.00	-1.82	-0.89	1.00	0.85	-0.57	-0.44
30-34	0.47	-1.31	0.54	0.34	0.80	-0.29	0.25	1.41	1.95	-0.65
35-39	-0.09	-0.40	-0.25	2.31	1.39	0.26	-0.40	1.86	-2.03	0.61
40-44	-0.06	-2.05	-1.83	1.12	-2.05	-1.19	-1.45	-0.13	-1.49	0.45
45-49	-0.95	0.50	-0.37	-0.93	-0.13	0.04	0.47	-1.60	-0.03	1.48
50-54	-0.12	-0.10	0.13	-0.05	-0.71	-0.64	0.94	0.54	0.71	-0.41
55-59	1.04	0.80	1.81	1.18	-0.33	0.57	2.07	-0.70	0.98	-0.32
60-64	0.03	-1.24	1.91	-1.65	0.38	-0.43	-1.84	-0.23	-0.16	-1.60

Deferred period 4 weeks

Table 5.2 (continued) Scaled deviance residuals by age and period

	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
18-24	-0.80	-0.08	-0.56	-0.79	0.63	-0.61	-2.08	-2.15	0.24	-0.29
25-29	0.20	-1.21	-0.05	-0.93	0.85	-0.56	0.69	-0.48	-0.63	1.34
30-34	-1.14	0.38	-1.33	-0.29	0.07	-0.41	1.15	-1.24	-0.39	-0.90
35-39	0.48	1.00	0.57	0.10	0.34	1.06	-1.02	-1.20	-0.72	0.56
40-44	0.87	0.65	-1.10	0.96	0.12	-1.24	0.27	-1.17	-0.11	0.93
45-49	0.87	-0.53	0.59	-0.14	-1.59	0.52	-2.18	-2.23	-0.17	0.62
50-54	-0.24	-1.22	0.89	0.43	1.12	-0.28	-0.35	-1.42	1.50	1.63
55-59	0.85	-0.67	-0.39	0.43	2.40	-0.10	-1.21	-1.02	0.90	-0.61
60-64	0.32	-0.32	1.34	0.45	1.53	1.33	-1.59	-0.81	-1.73	-0.53

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
18-24	-1.35	-0.20	0.90	0.47	1.73	-0.58	-0.42	-1.35	-1.17	-0.39
25-29	0.01	1.17	0.72	-0.03	0.01	0.25	1.10	0.09	-0.31	0.53
30-34	1.28	1.21	-1.32	0.76	0.42	0.34	-0.37	-0.48	1.14	0.53
35-39	0.52	0.09	0.01	1.04	-0.71	2.47	0.04	2.11	-0.99	1.07
40-44	1.30	0.42	0.15	-0.37	-0.51	-0.93	-0.76	-0.56	0.51	-1.84
45-49	-1.86	0.43	-1.25	-0.52	-0.62	-1.41	-0.36	-0.41	0.06	-2.09
50-54	1.29	1.15	0.17	-0.81	-1.44	0.94	-1.50	-0.87	1.21	2.40
55-59	1.00	1.60	0.77	-0.29	0.17	-0.50	0.65	1.89	0.43	0.67
60-64	0.84	-1.17	-0.42	-2.06	1.02	-1.85	-0.71	-0.22	0.28	-0.43

Deferred period 13 weeks

	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984
18-24	0.03	-1.59	-0.36	-1.21	0.11	-0.57	-1.20	-1.12	-0.94	-0.99
25-29	-0.46	0.69	1.88	-0.37	0.50	1.52	0.95	-1.75	0.05	-0.07
30-34	0.44	-0.83	-0.38	0.26	0.06	1.40	-0.87	-1.52	-0.49	-0.29
35-39	1.69	-0.06	-0.61	0.86	-0.73	1.12	0.28	-1.39	0.69	2.13
40-44	-1.25	0.58	0.90	1.01	0.36	-0.50	0.08	-0.56	-0.66	0.47
45-49	0.25	-1.16	-0.84	0.09	-1.94	-1.21	-0.80	-0.83	-0.51	-0.42
50-54	0.10	-0.02	-0.77	1.21	-0.35	-0.75	-1.60	-2.02	-0.65	-0.46
55-59	-0.03	0.79	1.18	-1.27	0.61	1.08	-0.23	0.65	0.01	0.95
60-64	0.12	0.33	1.08	0.20	0.45	1.15	0.46	-0.63	0.03	-1.70

	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994
18-24	-0.99	-1.02	-1.08	-1.08	-1.06	-1.03	0.32	-0.29	0.33	-1.54
25-29	-0.76	-0.72	-1.75	0.37	0.13	0.73	-0.33	1.81	-0.66	-1.97
30-34	0.23	0.58	1.33	0.11	0.92	0.01	-0.11	0.55	0.93	1.01
35-39	0.70	-0.41	0.08	1.28	-0.54	0.12	0.73	0.19	-0.31	0.67
40-44	-0.18	-1.46	1.16	-0.79	-1.39	0.29	1.73	0.40	0.07	0.96
45-49	0.48	-1.35	-0.44	-2.12	2.04	-1.35	-0.08	0.14	-0.68	1.24
50-54	-0.16	-0.25	-1.89	-1.44	-0.22	-0.42	1.96	-0.44	-0.29	0.07
55-59	1.82	3.21	1.02	1.02	-0.35	-0.30	1.22	0.39	1.25	-1.74
60-64	0.25	-0.35	1.58	-1.69	-1.64	-1.91	-0.44	-1.49	1.43	-0.65

Deferred period 26 weeks



Table 6.1 Values of  $(\hat{\sigma}_{tx}^* - \hat{\sigma}_{tx})/\hat{\sigma}_{tx}$  expressed as percentages

	1975	1977	1979	1981	1983	1985	1987	1989	1991	1993	1994
18-24	1.8	0.0	-2.7	-5.5	-8.0	-10.1	-11.7	-12.9	-14.0	-15.4	-16.3
25-29	2.0	2.1	1.2	0.2	-0.7	-1.0	-0.8	-0.3	0.4	0.9	0.9
30-34	0.5	1.2	1.0	0.6	0.5	0.7	1.5	2.6	3.8	4.7	5.0
35-39	-1.5	-0.6	-0.6	-0.9	-1.0	-0.8	-0.1	0.9	1.9	2.5	2.6
40-44	-2.2	-1.2	-1.2	-1.6	-1.9	-1.9	-1.6	-1.0	-0.5	-0.5	-0.8
45-49	-1.0	0.0	-0.0	-0.5	-1.0	-1.2	-1.1	-0.9	-0.9	-1.4	-2.0
50-54	0.9	2.1	2.1	1.7	1.2	0.9	0.9	0.9	0.7	-0.1	-0.9
55-59	0.8	2.3	2.6	2.4	2.1	2.0	2.1	2.1	1.9	1.0	0.2
60-64	-4.5	-2.6	-1.9	-1.7	-1.6	-1.4	-1.0	-0.7	-0.7	-1.3	-2.0

Deferred period 1 week

	1975	1977	1979	1981	1983	1985	1987	1989	1991	1993	1994
18-24	1.7	-1.3	-4.1	-6.6	-8.8	-10.8	-12.5	-13.9	-15.3	-16.4	-16.7
25-29	5.0	4.1	3.3	2.8	2.6	2.7	3.1	3.7	4.5	5.7	6.4
30-34	2.8	2.7	2.6	2.8	3.2	4.0	5.0	6.1	7.5	9.1	10.2
35-39	-1.6	-1.6	-1.6	-1.4	-1.0	-0.4	0.3	1.1	2.1	3.2	3.9
40-44	-3.5	-3.6	-3.7	-3.8	-3.8	-3.6	-3.3	-3.1	-2.9	-2.6	-2.3
45-49	-1.4	-1.5	-1.8	-2.0	-2.2	-2.3	-2.4	-2.7	-3.0	-3.4	-3.5
50-54	2.7	2.8	2.7	2.5	2.4	2.2	1.9	1.4	0.8	0.1	-0.2
55-59	3.1	3.6	3.9	4.1	4.3	4.4	4.4	4.1	3.6	2.9	2.6
60-64	-7.7	-6.5	-5.6	-4.7	-4.0	-3.3	-2.8	-2.6	-2.6	-2.8	-2.9

Deferred period 4 weeks

	1975	1977	1979	1981	1983	1985	1987	1989	1991	1993	1994
18-24	-28.4	-28.9	-28.6	-27.6	-25.9	-23.5	-20.3	-16.4	-11.8	-6.2	-3.1
25-29	-5.2	-6.1	-5.9	-4.8	-2.6	0.5	4.7	9.9	16.3	23.9	28.2
30-34	2.6	1.1	0.8	1.5	3.2	5.9	9.5	14.1	19.9	26.7	30.6
35-39	0.8	-1.2	-2.1	-2.2	-1.3	0.3	2.8	6.0	10.0	14.8	17.4
40-44	-1.5	-3.9	-5.3	-6.0	-6.0	-5.3	-4.0	-2.1	0.2	3.1	4.7
45-49	1.6	-1.3	-3.3	-4.5	-5.1	-5.2	-4.8	-4.0	-2.8	-1.3	-0.4
50-54	9.5	6.1	3.6	1.8	0.6	-0.1	-0.4	-0.4	-0.1	0.4	0.7
55-59	15.8	12.1	9.1	6.9	5.2	4.0	3.1	2.5	2.1	1.8	1.7
60-64	9.2	5.5	2.6	0.2	-1.6	-3.1	-4.3	-5.3	-6.1	-6.9	-7.2

Deferred period 13 weeks

	1975	1977	1979	1981	1983	1985	1987	1989	1991	1993	1994
18-24	-30.2	-50.0	-45.8	-40.0	-40.3	-44.4	-46.0	-40.5	-27.8	-24.8	-38.1
25-29	22.9	-13.3	-7.4	1.3	-0.2	-8.0	-11.4	-2.9	17.2	21.7	-0.0
30-34	46.2	2.6	8.8	18.2	15.7	5.9	1.3	10.1	31.9	35.7	11.0
35-39	42.8	0.2	6.1	15.0	12.1	2.1	-3.0	4.6	24.1	26.5	2.8
40-44	31.0	-7.7	-2.0	6.3	3.5	-6.0	-11.1	-4.7	12.1	13.1	-8.5
45-49	24.8	-11.5	-5.6	2.6	-0.0	-9.3	-14.5	-8.7	6.8	6.9	-14.0
50-54	26.3	-9.8	-3.3	5.4	2.9	-6.6	-12.1	-6.4	8.9	8.3	-13.1
55-59	30.5	-6.4	0.8	10.2	7.7	-2.2	-8.0	-2.3	13.3	12.3	-10.2
60-64	26.8	-8.7	-1.6	7.8	5.4	-4.3	-10.1	-4.7	10.3	9.0	-13.0



Deferred period 26 weeks

Table 6.2 Inceptions: 1991-94 quadrennial experience, parameter details



	targeting $\sigma$				targeting $\tau$			
	DP1	DP4	DP13	DP26	DP1	DP4	DP13	DP26
$\beta_0$	-1.507E-1	-3.030E-1	-4.624E-1	*	-1.760	-3.718	-5.285	-5.307
(s.e.)	6.282E-2	6.945E-2	7.106E-2		5.899E-2	4.868E-2	1.021E-1	7.612E-2
t-stat.	-2.40	-4.36	-6.51		-29.8	-76.4	-51.8	-69.7
$\beta_1$	-5.105E-1	-2.575	-3.104	-3.539	8.869E-1	7.440E-1	1.152	9.341E-1
(s.e.)	2.226E-1	3.866E-1	3.081E-1	5.382E-1	1.107E-1	3.776E-1	1.488E-1	5.279E-1
t-stat.	-2.29	-6.66	-10.1	-6.58	8.01	1.97	7.74	1.77
$\beta_2$	-1.256	-1.993	*	-3.111	-4.412E-1	*	2.334	*
(s.e.)	2.608E-1	2.505E-1		3.416E-1	2.309E-1	*	8.959E-1	*
t-stat.	-4.81	-7.96		-9.11	-1.91	*	2.60	*
$\beta_3$	1.123	8.971	5.379	1.233E+1	*	4.006	*	5.416
(s.e.)	5.193E-1	1.857	7.941	2.713		1.818		2.608
t-stat.	2.16	4.83	6.77	4.55		2.20		2.08
$\beta_4$	*	*	-4.060	*	*	*	-2.782	*
(s.e.)			6.094E-1			*	1.407	
t-stat.			-6.66			*	-1.98	
$\beta_5$	*	-7.305	*	-9.124	*	-5.381	*	-6.480
(s.e.)		2.038		3.073		1.994		2.965
t-stat.		-3.58		-2.97		-2.70		-2.19
$\bar{\phi}$	2.528	1.488	1.099	1.519	2.457	1.430	1.124	1.490

Table 6.3 Values of  $(\hat{\sigma}_x^* - \hat{\sigma}_x)/\hat{\sigma}_x$  expressed as percentages: 1991-94 quadrennium



age	DP1	DP4	DP13	DP26
18	-16.6	2.1	794	5.2
20	-19.7	-1.6	219	-4.8
22	-17.9	-3.1	65.7	-9.1
24	-13.0	-2.3	14.6	-8.6
26	-6.5	-0.2	-3.3	-4.9
28	-0.0	2.0	-7.8	0.1
30	5.2	3.3	-6.5	4.5
32	8.1	3.2	-3.4	7.1
34	8.5	1.9	-0.7	7.5
36	7.0	-0.0	0.6	6.1
38	4.2	-1.9	0.5	3.9
40	1.0	-3.1	-0.4	1.9
42	-1.7	-3.2	-1.2	0.8
44	-3.7	-2.2	-1.3	0.8
46	-4.6	-0.5	-0.6	1.6
48	-4.4	1.4	1.0	2.8
50	-3.3	3.0	3.0	3.7
52	-1.3	3.4	4.6	3.5
54	1.0	2.3	4.8	1.9
56	3.3	0.0	2.8	-0.7
58	4.9	-2.7	-2.4	-3.1
60	5.4	-4.3	-10.8	-3.2
62	4.2	-2.2	-22.1	3.0
64	0.8	7.8	-35.1	23.0





Log predicted inception intensities vs period, by age, DP1  
Figure 5.1(a)



Log predicted inception intensities vs period, by age, DP4  
Figure 5.1(b)



Log predicted inception intensities vs period, by age, DP13  
Figure 5.1(c)



Log predicted inception intensities vs period, by age, DP26  
 Figure 5.1(d)



DEPARTMENT OF ACTUARIAL SCIENCE AND STATISTICS

Actuarial Research Papers since 1992

32. England P.D. and Verrall R.J. Dynamic Estimation for Models of Excess Mortality. January 1992. 19 pages. ISBN 1 874 770 32 8
33. Verrall R.J. A State Space Formulation of Whittaker-Henderson Graduation, with Extensions. January 1992. 13 pages. ISBN 1 874 770 33 6
34. Verrall R.J. Graduation by Dynamic Regression Methods. January 1992. 37 pages. ISBN 1 874 770 34 4
35. Gerrard R.G. and Haberman S. Stability of Pension Systems when Gains/Losses are Amortized and Rates of Return are Autoregressive. March 1992. 12 pages. ISBN 1 874 770 35 2
36. Haberman S. HIV, AIDS and the Approximate Calculation of Life Insurance Functions, Annuities and Net Premiums. April 1992. 28 pages. ISBN 1 874 770 36 0
37. Cooper D.R. Savings and Loans: An Historical Perspective. May 1992. 29 pages. ISBN 1 874 770 37 9
38. Verrall R.J. Dynamic Bayes Estimation for AIDS Delay Tables. May 1992. 16 pages. ISBN 1 874 770 38 7
39. Kaye G.D. Data Sources Relating to Life Assurance Expenses. May 1992. 39 pages. Presented to the Staple Inn Actuarial Society and Royal Statistical Society - February 1992. ISBN 1 874 770 39 5
40. Renshaw A.E., and Haberman S. On the Graduation Associated with a Multiple State Model for Permanent Health Insurance. May 1992. ISBN 1 874 770 40 9
41. England P.D. Statistical Modelling of Excess Mortality Number 3. June 1992. 163 pages. ISBN 1 874 770 41 7
42. Bloomfield D.S.F. and Haberman S. Male Social Class Mortality Differences Around 1981: An Extension to Include Childhood Ages. 21 pages. June 1992. ISBN 1 874 770 42 5
43. Berg M.P and Haberman S. Trend Analysis and Prediction Procedures for Time Nonhomogeneous Claim Processes. 33 pages. June 1992. ISBN 1 874 770 43 3
44. Booth P.M. The Single Market for Insurance, Free Capital Movements and attempts to Fix Exchange Rates. October 1992. 28 pages. ISBN 1 874 770 44 1
45. Verrall R.J. Chain Ladder with Varying Run-off Evolutions. February 1993. 15 pages. ISBN 1 874 770 45 X

46. Gavin J., Haberman S. and Verrall R.J. Moving Weighted Average Graduation using Kernel Estimation. November 1992. 14 pages. ISBN 1 874 770 46 8
47. Gavin J., Haberman S. and Verrall R.J. On the Choice of Bandwidth for Kernel Graduation. November 1992. 21 pages. ISBN 1 874 770 47 6
48. S. Haberman. Pension Funding with Time Delays and the Optimal Spread Period. May 1993. 13 pages. ISBN 1 874 770 48 4
49. S. Haberman. Stochastic Investment Returns and the Present Value of Future Contributions in a Defined Benefit Pension Scheme. May 1993. 22 pages. ISBN 1 874 770 49 2
50. A. Zimbidis and S. Haberman. Delay, Feedback and Variability of Pension Contributions and Fund Levels. May 1993. 19 pages. ISBN 1 874 770 50 6
51. S. Haberman. Pension Funding: The Effect of Changing The Frequency Valuations. June 1993. 19 pages. ISBN 1 874 770 51 4
52. S Haberman. HIV, AIDS Markov Chains and PHI. June 1993. 15 pages. ISBN 1 874 770 52 2
53. S Haberman. A Consideration of Pension Credit and Termination Insurance. June 1993. 22 pages. ISBN 1 874 770 53 0
54. M Z Khorasanee. Survey of Actuarial Practice in the Funding of UK Defined Benefit Pension Schemes. July 1993. 19 pages. ISBN 1 874 770 54 9
55. P M Booth, R G Chadburn and A S K Ong. A Utility Maximisation Approach to Asset Allocation. September 1993. 40 pages. ISBN 1 874 770 55 7
56. R G Chadburn. Bias in Select Mortality Investigations. August 1993. 62 pages. ISBN 1 874 770 56 5
57. M Z Khorasanee. A Comparison of Pension Funding Strategies by means of Simulations for a Model Scheme. August 1993. 43 pages. ISBN 1 874 770 57 3
58. A E Renshaw, P Hatzopolous and S Haberman. Recent Mortality Trends in Male Assured Lives. June 1993. 23 pages. ISBN 1 874 770 58 1
59. E Pitacco. Disability Risk Models: Towards A Unifying Approach. September 1993. 33 pages. ISBN 1 874 770 59 X
60. M Boskov and R J Verrall. Premium Rating by Geographic Area Using Spatial Models. September 1993. 14 pages. ISBN 1 874 770 60 3
61. R G Chadburn. Managing Mutual Life Offices: The Use of an Expense Ratio in New Business Decision Making and Expense Control. October 1993. 21 pages. ISBN 1 874 770 61 1
62. Haberman S. Pension Funding Modelling and Stochastic Investment Returns. 56 pages. March 1994. ISBN 1 874 770 62 X
63. Renshaw A E and Verrall R J. The Stochastic Model Underlying the Chain-Ladder Technique. 25 pages. April 1994. ISBN 1 874 770 63 8



64. Haberman S and Sung J-H. Dynamic Approaches to Pension Funding. 22 pages. April 1994.  
ISBN 1 874 770 64 6
65. Renshaw A.E. On the Second Moment Properties and the Implementation of Certain GLIM Based Stochastic Claims Reserving Models. 36 pages. September 1994. ISBN 1 874 770 65 4
66. Booth P.M., J.N. Allan, and J.W. Jang. An Evaluation of the UK Life Insurance Mismatching Test. September 1994. ISBN 1 874 770 66 2
67. Booth P.M. and Stroinski K. Insurance and Investment Markets in Poland. September 1994. 35 pages. ISBN 1 874 770 67 0
68. Ong A. A Stochastic Model for Treasury-Bills: An Extension to Wilkie's Model. September 1994. 12 pages. ISBN 1 874 770 68 9
69. Bloomfield D.S.F. Moving on from Undergraduate Exams to Professional Exams: Actuaries. November 1994. 22 pages. ISBN 1 874 770 69 7
70. Huber P. A Review of Wilkie's Stochastic Investment Model. January 1995. 22 pages. ISBN 1 874 770 70 0
71. Renshaw A.E. On the Graduation of 'Amounts'. January 1995. 24 pages. ISBN 1 874 770 71 9
72. Renshaw A.E. Claims Reserving by Joint Modelling. December 1994. 26 pages. ISBN 1 874 770 72 7
73. Renshaw A.E. Graduation and Generalised Linear Models: An Overview. February 1995. 40 pages. ISBN 1 874 770 73 5
74. Khorasane M.Z. Simulation of Investment Returns for a Money Purchase Fund. June 1995. 20 pages. ISBN 1 874 770 74 3
75. Owadally M.I. and Haberman S. Finite-time Pension Fund Dynamics with Random Rates of Return. June 1995. 28 pages. ISBN 1 874 770 75 1
76. Owadally M.I. and Haberman S. Stochastic Investment Modelling and Optimal Funding Strategies. June 1995. 25 pages. ISBN 1 874 770 76 X
77. Khorasane M.Z. Applying the Defined Benefit Principle to a Defined Contribution Scheme. August 1995. 30 pages. ISBN 1 874 770 77 8
78. Sebastiani P. and Settini R. Experimental Design for Non-Linear Problems. September 1995. 13 pages. ISBN 1 874 770 78 6
79. Verrall R.J. Whittaker Graduation and Parametric State Space Models. November 1995. 23 pages. ISBN 1 874 770 79 4
80. Verrall R.J. Claims Reserving and Generalised Additive Models. November 1995. 17 pages. ISBN 1 874 770 80 8
81. Nelder J.A. and Verrall R.J. Credibility Theory and Generalized Linear Models. November 1995. 15 pages. ISBN 1 874 770 81 6

82. Renshaw A.E., Haberman S. and Hatzopoulos P. On The Duality of Assumptions Underpinning The Construction of Life Tables. December 1995. 17 Pages. ISBN 1 874 770 82 4
83. Chadburn R.G. Use of a Parametric Risk Measure in Assessing Risk Based Capital and Insolvency Constraints for With Profits Life Insurance. March 1996. 17 Pages. ISBN 1 874 770 84 0
84. Haberman S. Landmarks in the History of Actuarial Science (up to 1919). March 1996. 62 Pages. ISBN 1 874 770 85 9
85. Renshaw A.E. and Haberman S. Dual Modelling and Select Mortality. March 1996. 30 Pages. ISBN 1 874 770 88 3
86. Booth P.M. Long-Term Care for the Elderly: A Review of Policy Options. April 1996. 45 Pages. ISBN 1 874 770 89 1
87. Huber P.P. A Note on the Jump-Equilibrium Model. April 1996. 17 Pages. ISBN 1 874 770 90 5
88. Haberman S and Wong L.Y.P. Moving Average Rates of Return and the Variability of Pension Contributions and Fund Levels for a Defined Benefit Pension Scheme. May 1996. 51 Pages. ISBN 1 874 770 91 3
89. Cooper D.R. Providing Pensions for Employees with Varied Working Lives. June 1996. 25 Pages. ISBN 1 874 770 93 X
90. Khorasane M.Z. Annuity Choices for Pensioners. August 1996. 25 Pages. ISBN 1 874 770 94 8
91. Verrall R.J. A Unified Framework for Graduation. November 1996. 25 Pages. ISBN 1 874 770 99 9
92. Haberman S. and Renshaw A.E. A Different Perspective on UK Assured Lives Select Mortality. November 1996. 61 Pages. ISBN 1 874 770 00 X
93. Booth P.M. The Analysis of Actuarial Investment Risk. March 1997. 43 Pages. ISBN 1 901615 03 0
94. Booth P.M., Chadburn R.G. and Ong A.S.K. Utility-Maximisation and the Control of Solvency for Life Insurance Funds. April 1997. 39 Pages. ISBN 1 901615 04 9
95. Chadburn R.G. The Use of Capital, Bonus Policy and Investment Policy in the Control of Solvency for With-Profits Life Insurance Companies in the UK. April 1997. 29 Pages. ISBN 1 901615 05 7
96. Renshaw A.E. and Haberman S. A Simple Graphical Method for the Comparison of Two Mortality Experiences. April 1997. 32 Pages. ISBN 1 901615 06 5
97. Wong C.F.W. and Haberman S. A Short Note on Arma (1, 1) Investment Rates of Return and Pension Funding. April 1997. 14 Pages. ISBN 1 901615 07 3
98. Puzey A S. A General Theory of Mortality Rate Estimators. June 1997. 26 Pages. ISBN 1 901615 08 1

99. Puzey A S. On the Bias of the Conventional Actuarial Estimator of  $q_x$ . June 1997. 14 Pages.  
ISBN 1 901615 09 X
100. Walsh D. and Booth P.M. Actuarial Techniques in Pricing for Risk in Bank Lending. June 1997.  
55 Pages. ISBN 1 901615 12 X
101. Haberman S. and Walsh D. Analysis of Trends in PHI Claim Inception Data. July 1997.  
51 Pages. ISBN 1 901615 16 2
102. Haberman S. and Smith D. Stochastic Investment Modelling and Pension Funding: A Simulation  
Based Analysis. November 1997. 91 Pages. ISBN 1 901615 19 7
103. Rickayzen B.D. A Sensitivity Analysis of the Parameters used in a PHI Multiple State Model.  
December 1997. 18 Pages. ISBN 1 901615 20 0
104. Verrall R.J. and Yakoubov Y.H. A Fuzzy Approach to Grouping by Policyholder Age in General  
Insurance. January 1998. 18 Pages. ISBN 1 901615 22 7
105. Yakoubov Y.H. and Haberman S. Review of Actuarial Applications of Fuzzy Set Theory.  
February 1998. 88 Pages. ISBN 1 901615 23 5
106. Haberman S. Stochastic Modelling of Pension Scheme Dynamics. February 1998. 41 Pages.  
ISBN 1 901615 24 3
107. Cooper D.R. A Re-appraisal of the Revalued Career Average Benefit Design for Occupational  
Pension Schemes. February 1998. 12 Pages. ISBN 1 901615 25 1
108. Wright I.D. A Stochastic Asset Model using Vector Auto-regression. February 1998. 59 Pages.  
ISBN 1 901615 26 X
109. Huber P.P. and Verrall R.J. The Need for Theory in Actuarial Economic Models. March 1998.  
15 Pages. ISBN 1 901615 27 8
110. Booth P.M. and Yakoubov Y. Investment Policy for Defined Contribution Pension Scheme  
Members Close to Retirement. May 1998. 32 Pages ISBN 1 901615 28 6
111. Chadburn R.G. A Genetic Approach to the Modelling of Sickness Rates, with Application to Life  
Insurance Risk Classification. May 1998. 17 Pages. ISBN 1 901615 29 4
112. Wright I.D. A Stochastic Approach to Pension Scheme Funding. June 1998. 24 Pages.  
ISBN 1 901615 30 8
113. Renshaw A.E. and Haberman S. Modelling the Recent Time Trends in UK Permanent Health  
Insurance Recovery, Mortality and Claim Inception Transition Intensities. June 1998. 57 Pages.  
ISBN 1 901615 31 6
114. Megaloudi C. and Haberman S. Contribution and Solvency Risk in a Defined Benefit Pension  
Scheme. July 1998. 39 Pages ISBN 1 901615 32 4
115. Chadburn R.G. Controlling Solvency and Maximising Policyholders' Returns: A Comparison of  
Management Strategies for Accumulating With-Profits Long-Term Insurance Business. August  
1998. 29 Pages ISBN 1 901615 33 2

116. Fernandes F.N. Total Reward - An Actuarial Perspective. August 1998. 25 Pages.  
ISBN 1 901615 34 0
117. Booth P.M. and Walsh D. The Application of Financial Theory to the Pricing of Upward Only Rent Reviews. November 1998. 23 Pages.  
ISBN 1 901615 35 9
118. Renshaw A.E. and Haberman S. Observations on the Proposed New Mortality Tables Based on the 1991-94 Experience for Male Permanent Assurances. February 1999. 40 Pages.  
ISBN 1 901615 36 7
119. Velmachos D. And Haberman S. Moving Average Models for Interest Rates and Applications to Life Insurance Mathematics. July 1999. 27 Pages.  
ISBN 1 901615 38 3
120. Chadburn R.G. and Wright I.D. The Sensitivity of Life Office Simulation Outcomes to Differences in Asset Model Structure. July 1999. 58 Pages.  
ISBN 1 901615 39 1
121. Renshaw A.E. and Haberman S. An Empirical Study of Claim and Sickness Inception Transition Intensities (Aspects of the UK Permanent Health Insurance Experience). November 1999. 35 Pages.  
ISBN 1 901615 41 3

#### Statistical Research Papers

1. Sebastiani P. Some Results on the Derivatives of Matrix Functions. December 1995.  
17 Pages. ISBN 1 874 770 83 2
2. Dawid A.P. and Sebastiani P. Coherent Criteria for Optimal Experimental Design.  
March 1996. 35 Pages. ISBN 1 874 770 86 7
3. Sebastiani P. and Wynn H.P. Maximum Entropy Sampling and Optimal Bayesian Experimental Design. March 1996. 22 Pages.  
ISBN 1 874 770 87 5
4. Sebastiani P. and Settini R. A Note on D-optimal Designs for a Logistic Regression Model. May 1996. 12 Pages.  
ISBN 1 874 770 92 1
5. Sebastiani P. and Settini R. First-order Optimal Designs for Non Linear Models. August 1996.  
28 Pages. ISBN 1 874 770 95 6
6. Newby M. A Business Process Approach to Maintenance: Measurement, Decision and Control.  
September 1996. 12 Pages. ISBN 1 874 770 96 4
7. Newby M. Moments and Generating Functions for the Absorption Distribution and its Negative Binomial Analogue. September 1996. 16 Pages.  
ISBN 1 874 770 97 2
8. Cowell R.G. Mixture Reduction via Predictive Scores. November 1996. 17 Pages.  
ISBN 1 874 770 98 0
9. Sebastiani P. and Ramoni M. Robust Parameter Learning in Bayesian Networks with Missing Data. March 1997. 9 Pages.  
ISBN 1 901615 00 6
10. Newby M.J. and Coolen F.P.A. Guidelines for Corrective Replacement Based on Low Stochastic Structure Assumptions. March 1997. 9 Pages.  
ISBN 1 901615 01 4.

11. Newby M.J. Approximations for the Absorption Distribution and its Negative Binomial Analogue. March 1997. 6 Pages. ISBN 1 901615 02 2
12. Ramoni M. and Sebastiani P. The Use of Exogenous Knowledge to Learn Bayesian Networks from Incomplete Databases. June 1997. 11 Pages. ISBN 1 901615 10 3
13. Ramoni M. and Sebastiani P. Learning Bayesian Networks from Incomplete Databases. June 1997. 14 Pages. ISBN 1 901615 11 1
14. Sebastiani P. and Wynn H.P. Risk Based Optimal Designs. June 1997. 10 Pages. ISBN 1 901615 13 8
15. Cowell R. Sampling without Replacement in Junction Trees. June 1997. 10 Pages. ISBN 1 901615 14 6
16. Dagg R.A. and Newby M.J. Optimal Overhaul Intervals with Imperfect Inspection and Repair. July 1997. 11 Pages. ISBN 1 901615 15 4
17. Sebastiani P. and Wynn H.P. Bayesian Experimental Design and Shannon Information. October 1997. 11 Pages. ISBN 1 901615 17 0
18. Wolstenholme L.C. A Characterisation of Phase Type Distributions. November 1997. 11 Pages. ISBN 1 901615 18 9
19. Wolstenholme L.C. A Comparison of Models for Probability of Detection (POD) Curves. December 1997. 23 Pages. ISBN 1 901615 21 9
20. Cowell R.G. Parameter Learning from Incomplete Data Using Maximum Entropy I: Principles. February 1999. 19 Pages. ISBN 1 901615 37 5
21. Cowell R.G. Parameter Learning from Incomplete Data Using Maximum Entropy II: Application to Bayesian Networks. November 1999. 12 Pages. ISBN 1 901615 40 5

**Department of Actuarial Science and Statistics**

Actuarial Research Club

The support of the corporate members

CGU  
Government Actuary's Department  
Guardian Insurance  
Hymans Robertson  
KPMG  
Munich Reinsurance  
PricewaterhouseCoopers  
Swiss Reinsurance  
Watson Wyatt

is gratefully acknowledged.

ISBN 1 901615 39 1