December 1967

Brief 67-10590



AEC-NASA TECH BRIEF



AEC-NASA Tech Briefs describe innovations resulting from the research and development program of the U.S. AEC or from AEC-NASA interagency efforts. They are issued to encourage commercial application. Tech Briefs are published by NASA and may be purchased, at 15 cents each, from the Clearinghouse for Federal Scientific and Technical Information, Springfield, Virginia 22151.

Ultraviolet Microscopy Aids in Cytological and Biomedical Research

Ultraviolet microscopy has been used by cytologists and biochemists to study the morphological and physiological changes which occur in the living cell under varied culture conditions. The yeast cell was chosen as a suitable specimen for investigation because its content of ultraviolet-absorbing materials lies within the range of observation by ultraviolet microscopy, and its lack of motility is an important factor in the production of useful images with the available ultraviolet energy.

The yeast cells were cultured under conditions which produce a high concentration of the strongly ultraviolet-absorbing compound, S-adenosylmethionine, in the vacuole. The cells were induced to sporulate and ultraviolet photomicrographs were obtained. These showed that the vacuole is abolished at the onset of sporulation, and its contents may be observed temporarily within the old cell wall surrounding the spores. As sporulation progresses, the material is discharged into the culture medium and is not utilized. This is quite different from what occurs during budding, where the vacuolar material appears in the vacuole of the bud in a concentration indicating that parent and daughter share equally.

Ultraviolet photomicrographs of cells containing large, easily observed, cytoplasmic granules revealed that most of the granules disappeared during sporulation and that an ultraviolet-absorbing lipid substance lay between the spores, wetting them. It appeared, thus, that some of the granules were composed of, or contain, this lipid.

It is believed that the lipid material present in the granules takes part in the formation of the spore coat, especially the outer layer. Many more tests are being run to clarify this and other questions such as the role of deoxyribonucleic acid (DNA) during sporulation.

Notes:

- 1. This information would be of interest to biophysicists, microbiologists, biochemists, and physiologists, as it can be applied to other cytological and biochemical studies.
- 2. Additional details are contained in:
 - (a) Ultraviolet Microscopy of the Vacuole of Saccharomyces Cerevisiae during Sporulation, Journal of Bacteriology. 88, 449-456 (1964), by G. Svihla, J. L. Dainko, and F. Schlenk.
 - (b) Biological and Medical Research Division Annual Report, 1964, ANL-6971, pp. 21-22, Argonne National Laboratory, Argonne, Illinois. The Argonne report is available from the Clear-inghouse for Federal Scientific and Technical Information, Springfield, Virginia 22151; \$3.00 each (microfiche, \$0.65).
- 3. Inquiries concerning this innovation may be directed to:

Office of Industrial Cooperation Argonne National Laboratory 9700 South Cass Avenue Argonne, Illinois 60439 Reference: B67-10590

Source: B. Svihla and F. Schlenk Biological and Medical Research Division (ARG-178)

Patent status:

Inquiries about obtaining rights for commercial use of this innovation may be made to:

Mr. George H. Lee, Chief Chicago Patent Group U.S. Atomic Energy Commission Chicago Operations Office 9800 South Cass Avenue Argonne, Illinois 60439

Category 04

This document was prepared under the sponsorship of the Atomic Energy Commission and/or the National Aeronautics and Space Administration. Neither the United States Government nor any person acting on behalf of the United States Government assumes any liability resulting from the use of the information contained in this document, or warrants that the use of any information, apparatus, method, or process disclosed in this document may not infringe privately owned rights.

Place of States

Heermon 1991

AEC-NASA TECH BRIEF



vice, valia, if white deviate providing results from the rescuent and de playment prostant a the US A DS or them AirC MA SA interspective obtains. They are noted to choose any solution terrate more during. Foch Brief, and published by "Assa and man he play have and in consultant. Interhere (datasets) are first and published by "Assa and man he play have and the second terrate and the first public of the terral Sciences of the result of the play have anti-second to the second terrate

to the second second second strain and second a second s

An entropy of the first of a set of the set

and an an anna. An a tarrait that are hard a function of a second solution (a) that are hard a function of a second and any of a second fined with a second to be set a second the second second and a second second and a second to second second second a second second and a second to second second second second second second and a second second second second second second second and second second second second second second second second and second second second second second second second second and second and second se

Lord San is the base of the same transport of the second s