View metadata, citation and similar papers at core.ac.uk

June 1968

brought to you by 🐰 CORE

Brief 68-10228

NASA TECH BRIEF



NASA Tech Briefs are issued to summarize specific innovations derived from the U.S. space program, to encourage their commercial application. Copies are available to the public at 15 cents each from the Clearinghouse for Federal Scientific and Technical Information, Springfield, Virginia 22151.

Theory of Refined Earth Model

Refined equations have been derived relating the variations of the earth's gravity and radius as functions of longitude and latitude. Starting from the expansion in spherical harmonics of the geopotential of a rotating three-axial earth model, the three components of the gravity acceleration vector were obtained. The generalized equation for the surface of the earth (also expressed as an expansion in spherical harmonics) was inserted into the equation for the constant surface potential to derive quantitative mathematical relationships between the oblateness coefficients of the zonal and tesseral harmonics and the geometrical and gravitational parameters of the earth. These equations particularly relate the oblateness coefficients of the odd harmonics and the difference of the polar radii (respectively, ellipticities and polar gravity accelerations) in the northern and southern hemispheres. The new theory on which the equations are based may well replace the first-, second-, and third-order theories of the earth's figure developed by Clairaut, G. H. Darwin, Callandreau, Helmert, de Sitter, and Cook.

Starting from the newest values for the oblateness coefficients of the even and odd spherical harmonics

up to the 14th degree, given by Y. Kozai in 1964, the new theory gives a mean meridional reciprocal flattening 1/f = 298.17. The corresponding values for the northern and southern hemispheres are, respectively, 298.43 and 297.91. The difference between the polar radii is 37.8 meters, and the difference between the polar gravity accelerations is 13.4 milligals (1 gal = 1 centimeter per second per second). The equator has an ellipticity of 1.082×10^{-5} , corresponding to a difference of 69.0 meters between the largest and smallest equatorial radii, and a maximum difference of 31.8 milligals between the equatorial gravity accelerations.

Note:

Complete details may be obtained from: Technology Utilization Officer Marshall Space Flight Center Huntsville, Alabama 35812 Reference: B68-10228

Patent status:

No patent action is contemplated by NASA.

Source: H. G. L. Krause (MFS-14679)

Category 02

This document was prepared under the sponsorship of the National Aeronautics and Space Administration. Neither the United States Government nor any person acting on behalf of the United States

0

Government assumes any liability resulting from the use of the information contained in this document, or warrants that such use will be free from privately owned rights.