

N73-28367

**ERTS-1 APPLIED FOR STRUCTURAL AND MORPHOLOGICAL INVESTIGATIONS
CASE STUDIES: (1) LOS ANGELES, CALIFORNIA AND (2) COASTAL PLAIN,
NEW JERSEY**

Ervin Y. Kedar, Department of Geography, William Paterson College of New Jersey

This report addresses itself to a major earth's resources management problem; namely, the application of ERTS-1 imagery for geo-morpho-tectonics, and subsequently seismic-risk, earthquake, and mineral exploration applications.

Case Study No. 1 - Los Angeles, California

ERTS-1 image (MSS color composite of August 10, 1972, E-1018-501, centered at N 34.13/w 118-27), which covers the area from South Los Angeles to the Mojave Desert and from Eastern San Gabriel Mts. to Cuyama Valley and Central Santa Ynez Mts., of the Transverse Ranges, California, was analyzed (fig. 1). This image was studied in conjunction with the following conventional sources: (1) the Los Angeles Sheet of the geologic map of California (1); (2) the map of the Geology of the Central Santa Ynez Mountains, Santa Barbara County, California (2); (3) the geologic map of Transverse Ranges, Southern California (3).

A comparison of ERTS-1 image to the conventional sources indicated the usefulness of the image in studying structural geology.

The following are aspects of the comparisons:

- (1) Each conventional source failed to show the Verdugo Mts. lineament, the Simi Hills lineament and the Santa Monica Mts. lineament, all of which bound the San Fernando Valley (fig. 2). These structural lineaments which are seen readily on the ERTS-1 image are apparently the major morphotectonic features in the structural framework of the San Fernando Valley. The image also indicates that Point Dume is the seaward extension of the Simi Hills lineament across the Santa Monica Mts., and that Point Dume represents a post-Cretaceous/early Paleocene geotectonic episode.
- (2) Each conventional source fails to show the oval-shaped sedimentary basin between the Simi Hills lineament in the S.E. and the Oakridge fault in the N.W. This oval-shaped unit, which has no physiographic name, since it was not seen yet, is a lithological and structural unit separated from the adjacent region. It is inferred from ERTS-1 image that the Violin Canyon watershed, the upper part of the Santa Clara River system, is the major source of sedimentation in the oval basin.

1527

PRECEDING PAGE BLANK NOT FILMED

Original photography may be purchased from
EROS Data Center
10th and Dakota Avenue
Sioux Falls, SD 57198

- (3) Neither conventional source showed the major structural element in the Transverse Ranges; namely, the Ventura-Soledad Trough. This structural trough, which is readily seen on the ERTS-1 image, cuts across the Transverse Ranges from Antelop Valley to Santa Barbara Channel. It is believed that Ventura-Soledad Trough is the most seismic part of greater Los Angeles. Newhall, California, the epicenter of the San Fernando Valley earthquake in February 1971, is located on the intersection of the San Gabriel Fault, the Verdugo Mts. Fault, the Simi Hills Fracture Belt and the Ventura-Soledad Trough. This conclusion was derived from ERTS-1 image analysis (fig. 3).

The conventional sources do not show the above mentioned information because they are either overgeneralized or undergeneralized. The geologic map of the Transverse Range Province has only fault lines and rock age distribution and lacks drainage patterns and contour lines. The Los Angeles sheet of the geologic map of California has too much information (road system, railroads, power line, etc.) and makes no distinction between major faults and local faults. ERTS-1 image was found to be a generalized, yet selective, source of information.

Case Study No. 2 - Coastal Plain, New Jersey

ERTS-1 image (MSS color composite of August 17, 1972, E-1024-15071-501 centered at N 40-21/w 073.26), which covers the area from the Palisades, New York-New Jersey to Burlington, New Jersey, and from New Brunswick, New Jersey to the great Pocomic Bay, New York, was analyzed (fig. 4). This image was compared with: (1) the geologic map of New Jersey 1:1,000,000 (4); (2) the geologic map of New Jersey 1:250,000 (5); and (3) selected maps as published in the geology of selected areas in New Jersey and Eastern Pennsylvania (6).

In this case study, too, ERTS-1 image indicated its usefulness in studying the structural geology of the test site region.

The following are aspects of this case study:

- (1) The coastal plain is underlain almost entirely by unconsolidated marine and fluvial clay, silt, sand and gravel of Late Cretaceous and Tertiary Age. Much of this province is covered by Quarternary deposits, particularly in the inner and outer lowlands and the southern one-third of the central upland of the Coastal Plain. The difficulty of mapping the quarternary deposits was recognized a long time ago (Salisbury 1917, 7). He subdivided the Pleistocene deposits into three formations: (from the oldest to the youngest) (a) the Bridgeton; (b) Pensauken; and (c) Cape

May. The major portion of the Pensauken is confined to a 20-mile wide belt whose axis extends from Staten Island, N. Y., south westward to Salem, N. Y. This belt was first seen and traced on the ERTS-1 image. It was suggested that this belt is trough shaped. This trough was only roughly mapped (Campbell and Bascom 1933, 8). It was also concluded that the Pensauken was an alluvial deposit formed by the combined ancestral drainage system of the Hudson River (9), (fig. 5).

- (2) The Coastal Plain has potential economic ilmenite sands type ore bodies and other heavy mineral deposits (10). Today several companies mine these and other minerals in the Coastal Plain. Some of the mines can be seen and identified on the ERTS-1 image, such as the mines at Lakehurst, N. J. In order to determine the location of these mineral deposits, the ancient drainage patterns have to be determined. ERTS-1 imagery can be used for the hypothetical reconstruction of ancient drainage patterns (see (1)a on previous page), (fig. 6).

Conventional sources, including field investigation, have been found to be inadequate to supply all the required structural and paleohydrographic information needed.

It is, therefore, concluded that ERTS-1 was found to be an essential supplementary source of information for geologic explorations, much like and even better than space photography from manned spacecraft, and SLR imagery (11).

REFERENCES

1. Anon: Los Angeles Sheet of the Geologic Map of California, 1:50,000, 1955.
2. T. W. Dibblee, Jr., Geology of the Central Santa Ynez Mountains, Santa Barbara County, California, Bull. 186, California Division of Mines and Geology, San Francisco, 1966.
3. Olaf P. Jenkins, Geologic Map of Transverse Range Province, Southern California, Department of Natural Resources, State of California, Bull. 170, Chap. 2, Cont. 3, Plate 4, 1954.
4. Geologic Map of New Jersey, 1:1,000,000, New Jersey Geographical Press, Little Falls, N. J., 1962.
5. J. V. Lewis and H. B. Kummel, The Geology of New Jersey, Geologic Map 1:250,000, Geological Survey of New Jersey, 1915.

6. **Geology of Selected Areas in New Jersey and Eastern Pennsylvania and Guidebook of Excursions**, edit. by Seymour Subitzky, Rutgers University Press, New Jersey, 1969.
7. **R. D. Salisbury and G. N. Knapp, The Quarternary Formations of Southern New Jersey: The Final Report Series of the State Geologist, Vol. VIII** Department of Conservation and Economic Development, Trenton, New Jersey, 1917.
8. **M. R. Campbell and F. Bascom, Origin and Structure of the Pensauken Gravel: Am. Your. Sci., Vol. 26, p. 300-318.**
9. **R. R. Jordan, Columbia Sediments of Delaware: Delaware Geological Survey, Bull. No. 12, Newark, 1964.**
10. **Frank J. Markewicz, Ilmenite Deposits of the New Jersey Coastal Plain, in Geology of Selected Areas in New Jersey, pp. 363-382, 1969.**
11. **The following are publications by E. Y. Kedar:**
 - Side-Looking Radar (SLR) Imagery Applied to Seismic-Risk Mapping, Eighth International Symposium on Remote Sensing of Environment, October 2-6, 1972, Ann Arbor, Michigan.**
 - Clustering Phenomena in Side-Looking Radar (SLR) Micro-Texture, Electro-Optics 1972 Conference, September 12-14, 1972, New York City.**
 - Space Photographs of the Earth in the Study of Plate Tectonics, 22nd International Geographical Congress, Montreal, Canada, August 10-15, 1972.**
 - Earth Photographs from Space Applied for Geomorphotectonic References presented at the Association of the American Geographers, Kansas meeting, April 1972.**
 - Earth Photographs from Space for Plate Tectonics and Earthquake-Risk Mapping, American Society of Photogrammetry, March 12-17, 1972, Washington, D. C., Proceedings of the 38th Annual Meeting, ASP.**
 - Application of the Discriminant Function in Automatic Pattern Recognition of Side-Looking Radar, presented to the Seminar of the Society of Photo-Optical Instrumentation Engineers, Redondo Beach, California, Feb., 1972.**

D

Plate Tectonics in the Israel-Sinai Region, The Geological Society of America, Abstracts, the 1971 Annual Meeting, Washington, D. C., November, 1971, Vol. 3, No. 7, p. 476, 619.

An Automatic Analytical System of Side-Looking Radar (SLR) Imagery and its Applications in Environmental Discrimination, Proceedings of the Technical Program, Electro-Optical Systems, N. Y., September 5-6, 1971, pp. 353-356.

Manual and Automatic-Computerized Systems of Side-Looking Radar Imagery Analysis for Crop Discrimination, in the American Society of Photogrammetry, Proceedings, ASP-ACSM, Fall Convention, San Francisco, September 1971, pp. 190-199.

Manual and Automatic-Computerized System in SLR Application for Land Use Survey, Technology, Utilization Ideas for the 70's and Beyond, American Astronautical Society, Vol. 26, Science and Technology, May 1971, pp. 239-262.

Plate Tectonics in the Red Sea Region as Inferred from Space Photography, NASA Technical Note, TN D-6261, NASA, Washington, D. C., April 1971 and presented at the Seventh International Symposium on Remote Sensing of Environment, May 1971, Ann Arbor, Michigan.

Plate Tectonics in the Red Sea Region as Inferred from Space Photography, Earth-Observation Division, Johnson Space Center, NASA, S-257.

Transverse Range: Tectonic Inferences from Earth Photographs from Space. Earth-Observation Division, Johnson Space Center, NASA, TF7-2, June 1970. NR-E9-00-000-01056.

Texas Lineament Traced on Earth Photographs Taken from Space. Earth-Observation Division, Johnson Space Center, NASA TF7-3, June 1970. NR-E9-09-998-00407.

Space Photographs of the Earth in the Study of Plate Tectonics. Earth-Observation Division, Johnson Space Center, NASA, TF7-4, June 1970. NR-E9-EJ-998-01055.

Plate-Margin Morphotectonics Case Study: The Israel-Sinai Section. Earth-Observation Division, Johnson Space Center, NASA, TF7-6, June 1970. NR-E9-00-998-01131.

Space Photographs of the Earth in the Study of Geotectonics, Science and Earth Technology, Vol. 23, American Astronautical Society 69-579, October 1969, pp. 331-332.

Application of Space Photographs for Geomorphological Study; Taurus Mountains, Turkey, Earth Observation Division, Johnson Space Center, NASA, TF7-1, August 1969, NR-E9-03-000-00634.

Geomorphotectonic Inferences from Apollo and Gemini Photographs: A Case Study of the Southwestern United States, presented at the 32nd Annual Meeting of the Association of Pacific Coast Geographers, San Fernando Valley State College, June 11-14, 1969.



Fig. 1 ERTS MSS Color Composite of August 10, 1972
Centered at N 34.13/W 118.27.




Fig. 2 Geologic Map, Transverse Range Province.




Fig. 3 ERTS-1 MSS Color Composite of August 10, 1972, Structural Analysis (See fig. 1).




Fig. 4 ERTS-1 MSS Color Composite of August 16, 1972,
Centered at N 40.21/W 073.26.





Fig. 5 ERTS-1 MSS of Eastern New Jersey Paleo-Hydrology.



AUG72 C N40-21/W073-26 N N40-21/W073-23 MSS 6 D SUN EL53 AZ130 191-0333- -1-N-D-2L NASA ERTS E-1224-1521-8

Fig. 6 ERTS-1 MSS Structural and Hydrographic Interpretations.