

320

5.3A CHARACTERISTICS OF SUNSET RADAR

J. L. Green

NOAA, Aeronomy Laboratory
325 Broadway
Boulder, CO 80303

The Sunset radar is a VHF, pulsed Doppler radar located in a narrow canyon near the Sunset townsite 15 km west of Boulder, CO. This facility is operated by the Aeronomy Laboratory, ERL, NOAA, exclusively for meteorological research and the development of the MST and ST radar technique.

ANTENNA

The antenna system consists of two colocated arrays, each containing 16 lines of coaxial-colinear dipoles. One of these arrays can be steered in the N-S vertical plane and the other in the E-W plane. The phase progression necessary for steering the antenna beam is provided by the insertion of remotely controlled phase shifters in the transmission line to each line of dipoles. The phase shifters have binary format (i.e., phase increments of $\lambda/2$, $\lambda/4$,..... $\lambda/256$ can be summed). The practical range of beam steering for meteorological experiments is $\pm 45^\circ$ due to the loss of antenna gain with increasing zenith angle. The beam steering increment near the zenith is 0.4° to take advantage of the sharp nulls in the antenna pattern. The phase shifters also perform the switching between the two arrays. Each phase shifter is controlled by the on-line computer.

DATA PROCESSING

On-line data processing is performed by a Nova 800 computer. This computer provides radar control, antenna steering control, spectral analysis by a FFT algorithm, spectral averaging, on-line spectral plotting, and output to a tape recorder and a hard copy unit. Control of the radar is by a "menu" subroutine that allows immediate changes in range gate altitudes, range gate length, number of spectra averaged, and antenna beam position. Sequences of antenna positions may also be specified.

Coherent averaging is provided by analogue filters on each of the 16 range gates. The effects of ground clutter are reduced by a 0 - 0.1 Hz notch in the response of these filters. Coherent averaging is especially effective because of the high pulse repetition frequency of the radar (typically 6 kHz).

Since March, 1981, calibration signals from a stable noise generator have been placed on each data tape for each mode of operation used. These calibrations, along with the recorded transmitted power, are used to determine absolute radar reflectivity.

Magnetic data tapes are further processed by a scaling program on the central computing facility of the Boulder Laboratories of NOAA. The output of this program contains radial velocity, velocity width, received power, C_n^2 , and the magnitude of the reflection coefficient for each range gate.

HISTORY AND SIGNIFICANT ACCOMPLISHMENTS

The Sunset radar was the first radar designed and constructed specifically

 Summary of Sunset radar characteristics

Frequency	40.475 MHz
Wavelength	7.41 m
Transmitter power	
Peak	100 kW
Average (maximum)	16 kW
(typical)	2.5 kW
Antenna	
Phased array of coaxial-colinear dipoles with computer-controlled phase shifters for each line of dipoles	
Area	3600 m ²
Aperture efficiency	0.58
Resistive loss	0.30
Steerability	
(E-W or N-S vertical plane only)	± 60°
Beam width H plane (steerable)	4.8°
E plane	4.4°
Steering increment	0.2°
Practical steering rate	1 record/minute/position to any arbitrary antenna beam position

as a VHF ST radar. Construction was begun in 1973 and the first clear-air echoes and wind velocity measurements were obtained in 1974 (GREEN et al., 1975). The Sunset radar has been used to demonstrate the feasibility of using an ST radar to measure wind velocity (WARNOCK et al., 1978), all three vectors of wind velocity (GREEN et al., 1978a), turbulence (VANZANDT et al., 1978), wind variability (GAGE and CLARK, 1978), variation of clear air turbulence near and above thunderstorms (GREEN et al., 1978b), simultaneous measurement of hydrometeor and air velocity (GREEN et al., 1978b), turbulence intensity using velocity width (1978c) specular reflection from stable layers (GAGE and GREEN, 1978), gravity (buoyancy) waves (VANZANDT et al., 1979), the height of the tropopause (GAGE and GREEN, 1979b), turbulence parameters (GAGE et al., 1980); devise a practical and economical method of steering ST radar antennas (GREEN et al., 1981); and estimate the gradient of temperature in the stratosphere (GAGE and GREEN, 1982).

(References in this paper are included in the Publications listed below.)

PUBLICATIONS RELATED TO THE SUNSET RADAR

- Green, J. L., J. M. Warnock, R. H. Winkler and T. E. VanZandt (1975), Studies of winds in the upper troposphere with a sensitive VHF radar, Geophys. Res. Lett., **2**, 19.
- Green, J. L., J. M. Warnock, R. H. Winkler and T. E. VanZandt (1975), A sensitive VHF radar for the study of winds, waes, and turbulence in the troposphere, stratosphere, and mesosphere, 16th Radar Meteorol. Conf., Houston, Am. Meteorol. Soc.
- VanZandt, T. E., J. L. Green, J. M. Warnock and R. W. Winkler (1975), Studies of winds in the upper troposphere with a sensitive VHF radar, 16th Radar Meteorol. Con., Houston, AMS, April 22-24.

- VanZandt, T. E., J. L. Green, J. M. Warnock, K. S. Gage and W. L. Clark (1977), Reflectivity studies with the Sunset VHF pulsed Doppler radar, Proc. Symp. on Radio Prop. and Non-ionized Media, 28 April - 6 May, Centre National d'Etudes des Telecommunications, Paris, France, 519-524.
- B. B. Balsley and J. L. Green (1978), Coherent radar systems for probing the troposphere, stratosphere, and mesosphere, Proc. 4th Symp. on Meteorol. Observations and Instrumentation, April 10-14, 1978, Denver, CO, 227-230.
- Gage, K. S. (1978), Comparison of sensitive Doppler radar observations of winds and turbulence with in situ observations, Proc. 4th Symp. on Meteorol. Observations and Instrumentation, April 10-14, Denver, CO, 231-238.
- Gage, K. S. and J. L. Green (1978), Evidence for specular reflection from monostatic VHF radar observations of the stratosphere, Radio Sci., 13, 991-1001.
- Gage, K. S., J. L. Green and T. E. VanZandt (1978), Application of the VHF pulsed Doppler radar to cloud physics research, Proc. Conf. on Cloud Physics and Atmospheric Electricity, July 31 - August 4, 1978, Issaquah, Wash. 394-401.
- Gage, K. S. and B. B. Balsley (1978), Doppler radar probing of the clear atmosphere, Bull. Amer. Meteorol. Soc., 59, 1074-1093.
- Gage, K. S. and W. L. Clark (1978), Mesoscale variability of jet stream winds observed by the Sunset VHF Doppler radar, J. Applied Meteorol., 17, 1412-1416.
- Gage, K. S., T. E. VanZandt and J. L. Green (1978), Vertical profiles of C in the free atmosphere: Comparison of model calculations with radar observations Proc. 18th Conf. on Radar Meteorol., March 28-31, 1978, Atlanta, GA, 184-189.
- Green, J. L., K. S. Gage and T. E. VanZandt (1978c), VHF Doppler studies of CAT in the troposphere and lower stratosphere, Preprint Vol, Conf. on Atmos. Environment of Aerospace Systems and Appl. Meteorol., Nov. 14-16, New York, NY, 159-164.
- Green, J. L., K. S. Gage and T. E. VanZandt (1978a), Three dimensional wind observations of a jet stream using a VHF Doppler radar, Proc. 18th Conf. on Radar Meteorol., March 28-31, Atlanta, GA, 184-189.
- Green, J. L., R. H. Winkler, J. M. Warnock, W. L. Clark, K. S. Gage and T. E. VanZandt (1978b), Observations of enhanced clear air reflectivity associated with convective clouds, Proc. 18th Conf. on Radar Meteorol., March 28-31, Atlanta, GA, 88-93.
- VanZandt, T. E., J. L. Green, K. S. Gage and W. L. Clark, (1978), Vertical profiles of refractivity turbulence structure constant: comparison of observations by the Sunset radar with a new theoretical model, Radio Sci., 13, 818-829.
- Warnock, J. M., T. E. VanZandt, J. L. Green and R. H. Winkler (1978), Comparisons between wind profiles measured by Doppler radar and by rawinsonde balloons, Geophys. Res. Lett., 5, 109-112.
- Gage, K. S., J. L. Green, W. L. Clark and T. E. VanZandt (1979), Doppler radar measurement of turbulence in the clear atmosphere, Proc. Conf. Fourth Symp. on Turbulence, Diffusion and Air Pollution, January 15-18, Reno, Nevada, 522-529.

- Gage, K. S. and J. L. Green (1979b), Tropopause detection by partial specular reflection using VHF radar, Science, 203, 1238-1240.
- Green, J. L., K. S. Gage and T. E. VanZandt (1979), Atmospheric measurements by VHF pulsed Doppler radar, IEEE Trans. on Geosci. Elec., GE-17, 262-280.
- VanZandt, T. E., K. S. Gage and J. M. Warnock (1979), A statistical model for the probability of turbulence and the calculation of vertical profiles of turbulence parameters, Proc. Conf. Fourth Symp. on Turbulence, Diffusion, and Air Pollution, January 15-18, Reno, Nevada, 52-55.
- VanZandt, T. E., J. L. Green, W. L. Clark and J. R. Grant (1979), Buoyancy waves in the troposphere: Doppler radar observations and a theoretical model, Geophys. Res. Lett., 6, 429-432.
- Gage, K. S., J. L. Green and T. E. VanZandt (1980), Use of Doppler radar for the measurement of atmospheric turbulence parameters from the intensity of clear-air echoes, Radio Sci., 15, 407-416.
- Green, J. L. and K. S. Gage (1980), Observations of stable layers in the troposphere and stratosphere using VHF radar, Radio Sci., 15, 395-405.
- VanZandt, T. E., K. S. Gage, and J. M. Warnock (1980), Progress in modelling turbulence parameters from profiles of background winds, temperature, and humidity, Proc. AMS 19th Conf. on Radar Meteorol., April 15-18, 613-615.
- Gage, K. S. and J. L. Green (1980), Prospects for a quantitative measurement of atmospheric stability using VHF radar, Proc. AMS 19th Conf. on Radar Meteorol., April 15-18, Miami Beach, FL, 605-612.
- Green, J. L., J. M. Warnock, W. L. Clark and F. J. Eggert (1980), Modifications to the Sunset radar to provide beam steering, Proc. AMS 19th Conf. on Radar Meteorol., April 15-18, Miami Beach, FL, 575-576.
- Green, J. L., K. S. Gage, W. L. Clark, T. E. VanZandt and P. H. Hildebrand (1980), Joint instrumented aircraft and VHF Doppler radar measurements of wind near Boulder, CO, Proc. AMS 19th Conf. on Radar Meteorol., April 15-18, Miami Beach, FL, 624-628.
- VanZandt, T. E., K. S. Gage and J. L. Green (1979), The use of Doppler radar measurement of atmospheric turbulence parameters in the clear atmosphere, Collection of Extended Abstracts Presented at ICMUA Sessions and IUGG Symp. 18, XVII IUGG Gen. Assemb., Canberra, Australia, 285-292.
- Gage, K. S. and J. L. Green (1979a), Observations of stable layers in the middle atmosphere using VHF radar, Collection of Extended Abstracts Presented at ICMUA Sessions and IUGG Symp. 18, XVII IUGG Gen. Assemb., Canberra, Australia, 325-332.
- Gage, K. S., B. B. Balsley and J. L. Green (1981), Fresnel scattering model for the specular echoes observed by VHF radar, Radio Sci., 16, 1447-1453.
- Gage, K. S. and J. L. Green (1982), A technique for determining the temperature profile from VHF radar observations, J. Appl. Meteorol., 21, 1155-1158.
- Balsley, B. B., J. L. Green, W. L. Ecklund, W. L. Clark, R. G. Strauch and A. C. Riddle (1981), Joint observations of gravity wave activity in vertical winds in the troposphere and lower stratosphere over a 63 km baseline obtained with clear-air VHF radars at Platteville and Sunset, Colorado, Preprint Vol, 20th Conf. on Radar Meteorol., Boston, Nov 30 - Dec 3, AMS, 110-115.

Green, J. L., J. M. Warnock, W. L. Clark, F. J. Eggert and K. J. Ruth (1981), Modifications to the Sunset radar to provide antenna beam steering, Preprint Vol., 20th Conf. on Radar Meteorol., Boston, Nov 30 - Dec 3, AMS, 7-10.

VanZandt, T. E., K. S. Gage and J. M. Warnock (1981), An improved model for the calculation of profiles of C and in the free atmosphere from background profiles of wind, temperature and humidity, Preprint Vol., 20th Conf. on Radar Meteorol., Boston, MA, Nov 30 - Dec 3, AMS, 129-135.

Ecklund, W. L., K. S. Gage, B. B. Balsley, R. G. Strauch and J. L. Green (1982), Vertical wind variability observed by VHF radar in the lee of the Colorado Rockies, Mon. Wea. Rev., 110, 1451-1457.