



Horn, D. and McCulloch, R. (2010) *Molecular mechanisms underlying the control of antigenic variation in African trypanosomes*. *Current Opinion in Microbiology*, 13 (6). pp. 700-705. ISSN 1369-5274.

<http://eprints.gla.ac.uk/48025/>

Deposited on: 20 January 2011

**Antigenic variation in the African trypanosome: molecular mechanisms and phenotypic complexity**

Liam J. Morrison, Lucio Marcello and Richard McCulloch\*

University of Glasgow,

Wellcome Centre for Molecular Parasitology and Division of Infection and Immunity,

Glasgow Biomedical Research Centre,

120 University Place,

Glasgow,

G12 8TA.

\*Correspondence:

e-mail: [rmc9z@udcf.gla.ac.uk](mailto:rmc9z@udcf.gla.ac.uk)

phone: 44 (0)141 330 5946

fax.: 44 (0)141 330 5422

## **Abstract**

Antigenic variation is an immune evasion strategy that has evolved in viral, bacterial and protistan pathogens. In the African trypanosome this involves stochastic switches in the composition of a Variant Surface Glycoprotein (VSG) coat, using a massive archive of silent *VSG* genes to change the identity of the single VSG expressed at a time. VSG switching is driven primarily by recombination reactions that move silent *VSGs* into specialised expression sites, though transcription-based switching can also occur. Here we discuss what is being revealed about the machinery that underlies these switching mechanisms, including what parallels can be drawn with other pathogens. In addition, we discuss how such switching reactions act in a hierarchy and contribute to pathogen survival in the face of immune attack, including the establishment and maintenance of chronic infections, leading to host-host transmission.

## **Antigenic variation in African trypanosomes**

African trypanosomes are protistan parasites that infect a range of mammals and are transmitted between successive hosts by the tsetse fly. In all mammals, the parasites replicate extracellularly in the tissue fluids and bloodstream. Despite continuous exposure to immune attack trypanosome infections can be very prolonged, causing a range of pathological manifestations (Barrett *et al.*, 2003). Trypanosome persistence in the mammal is due to antigenic variation, which involves changes in the identity of the Variant Surface Glycoprotein (VSG), which forms a dense cell surface coat that shields invariant surface antigens from immune recognition. Antibodies against the VSG kill the trypanosome, but the population survives due to stochastic switching to a coat composed of an antigenically distinct VSG. The rate of VSG switching (up to  $10^3$  switchers/cell/division)(Turner and Barry, 1989) is typical of what Moxon and colleagues termed ‘enhanced phenotypic variation’ (Moxon *et al.*, 1994) in subset of genes in viral, bacterial and eukaryotic pathogens (Deitsch *et al.*, 2009).

Trypanosomes have made a huge investment in antigenic variation. The genome of *T. brucei* contains >1600 *VSG* genes (Marcello and Barry, 2007), hugely in excess of antigenic variation gene families elsewhere (Table 1). Most (80-90%) of this *VSG* archive is in 11 diploid, megabase-sized chromosomes; the subtelomeres harbour

silent *VSG* arrays (Berriman *et al.*, 2005), while sites for *VSG* transcription, termed the *VSG* expression sites (ES) (Fig.2), are found directly proximal to the telomeres (Hertz-Fowler *et al.*, 2008). Only a minority (~5%) of *VSG* array genes are functional, with the large majority (~85%) being pseudogenes (Marcello and Barry, 2007), which (in common with other pathogens) provide a critical resource during antigenic variation. *T. brucei* has also evolved novel, aneuploid chromosomes to facilitate antigenic variation. Minichromosomes (Wickstead *et al.*, 2004) are present in ~100 copies per cell and harbour *VSGs* at their ends, suggesting they evolved to expand the telomeric pool of *VSGs* (Barry *et al.*, 2003). Intermediate chromosomes are related to minichromosomes (Wickstead *et al.*, 2004) and contain ES. These chromosomes may also be a product of the evolutionary drive to expand the *VSG* archive, or may indicate selection for increased flexibility in ES-associated gene (*ESAG*) expression (see below). Two modes of *VSG* switching occur: recombination reactions that move silent *VSGs* into the ES, or transcriptional switches in which the single actively transcribed ES is silenced and a silent ES becomes actively transcribed. Given that the overwhelming number of archival *VSGs* are outwith the ES, recombinational switching is the main player in *T. brucei* antigenic variation.

### **Recombination mechanisms in *T. brucei* antigenic variation**

Three recombination pathways contribute to *VSG* switching. The extent to which the underlying recombination machinery is shared by these pathways, and whether or not each can occur by different recombination reactions, is being dissected. Additionally, *VSG* switching by recombination operates in a hierarchy (Capbern *et al.*, 1977; Morrison *et al.*, 2005), the basis for which is not fully understood. In the hierarchy, telomeric *VSGs* are activated earliest in infections, followed by intact subtelomeric array *VSGs*, and finally pseudogenes, which are recombined to form functional *VSG* ‘mosaics’ (Marcello and Barry, 2007; Morrison *et al.*, 2005; Thon *et al.*, 1990).

The most commonly described recombination pathway is gene conversion, where a silent, functional *VSG* is copied and transferred to the ES, replacing the resident *VSG* (Fig.1). The sequence copied in this reaction normally extends beyond the *VSG* ORF. Upstream, the boundary is often degenerate 70 bp repeats (Liu *et al.*, 1983), which flank virtually all *VSGs* (~90%) (Marcello and Barry, 2007), whereas gene conversions in which a silent ES acts as *VSG* donor can extend much further upstream

(Pays *et al.*, 1983), even including the *VSG* promoter ~50 kbp distant (Hertz-Fowler *et al.*, 2008). Downstream, gene conversion of array *VSGs* can extend to the 3' coding or non-coding parts of the gene (Bernards *et al.*, 1981), or to the chromosome end (de Lange *et al.*, 1983). A number of recombination models can account for *VSG* gene conversion. Synthesis-dependent strand-annealing (SDSA) does not generate crossovers and has been invoked (Barry, 1997; Borst *et al.*, 1996) as it avoids potentially lethal translocations when array *VSGs* are copied into the ES (Fig. 1B). However, given that a substantial minority of *VSGs* are telomeric, break-induced replication (BIR) has also been suggested (Barry and McCulloch, 2001; Dreesen *et al.*, 2007). BIR (Fig. 1D) is enzymatically and kinetically distinct from SDSA (Lydeard *et al.*, 2007), primarily in that involves repair by a processive DNA replication fork established from one end of a DNA double strand break (DSB). In yeast, BIR can stabilise telomeric repeats using subtelomeric sequences (Lydeard *et al.*, 2007), and this may be promoted in *T. brucei* *VSG* switching by the preponderance of conserved subtelomeric sequences, such as the 70 bp repeats. In addition, long-range *VSG* gene conversions (Hertz-Fowler *et al.*, 2008) may be better explained by BIR than SDSA. Nevertheless, no genetic or mechanistic dissection has shown that SDSA and BIR both act in *VSG* switching. Moreover, if both reactions do act, it is highly likely that BIR is important only for activation of telomeric *VSGs*, while SDSA can activate any gene in the archive.

A second recombination pathway has been termed reciprocal, telomeric *VSG* exchange (Pays *et al.*, 1985). Here, chromosome ends are exchanged by crossover, moving a silent telomeric *VSG* into the active ES and retaining the previously active *VSG* on the other chromosome, most likely by double strand break repair (Fig. 1B). It seems this is a minor pathway of *VSG* switching. Indeed, recent work has suggested that the ES undergo extensive recombination within the *ESAGs* upstream of the *VSG* (Hertz-Fowler *et al.*, 2008), and it is possible that *VSG* reciprocal exchange might be a by-product of this activity.

A final recombination pathway is used to generate novel, functional *VSGs* by combining portions of the ORFs from at least two pseudogenes to yield *VSG* mosaics (Thon *et al.*, 1990). A number of observations suggest this is a distinct pathway to intact *VSG* gene conversion. First, this appears to be a relatively inefficient reaction, with *VSG* mosaics observed late in infections (Marcello and Barry, 2007).

Nevertheless, the huge number of VSG pseudogenes suggests this is very important (Barbet and Kamper, 1993), expanding the VSG coat repertoire beyond that encoded directly from the genome. Second, in mosaic gene conversion recombination occurs within the VSG ORFs, rather than flanking homology, despite VSGs sharing very limited overall primary sequence homology. Third, mosaic VSG formation involves multiple donor genes, with a functional gene most likely built up sequentially by segmental gene conversions. Finally, activation of intact VSGs almost certainly targets the ES (Boothroyd *et al.*, 2009), but it is unclear if this is also the case for mosaic VSG formation. Sequencing the bloodstream stage ES repertoire in one *T. brucei* strain revealed that each site possesses a functional VSG adjacent to the telomere (Hertz-Fowler *et al.*, 2008) and it is unclear if the VSG pseudogenes sometimes observed upstream are assembly intermediates.

### **Molecular players in antigenic variation by recombination**

VSG switching, at least of intact VSGs, occurs by exploiting homologous recombination (HR), a universally conserved DNA repair mechanism (San Filippo *et al.*, 2008). Mutation of three key factors of HR has been shown to impair (not abrogate) VSG switching: RAD51, the central enzyme of HR (McCulloch and Barry, 1999), and BRCA2 (Hartley and McCulloch, 2008) and RAD51-3 (Proudfoot and McCulloch, 2005), distinct proteins that mediate RAD51 activity (San Filippo *et al.*, 2008). These data show striking parallels with *Neisseria* type IV pili antigenic variation (Hill and Davies, 2009), which occurs by gene conversion of silent, non-functional *pilS* genes to a *pilE* expression locus. This is dependent on RecA, the bacterial homologue of eukaryotic RAD51, exploiting each available subpathway of RecA-dependent HR, initiated by a RecF-like route (Mehr and Seifert, 1998) or by recBCD (though the evidence for this is controversial, and may be strain-specific)(Hill *et al.*, 2007). *T. brucei* VSG gene conversion also appears to exploit different recombination sub-pathways: residual VSG gene conversion in RAD51 mutants suggests RAD51-independent recombination can act, while four RAD51 paralogues appear to make non-equivalent contributions to VSG switching (R. Dobson, C. Stockdale and R. McCulloch, unpublished)(Proudfoot and McCulloch, 2005). Intriguingly, the contribution of mismatch repair (MMR) to antigen gene conversion in the two organisms also shows parallels. MMR recognises and repairs base mismatches, which can occur during recombination between sequence-diverged

DNA molecules, in which context MMR suppresses recombination. Surprisingly, mutation of neither *mutS* in *N. gonorrhoeae* nor the eukaryotic orthologue, MSH2, in *T. brucei* alters antigenic variation frequency, despite elevating rates of mutation and HR (Bell and McCulloch, 2003; Hill and Davies, 2009). This could suggest the gene conversion reactions are not subject to normal MMR surveillance, which for *T. brucei* could implicate an MMR-independent HR reaction (Barnes and McCulloch, 2007). Alternatively, MMR may function both to initiate and regulate antigen recombination, nullifying the phenotype of any mutation.

The use of both antigen pseudogenes and intact genes for gene conversion in a single pathogen is rare (Table 1). To date, approaches in *T. brucei* have only assayed for the recombination factors that contribute to, or have modelled experimentally (Boothroyd *et al.*, 2009), recombination of intact VSGs. Nevertheless, comparison with other pathogens may provide clues regarding mosaic VSG formation. *B. burgdorferi* antigenic variation involves recombining silent *vls* segments into a *vlsE* ES (Norris, 2006) and looks superficially similar to that of *Neisseria*, but is RecA-independent (Liveris *et al.*, 2008). The explanation for this difference may reside in the fact that *VlsE* antigenic variation involves apparently random segmental conversions from multiple *vls* donors, without any clear homology sequences guiding the reaction (Zhang and Norris, 1998). This contrasts with the ‘mini-cassette’ process of *Neisseria pilS* gene conversion (Hill and Davies, 2009), which is normally guided by upstream and downstream flanking sequence homology, similar to the recombination of intact *vlp/vsp* genes in *B. hermsii* (Dai *et al.*, 2006). In principle, therefore, *B. burgdorferi* antigenic variation appears more similar to *A. marginale msp2* gene conversion, which employs only a very small donor archive and uses segmental gene conversion to generate huge numbers of variants (Futse *et al.*, 2005). This reaction appears to be ‘anchored’ by sequence homology at one flank, but is resolved in regions of limited homology, allowing huge recombinatorial flexibility (Futse *et al.*, 2005). Two speculations seem warranted. First, it is likely that *B. hermsii* antigenic variation compares with gene conversion of intact *T. brucei* VSGs and *Neisseria pilS* genes in being driven by RecA/Rad51-dependent recombination. Second, the segmental gene conversion reactions that underlie *T. brucei* VSG and *A. marginale msp2* mosaic gene formation, like *B. burgdorferi vlsE* switching, are (at least partly) RecA/Rad51-independent. Segmental gene conversion pathways may have evolved to elaborate

antigenic variation, allowing highly protracted infections marked by huge antigen variability and allowing superinfection in partly immune hosts. What features are common to those pathogens that employ only intact genes are less clear; this is not limited to intracellular pathogens (Table 1), but may be used where a relatively precise surface antigen activation hierarchy is seen, such as in *B. hermsii* (Barbour *et al.*, 2006).

### **Initiation of VSG switching**

A question long unanswered in any pathogen, is how are recombination-based switching events initiated? DSBs within the antigen ES have been suggested as the likeliest culprit (Barry, 1997; Borst *et al.*, 1996; Hill and Davies, 2009) and recently Boothroyd *et al.* (Boothroyd *et al.*, 1980) provided evidence that that this may be so for *T. brucei* VSG switching. Controlled generation of a DSB adjacent to the 70 bp repeats in the active ES increased the rate of VSG switching and initiated VSG gene conversion. In addition, using ligation-mediated PCR, DNA breaks could be detected within the 70 bp repeats upstream of the VSG in the active ES, but much less readily (if at all) in an inactive ES. The 70 bp repeats have been predicted as the site of initiation previously (Barry, 1997; Liu *et al.*, 1983), but further work is needed to determine how they act (Barry and McCulloch, 2009). This may not be straightforward, as several observations appear at odds with the suggestion that the 70 bp repeat breaks always initiate VSG switching. First, both previous work (McCulloch *et al.*, 1997) and Boothroyd *et al.* (Boothroyd *et al.*, 2009) examined VSG switching in *T. brucei* cells expressing VSG from an ES engineered to lack 70 bp repeats. Neither study could detect differences in switch frequency compared with *T. brucei* expressing VSG from the same ES when a large array of 70 bp repeats was retained. Second, in all eukaryotes DNA breaks elicit a DNA damage response, allowing the cell to co-ordinate repair with cell cycle progression. For DSBs, a key factor is the Mre11-Rad50-Nbs1/Xrs2 complex (Harper and Elledge, 2007), but mutation of MRE11 in *T. brucei* has no discernible effect on VSG switching (Robinson *et al.*, 2002), even though the protein is clearly important for genome stability (Robinson *et al.*, 2002; Tan *et al.*, 2002). Finally, generation of a DSB appeared not to recapitulate the full diversity of VSG switching events, in that all switching reactions used telomeric VSGs as donors (>80% residing in silent ES). This raises the question of whether DSBs also elicit switching of intact array VSGs or VSG

pseudogenes. Indeed, we do not yet know if DSBs in the ES recruit RAD51 for HR, as seen elsewhere in the genome (Glover *et al.*, 2008).

### **What is the role of VSG transcriptional switching, and how is it catalysed?**

Antigenic variation relies on the expression of a single ES from many, a process termed allelic exclusion (Borst and Genest, 2006). An important context to understand this process is the observation (Navarro and Gull, 2001) that the active ES, which is transcribed by RNA polymerase (Pol) I, does not localise to the nucleolus (as expected for PolII-transcribed genes such as rRNA) but to a putative subnuclear site termed the expression site body (ESB). A model for the functioning of the ESB (whose constitution remains unknown) is that it accommodates only a single ES, explaining why attempts to select for trypanosomes expressing >1 ES yield only unstable, rapidly switching intermediates (Chaves *et al.*, 1999). Recently, a number of factors have been described whose mutation or RNAi-mediated knockdown results in derepression of transcription from the silent ES. Each factor influences chromatin structure, and in each case derepression affected differing amounts of the ES transcription units (Fig. 2). RNAi of TbISWII, a SWI2/SNF-2-related chromatin remodelling factor, leads to a ~30 fold increase in transcripts of genes proximal to the ES promoter (Hughes *et al.*, 2007). Conversely, RNAi of TbRAP1, which in yeast binds telomeres and contributes to heterochromatin formation, causes a 2-50 fold increases in *VSG* transcripts from the silent ES, but lesser derepression proximal to the promoter (Yang *et al.*, 2009). A similar telomere-mediated effect is seen in histone deacetylase TbSir2 mutants (Alsford *et al.*, 2007), but limited to the region downstream of the *VSGs*. Finally, RNAi of DOT1B, one of two *T. brucei* histone H3 methyltransferases, results in ~10 fold depression of the whole silent ES transcription unit (Figueiredo *et al.*, 2008). In addition, DOT1B RNAi caused accumulation of cells expressing two *VSGs* on the cell surface, indicating a perturbation in *VSG* coat transition during switching. These studies reveal that a number of epigenetic processes contribute to singular ES expression, but it is striking that the extent of silent ES de-repression never reveals transcription to the levels from the active site, and the frequency of *VSG* switching appears not to change (where tested). It seems, likely, therefore, that these are secondary processes that follow from the establishment of singular expression. Moreover, the factors, and potential triggers, that mediate

transcriptional switching remain elusive for trypanosomes, as for pathogens exclusively reliant on this reaction, such as *P. falciparum* (Scherf *et al.*, 2008).

A broader question about transcriptional switching in trypanosomes is the purpose of this reaction. Given the overwhelming numbers of silent *VSGs* that must be activated by recombination, does the presence of ~20 ES add much to antigenic variation, or does it contribute another function? It has been suggested that the reason for possessing multiple ES may lie in the *ESAGs*, not the *VSGs* (Bitter *et al.*, 1998; Young *et al.*, 2008). *ESAG6* and *ESAG7* form a heterodimeric receptor for host transferrin, supplying the parasite with iron. A host-range hypothesis has stemmed from this, suggesting that transcriptional switching reveals different transferrin receptors adapted to the different mammalian hosts trypanosomes infect (Bitter *et al.*, 1998). This has been extended to other *ESAGs* and, though questioned by others (Steverding, 2006), would suggest that transcriptional switching provides a means for infection establishment. However, we cannot yet exclude that transcriptional switching provides antigenic variation functions throughout a chronic infection: the silent ES may provide sites for the generation of *VSG* mosaics (Barry and McCulloch, 2001), or might provide a failsafe switch mechanism when *VSG* pseudogenes are switched into the active ES. Understanding the evolution of the *VSG* system for antigenic variation would shed light on this (Barry and McCulloch, 2001). For instance, do transcriptional and recombinational switching reactions share a common trigger?

### **Infection dynamics and antigenic variation**

The essence of the antigenic variation system in trypanosomes is to ensure survival within the mammal to allow transmission to the next host, via the tsetse vector. The system is clearly efficient, since infections in experimental cattle can last for hundreds of days (Luckins and Mehlitz, 1976), while human infections can have a prepatent period of years before clinical signs emerge (Barrett *et al.*, 2003). This picture of a chronic disease is not mirrored by the majority of the trypanosome antigenic variation literature, which, largely due to experimental restrictions, has concentrated on the first peaks of parasitaemia, essentially the initiation and establishment of infection. Despite these shortcomings, studies that have ventured into the chronic phase of infection (Capbern *et al.*, 1977; Kamper and Barbet, 1992; Marcello and Barry, 2007; Morrison *et al.*, 2005; Thon *et al.*, 1990) have revealed some intriguing insights into the dynamics and functioning of antigenic variation.

Antigenic variation occurs independently of host immune responses as a stochastic process. However, it is antibody selection that results in the observed hierarchy of VSG expression. Experimental infections and mathematical modelling suggest the first peak of parasitaemia is controlled by a combination of density dependent growth arrest and immunity (Lythgoe *et al.*, 2007; Morrison *et al.*, 2005). The sequential expression of VSGs that follows is a product of cumulative antibody responses that select against the more frequently activated VSGs as the infection progresses. As we have argued above, VSG switching is two-tiered, using one form of recombination (mosaic VSG formation) that is VSG-sequence homology-dependent, and another (gene conversion of intact VSGs) driven by sequence homology in the flanking regions. Mosaic VSG formation is relatively rare (Barry, 1997; Kamper and Barbet, 1992; Thon *et al.*, 1990) and predominates later in infection (Marcello and Barry, 2007). This difference in rate of switching between VSG classes provides some basis for understanding the VSG hierarchy, but a modelling analysis (Lythgoe *et al.*, 2007) suggests further subtleties. Within the limitations of the model, only by grouping genes into subsets with distinct ranges of switching probabilities could a realistic representation of *in vivo* infection dynamics be generated. These groupings may correspond to (a) those genes that comprise intact and functional open reading frames that are switched to by a simple single-step recombination reaction, and (b) pseudogenes requiring more than a single recombination event to create an intact ORF and functional protein. Within these probability clusters there is further hierarchy; for example, within (a), probability of switching has been experimentally demonstrated to be dependent upon the locus position of the donor sequence (Aline, Jr. *et al.*, 1985; Liu *et al.*, 1985; Morrison *et al.*, 2005; Robinson *et al.*, 1999) or (b), dependent upon the number of recombination events required to create a functionally expressed VSG, from involving few sequences (Pays *et al.*, 1983) to multiple and complex recombinations (Kamper and Barbet, 1992; Marcello and Barry, 2007). Several aspects remain unclear, however. Most prominently, we do not yet know if the transition to mosaic VSG formation involves a change in dominance of switching mechanisms *per se* (i.e. specific recombination pathway differences), mainly because recombination pathway responsible has not been characterised. The demonstration of RAD51-independent recombination in *T. brucei* requiring as little as 7-13 bp of homology (Conway *et al.*, 2002; Glover *et al.*, 2008) suggests a pathway, but this remains to be tested. There may be further mechanistic differences in VSG switching

at the species level. The VSG archive of *T. brucei* is now fairly well characterised, but those of *T. congolense* and *T. vivax* have not been similarly analysed. Genome projects are present for both (<http://www.genedb.org/>), and preliminary analysis suggests that *T. congolense* does not have the 70-bp repeat system of *T. brucei* (L. Marcello, PhD thesis), hinting that phenotypic similarities in infection profiles may hide mechanistic differences in VSG switching.

Recombination-based determinants of switch hierarchy have been described during antigenic variation in other pathogens. In *B. hermsii* the activation probability of silent (functionally intact) *vlp/vsp* genes appears to be dependent on the sequence homology and positioning, respectively, of upstream and downstream homology elements (Barbour *et al.*, 2006). *A. marginale* also displays a hierarchy, but using only pseudogenes: simple *msp2* mosaics appear preferentially early in infection and more complex mosaic genes predominate later, in the presence of a patent specific immune response to the earlier, simple mosaics (Palmer *et al.*, 2007). It has been suggested that the simple mosaic confer a fitness advantage over the complex mosaics, but this fitness advantage is balanced by the simple mosaics being more immunogenic (Zhuang *et al.*, 2007). A potential explanation for antigens expressed early in infection being more immunogenic is that if the switching probability matrix is maintained throughout infection, they will be continuously switched to, giving rise to a persistent stimulation of the immune response to those epitopes. Once the antibody response against these epitopes is patent, further switches to these commonly switched to antigens will be redundant, but the maintenance of this spectrum of switching probability throughout infection will result in the characteristic profile (in trypanosomes and *Anaplasma*) of large, early parasitaemic peaks followed by intermittent and smaller peaks (Lythgoe *et al.*, 2007)(i.e. a reduction in the effective switching rate). Variant fitness requires addressing in trypanosome chronic infections, as this may be due to cross-reactive epitopes, as suggested in *Anaplasma*, but also potentially the creation of non- (or less-) functional VSGs as the more complex (and less probable) mosaic formation occurs, and consequently reduced fitness of the trypanosome expressing that VSG. In addition, homology-driven switching is likely to generate VSGs with epitopes that are cross-reactive with patent immune responses, and such cross-reactivity has been suggested to play a vital role in determining *Plasmodium* infection dynamics (Recker *et al.*, 2004). Cross reaction between

immune responses to VSGs has not been observed in trypanosome infections (Capbern *et al.*, 1977; Morrison *et al.*, 2005; Robinson *et al.*, 1999; Van Meirvenne *et al.*, 1975), although it is a phenomenon that has not specifically been searched for. Given the recent description of loss of memory B cells during trypanosome infections (Radwanska *et al.*, 2008), it might be expected that this immunosuppression will allow increased tolerance of cross-reactive epitopes generated as the infection progresses. Loss of immunological memory may explain the reappearance of early variants late in infection observed in mice, rabbits and goats (but not cattle) (Barry, 1986). Immune selection at a population level, combined with homology-driven switching, may drive VSG archive evolution down homology-mediated routes, explaining how strain divergence in VSG repertoire develops (Hutchinson *et al.*, 2007). The continual generation of antigenic diversity at the repertoire level is essential to allow the parasite to overcome herd immunity (Futse *et al.*, 2008).

Antigenic variation undoubtedly combines with other parasite- and host-driven mechanisms to maintain chronicity. Probably the most important parasite determinant, other than the antigenic variation system, for maintaining chronic infections is the self-regulation of growth by the density-dependent differentiation of mitotically dividing long slender bloodstream form cells to non-dividing short stumpy form cells, which are pre-adapted to initiate infection in the tsetse fly midgut (Barry and McCulloch, 2001). The interplay of differentiation and antigenic variation are key (Lythgoe *et al.*, 2007; Tyler *et al.*, 2001), and the recent identification of a key marker of differentiation will allow this relationship to be examined in detail (Dean *et al.*, 2009). Further fitness advantages that allow an increased time window for a successful switch to novel variants may be at the population level and more generic, such as hydrodynamic uptake and removal of bound antibody (Engstler *et al.*, 2007), or the advantages conferred by different trypanosome transferrin receptors (Bitter *et al.*, 1998). These multiple mechanisms, along with the VSG switching dynamics, will contribute to the evolution of fitness within the host. Certainly, when superinfections with a different strain of trypanosome are attempted in an animal with an established infection, the strain that is established does seem to have a fitness advantage (Morrison *et al.*, 1982).

In summary, the availability of post-genomic tools has begun the exploration of several key aspects of antigenic variation, revealing underlying mechanisms and VSG

switching dynamics in the chronic stage of infection. Understanding this stage may explain the apparently uniquely enormous antigen repertoire in trypanosomes, and identification of the molecular mechanism of mosaic VSG formation will inform how infection time and potential for transmission is maximised. Appreciating the diversity of host-parasite interactions is important. Combining markers for both parasite (VSG switching and stumpy formation) and host (immune response and antibodies) will result in better overall understanding of how trypanosomiasis manifests as a chronic disease, and potentially provide opportunities for disease intervention.

### **Acknowledgments.**

The authors thank the Medical Research Council, Wellcome Trust and Royal Society for funding.

### **Figure legends**

#### **Figure 1. Recombination models for VSG switching by gene conversion. A**

Recombination is shown initiated by a DNA double strand break (DSB) in the 70 bp repeats (hatched) upstream of the VSG (black arrow) in the active expression site. DSB processing reveals 3' single-stranded ends, a reaction in which MRE11-RAD50-XRS2/NBS1 acts, providing a substrate for RAD51 to form a nucleoprotein filament. RAD51 function is aided by a number of factors, including BRCA2 and multiple RAD51 paralogues (not shown). RAD51 catalyses homology-dependent invasion of the single-stranded end into intact DNA duplex (grey lines) containing a silent VSG (grey arrow); mismatch repair (MMR) limits recombination to sufficiently homologous sequences. Three pathways for DSB repair can be used. **B** shows double strand break repair, where newly synthesised DNA is copied from the intact DNA duplex and remains base-paired, leading to Holliday junction recombination intermediates, whose enzymatic resolution can lead to gene conversion with (not shown) or without (shown) cross-over of flanking sequences. **C** shows synthesis-dependent strand annealing, where newly replicated DNA is displaced from the intact duplex, re-anneals with homologous sequence at the DSB, allowing synthesis of the other strand. In **D**, break-induced replication, an origin-independent replication fork,

with leading and lagging strand synthesis, forms on the strand invasion intermediate allowing replication to the chromosome end.

**Figure 2. VSG mosaic formation by segmental gene conversion.** A VSG gene (blue box) is shown within an expression site (ES), which is represented as an archetype of such a transcription unit: expression site associated genes are shown as functional (black box) or pseudogenes (grey box) and are identified by number; repeat sequences within (70 bp repeats) or upstream (50 bp repeats) of the ES are indicated by hatched boxes; the promoter is indicated by an arrow; and the telomere by a vertical line. Segmental gene conversion of three silent VSG genes from the subtelomeric array archive is diagrammed, indicating segments from each ORF being combined (arrows) to replace the transcribed VSG. The inset shows details of an experimentally verified VSG mosaic (adapted from Marcello and Barry, 2007). Three silent VSG pseudogene donor ORFs are shown below the mosaic sequence of the functional VSG (identified as an expressed cDNA): colours indicate regions of sequence homology between the donors and the cDNA, and the potential junctions in the gene conversion(s); overall sequence identity is shown.

**Figure 3. VSG switching hierarchy in *T. brucei*.** The graph shows the numbers of *T. brucei* cells (parasitaemia) measured in a cow up to 70 days post-infection (plotted as the prepatent period, in days, that a 0.2 ml inoculum of cattle blood achieves a parasitaemia of  $1 \times 10^{8.1}$  trypanosomes.ml<sup>-1</sup> units in a mouse; adapted from Morrison *et al*, 2005). Below is a depiction of the timing of VSG switch mechanism usage and gene activation. Transcriptional switching is proposed to occur at the outset of an infection, whereas recombination switching is used throughout; note, the former proposal is not experimentally verified. Silent telomeric VSGs are activated more frequently than intact, subtelomeric array VSGs, which are activated more frequently than VSG pseudogenes. During recombinational switching, VSG flanking sequence (e.g. 70 bp repeats) is proposed to mediate the VSG switching reactions of intact genes early in infections, whereas the formation of VSG mosaics relies on recombination using the genes' ORF sequences later in infections. The intensity of shading in each bar represents the effective switching rate, i.e. the proportion of switches that give rise to novel VSGs.

**Table 1. Antigenic variation systems in eukaryotic and bacterial pathogens.**

<sup>a</sup>VSG, variant surface glycoprotein; MSG, major surface glycoprotein; VSP, variant specific surface protein; VES1, variant erythrocyte surface antigen, of both  $\alpha$  and  $\beta$  types; VAR encodes *P. falciparum* erythrocyte membrane protein 1; VLP/VSP, variable large protein/variable small protein; VLS, Vmp-like sequence; PIL encodes the type IV pilus; MSP2, major surface protein 2.

<sup>b</sup> N/A; not applicable

## References

- Aline, R. F., Jr., Scholler, J. K., Nelson, R. G., Agabian, N., and Stuart, K. (1985) Preferential activation of telomeric variant surface glycoprotein genes in *Trypanosoma brucei*. *Mol Biochem Parasitol* **17**: 311-320.
- Alsford, S., Kawahara, T., Isamah, C., and Horn, D. (2007) A sirtuin in the African trypanosome is involved in both DNA repair and telomeric gene silencing but is not required for antigenic variation. *Mol Microbiol* **63**: 724-736.
- Barbet, A. F. and Kamper, S. M. (1993) The importance of mosaic genes to trypanosome survival. *Parasitol Today* **9**: 63-66.
- Barbour, A. G., Dai, Q., Restrepo, B. I., Stoenner, H. G., and Frank, S. A. (13-11-2006) Pathogen escape from host immunity by a genome program for antigenic variation. *Proc Natl Acad Sci U S A*.
- Barnes, R. L. and McCulloch, R. (2007) *Trypanosoma brucei* homologous recombination is dependent on substrate length and homology, though displays a differential dependence on mismatch repair as substrate length decreases. *Nucleic Acids Res* **35**: 3478-3493.
- Barrett, M. P., Burchmore, R. J., Stich, A., Lazzari, J. O., Frasch, A. C., Cazzulo, J. J., and Krishna, S. (1-11-2003) The trypanosomiases. *Lancet* **362**: 1469-1480.
- Barry, D. and McCulloch, R. (14-5-2009) Molecular microbiology: A key event in survival. *Nature* **459**: 172-173.
- Barry, J. D. (1986) Antigenic variation during *Trypanosoma vivax* infections of different host species. *Parasitology* **92** ( Pt 1): 51-65.
- Barry, J. D. (1997) The relative significance of mechanisms of antigenic variation in African trypanosomes. *Parasitol Today* **13**: 212-218.

- Barry, J. D., Ginger, M. L., Burton, P., and McCulloch, R. (2003) Why are parasite contingency genes often associated with telomeres? *Int J Parasitol* **33**: 29-45.
- Barry, J. D. and McCulloch, R. (2001) Antigenic variation in trypanosomes: enhanced phenotypic variation in a eukaryotic parasite. *Adv Parasitol* **49**: 1-70.
- Bell, J. S. and McCulloch, R. (14-11-2003) Mismatch repair regulates homologous recombination, but has little influence on antigenic variation, in *Trypanosoma brucei*. *J Biol Chem* **278**: 45182-45188.
- Bernards, A., Van der Ploeg, L. H., Frasnich, A. C., Borst, P., Boothroyd, J. C., Coleman, S., and Cross, G. A. (1981) Activation of trypanosome surface glycoprotein genes involves a duplication-transposition leading to an altered 3' end. *Cell* **27**: 497-505.
- Berriman, M., Ghedin, E., Hertz-Fowler, C., Blandin, G., Renaud, H., Bartholomeu, D. C., Lennard, N. J., Caler, E., Hamlin, N. E., Haas, B., Bohme, U., Hannick, L., Aslett, M. A., Shallom, J., Marcello, L., Hou, L., Wickstead, B., Alsmark, U. C., Arrowsmith, C., Atkin, R. J., Barron, A. J., Bringaud, F., Brooks, K., Carrington, M., Cherevach, I., Chillingworth, T. J., Churcher, C., Clark, L. N., Corton, C. H., Cronin, A., Davies, R. M., Doggett, J., Djikeng, A., Feldblyum, T., Field, M. C., Fraser, A., Goodhead, I., Hance, Z., Harper, D., Harris, B. R., Hauser, H., Hostetler, J., Ivens, A., Jagels, K., Johnson, D., Johnson, J., Jones, K., Kerhornou, A. X., Koo, H., Larke, N., Landfear, S., Larkin, C., Leech, V., Line, A., Lord, A., MacLeod, A., Mooney, P. J., Moule, S., Martin, D. M., Morgan, G. W., Mungall, K., Norbertczak, H., Ormond, D., Pai, G., Peacock, C. S., Peterson, J., Quail, M. A., Rabbinowitsch, E., Rajandream, M. A., Reitter, C., Salzberg, S. L., Sanders, M., Schobel, S., Sharp, S., Simmonds, M., Simpson, A. J., Tallon, L., Turner, C. M., Tait, A., Tivey, A. R., Van Aken, S., Walker, D., Wanless, D., Wang, S., White, B., White, O., Whitehead, S., Woodward, J., Wortman, J., Adams, M. D., Embley, T. M., Gull, K., Ullu, E., Barry, J. D., Fairlamb, A. H., Opperdoes, F., Barrell, B. G., Donelson, J. E., Hall, N., Fraser, C. M., Melville, S. E., and El Sayed, N. M. (15-7-2005) The genome of the African trypanosome *Trypanosoma brucei*. *Science* **309**: 416-422.
- Bitter, W., Gerrits, H., Kieft, R., and Borst, P. (29-1-1998) The role of transferrin-receptor variation in the host range of *Trypanosoma brucei*. *Nature* **391**: 499-502.
- Boothroyd, C. E., Dreesen, O., Leonova, T., Ly, K. I., Figueiredo, L. M., Cross, G. A., and Papavasiliou, F. N. (14-5-2009) A yeast-endonuclease-generated DNA break induces antigenic switching in *Trypanosoma brucei*. *Nature* **459**: 278-281.
- Boothroyd, J. C., Cross, G. A., Hoeijmakers, J. H., and Borst, P. (1980) A variant surface glycoprotein of *Trypanosoma brucei* synthesized with a C-terminal hydrophobic 'tail' absent from purified glycoprotein. *Nature* **288**: 624-626.
- Borst, P. and Genest, P. A. (23-2-2006) Parasitology: switching like for like. *Nature* **439**: 926-927.
- Borst, P., Rudenko, G., Taylor, M. C., Blundell, P. A., van Leeuwen, F., Bitter, W., Cross, M., and McCulloch, R. (1996) Antigenic variation in trypanosomes. *Arch Med Res* **27**: 379-388.

Capbern, A., Giroud, C., Baltz, T., and Mattern, P. (1977) [Trypanosoma equiperdum: antigenic variations in experimental trypanosomiasis of rabbits]. *Exp Parasitol* **42**: 6-13.

Chaves, I., Rudenko, G., Dirks-Mulder, A., Cross, M., and Borst, P. (1-9-1999) Control of variant surface glycoprotein gene-expression sites in Trypanosoma brucei. *EMBO J* **18**: 4846-4855.

Conway, C., Proudfoot, C., Burton, P., Barry, J. D., and McCulloch, R. (2002) Two pathways of homologous recombination in Trypanosoma brucei. *Mol Microbiol* **45**: 1687-1700.

Dai, Q., Restrepo, B. I., Porcella, S. F., Raffel, S. J., Schwan, T. G., and Barbour, A. G. (2006) Antigenic variation by Borrelia hermsii occurs through recombination between extragenic repetitive elements on linear plasmids. *Mol Microbiol* **60**: 1329-1343.

de Lange, T., Kooter, J. M., Michels, P. A., and Borst, P. (10-12-1983) Telomere conversion in trypanosomes. *Nucleic Acids Res* **11**: 8149-8165.

Dean, S., Marchetti, R., Kirk, K., and Matthews, K. R. (14-5-2009) A surface transporter family conveys the trypanosome differentiation signal. *Nature* **459**: 213-217.

Deitsch, K. W., Lukehart, S. A., and Stringer, J. R. (2009) Common strategies for antigenic variation by bacterial, fungal and protozoan pathogens. *Nat Rev Microbiol* **7**: 493-503.

Dreesen, O., Li, B., and Cross, G. A. (2007) Telomere structure and function in trypanosomes: a proposal. *Nat Rev Microbiol* **5**: 70-75.

Engstler, M., Pfohl, T., Herminghaus, S., Boshart, M., Wiegertjes, G., Heddergott, N., and Overath, P. (2-11-2007) Hydrodynamic flow-mediated protein sorting on the cell surface of trypanosomes. *Cell* **131**: 505-515.

Figueiredo, L. M., Janzen, C. J., and Cross, G. A. (1-7-2008) A histone methyltransferase modulates antigenic variation in African trypanosomes. *PLoS Biol* **6**: e161.

Futse, J. E., Brayton, K. A., Dark, M. J., Knowles, D. P., Jr., and Palmer, G. H. (12-2-2008) Superinfection as a driver of genomic diversification in antigenically variant pathogens. *Proc Natl Acad Sci U S A* **105**: 2123-2127.

Futse, J. E., Brayton, K. A., Knowles, D. P., Jr., and Palmer, G. H. (2005) Structural basis for segmental gene conversion in generation of Anaplasma marginale outer membrane protein variants. *Mol Microbiol* **57**: 212-221.

Glover, L., McCulloch, R., and Horn, D. (2008) Sequence homology and microhomology dominate chromosomal double-strand break repair in African trypanosomes. *Nucleic Acids Res* **36**: 2608-2618.

- Harper, J. W. and Elledge, S. J. (14-12-2007) The DNA damage response: ten years after. *Mol Cell* **28**: 739-745.
- Hartley, C. L. and McCulloch, R. (2008) Trypanosoma brucei BRCA2 acts in antigenic variation and has undergone a recent expansion in BRC repeat number that is important during homologous recombination. *Mol Microbiol* **68**: 1237-1251.
- Hertz-Fowler, C., Figueiredo, L. M., Quail, M. A., Becker, M., Jackson, A., Bason, N., Brooks, K., Churcher, C., Fahkro, S., Goodhead, I., Heath, P., Kartvelishvili, M., Mungall, K., Harris, D., Hauser, H., Sanders, M., Saunders, D., Seeger, K., Sharp, S., Taylor, J. E., Walker, D., White, B., Young, R., Cross, G. A., Rudenko, G., Barry, J. D., Louis, E. J., and Berriman, M. (2008) Telomeric expression sites are highly conserved in Trypanosoma brucei. *PLoS ONE* **3**: e3527.
- Hill, S. A. and Davies, J. K. (2009) Pilin gene variation in Neisseria gonorrhoeae: reassessing the old paradigms. *FEMS Microbiol Rev* **33**: 521-530.
- Hill, S. A., Woodward, T., Reger, A., Baker, R., and Dinse, T. (2007) Role for the RecBCD recombination pathway for pilE gene variation in repair-proficient Neisseria gonorrhoeae. *J Bacteriol* **189**: 7983-7990.
- Hughes, K., Wand, M., Foulston, L., Young, R., Harley, K., Terry, S., Ersfeld, K., and Rudenko, G. (2-5-2007) A novel ISWI is involved in VSG expression site downregulation in African trypanosomes. *EMBO J* **26**: 2400-2410.
- Hutchinson, O. C., Picozzi, K., Jones, N. G., Mott, H., Sharma, R., Welburn, S. C., and Carrington, M. (2007) Variant Surface Glycoprotein gene repertoires in Trypanosoma brucei have diverged to become strain-specific. *BMC Genomics* **8**: 234.
- Kamper, S. M. and Barbet, A. F. (1992) Surface epitope variation via mosaic gene formation is potential key to long-term survival of Trypanosoma brucei. *Mol Biochem Parasitol* **53**: 33-44.
- Liu, A. Y., Michels, P. A., Bernards, A., and Borst, P. (5-4-1985) Trypanosome variant surface glycoprotein genes expressed early in infection. *J Mol Biol* **182**: 383-396.
- Liu, A. Y., Van der Ploeg, L. H., Rijsewijk, F. A., and Borst, P. (15-6-1983) The transposition unit of variant surface glycoprotein gene 118 of Trypanosoma brucei. Presence of repeated elements at its border and absence of promoter-associated sequences. *J Mol Biol* **167**: 57-75.
- Liveris, D., Mulay, V., Sandigursky, S., and Schwartz, I. (2008) Borrelia burgdorferi vlsE antigenic variation is not mediated by RecA. *Infect Immun* **76**: 4009-4018.
- Luckins, A. G. and Mehlitz, D. (1976) Observations on serum immunoglobulin levels in cattle infected with Trypanosoma brucei, T. vivax and T. congolense. *Ann Trop Med Parasitol* **70**: 479-480.
- Lydeard, J. R., Jain, S., Yamaguchi, M., and Haber, J. E. (16-8-2007) Break-induced replication and telomerase-independent telomere maintenance require Pol32. *Nature* **448**: 820-823.

- Lythgoe, K. A., Morrison, L. J., Read, A. F., and Barry, J. D. (8-5-2007) Parasite-intrinsic factors can explain ordered progression of trypanosome antigenic variation. *Proc Natl Acad Sci U S A* **104**: 8095-8100.
- Marcello, L. and Barry, J. D. (2007) Analysis of the VSG gene silent archive in *Trypanosoma brucei* reveals that mosaic gene expression is prominent in antigenic variation and is favored by archive substructure. *Genome Res* **17**: 1344-1352.
- McCulloch, R. and Barry, J. D. (1-11-1999) A role for RAD51 and homologous recombination in *Trypanosoma brucei* antigenic variation. *Genes Dev* **13**: 2875-2888.
- McCulloch, R., Rudenko, G., and Borst, P. (1997) Gene conversions mediating antigenic variation in *Trypanosoma brucei* can occur in variant surface glycoprotein expression sites lacking 70- base-pair repeat sequences. *Mol Cell Biol* **17**: 833-843.
- Mehr, I. J. and Seifert, H. S. (1998) Differential roles of homologous recombination pathways in *Neisseria gonorrhoeae* pilin antigenic variation, DNA transformation and DNA repair. *Mol Microbiol* **30**: 697-710.
- Morrison, L. J., Majiwa, P., Read, A. F., and Barry, J. D. (2005) Probabilistic order in antigenic variation of *Trypanosoma brucei*. *Int J Parasitol* **35**: 961-972.
- Morrison, W. I., Wells, P. W., Moloo, S. K., Paris, J., and Murray, M. (1982) Interference in the establishment of superinfections with *Trypanosoma congolense* in cattle. *J Parasitol* **68**: 755-764.
- Moxon, E. R., Rainey, P. B., Nowak, M. A., and Lenski, R. E. (1-1-1994) Adaptive evolution of highly mutable loci in pathogenic bacteria. *Curr Biol* **4**: 24-33.
- Navarro, M. and Gull, K. (13-12-2001) A pol I transcriptional body associated with VSG mono-allelic expression in *Trypanosoma brucei*. *Nature* **414**: 759-763.
- Norris, S. J. (2006) Antigenic variation with a twist--the *Borrelia* story. *Mol Microbiol* **60**: 1319-1322.
- Palmer, G. H., Futse, J. E., Leverich, C. K., Knowles, D. P., Jr., Rurangirwa, F. R., and Brayton, K. A. (2007) Selection for simple major surface protein 2 variants during *Anaplasma marginale* transmission to immunologically naive animals. *Infect Immun* **75**: 1502-1506.
- Pays, E., Guyaux, M., Aerts, D., Van Meirvenne, N., and Steinert, M. (8-8-1985) Telomeric reciprocal recombination as a possible mechanism for antigenic variation in trypanosomes. *Nature* **316**: 562-564.
- Pays, E., Van Assel, S., Laurent, M., Dero, B., Michiels, F., Kronenberger, P., Matthyssens, G., Van Meirvenne, N., Le Ray, D., and Steinert, M. (1983) At least two transposed sequences are associated in the expression site of a surface antigen gene in different trypanosome clones. *Cell* **34**: 359-369.
- Proudfoot, C. and McCulloch, R. (2005) Distinct roles for two RAD51-related genes in *Trypanosoma brucei* antigenic variation. *Nucleic Acids Res* **33**: 6906-6919.


- Radwanska, M., Guirnalda, P., De Trez, C., Ryffel, B., Black, S., and Magez, S. (2008) Trypanosomiasis-induced B cell apoptosis results in loss of protective anti-parasite antibody responses and abolishment of vaccine-induced memory responses. *PLoS Pathog* **4**: e1000078.
- Recker, M., Nee, S., Bull, P. C., Kinyanjui, S., Marsh, K., Newbold, C., and Gupta, S. (3-6-2004) Transient cross-reactive immune responses can orchestrate antigenic variation in malaria. *Nature* **429**: 555-558.
- Robinson, N. P., Burman, N., Melville, S. E., and Barry, J. D. (1999) Predominance of duplicative VSG gene conversion in antigenic variation in African trypanosomes. *Mol Cell Biol* **19**: 5839-5846.
- Robinson, N. P., McCulloch, R., Conway, C., Browitt, A., and Barry, J. D. (19-7-2002) Inactivation of Mre11 Does Not Affect VSG Gene Duplication Mediated by Homologous Recombination in *Trypanosoma brucei*. *J Biol Chem* **277**: 26185-26193.
- San Filippo, J., Sung, P., and Klein, H. (14-2-2008) Mechanism of Eukaryotic Homologous Recombination. *Annu Rev Biochem*.
- Scherf, A., Lopez-Rubio, J. J., and Riviere, L. (2008) Antigenic variation in *Plasmodium falciparum*. *Annu Rev Microbiol* **62**: 445-470.
- Steverding, D. (2006) On the significance of host antibody response to the *Trypanosoma brucei* transferrin receptor during chronic infection. *Microbes Infect* **8**: 2777-2782.
- Tan, K. S., Leal, S. T., and Cross, G. A. (2002) *Trypanosoma brucei* MRE11 is non-essential but influences growth, homologous recombination and DNA double-strand break repair. *Mol Biochem Parasitol* **125**: 11-21.
- Thon, G., Baltz, T., Giroud, C., and Eisen, H. (1990) Trypanosome variable surface glycoproteins: composite genes and order of expression. *Genes Dev* **4**: 1374-1383.
- Turner, C. M. and Barry, J. D. (1989) High frequency of antigenic variation in *Trypanosoma brucei rhodesiense* infections. *Parasitology* **99 Pt 1**: 67-75.
- Tyler, K. M., Higgs, P. G., Matthews, K. R., and Gull, K. (7-11-2001) Limitation of *Trypanosoma brucei* parasitaemia results from density-dependent parasite differentiation and parasite killing by the host immune response. *Proc Biol Sci* **268**: 2235-2243.
- Van Meirvenne, N., Janssens, P. G., Magnus, E., Lumsden, W. H., and Herbert, W. J. (1975) Antigenic variation in syringe passaged populations of *Trypanosoma* (Trypanozoon) *brucei*. II. Comparative studies on two antigenic-type collections. *Ann Soc Belg Med Trop* **55**: 25-30.
- Wickstead, B., Ersfeld, K., and Gull, K. (2004) The small chromosomes of *Trypanosoma brucei* involved in antigenic variation are constructed around repetitive palindromes. *Genome Res* **14**: 1014-1024.



Yang, X., Figueiredo, L. M., Espinal, A., Okubo, E., and Li, B. (3-4-2009) RAP1 is essential for silencing telomeric variant surface glycoprotein genes in *Trypanosoma brucei*. *Cell* **137**: 99-109.



Young, R., Taylor, J. E., Kurioka, A., Becker, M., Louis, E. J., and Rudenko, G. (2008) Isolation and analysis of the genetic diversity of repertoires of VSG expression site containing telomeres from *Trypanosoma brucei gambiense*, *T. b. brucei* and *T. equiperdum*. *BMC Genomics* **9**: 385.

Zhang, J. R. and Norris, S. J. (1998) Genetic variation of the *Borrelia burgdorferi* gene *vlsE* involves cassette-specific, segmental gene conversion. *Infect Immun* **66**: 3698-3704.

Zhuang, Y., Futse, J. E., Brown, W. C., Brayton, K. A., and Palmer, G. H. (2007) Maintenance of antibody to pathogen epitopes generated by segmental gene conversion is highly dynamic during long-term persistent infection. *Infect Immun* **75**: 5185-5190.







Pathogen	Infection strategy	Surface antigen <sup>a</sup>	Gene no. in archive	% of archive pseudogene	Expression site no.	Switch mechanism	RecA/Rad51 dependency <sup>b</sup>
<i>T. brucei</i>	extracellular	VSG	~1600	85	14-23	recombination and transcription	Rad51-dependent and -independent
<i>P. carinii</i>	intracellular	MSG	~80	0	1	recombination	?
<i>G. lamblia</i>	extracellular	VSP	~150	0	~150	transcription	N/A
<i>B. bovis</i>	intracellular	VES1	~150	~50	24-42	recombination and transcription	?
<i>P. falciparum</i>	intracellular	VAR	~60	0	~60	transcription	N/A
<i>B. hermsii</i>	extracellular	VLP/VSP	~60	0	1	recombination	?
<i>B. burgdorferi</i>	extracellular	VLS	~15	100	1	recombination	RecA-independent
<i>N. gonorrhoeae</i>	intracellular	PIL	~20	100	1-2	recombination	RecA-dependent
<i>A. marginale</i>	intracellular	MSP2	<10	100	1	recombination	?