

Long-Term Variations of the Electron Slot Region and Global Radiation Belt Structure

Shing F. Fung¹, Xi Shao^{1,2}, and Lun C. Tan^{1,3}

¹Space Physics Data Facility, NASA Goddard Space Flight Center, Greenbelt, MD 20771

²NAS National Research Council, Washington DC

³QSS Group, Inc., Lanham, MD

Submitted to the *Geophysical Research Letters*

October 7, 2005

Abstract

We report the observations of changes of the nominal position of the quiet-time radiation belt slot over the solar cycles. It has been found that the slot region, believed to be a result of enhanced precipitation losses of energetic electrons due to their interactions with VLF waves in the magnetosphere, tends to shift to higher L (~ 3) during a solar maximum compared to its canonical L value of ~ 2.5 , which is more typical of a solar minimum. The solar-cycle migration of the slot can be understood in terms of the solar-cycle changes in ionospheric densities, which may cause the optimal wave-particle interaction region during higher solar activity periods to move to higher altitudes *and* higher latitudes, thus higher L . Our analysis also suggests that the primary wave-particle interaction processes that result in the slot formation are located off of the magnetic equator.

Introduction

The *Van Allen* radiation belts are permanent features of Earth's inner magnetosphere, although their structures are known to vary with geomagnetic activities. Enhancements of the radiation belts have been observed to occur during geomagnetic storms. New belt formations have also been observed during strong storms. Subsequent to the main phase of a storm, the electron slot region, believed to result from losses of trapped electrons caused by precipitation induced by pitch-angle scattering by VLF waves in the plasmasphere, can be completely filled with energetic electrons, leading to the disappearance of the slot and the overall enhancement of the outer-belt electron fluxes. The slot then recovers after the storm has decayed sufficiently, typically a few days after the main phase. The slot region, nominally located at $2 < L < 3$, thus is a salient feature of the radiation belts, separating the inner and outer belts.

Due to the potential hazards of energetic electrons on orbiting space systems, studies of radiation belt have focused on the sources and acceleration mechanisms that might lead to enhancements of energetic electron fluxes. Much less attention has been paid to the particle loss mechanisms or the slot region. The presence of the slot region is important for spacecraft operations as it provides a region in space with a range of L values that is relatively free of hazardous energetic electrons. In addition, the processes involved in the creation of the slot are also connected intimately to the decay of a geomagnetic storm. Understanding the development and maintenance of the slot, *i.e.* processes leading to the production of VLF waves in the slot L range, are therefore important for delineating the physics of post-storm magnetospheric dynamics.

We have investigated the long-term (solar-cycle) behavior of the electron slot region in order to elucidate the magnetospheric conditions controlling the long-term slot structure. The

main topic of this paper is to describe the apparent shifts of the slot region observed at low altitudes (~ 850 km) as observed by different *NOAA* satellites over the solar-cycle time scale [*Fung et al.*, 2005], investigate the probable causes of that shift and its relationship with changes in ionospheric densities and the VLF wave pattern observed at high altitudes.

Formation of the Quiet-Time Electron Slot Region

There may be little disagreement with the notion that the radiation belt slot region is formed by losses of energetic electrons by enhanced pitch-angle scattering by VLF waves, most likely whistler waves associated with plasmaspheric hiss emission [e.g., *Lyons and Thorne*, 1973; *Lyons and Williams*, 1984, *Imhof et al.*, 1982]. There have been many discussions, however, on the sources and distributions of the wave activities responsible for the scattering of energetic electrons in the slot [e.g., *Abel and Thorne*, 1984a,b; *Imhof et al.*, 1986; *Bortnik et al.*, 2002; 2003; *Green et al.*, 2005a,b]. It is remarkable that they all seem to predict the canonical position of the slot ($2 < L < 3$), making it difficult to observationally distinguish the dominant slot formation process.

Abel and Thorne [1998a,b] conducted relatively extensive studies of pitch angle diffusion and calculated electron precipitation lifetimes resulting from various scattering processes due to Coulomb interactions with thermal plasma, plasmaspheric hiss, lightning-generated whistlers, and VLF transmitter waves. They found that the slot occurrence is basically accounted for by some appropriate combinations of the various scattering processes. They also found, however, that these processes can also be affected by the average propagation characteristics of the scattering waves. As the plasma and wave characteristics are likely to be affected by magnetospheric conditions, it is then of interest to determine the quiet-time equilibrium structure

of the slot without the transient effects of dynamical processes, so that the “ground-state” structure of the radiation belt structure can be ascertained.

Solar-Cycle Variations of the Electron Slot Region

Except during strong storms when the slot region can be filled temporarily or new belts can form, the slot is a salient feature of the *Van Allen* radiation belts, most noticeable during geomagnetically quiet times. While most radiation belt studies tend to focus on the outer or inner belt [e.g., *Li et al.*, 2001; *Friedel and Korth*, 1995], long-term variations of the slot region have not received much attention. Since the slot region is relatively devoid of hazardous radiation fluxes and can thus be a safe zone for Earth orbiting satellites, we need to examine if there exist any long-term variations of the slot that may shed light on the maintenance of the global radiation belt structures.

Fung et al [2005] show recently that there is a noticeable shift in the average slot location from $L \sim 2.5$ during solar minimum to $L \sim 3$ during solar maximum. A similar shift is noted in *Green et al* [2005]. Figure 1 shows the L - B/B_0 profiles of omnidirectional electron fluxes observed during geomagnetically quiet times in the months of June, July and August of 1980 and 1990 for solar maximum, and of 1986 and 1996 for solar minimum. The particle observations were obtained by the *Medium Energy Proton Electron Detectors (MEPED)* [*Raben et al.*, 1995] aboard different polar-orbiting, low-altitude *NOAA* satellites. The quiet intervals were obtained by using the *Magnetospheric State Query System (MSQS)*, radbelts.gsfc.nasa.gov/RB_model_int/Psi_database.html; *Fung*, [2004]) for the conditions of $K_p \leq 2$ and solar wind speeds $\leq 400 \text{ km s}^{-1}$.

As shown clearly in the energetic electron (> 300 keV) observations in Figure 1, there is an apparent shift in the nominal slot location between higher and lower L as the solar cycle phase changes from solar maximum to solar minimum condition, and vice versa. While a slot-like feature is seen at lower energies (> 100 keV), no apparent solar cycle-phase shift is seen in the less energetic electron observations.

Figure 2 shows the count rates, instead of omnidirectional fluxes, observed by the *NOAA* satellites years three consecutive years during a solar maximum (1979, 1989, and 1981) and solar minimum (1984, 1985, and 1986) period. Again to avoid seasonal variations, only observations from the same summer months: May, June and July from each year are plotted. The consistency in the quiet-time slot locations in the three consecutive years in a given solar cycle phase (solar minimum or solar maximum) is striking. The shift in the slot locations due to solar cycle changes, however, is quite apparent in the >300 keV observations. Figure 2 also shows that the slot at successively lower energies tends to occur at higher and higher L , although they are not affected by solar cycle phases.

Solar-cycle variations of the quiet-time minimum electron flux location (> 300 keV) in the slot and the corresponding minimum fluxes are shown in the lower and upper panels in Figure 3, respectively. The data are all taken in the months of June, July, and August during each year except that in 1988, the data is taken from the entire year since NOAA10 has poor coverage. The quiet time intervals were again obtained by using the *MSQS* as above. As seen from the lower panel of Figure 3, the L value of the minimum flux in the slot correlates nearly perfectly with solar activity. On the other hand, the upper panel shows that the slot minimum flux reaches a relative minimum near the beginning of the rising phase of a solar cycle, and it tends to maximize at the beginning of the declining phase of the solar cycle.

Discussions

The quiet-time slot structures (Figure 3) found in this paper represent the equilibrium states of the radiation belt slot during solar maximum and solar minimum conditions, when all transient effects have subsided. While it may be expected that similar pitch angle scattering processes are responsible for the creation of the slot during different solar cycle phases, the degree to which the slot is evacuated and the primary location of the electron-loss processes are apparently solar-cycle dependent.

From the results shown in Figures 1-3, it can be inferred that the primary site of precipitation losses of energetic electrons (>300 keV), hence the slot, must shift from $L \sim 2.5$ to slightly higher L (to ~ 3) as solar activity level changes from a minimum to a maximum. This seemingly oscillatory motion of the slot appears cyclical and is nearly in-phase with the 11-year solar cycle. This strong correlation between the slot position in L and solar activity suggests that the distributions of the whistler waves responsible for scattering and of the energetic electrons being scattered must somehow be controlled by the ionospheric density, which is higher and has a larger scale height during solar maximum than solar minimum [e.g., *Jursa*, 1985].

As shown in Figures 1 and 2, the outer-belt energetic electron (>300 keV) distributions during solar minimum (quiet) conditions tend to be broader in L and have less well-defined peaks. But the distribution in 1984 (top right panel for solar minimum in Figure 2) does show evidence of an outer-belt peak location that is consistent with those observed during solar maximum, such as in 1979 (top left panel in Figure 2). The inner-belt peak location ($L \sim 1.8$ for > 300 keV) is relatively insensitive to solar cycle phase variations. Therefore, the observed shift of the slot as a function of solar cycle phase is not likely due to any major shifts in the spatial

distributions of trapped electrons in the radiation belts, but is probably due to the shifts in the wave distributions.

It may be instructive to consider how increases in ionospheric densities and scale heights during a solar activity maximum may lead to shifting of the slot to higher L values. Figure 4 shows a schematic of a given density level ($N = N_f$) during a solar minimum (inner dotted contour) and a solar maximum (outer dotted contour) superposed over the Earth's magnetic field (solid lines). Points A and B are the sites where precipitation losses of energetic electrons are effective due to pitch angle scattering by waves in a solar minimum and solar maximum period, respectively.

For VLF waves that are effective in pitch angle scattering the slot electrons [e.g., *Abel and Thorne, 1998a,b; Bortnik et al., 2002; Green et al., 2005a,b*], their propagations are mediated by the background magnetized plasma that is generally characterized by its electron plasma frequency ω_{pe} and gyro frequency Ω_{ce} . Therefore the specific ratio of ω_{pe}/Ω_{ce} at A may be considered as the plasma condition appropriate for the occurrence of enhanced wave-particle interactions during a solar minimum condition. Due to the overall increases in the ionospheric electron densities and scale heights at solar maximum, the region where the same plasma condition occurs must migrate to point B at higher altitudes and on a field line with slightly higher L in order to maintain the same Ω_{ce} , consistent with the observed shift of the slot between a solar minimum and a solar maximum.

One point to note from Figure 4 is that the regions of enhanced scattering losses, A and B , cannot be too close to the magnetic equator (at least for quiet times) as the magnetic field strengths there only decrease with radial distances. The finite latitudinal change between A and B is a direct consequence of maintaining the same plasma and magnetic field conditions in order

for the optimal occurrences of the same scattering processes to form the slot in different solar cycles. In addition, the scattering losses seem to be more effective at the beginning of a rising phase of a solar cycle and are least effective at the beginning of a declining phase, as suggested by the observed long-term variations of the residual energetic electron fluxes in the slot over the solar activity cycles (lower panel of Figure 3).

Finally, we need to point out that movements of the slot over the solar cycle time scale in a sense *opposite* to what is described in this paper has been noted previously. *Vernov et al.* [1969] analyzed multiple short-term datasets obtained by different platforms in ~1958-1965 with various threshold energies in the range of >100 keV to > 1 MeV and noted that the radiation belt gap tends to migrate to higher L as solar activity *decreases*. This result is clearly inconsistent with what is being reported here, although *Vernov et al.* [1969] also pointed out that “some scatter of points may be due to time variations and to the differences between electron energies measured in various experiments.” The long-term *NOAA* data sets used in the present study may be better suited for the type of study being reported and therefore may provide a more accurate characterization of the solar-cycle changes of the radiation belt slot.

Conclusions

After analyzing the long-term *NOAA* satellite (5, 6, 8, 10, and 15) observations at low altitude (~850 km), we have found that the quiet-time slot region of the Earth’s radiation belts exhibits an oscillatory motion in L that is connected to the solar-cycle variations. The slot is found to be located at slightly higher L (~ 3) during a solar maximum with a higher residual slot energetic electron fluxes (> 300 keV) occurring at the end of the corresponding solar maximum.

By comparison, at solar minimum, the slot is located at lower L (~ 2.5) and is more depleted with energetic electrons near the end of the solar minimum.

The reported findings can be understood in terms of the migrations of the enhanced precipitation loss region with the changes of ionospheric densities and scale heights between solar cycle phases. They suggest that the primary wave-particle interaction region is situated along the mid-latitude field lines and not at the magnetic equator. This supports the importance of incorporating wave activity distributions along the geomagnetic field lines when modeling the geomagnetic cutoffs of trapped particles, as demonstrated by *Boscher et al.* [1997].

Acknowledgements

We thank the *NOAA National Geophysical Data Center (NGDC)* for making the *NOAA* datasets publicly accessible. This work was partially supported by *NASA* contract NAS 5-97057 (LCT) and the *National Research Council Research Associate Program (XS)*.

References

- Abel B. and R. M. Thorne, Electron scattering loss in Earth's inner magnetosphere: 1, dominant physical processes, *J. Geophys. Res.*, **103**, 2385-2396, 1998a.
- Abel, B., and R. M. Thorne, Electron scattering loss in Earth's inner magnetosphere: 2, sensitivity to model parameters, *J. Geophys. Res.*, **103**, 2397-2407, 1998b.
- Boscher, D. S. F. Fung, and L. C. Tan, Spatial distributions of the inner radiation belt electrons: a comparison between observations and radial diffusion theory predictions, *Adv. Space Res.*, **20**, 369-372, 1997.

- Bortnik, J., U. S. Inan, and T. F. Bell, Energy distribution and lifetime of magnetospherically reflecting whistlers in the plasmasphere, *J. Geophys. Res.*, **108**(A5), 1199, doi:10.1029/2002JA009316, 2003.
- Bortnik, J., U. S. Inan, and T. F. Bell, L-dependence of energetic electron precipitation driven by magnetospherically reflecting whistle waves, *J. Geophys. Res.*, **107**(A8), 1150, doi:10.1029/2001JA000303, 2002.
- Friedel, R. H. W., and A. Korth, Long-term observations of keV ion and electron variability in the outer radiation belt from CRRES, *Geophys. Res., Letts.*, **22**, pp. 1853-1856, 1995.
- Fung, S. F., Announcement of a "Magnetospheric-State Query System", *American Geophysical Union SPA Section Newsletter*, vol. XI, Issue 9, ftp://igpp.ucla.edu/scratch/aguspa/volume11_2004/vol11no009, January 28, 2004.
- Fung, S. F., E. V. Bell, L. C. Tan, R. M. Candey, M. J. Golightly, S. L. Huston, J. H. King, and R. E. McGuire, Development of A Magnetospheric State-Based Trapped Radiation Data Base, *Adv. Space Research*, in press, 2005.
- Green, J. L., S. F. Fung, S. Boardsen, and H. J. Christian, Distribution and origin of plasmaspheric plasma waves, in *Inner Magnetosphere Interactions: New Perspectives from Imaging*, edited by James L. Burch, Michael Schulz, and Harlan Spence, Geophysical Monograph Series, **159**, Chapter 6, AGU Washington D. C., 2005a.
- Green, J. L., S. Boardsen, L. Garcia, W. W. L. Taylor, S. F. Fung, and B. W. Reinisch (2005b), On the origin of whistler mode radiation in the plasmasphere, *J. Geophys. Res.*, **110**, A03201, doi:10.1029/2004JA010495.
- Imhof, W. L., Reagan, J. B., Gaines, E. E. and Anderson, R. R., Coordinated measurements of slot region electron precipitation by plasmaspheric wave bands, *J. Geophys Res.*, **87**, p. 4418-4426, 1982.

- Imhof, W. L., Voss, H. D., Walt, M., Gaines, E. E., and Mobilia, J., Slot region electron precipitation by lightning, VLF chorus, and plasmaspheric hiss, *J. Geophys. Res.*, **91**, p. 8883-8894, 1986.
- Jursa, A. S., *Handbook of Geophysics and the Space Environment*, Air Force Geophysics Laboratory, United States Air Force, pp. 9-1, 10-47-10-50, 1985.
- Li, Xinlin, D. N. Baker, S. G. Kanekal, M. Looper, and M. Temerin, Long Term Measurements of Radiation Belts by SAMPEX and Their Variations, *Geophys. Res. Letts.*, **28**, pp. 3827-3830, 2001.
- Lyons L. R., and D. J. Williams, *Quantitative Aspects of Magnetospheric Physics*, D. Reidel Pub. Co., Dordrecht, pp.168-180, 1984.
- Lyons, L. R. and R. M. Thorne, Equilibrium structure of radiation belt structure, *J. Geophys. Res.*, **78**, 2142-2149, 1973.
- Raben, V. J., D. S. Evans, H. H. Sauer, S. R. Sahn, and M. Huynh, *TIRO/NOAA Satellite Space Environment Monitor Data Archive Documanetation: 1995 Update*, NOAA Technical Memorandum ERL SEL-86, Space Environment Laboratory, Boulder CO, 1995.
- Vernov S.N., E.V.Gorchakov, S.N.Kuznetsov, Yu.I.Logachev, E.N.Sosnovets, and V.G.Stolpovsky, Particle Fluxes in the Outer Geomagnetic Field, *Reviews of Geophys.*, **7**, pp. 257-280, 1969.




Figure 1. L - B/B_{min} profiles of quiet-time ($K_p \leq 2$ and $V_{sw} \leq 400$ km/s) omnidirectional electron fluxes [> 100 keV (in blue) and > 300 keV (in red)] measured by *MEPED* [Raben et al., 1995] aboard the *NOAA 5, 6, 7, 8, 10, and 12* satellites during 3 months (June, July and August) in solar maximum (1980, 1990) and solar minimum (1986,1996) years. A shift in the slot location between solar minimum and solar maximum is clearly seen at > 300 keV.




Figure 2. Radiation belt electron count rates at >30 (red), >100 (green), and >300 (blue) keV observed at low altitudes (at $0.55L^{3.29} < B/B_0 < 0.65L^{3.29}$) by the *MEPED* instruments [Raben *et al.*, 1995] aboard the *NOAA TIROS 5, 6, 7, and 8* satellites during quiet conditions ($K_p < 2$ and solar wind speeds $< 400 \text{ km s}^{-1}$) in the months of May, June and July of three consecutive years during the solar maximum (left panels) and solar minimum (right panels) periods. The quiet-time slot feature is best seen in the energetic electron data ($> 300 \text{ keV}$, blue). The black vertical lines indicate the nominal slot positions in L during the solar minimum and solar maximum intervals. A small shift in the slot location ($\Delta L \sim 0.5$) is apparent between the two solar cycle phases [After Fung *et al.*, 2005].




Figure 3. Solar-cycle variations of the minimum flux in the slot region (top panel) and the L -value of minimum flux (bottom panel) for energetic electrons (> 300 keV) during quiet-time conditions (see Figure 1). The data are all taken in the months of June, July, and August of each year except that in 1988, the data is taken from the entire year since NOAA10 has poor coverage.




Figure 4. A schematic depicting the migration of the location of optimal wave-particle interaction region characterized by a specific electron plasma frequency-to-gyro frequency ratio, ω_{pe}/Ω_{ce} . Due to the increases of ionospheric densities and scale heights from a solar minimum to a solar maximum, the slot region is expected to move correspondingly to a slightly high L value as observed, and vice versa.