

Paper, Presentation, Short Abstract, & Extended Abstract: Overview of the Field Phase of the NASA Tropical Cloud Systems and Processes (TCSP) Experiment

Authors: Robbie E. Hood, NASA/MSFC, Huntsville, AL; E. Zipser, University of Utah, Salt Lake City, Utah; G. M. Heymsfield, NASA Goddard Space Flight Center, Greenbelt, Maryland; R. Kakar, NASA Headquarters, Washington, DC; J. Halverson, Joint Center for Earth Systems Technology, Greenbelt, Maryland; R. Rogers, NOAA Hurricane Research Division, Miami, FL; and M. Black, NOAA Hurricane Research Division, Miami, FL

Conference: American Meteorological Society 27th Conference on Hurricanes and Tropical Meteorology

Date & Location: April 24-28, 2006; Monterey, CA

The Tropical Cloud Systems and Processes experiment is sponsored by the National Aeronautics and Space Administration (NASA) to investigate characteristics of tropical cyclone genesis, rapid intensification and rainfall using a three-pronged approach that emphasizes satellite information, suborbital observations and numerical model simulations. Research goals include demonstration and assessment of new technology, improvements to numerical model parameterizations, and advancements in data assimilation techniques.

The field phase of the experiment was based in Costa Rica during July 2005. A fully instrumented NASA ER-2 high altitude airplane was deployed with Doppler radar, passive microwave instrumentation, lightning and electric field sensors and an airborne simulator of visible and infrared satellite sensors. Other assets brought to TCSP were a low flying uninhabited aerial vehicle, and a surface-based radiosonde network. In partnership with the Intensity Forecasting Experiment of the National Oceanic and Atmospheric Administration (NOAA) Hurricane Research Division, two NOAA P-3 aircraft instrumented with radar, passive microwave, microphysical, and dropsonde instrumentation were also deployed to Costa Rica.

The field phase of TCSP was conducted in Costa Rica to take advantage of the geographically compact tropical cyclone genesis region of the Eastern Pacific Ocean near Central America. However, the unusual 2005 hurricane season provided numerous opportunities to sample tropical cyclone development and intensification in the Caribbean Sea and Gulf of Mexico as well. Development of Hurricane Dennis and Tropical Storm Gert were each investigated over several days in addition to Hurricane Emily as it was close to Saffir-Simpson Category 5 intensity. An overview of the characteristics of these storms along with the pre-genesis environment of Tropical Storm Eugene in the Eastern Pacific will be presented.

1A.1 OVERVIEW OF THE FIELD PHASE OF THE NASA TROPICAL CLOUD SYSTEMS AND PROCESSES (TCSP) EXPERIMENT

Robbie E. Hood*
NASA Marshall Space Flight Center, Huntsville, Alabama

Edward Zipser
University of Utah, Salt Lake City, Utah

Gerald M. Heymsfield
NASA Goddard Space Flight Center, Greenbelt, Maryland

Ramesh Kakar
NASA Headquarters, Washington, DC

Jeffery Halverson
Joint Center for Earth Systems Technology, Greenbelt, Maryland

Robert Rogers and Michael Black
NOAA Hurricane Research Division, Miami, Florida

1. INTRODUCTION

Although many who reside in the countries bordered by the Caribbean Sea and Gulf of Mexico can never be truly compensated for the heartbreaking losses they suffered during the highly active 2005 hurricane season, one glimmer of hope for the future is that the numerous tropical cyclones which occurred during this season were extensively studied during three Federally funded aircraft field experiments. The National Aeronautics and Space Administration (NASA), The National Oceanic and Atmospheric Administration (NOAA), and the National Science Foundation (NSF) each sponsored research experiments which used aircraft, satellite, oceanic sensors, and research models to collect an extensive body of information that will assist in improving the understanding and predictability of future tropical cyclones.

2. EXPERIMENT DESCRIPTION

The unusually early start of the 2005 Atlantic hurricane season was jointly captured by the NASA Tropical Cloud Systems and Processes (TCSP) Experiment and the initial phase of the

NOAA Intensity Forecasting Experiment (IFEX) which were conducted in collaboration from a base in San Jose, Costa Rica during July 2005. The TCSP experiment was planned by NASA to investigate characteristics of tropical cyclone genesis, rapid intensification and rainfall using a three-pronged approach that emphasizes satellite information, suborbital observations and numerical model simulations. Research goals include demonstration and assessment of new technology, improvements to numerical model parameterizations, and advancements in data assimilation techniques. The goals of the NOAA IFEX are described in Rogers et al. 2006 of this conference.

The field phase of the experiment was based in Costa Rica during July 2005. A fully instrumented NASA ER-2 high altitude airplane was deployed with Doppler radar, passive microwave instrumentation, lightning and electric field sensors and an airborne simulator of visible and infrared satellite sensors. The sensor names and observations of the ER-2 instrument payload are depicted in Fig. 1. Other TCSP assets included near real-time satellite imagery and products, a low flying uninhabited aerial vehicle to sample the eastern Pacific boundary layer during both pre-genesis and inactive phases of TCSP, and a surface-based radiosonde network releasing six-hourly GPS radiosondes from Juan Santamaria International Airport to support

* Corresponding author address: Robbie E. Hood,
NASA/Marshall Space Flight Center, Code:VP61,
320 Sparkman Dr., Huntsville, AL 35805
E-mail: Robbie.Hood@nasa.gov.

scientific and aircraft operations. Costa Rican scientists and forecasters familiar with local weather also contributed to the TCSP weather forecasting and mission planning activities.

Costa Rica was chosen as a deployment site to take advantage of the geographically compact tropical cyclone genesis region of the Eastern Pacific Ocean near Central America. However, the unusual start to the Atlantic basin 2005 hurricane season provided numerous opportunities to sample tropical cyclone development and intensification in the Caribbean

Sea and Gulf of Mexico as well. Development of Hurricane Dennis and Tropical Storm Gert were each investigated over several days in addition to Hurricane Emily as it intensified to nearly Saffir-Simpson Category 5 status. Overviews of the characteristics of these storms along with the pre-genesis environment of Tropical Storm Eugene in the Eastern Pacific are described in more detail in Halverson et al. 2006.

NASA ER-2 Instrument Payload for TCSP




Figure 1. Scientific instrumentation flown on board the NASA ER-2 aircraft during TCSP.

3. CONCLUDING REMARKS

Detailed information about the TCSP campaign, daily mission summaries, satellite animations, flight tracks, instrument data and quicklook images of the data can be found by visiting the TCSP website at <http://tcsp.nsstc.nasa.gov>. This information has considerable merit for research pertaining to tropical cyclone life cycle and intensity changes. However, the true value of the July TCSP observations increase significantly when combined with NOAA IFEX information collected

during July-November and the NSF Rainband and Intensity Experiment (RAINEX) observations collected during August-September to characterize the unique behavior of the unusually long and active 2005 hurricane season in the Atlantic Ocean, Caribbean Sea, and Gulf of Mexico.. The TCSP observations also increase the value of the tropical cyclone data collected during the previous NASA Convection and Moisture Experiments (CAMEX) of 1998 and 2001 (described in Kakar et al. 2006) which provided an unprecedented body of 3-dimensional information of tropical cyclone behavior, structure, and life cycle collected by

high altitude NASA aircraft and lower altitude NOAA aircraft. The comprehensive scope and diverse variety of the NASA, NOAA, and NSF tropical cyclone field data provides a very important opportunity for collaborative research across Federal agencies, academia, and private industry to improve the understanding and predictability of future tropical cyclones.

4. REFERENCES

Halverson J., P. L. Azofeifa, M. Black, S. Braun, D. Cecil, M. Goodman, A. Heymsfield, G. Heymsfield, R. Hood, T. Krishnamurti, G. McFarquhar, J. Molinari, R. Rogers, J. Turk, C. Velden, D.-L. Zhang, E. Zipser, and R. Kakar, 2006: NASA's Tropical Cloud Systems and Processes (TCSP) Experiment: Investigating Tropical Cyclogenesis and Hurricane Intensity Change. *Submitted to Bull. Amer. Meteor. Soc.*

Kakar, R, M. Goodman, R. Hood, A. Guillory, 2006: Overview of the Convection And Moisture Experiment, *J. of Atmos. Sci.* **63**, 5-18

Rogers, R., M. Black, R. Hood, J. Halverson, E. Zipser, and G. Heymsfield, 2006: THE INTENSITY FORECASTING EXPERIMENT (IFEX): A NOAA MULTI-YEAR FIELD PROGRAM FOR IMPROVING TROPICAL CYCLONE INTENSITY FORECASTING, 27th Conference on Hurricane and Tropical Meteorology, 24-28 April, Monterey, CA.

5. ACKNOWLEDGEMENTS

The authors are grateful to Drs. Mary Cleave and Jack Kaye of the NASA Science Mission Directorate for their support of the NASA TCSP mission. The authors also acknowledge Dr. Frank Marks of the NOAA Hurricane Research Division and other members of NOAA management who have supported this NASA/NOAA research collaboration. The authors are deeply indebted to the aircraft pilots and crew, the NASA TCSP Project Office and NOAA Aircraft Operations Center and the Costa Rican government, weather service, and scientific community who significantly contributed to the field phase of the TCSP experiment