



NASA/TM-2001-210926



# Effect of Exercise Training and +Gz Acceleration Training on Men

*John E. Greenleaf, Shawn R. Simonson, Jodie M. Stocks, Joyce Evans, Charles F. Knapp, Stephenie A. Cowell, Kendra N. Pemberton, Heather W. Wilson, Jamie M. Vener, Simon N. Evetts, Peter A. Hardy, Richard E. Grindeland, Helmut Hinghofer-Szalkay, Scott M. Smith, Michael G. Ziegler, David R. Brown, David G. Evans, Fritz B. Moore, and David T. Quach*



## The NASA STI Program Office . . . in Profile

Since its founding, NASA has been dedicated to the advancement of aeronautics and space science. The NASA Scientific and Technical Information (STI) Program Office plays a key part in helping NASA maintain this important role.



The NASA STI Program Office is operated by Langley Research Center, the Lead Center for NASA's scientific and technical information. The NASA STI Program Office provides access to the NASA STI Database, the largest collection of aeronautical and space science STI in the world. The Program Office is also NASA's institutional mechanism for disseminating the results of its research and development activities. These results are published by NASA in the NASA STI Report Series, which includes the following report types:

- **TECHNICAL PUBLICATION.** Reports of completed research or a major significant phase of research that present the results of NASA programs and include extensive data or theoretical analysis. Includes compilations of significant scientific and technical data and information deemed to be of continuing reference value. NASA's counterpart of peer-reviewed formal professional papers but has less stringent limitations on manuscript length and extent of graphic presentations.
- **TECHNICAL MEMORANDUM.** Scientific and technical findings that are preliminary or of specialized interest, e.g., quick release reports, working papers, and bibliographies that contain minimal annotation. Does not contain extensive analysis.
- **CONTRACTOR REPORT.** Scientific and technical findings by NASA-sponsored contractors and grantees.
- **CONFERENCE PUBLICATION.** Collected papers from scientific and technical conferences, symposia, seminars, or other meetings sponsored or cosponsored by NASA.
- **SPECIAL PUBLICATION.** Scientific, technical, or historical information from NASA programs, projects, and missions, often concerned with subjects having substantial public interest.
- **TECHNICAL TRANSLATION.** English-language translations of foreign scientific and technical material pertinent to NASA's mission.

Specialized services that complement the STI Program Office's diverse offerings include creating custom thesauri, building customized databases, organizing and publishing research results . . . even providing videos.

For more information about the NASA STI Program Office, see the following:

- Access the NASA STI Program Home Page at <http://www.sti.nasa.gov>
- E-mail your question via the Internet to [help@sti.nasa.gov](mailto:help@sti.nasa.gov)
- Fax your question to the NASA Access Help Desk at (301) 621-0134
- Telephone the NASA Access Help Desk at (301) 621-0390
- Write to:  
NASA Access Help Desk  
NASA Center for AeroSpace Information  
7121 Standard Drive  
Hanover, MD 21076-1320



## Effect of Exercise Training and +Gz Acceleration Training on Men

*John E. Greenleaf<sup>1</sup>, Shawn R. Simonson<sup>2</sup>, Jodie M. Stocks<sup>1</sup>, Joyce M. Evans<sup>3</sup>, Charles F. Knapp<sup>3</sup>, Stephenie A. Cowell<sup>2</sup>, Kendra N. Pemberton<sup>4</sup>, Heather W. Wilson<sup>1</sup>, Jamie M. Vener<sup>5</sup>, Simon N. Evetts<sup>6</sup>, Peter A. Hardy<sup>3</sup>, Richard E. Grindeland<sup>1</sup>, Helmut Hinghofer-Szalkay<sup>7</sup>, Scott M. Smith<sup>8</sup>, Michael G. Ziegler<sup>9</sup>, David R. Brown<sup>3</sup>, David G. Evans<sup>4</sup>, Fritz B. Moore<sup>10</sup>, and David T. Quach<sup>1</sup>*

*<sup>1</sup>NASA Ames Research Center, Moffett Field, CA 94035-1000; <sup>2</sup>Lockheed Martin Space Operations Corporation, NASA Ames Research Center, Moffett Field, CA 94035-1000; <sup>3</sup>University of Kentucky, Lexington, KY 40506; <sup>4</sup>San Francisco State University, San Francisco, CA 94132; <sup>5</sup>California State University, Fresno, CA 93740-8018; <sup>6</sup>King's College London, London W8 7AH, England; <sup>7</sup>Institute for Adaptive and Spaceflight Physiology, A-8010 Graz, Austria; <sup>8</sup>NASA, Johnson Space Center, Houston, TX 77058-3696; <sup>9</sup>University of California, San Diego, CA 92103-8341; <sup>10</sup>University of California, Davis, CA 95616-8741*

National Aeronautics and  
Space Administration

Ames Research Center  
Moffett Field, California 94035-1000

## Acknowledgments

The authors thank: for management and technical assistance – B.M. Asagi, P.R. Barnes, A. Bognar, T. Burchell, M. Crellin, J.B. Griffith, D.P. Gundo, J.F. Kaumeyer, J.A. Klem, M. Ludgate, S.G. Lumidao, C.M. McIntosh, R.T. Merlo, A.R. Patwardhan, T. Purcell, M.D. Smith, M. Stenger, and R.O. Talavera; for blood drawing – K. Campbell, J.D. Pavloff, and G. Shinn; for medical monitoring – B. Kharrazi, R. Pelligra, and S. Sastry; for manuscript preparation and editing – C.A. Barton, J.M. Donald, and L.N. Popish; and the test subjects for their splendid cooperation.

## Extant Presentations and Publications

- Evans, J.M., S.R. Simonson, C.K. Knapp, J.M. Stocks, H.W. Biagini, S.A. Cowell, K.N. Bailey, J.M. Vener, S.N. Evetts, F.B. Moore, M.B. Stenger, C.M. McIntosh, and J.E. Greenleaf. Differences in acceleration training and exercise training on resting cardiovascular variables. FASEB J. 14: A616, 2000. Abstract.
- Evans, J.M., M.B. Stenger, E. Kwong, C.M. McIntosh, D.R. Brown, A.R. Patwardhan, C.F. Knapp, SR Simonson, JM Stocks, SA Cowell, KN Bailey, JM Vener, SN Evetts, FB Moore, and MG Ziegler.. Human powered centrifuge training on cardiovascular responses to head up tilt. FASEB J. 15: A795, 2001. Abstract.
- Knapp, C.F., M.B. Stenger, E. Kwong, C.M. McIntosh, D.R. Brown, A.R. Patwardhan, J.M. Evans, S.R. Simonson, J.M. Stocks, H.W. Biagini, S.A. Cowell, K.N. Bailey, J.M. Vener, S.N. Evetts, F.B. Moore, and M.G. Ziegler. Passive acceleration training and cardiovascular responses to head up tilt. FASEB J. 15: A795, 2001. Abstract.
- Pemberton, K.N.B. The Effects of Passive +Gz and Exercise Plus +Gz Training on Orthostatic Tolerance. M.A. Thesis; Department of Biology: Physiology and Behavioral Biology, San Francisco State University, September 2000. 84p.
- Simonson, S.R., S.A. Cowell, J.M. Stocks, H.W. Biagini, J.M. Vener, S.N. Evetts, K.N. Bailey, J.M. Evans, C.F. Knapp, and J.E. Greenleaf. The impact of passive acceleration and exercise plus acceleration on work capacity and orthostasis. 13<sup>th</sup> I.A.A. Humans in Space Symposium: Exploring Space. Santorini, Greece, May 20-26, 2000.
- Vener, J.M. Cardiopulmonary Responses to Incremental Supine Cycle Ergometry with Concomitant +Gz Acceleration. M.A. Thesis; Department of Kinesiology, California State University, Fresno, May 2000. 92p.

Available from:

NASA Center for AeroSpace Information  
7121 Standard Drive  
Hanover, MD 21076-1320  
(301) 621-0390

National Technical Information Service  
5285 Port Royal Road  
Springfield, VA 22161  
(703) 487-4650

# CONTENTS

|                                                          |    |
|----------------------------------------------------------|----|
| ACKNOWLEDGMENTS.....                                     | iv |
| SUMMARY .....                                            | 1  |
| INTRODUCTION.....                                        | 3  |
| METHODS.....                                             | 4  |
| Approval .....                                           | 4  |
| Subjects .....                                           | 4  |
| Human-Powered Centrifuge .....                           | 4  |
| Design, construction, operation, and instrumentation     |    |
| Protocol .....                                           | 6  |
| Tests And Measurements.....                              | 11 |
| Maximal oxygen uptake ( $\dot{V}O_2$ max) protocol ..... | 11 |
| Maximal human-powered centrifuge protocol.....           | 14 |
| Tilt-table (orthostasis) protocol.....                   | 15 |
| Blood sampling and analyses.....                         | 18 |
| Urine collection and analyses.....                       | 21 |
| Magnetic resonance imaging (MRI) .....                   | 21 |
| DATA ANALYSES.....                                       | 22 |
| Hemodynamic.....                                         | 22 |
| Statistical.....                                         | 22 |
| RESULTS .....                                            | 23 |
| Maximal Exercise (Passive, Exercise, Combined) Data..... | 23 |
| Exercise load.....                                       | 23 |

|                                                   |    |
|---------------------------------------------------|----|
| Oxygen uptake.....                                | 23 |
| Heart rate .....                                  | 23 |
| Exercise tolerance .....                          | 23 |
| Orthostatic (Tilt-Table) Cardiovascular Data..... | 23 |
| Resting pre-tilt heart rate.....                  | 23 |
| Resting pre-tilt systolic blood pressure.....     | 23 |
| Resting pre-tilt diastolic blood pressure.....    | 23 |
| Resting pre-tilt mean arterial pressure.....      | 23 |
| Tilt-tolerance time .....                         | 24 |
| Heart rate at tolerance.....                      | 24 |
| Mean arterial pressure at tolerance.....          | 24 |
| Cardiac R-R interval with training.....           | 24 |
| Stroke volume with training .....                 | 24 |
| End diastolic volume with training .....          | 24 |
| Cardiac output with training.....                 | 24 |
| Cuff arterial pressure with training.....         | 24 |
| Total peripheral resistance with training.....    | 24 |
| Cardiovascular variables.....                     | 24 |
| Orthostatic (Tilt-Table) Biochemical Data.....    | 24 |
| Blood hemoglobin .....                            | 24 |
| Raw hematocrit .....                              | 25 |
| Plasma volume.....                                | 25 |
| Plasma sodium.....                                | 25 |

|                                               |    |
|-----------------------------------------------|----|
| Plasma potassium .....                        | 25 |
| Plasma osmolality .....                       | 25 |
| Plasma albumin.....                           | 25 |
| Plasma total protein.....                     | 25 |
| Plasma renin activity .....                   | 26 |
| Plasma aldosterone .....                      | 26 |
| Plasma vasopressin.....                       | 26 |
| Plasma epinephrine.....                       | 26 |
| Plasma norepinephrine.....                    | 26 |
| Plasma dopamine.....                          | 26 |
| Plasma growth hormone .....                   | 26 |
| Urine Data (24 hour) .....                    | 27 |
| Volume and rate (24 hr).....                  | 27 |
| Creatinine.....                               | 27 |
| Deoxypyridinoline.....                        | 27 |
| Deoxypyridinoline / creatinine ratio.....     | 27 |
| n-Telopeptide.....                            | 27 |
| n-Telopeptide / creatinine ratio.....         | 27 |
| Pyridinium crosslinks.....                    | 27 |
| Pyridinium crosslinks / creatinine ratio..... | 27 |
| Hydroxyproline .....                          | 27 |
| Hydroxyproline / creatinine ratio .....       | 27 |
| Calcium.....                                  | 27 |

|                                                    |    |
|----------------------------------------------------|----|
| Magnetic Resonance Imaging.....                    | 27 |
| Volume .....                                       | 27 |
| Excitation .....                                   | 27 |
| SUMMARY OF RESULTS.....                            | 28 |
| REFERENCES.....                                    | 30 |
| APPENDICES .....                                   | 36 |
| A: A1 – A4. Protocol Approvals .....               | 36 |
| B: B1 – B2. G-level vs. RPM .....                  | 40 |
| C: Instruments and Equipment Inventory.....        | 42 |
| D: Borg Intensity of Perceived Exertion Scale..... | 46 |
| E: CPX Oxygen Analyzer Calibration.....            | 47 |
| F: Raw Data.....                                   | 48 |
| G: Figures (12 to 57).....                         | 93 |



## EFFECT OF EXERCISE TRAINING AND +Gz ACCELERATION TRAINING ON MEN

John E. Greenleaf<sup>1</sup>, Shawn R. Simonson<sup>2</sup>, Jodie M. Stocks<sup>1</sup>, Joyce M. Evans<sup>3</sup>, Charles F. Knapp<sup>3</sup>, Stephenie A. Cowell<sup>2</sup>, Kendra N. Pemberton<sup>4</sup>, Heather W. Wilson<sup>1</sup>, Jamie M. Vener<sup>5</sup>, Simon N. Evetts<sup>6</sup>, Peter A. Hardy<sup>3</sup>, Richard E. Grindeland<sup>1</sup>, Helmut Hinghofer-Szalkay<sup>7</sup>, Scott M. Smith<sup>8</sup>, Michael G. Ziegler<sup>9</sup>, David R. Brown<sup>3</sup>, David G. Evans<sup>4</sup>, Fritz B. Moore<sup>10</sup>, and David T. Quach<sup>1</sup>

### SUMMARY

Reduction in work capacity (maximal oxygen uptake) during flight and enhanced orthostatic intolerance during reentry, landing, and egress from the return vehicle are continuing problems that have not been solved. Intermittent, high-intensity, short-duration isotonic cycle ergometer exercise training can maintain work capacity at ambulatory control levels over 1 month of bed-rest (BR)-deconditioning, and short-arm (< 2-meter radius) +Gz (head-to-foot) acceleration training without and with concomitant exercise can attenuate the usual orthostatic intolerance resulting from water-immersion or bed-rest deconditioning. Thus, the purpose for this study was to test the hypothesis that (1) passive-acceleration training; supine-interval-exercise plus acceleration training; and exercise combined with acceleration training would improve orthostatic tolerance in ambulatory men; and that (2) addition of the aerobic exercise conditioning would not alter this improved tolerance from that of passive-acceleration training.

Seven men (24–38 years) were test subjects. Three men underwent “Passive” training on the Ames human-powered centrifuge (HPC) for 30 min (warm-up, then 24 min of 2-min acceleration intervals (+1.0 Gz to 50%  $G_{z_{max}}$  at  $+2.4 \pm 0.1$  Gz), and cool-down) for 5 days/week for 3 weeks. Three other subjects underwent constant +Gz acceleration (50% of HPC maximal acceleration at  $+2.3 \pm 0.2$  Gz) while performing “Exercise” training on the cycle ergometer at 40% of maximal oxygen

---

<sup>1</sup>Laboratory for Human Environmental Physiology, NASA, Ames Research Center, Moffett Field, Calif. 94035-1000; <sup>2</sup>Lockheed Martin Space Operations Corporation, NASA, Ames Research Center, Moffett Field, Calif. 94035-1000; <sup>3</sup>Center for Biomedical Engineering, University of Kentucky, Lexington, Ky. 40506; <sup>4</sup>Department of Biology: Physiology and Behavioral Biology, San Francisco State University, San Francisco, Calif. 94132; <sup>5</sup>Department of Kinesiology, California State University, Fresno, Calif. 93740-8018; <sup>6</sup>Biomedical Sciences Division, King’s College London, London W8 7AH, England; <sup>7</sup>Institute for Adaptive and Spaceflight Physiology, A-8010 Graz, Austria; <sup>8</sup>Life Sciences Research Laboratories, NASA, Johnson Space Center, Houston, Tex. 77058-3696; <sup>9</sup>GCRC, UCSD Medical Center, University of California, San Diego, Calif. 92103-8341; <sup>10</sup>Department of Neurology, Physiology, and Behavior, University of California, Davis, Calif. 95616-8741.

uptake ( $\dot{V}O_{2\max}$ ), then 24 min of 2-min intervals (40% - 90%  $\dot{V}O_{2\max}$ ) for 5 days/week for 3 weeks. A crossover design utilized 4 weeks of ambulatory deconditioning between sessions. Six subjects also underwent similar “Combined” exercise training at 40% to 90% of the HPC +Gz<sub>max</sub> exercise level. Before and after each training session the maximal  $\dot{V}O_2$ , workload, and heart rate (HR) were determined supine using a ramped cycle ergometer protocol. Resting HR and blood pressures (systolic, diastolic, and mean arterial) were measured, pre- and post-training, after 40 min of supine rest and for 1 min before determination of orthostatic tolerance to 70° head-up tilt.

Maximal human-powered (Passive, Exercise, Combined) data. Maximal supine exercise loads increased significantly ( $P < 0.05$ ) by 8.3% (Passive Phase), 12.6% (Exercise Phase), and 15.4% (Combined Phase) after training, but the subjects’ post-training maximal oxygen uptakes and maximal heart rates were unchanged from their respective pre-training levels. Maximal time to fatigue (endurance) was unchanged with Passive, but was also increased ( $P < 0.05$ ) with Exercise and Combined training. Thus, the exercise in the Exercise and Combined training Phases resulted in greater maximal loads and endurance without effect on maximal oxygen uptake or heart rate.

Orthostatic (tilt-table) cardiovascular data. Resting pre-tilt heart rate was elevated by 12.9% ( $P < 0.05$ ) only after Passive training, suggesting that the exercise training attenuated the HR response. Resting pre-tilt blood pressures (SBR, DBP, MAP) were not different pre- or post-training in any Phase. Post-training tilt-tolerance time and heart rate were increased ( $P < 0.05$ ) only with Passive training, by 37.8% and by 29.1%, respectively. Thus, addition of exercise training appeared to attenuate the increased Passive tolerance. Resting (pre-tilt) and post-tilt cardiac R-R interval, stroke volume, end-diastolic volume, and cardiac output were all uniformly reduced ( $P < 0.05$ ), and peripheral resistance was uniformly increased ( $P < 0.05$ ) pre- and post-training for the three Phases, indicating no effect of the exercise training on these cardiovascular variables.

Orthostatic (tilt-table) biochemical data. Plasma volume (percent change) was uniformly decreased by 8% to 14% ( $P < 0.05$ ) at tilt-tolerance pre-training versus post-training, indicating essentially no effect of training on the level of hypovolemia. The latter was reflected in the 6% to 12% ( $P < 0.05$ ) increase in plasma aldosterone [PA] and plasma total protein [PTP] concentrations. Percent changes in plasma sodium concentration [PNa] pre-training versus post-training were minimal (less than -0.8%) as was plasma osmolality [POsm] (less than -0.4%) indicating essentially isotonic plasma shifts during tilting. Pre- and post-training percent changes in plasma renin activity (PRA), plasma aldosterone concentration [PA], plasma epinephrine concentration [PE], and [PNa] exhibited similar characteristic increases at tolerance; the usual increase in plasma vasopressin concentration [PVP] was greatly attenuated post-training with Exercise and also pre- and post-training with Combined. The explanation for the latter is not obvious but is unlikely a result of technical errors.

Urine data (24-hr). Urinary volumes were within normal limits (1.2 to 1.5 ml•min<sup>-1</sup>) between and among pre- and post-training samples for the three Phases. There were no significant differences between or among the 10 urinary variables pre-training and post-training for the three Phases.

## INTRODUCTION

Long-duration (> 1-yr) human spaceflight will require refinement of current physiological countermeasures, as well as implementation of others to allay deconditioning of crew members – defined as attenuation of their physical fitness. Crew exposure to weightlessness in the spaceflight environment of moderate confinement, restricted mobility, and enhanced ionizing radiation affects every organ system in the body that contributes to deconditioning (Convertino, 1990; Nicogossian, 1994; Sonnenfeld, 1998; Zerath, 1998). Effects of deconditioning such as reduction of maximal work capacity (Greenleaf et al., 1989), bone density and strength (LeBlanc et al., 1990, 1996), muscle mass and strength (Ellis et al., 1993; Greenleaf et al., 1994), orthostatic tolerance (Buckey et al., 1996), and neurovestibular sensitivity (Collins et al., 1995; Paloski et al., 1992) can lead to decreased crew health, safety, and productivity during flight, especially during and immediately after landing on a planet with a physiologically significant gravitational force (Buckey et al., 1996; Burton, 1988; Kotovskaya et al., 1977; Nicogossian, 1994).

Short arm (< 2-m radius) +Gz (head-to-foot) acceleration training without (Shulzhenko et al., 1976, 1979; Vil-Vilyams, 1994) and with (Vil-Vilyams and Shulzhenko, 1980) concomitant exercise training has been reported to significantly attenuate the usual orthostatic intolerance resulting from water-immersion deconditioning. In addition, the data of White et al. (1966) indicated that the consistent intolerance to 20 min of 70° head-up tilt after prolonged BR-deconditioning was reduced or alleviated by daily intermittent +1.75 Gz (4.7 G-hr) training periods without exercise during 10 days of horizontal BR as reviewed by Stone et al. (1966). Also, intensive, intermittent cycle ergometer exercise training can maintain aerobic exercise capacity (maximal oxygen uptake) at ambulatory control levels during 30 days of 6° head-down BR (Greenleaf et al., 1989; Kakurin et al., 1978). Performing exercise with acceleration has been proposed as a time-efficient countermeasure to attenuate the reduction in both exercise capacity and orthostatic tolerance simultaneously during deconditioning (Burton, 1988; Greenleaf et al., 1999; Shulzhenko et al., 1976; Vernikos, 1997; Vil-Vilyams and Shulzhenko, 1980).

The efficacy of these two countermeasures, singly or in combination, has not been confirmed on most physiological systems after prolonged BR-deconditioning. As a result, there are many questions to be addressed. For example: What is the most effective duration and intensity of exercise and acceleration? When performed simultaneously, do these two treatments interfere with each other? Will the muscle pumping action of exercising legs interfere with acceleration-induced caudal fluid shifts? Does daily exercise-acceleration training have salutary or adverse effects on the muscular, neurovestibular, or skeletal systems?

The first studies utilizing the human-powered centrifuge (HPC) (Mulenburg and Vernikos, 1997) at Ames Research Center (ARC) began in 1995 (Chou, 1997; Chou et al., 1998; Greenleaf et al., 1997, 1999; Stad, 1998; Vener, 2000) and were designed to familiarize the investigators with its operating characteristics, procedures, and linearity of its loads as an exercise ergometer. Chou et al. (1997) have annotated most of the literature concerning the effects of exercise and acceleration training on deconditioning through 1996.

The purpose of this study was to investigate exercise and acceleration training (singly and in combination) on metabolic, orthostatic, and blood and urine factors to provide background data in

preparation for a more extensive BR-deconditioning-training study. It was hypothesized that the three training protocols (Passive, Exercise, and Combined, all utilizing head-to-foot acceleration) would preserve normal orthostatic responses during tilt, and that the aerobic exercise conditioning stimuli would not alter those +Gz training responses.

## METHODS

### Approval

This experimental protocol entitled “Exercise Training on the Short-Arm Centrifuge” was approved by the San Francisco State University Committee for the Protection of Human Subjects on 8 May 1998, and by the ARC Human Research Institutional Review Board on 4 January 1999 (H.R. No. 158 was replaced by No. 191). Approval of Dr. R. E. Grindeland’s addendum was granted on 16 February 1999, and for a 1-month extension of H.R. No. 191 on 8 November 1999 (appendixes A1 – A4). Study data were collected from 14 June through 19 December 1999.

### Subjects

The male test subjects were recruited from San Jose State University students and ARC employees and contractors. Seven men (table 1) were selected who were not currently involved in a regular physical training program, and who had no tobacco or nonprescriptive drug use. After extensive presentations of the experimental requirements and potential hazards by the investigators and medical monitor, the subjects signed informed consent forms and passed a comprehensive medical examination including history, electrocardiogram, blood and urine chemistry panels, and an exercise stress test prior to data collection.

### Human-Powered Centrifuge

Design, construction, operation, and instrumentation. The HPC, designed and constructed at Ames Research Center, is a short-arm (1.9 m ) dual-couch machine powered by a chain-linked cycle driven by the subject’s legs (fig. 1)\*. The revolving circular platform assembly consists of three lightweight aluminum honeycomb and bonded aluminum sections – a center section and two wings weighing a total of 295 kg. The center section, which rides on tapered roller bearings (Timken), supports the two couches and the chain-drive pedal mechanism. Bevel gears (Boston, 4:1 gear ratio) transform pedaling power to platform rotation. The two wings, bolted to the center section, provide areas for walking and

---

\* Figures 1–11 are in the main text at the point of citation; figures 12-57 appear at the end of the report. In text, bracketed abbreviations designate concentrations; for example, plasma total protein [PTP]. Simple initials are enclosed in the conventional parentheses following their definitions; for example, plasma renin activity (PRA).

Table 1. Test subject characteristics.

| Subject   | Phase    | Age, yr | Ht, cm | Wt, kg | S.A., m <sup>2</sup> | PV, ml | TBV, ml | VO <sub>2</sub> max,  |                                           | HR <sub>max</sub> ,   | +Gz <sub>max</sub> , | RH <sub>max</sub> ,   |
|-----------|----------|---------|--------|--------|----------------------|--------|---------|-----------------------|-------------------------------------------|-----------------------|----------------------|-----------------------|
|           |          |         |        |        |                      |        |         | L • min <sup>-1</sup> | ml • min <sup>-1</sup> • kg <sup>-1</sup> | b • min <sup>-1</sup> | G                    | b • min <sup>-1</sup> |
| FLE       | III      | 32      | 172.0  | 71.7   | 1.82                 | 3300   | 5172    | 2.62                  | 37                                        | 166                   | 4.40                 | 174                   |
| FRE       | I II III | 38      | 176.0  | 89.8   | 2.10                 | 3781   | 5609    | 3.44                  | 38                                        | 170                   | 4.51                 | 180                   |
| HUN       | I II III | 38      | 186.0  | 96.8   | 2.14                 | 3616   | 6092    | 2.69                  | 28                                        | 174                   | 4.50                 | 186                   |
| JAG       | I II III | 24      | 180.5  | 85.6   | 2.07                 | 3493   | 5829    | 3.18                  | 38                                        | 186                   | 4.94                 | 192                   |
| RAY       | I II III | 27      | 178.0  | 83.0   | 2.01                 | 3457   | 5510    | 3.31                  | 40                                        | 188                   | 5.47                 | 179                   |
| RUI       | I II III | 38      | 177.0  | 86.0   | 2.03                 | 4486   | 7437    | 3.16                  | 37                                        | 159                   | 4.90                 | 161                   |
| SCH       | I II     | 30      | 172.5  | 76.5   | 1.88                 | 2864   | 4535    | 2.86                  | 37                                        | 188                   | 5.03                 | 186                   |
| $\bar{X}$ |          | 32      | 177.4  | 84.2   | 2.01                 | 3571   | 5741    | 3.04                  | 36                                        | 176                   | 4.82                 | 180                   |
| SD        |          | 6       | 4.8    | 8.3    | 0.12                 | 496    | 900     | 0.32                  | 4                                         | 12                    | 0.38                 | 10                    |

Phase I – Passive acceleration; Phase II – Exercise acceleration; Phase III – Combined acceleration.  
 PV = plasma volume; TBV = total blood volume.




Figure 1. The Human Powered Centrifuge

for mounting instrumentation. The three pedaling stations, two on-board at the end of the couches and the third located on the off-board operator's stationary cycle, are linked by standard bicycle chains and sprockets to the center hub. The top of the subject's head is located about 26 cm from the center of rotation, and the level of acceleration is calculated 1.9 m (6 ft) from the center of rotation. One 360° rotation of the platform requires 1.6 pedal sprocket rotations. Tables of acceleration as a function of angular velocity are given in appendixes B1 and B2.

The on-board subject and the off-board operator can actuate and control platform angular velocity (revolutions/min, rpm), but only the operator can stop the platform with a spring-set disc brake (Stearns series 87,300, model 1-087-352-00, Rexnord Corp., Milwaukee, Wis.) activated electronically by the operator or test subject. Centrifuge parameters, such as rpm (model H25D-SB encoder, BEI Motions Systems, Co., Goleta, Calif., and model P6020 tachometer, Newport Electronics, Inc., Santa Ana, Calif.; total accuracy of 0.0002%) and G-level, are displayed on the control panel of the monitoring station. The center hub-drive differential has a slip-ring assembly (model 1067, Fabricast, South El Monte, Calif.) that transmits physiological data electrically from the on-board test subjects to instruments in the adjoining control room (fig. 1).

In the current configuration, the second on-platform pedaling station is a standard independent cycle ergometer (model 845, Quinton Ergometer, Seattle, Wash.) that is not connected for platform rotation (fig. 2). For rotation while pedaling this ergometer, one of the other two pedaling stations must be engaged. There are three video cameras: one by each couch aimed at the subject's head, and the third covering the entire centrifuge. Subject energy output was measured with a metabolic (oxygen) analyzer (CPX Express, MedGraphics Corp., St. Paul, Minn.) with data downloaded and stored at the end of each run with a model H-1330 (Quantax Microsystems Corp., Somerset, N.J.) laptop computer, and then printed on the MedGraphics printer (model BJC, Canon Hi-Tech, Thailand). Heart-rate data were taken from the electrocardiogram (model 78202, Hewlett-Packard, Palo Alto, Calif.) and displayed in the control room. Information about all instruments and equipment is presented in appendix C.

## Protocol

This study consisted of a 2-week pre-training period, three 3-week centrifuge training periods separated by two 4-week ambulatory recovery (deconditioning) periods, and a 4-week post-training ambulatory recovery (deconditioning) period (fig. 3). Daily protocols are presented in appendix D. The exercise and acceleration training regimens were conducted in three Phases:

Phase I (Passive acceleration) required the supine, resting subjects (no exercise) to ride the centrifuge at relative, intermittent loads that stepped between 25% and 50% of their maximal +Gz acceleration (+Gz<sub>max</sub>), provided by the off-platform cycle operator. Each subject underwent a 6-min warm-up at 25% followed by alternating 25% and 50% loads at 2-min intervals (0.008 Hz) for 30 min (fig. 4). The range of 50% +Gz<sub>max</sub> levels was 2.2 G to 2.5 G at the foot (table 2).





Figure 2. The Human Powered Centrifuge, top, bottom.

Figure 3(a). Experimental Protocol (Overview).

| Baseline                     |                         |           | I: Centrifuge Training<br>Passive + Exercise                                   |         |        | Recovery<br>(deconditioning) |                         |           | II: Centrifuge Training<br>Exercise + Passive                                    |         |       |
|------------------------------|-------------------------|-----------|--------------------------------------------------------------------------------|---------|--------|------------------------------|-------------------------|-----------|----------------------------------------------------------------------------------|---------|-------|
| July 12                      | 2 weeks                 | July 23   | July 26                                                                        | 3 weeks | Aug 13 | Aug 16                       | 4 weeks                 | Sept 10   | Sept 13                                                                          | 3 weeks | Oct 1 |
| ↑<br>(BS) Tilt               | (BS) TILT (BS)          | ↑<br>(BS) | (14 training sessions)<br>(1 tilt test on day 15)                              |         |        | ↑<br>maxVO <sub>2</sub>      | (BS) TILT (BS)          | ↑<br>(BS) | (14 training sessions)<br>(1 tilt test on day 15)                                |         |       |
| ↑<br>UA                      |                         | ↑<br>UA   | (BS) TILT (BS)                                                                 |         |        | ↑<br>MRI                     |                         | ↑<br>MRI  | (BS) TILT (BS)                                                                   |         |       |
|                              | ↑<br>maxVO <sub>2</sub> | ↑<br>UA   | ↑<br>UA                                                                        |         |        |                              | ↑<br>maxVO <sub>2</sub> | ↑<br>UA   | ↑<br>UA                                                                          |         |       |
|                              | ↑<br>+GzVO <sub>2</sub> |           |                                                                                |         |        |                              | ↑<br>+GzVO <sub>2</sub> |           |                                                                                  |         |       |
|                              |                         |           | Group A; N=3 (exercise acceleration)*<br>Group B; N=3 (passive acceleration)** |         |        |                              |                         |           | Group A; N=3 (passive acceleration)**<br>Group B; N=3 (exercise + acceleration)* |         |       |
| Recovery<br>(deconditioning) |                         |           | III: Centrifuge<br>Training Combined<br>(HPC)                                  |         |        | Recovery<br>(deconditioning) |                         |           |                                                                                  |         |       |
| Oct 4                        | 4 weeks                 | Oct 29    | Nov 1                                                                          | 3 weeks | Nov 19 | Nov 22                       | 4 weeks                 | Dec 17    |                                                                                  |         |       |
| ↑<br>maxVO <sub>2</sub>      | (BS) TILT (BS)          | ↑<br>(BS) | (14 training sessions)<br>(1 tilt test on day 15)                              |         |        | ↑<br>maxVO <sub>2</sub>      | (BS) TILT (BS)          | ↑<br>(BS) |                                                                                  |         |       |
| ↑<br>MRI                     |                         | ↑<br>MRI  | (BS) TILT (BS)                                                                 |         |        | ↑<br>MRI                     |                         | ↑<br>MRI  |                                                                                  |         |       |
|                              | ↑<br>UA                 | ↑<br>UA   | ↑<br>UA                                                                        |         |        |                              | ↑<br>UA                 | ↑<br>UA   |                                                                                  |         |       |
|                              | ↑<br>maxVO <sub>2</sub> |           | ↑<br>UA                                                                        |         |        |                              | ↑<br>maxVO <sub>2</sub> |           |                                                                                  |         |       |
|                              | ↑<br>+GzVO <sub>2</sub> |           |                                                                                |         |        |                              | ↑<br>+GzVO <sub>2</sub> |           |                                                                                  |         |       |
|                              |                         |           | Groups A + B, N=6<br>(combined acceleration)***                                |         |        |                              |                         |           |                                                                                  |         |       |

Tilt tests will be performed 1–2 days before pre-training and on the last day of the centrifuge training periods.

MRI tests will be performed 2–3 days before pre-training and 1 day after the centrifuge training periods.

Max VO<sub>2</sub> will be measured within a week before and 2 days after the centrifuge training periods.

Max +Gz VO<sub>2</sub> will be measured within a week before the centrifuge training periods.

UA: Urine analysis for markers of bone remodeling will be made from a 24-hr collection prior to tilt.

BS: Blood samples will be drawn before and after each tilt test.

\* Exercise + acceleration = alternating ergometer exercise + constant off-platform acceleration.

\*\* Passive acceleration = alternating off-platform acceleration.

\*\*\* Combined acceleration = human-powered centrifuge acceleration alternating exercise and accompanying acceleration.



Buildup and Phase I: Passive (P) and Exercise (E)  
Acceleration Training Schedule.

|          | Wk             | Mon                                  | Tue                                      | Wed                       | Thu                                | Fri                                   | Sat | Sun |
|----------|----------------|--------------------------------------|------------------------------------------|---------------------------|------------------------------------|---------------------------------------|-----|-----|
| Build Up | June 14 - 20   |                                      | Supine Practice Part 1                   |                           | Supine Practice Part 2             | Supine Practice Part 2                |     |     |
|          | June 21 - 27   | 9am Pract VO <sub>2</sub> max Part 2 |                                          | 9am Pract HPCmax Part 1   |                                    | 9am Pract HPCmax Part 2               |     |     |
|          | Jun/Jul 28 - 4 |                                      |                                          |                           |                                    |                                       |     |     |
|          | July 5 - 11    | Lab Closed                           |                                          | Subject Orientation Day * | Upright VO <sub>2</sub> max Part 2 | Practice Tilt <sup>B</sup> Urine Coll |     |     |
| I        | July 12 - 18   | Supine VO <sub>2</sub> max Part 1    | Supine VO <sub>2</sub> max Part 2: Trial | Prelim HPCmax Part 1      | Pract Tilt                         | Prelim Tilt <sup>B</sup>              | MRI |     |
|          | July 19 - 25   | Study VO <sub>2</sub> max Part 2     | Practice HPC P & E                       | Study HPCmax Part 2       |                                    | Study Tilt <sup>B</sup> Urine Coll    |     |     |
| Train    | Jul/Aug 26 - 1 | 30min P<br>E                         | 30min P<br>E                             | 30min P<br>E              | 30min P<br>E                       | 30min P<br>E                          |     |     |
|          | Aug 2 - 8      | 30min P<br>E                         | 30min P<br>E                             | 30min P<br>E              | 30min P<br>E                       | 30min P<br>E                          |     |     |
|          | Aug 9 - 15     | 30min P<br>E                         | 30min P<br>E                             | 30min P<br>E              | 30min P<br>E                       | Study Tilt <sup>B</sup> Urine Coll    | MRI |     |
| Recov    | Aug 16 - 22    | Study VO <sub>2</sub> max Part 2     |                                          |                           |                                    |                                       |     |     |
|          | Aug 23 - 29    |                                      |                                          |                           |                                    |                                       |     |     |
|          | Aug/Sep 30 - 5 |                                      |                                          |                           |                                    |                                       |     |     |
|          | Sep 6 - 12     | Study VO <sub>2</sub> max Part 2     |                                          | Study HPCmax Part 2       |                                    | Study Tilt <sup>B</sup> Urine Coll    | MRI |     |

Phase II: Exercise and Passive Acceleration Training Schedule.

|       | Wk             | Mon                              | Tue          | Wed                 | Thu          | Fri                                | Sat | Sun |
|-------|----------------|----------------------------------|--------------|---------------------|--------------|------------------------------------|-----|-----|
| II    | Sep 13 - 19    | 30min E<br>P                     | 30min E<br>P | 30min E<br>P        | 30min E<br>P | 30min E<br>P                       |     |     |
|       | Sep 20 - 26    | 30min E<br>P                     | 30min E<br>P | 30min E<br>P        | 30min E<br>P | 30min E<br>P                       |     |     |
|       | Sep/Oct 27 - 3 | 30min E<br>P                     | 30min E<br>P | 30min E<br>P        | 30min E<br>P | Study Tilt <sup>B</sup> Urine Coll | MRI |     |
| Recov | Oct 4 - 10     | Study VO <sub>2</sub> max Part 2 |              |                     |              |                                    |     |     |
|       | Oct 11 - 17    |                                  |              |                     |              |                                    |     |     |
|       | Oct 18 - 24    |                                  |              |                     |              |                                    |     |     |
|       | Oct 25 - 31    | Study VO <sub>2</sub> max Part 2 |              | Study HPCmax Part 2 |              | Study Tilt <sup>B</sup> Urine Coll | MRI |     |

Phase III: Combined (HPC) Acceleration Training Schedule.

|       | Wk             | Mon                              | Tue       | Wed                 | Thu       | Fri                                | Sat | Sun |
|-------|----------------|----------------------------------|-----------|---------------------|-----------|------------------------------------|-----|-----|
| III   | Nov 1 - 7      | 30min HPC                        | 30min HPC | 30min HPC           | 30min HPC | 30min HPC                          |     |     |
|       | Nov 8 - 14     | 30min HPC                        | 30min HPC | 30min HPC           | 30min HPC | 30min HPC                          |     |     |
| Train | Nov 15 - 21    | 30min HPC                        | 30min HPC | 30min HPC           | 30min HPC | Study Tilt <sup>B</sup> Urine Coll | MRI |     |
|       | Nov 22 - 28    | Study VO <sub>2</sub> max Part 2 |           |                     |           |                                    |     |     |
| Recov | Nov/Dec 29 - 5 |                                  |           |                     |           |                                    |     |     |
|       | Dec 6 - 12     |                                  |           |                     |           |                                    |     |     |
|       | Dec 13 - 19    | Study VO <sub>2</sub> max Part 2 |           | Study HPCmax Part 2 |           | Study Tilt <sup>B</sup> Urine Coll | MRI |     |

LEGEND



- = No test subjects involved.
- = Test subjects involved.

- \* Orientation = Subject familiarization with HPC, tilt, supine ergometer and VO<sub>2</sub>max
- Tilt<sup>B</sup> = Tilt to +70° from horizontal for approximately 1 hr + blood samples.
- Study = Measures to be used for study analysis.
- VO<sub>2</sub>max Pt 1 = Maximal oxygen uptake preliminary assessment.
- VO<sub>2</sub>max Pt 2 = Maximal oxygen uptake main assessment.
- MRI = Magnetic resonance imaging (noninvasive scan).
- HPCmax Pt 1 = Maximal oxygen uptake + centrifugation preliminary assessment .
- HPCmax Pt 2 = Maximal oxygen uptake + centrifugation main assessment.
- P = Subjects undertake passive centrifugation.
- E = Subjects undertake exercise during centrifugation.
- ↔ Urine Coll = 24-hr urine collection.

Figure 3(b).

**Phase I. Centrifuge Training - Passive. Acceleration and Workload Over Time.**




Figure 4,

The Phase II (Exercise acceleration) protocol consisted of supine, oscillatory leg ergometer exercise varying from 40% of maximal oxygen uptake ( $\dot{V}O_2$ max) to 90%  $\dot{V}O_2$ max in alternating 2-min intervals (fig. 5). The constant 50% +Gz<sub>max</sub> acceleration load was provided by the off-board operator.

Phases I and II were conducted using a cross-over design with three subjects (Group A) undergoing exercise acceleration and three subjects (Group B) undergoing passive acceleration first; followed by the reverse in Phase II (fig. 1). The ranges of exercise loads (kg·m·min<sup>-1</sup>) in Phase II were: 40% (600-800), 50% (750-1,000), 60% (900-1,200), 70% (1,050-1,400), 80% (1,200-1,600), and 90% (1,350-1,800) (table 2).

Phase III (Combined acceleration) had the subjects exercising on the HPC arm of the centrifuge where their leg exercise drives the centrifuge; that is, the exercise and the acceleration were performed by the subjects. Here the 40% to 90% loads were determined previously from the +Gz<sub>max</sub> on the HPC arm expressed in revolutions per minute (fig. 6). The ranges of relative exercise acceleration (G-levels) were: 40% (1.8-2.2), 50% (2.2-2.8), 60% (2.6-3.3), 70% (3.1-3.9), 80% (3.5-4.4), and 90% (4.0-5.0) (table 2 and appendix B).

**Table 2. Individual +Gz<sub>max</sub> levels for the seven men during the three Phases.**

|     |                                      |            |            |            |            |            |            |            |          |
|-----|--------------------------------------|------------|------------|------------|------------|------------|------------|------------|----------|
| I   |                                      | <b>FLE</b> | <b>FRE</b> | <b>HUN</b> | <b>JAG</b> | <b>RAY</b> | <b>RUI</b> | <b>SCH</b> |          |
|     | <b>100% +Gz max</b>                  | 4.40       | 4.51       | 4.50       | 4.94       | 5.47       | 4.90       | 5.03       | <b>G</b> |
|     | <b>Passive</b>                       | —          | 13<br>Sept | 13<br>Sept | 26<br>July | 26<br>July | 26<br>July | 13<br>Sept |          |
|     | <b>50% +Gz max</b>                   | —          | 2.2        | 2.2        | 2.4        | 2.5        | 2.5        | 2.5        | <b>G</b> |
| II  | <b>Exercise (ergometer)</b>          | —          | 26<br>July | 26<br>July | 13<br>Sept | 13<br>Sept | 13<br>Sept | 26<br>July |          |
|     | <b>50% +Gz max</b>                   | —          | 2.4        | 1.8        | 2.5        | 2.8        | 2.4        | 2.2        | <b>G</b> |
|     | <b>Load in kg·m·min<sup>-1</sup></b> |            |            |            |            |            |            |            |          |
|     | <b>40%</b>                           | —          | 600        | 600        | 800        | 800        | 800        | 600        |          |
|     | <b>50%</b>                           | —          | 800        | 750        | 1000       | 1000       | 1000       | 750        |          |
|     | <b>60%</b>                           | —          | 1000       | 900        | 1200       | 1200       | 1100       | 900        |          |
|     | <b>70%</b>                           | —          | 1100       | 1050       | 1400       | 1400       | 1300       | 1050       |          |
|     | <b>80%</b>                           | —          | 1300       | 1200       | 1600       | 1600       | 1500       | 1200       |          |
|     | <b>90%</b>                           | —          | 1400       | 1350       | 1800       | 1800       | 1700       | 1350       |          |
| III | <b>Combined (HPC)</b>                | 1 Nov      | 1 Nov      | 1 Nov      | 1 Nov      | 1 Nov      | 1 Nov      |            |          |
|     | <b>Load in +Gz units</b>             |            |            |            |            |            |            |            |          |
|     | <b>40%</b>                           | 1.8        | 1.9        | 1.8        | 1.9        | 2.2        | 2.1        | —          |          |
|     | <b>50%</b>                           | 2.2        | 2.4        | 2.2        | 2.4        | 2.8        | 2.6        | —          |          |
|     | <b>60%</b>                           | 2.6        | 2.8        | 2.6        | 2.9        | 3.3        | 3.1        | —          |          |
|     | <b>70%</b>                           | 3.1        | 3.3        | 3.1        | 3.4        | 3.9        | 3.6        | —          |          |
|     | <b>80%</b>                           | 3.5        | 3.8        | 3.5        | 3.8        | 4.4        | 4.2        | —          |          |
|     | <b>90%</b>                           | 4.0        | 4.3        | 4.0        | 4.3        | 5.0        | 4.7        | —          |          |

The subjective intensity of perceived stress during the training sessions was noted by the subjects on the Borg (1982) scale (appendix D).

### Tests and Measurements

Maximal oxygen uptake ( $\dot{V}O_{2max}$ ) protocol. Maximal working capacity was measured in the pre-training period with the subjects in the upright (sitting) and supine body positions (table 3) on calibrated electronically-braked ergometers (models 845 sitting and 846T supine, Quinton, Seattle, Wash.) where work output was independent of pedal rpm. Sitting and supine exercise were used prior to exercise-training data collection to familiarize the subjects with the protocol and to determine their maximal exertion data. The supine position was then used for all subsequent maximal exercise protocols at both positions on the centrifuge: the isolated Quinton (model 845) ergometer at one couch

Phase II. Centrifuge Training - Exercise, Acceleration and Workload Over Time.




Figure 5.

Phase III. Combined Centrifuge Training. Acceleration Plus Workload Over Time.




Figure 6.

Table 3. Individual pre-training supine maximal ergometer exercise data on the seven men.

|           | Oxygen uptake,        |                                           | RER  | Heart rate,<br>b • min <sup>-1</sup> | V <sub>E</sub> ,<br>L • min <sup>-1</sup> | Exerc. load,<br>kg-m • min <sup>-1</sup> | RPE |
|-----------|-----------------------|-------------------------------------------|------|--------------------------------------|-------------------------------------------|------------------------------------------|-----|
|           | L • min <sup>-1</sup> | ml • kg <sup>-1</sup> • min <sup>-1</sup> |      |                                      |                                           |                                          |     |
| FLE       | 2.62                  | 37.0                                      | 1.36 | 166                                  | 104                                       | 1600                                     | 10  |
| FRE       | 3.44                  | 38.3                                      | 1.21 | 170                                  | 115                                       | 1800                                     | 10  |
| HUN       | 2.69                  | 27.9                                      | 1.21 | 174                                  | 102                                       | 1500                                     | 10  |
| JAG       | 3.18                  | 37.9                                      | 1.28 | 186                                  | 111                                       | 1800                                     | 10  |
| RAY       | 3.31                  | 39.6                                      | 1.28 | 188                                  | 138                                       | 1700                                     | 10  |
| RUI       | 3.16                  | 36.7                                      | 1.25 | 159                                  | 122                                       | 1900                                     | 9   |
| SCH       | 2.86                  | 37.2                                      | 1.26 | 188                                  | 99                                        | 1500                                     | 10  |
| $\bar{X}$ | 3.04                  | 36.4                                      | 1.26 | 176                                  | 113                                       | 1686                                     | 10  |
| SD        | 0.32                  | 3.9                                       | 0.05 | 12                                   | 14                                        | 157                                      | 0   |
| SE        | 0.12                  | 1.5                                       | 0.02 | 4                                    | 5                                         | 59                                       | 0   |
|           |                       |                                           |      |                                      |                                           |                                          |     |
|           |                       |                                           |      |                                      |                                           |                                          |     |

RER = respiratory exchange ratio;  $\dot{V}_E$  BTPS = ventilation; RPE = rated perceived exertion.

and the combined HPC exercise + acceleration station at the other couch (fig. 7). The subjects were secured to the couches with shoulder braces and a four-point shoulder and lap harness; handgrips were used for body stabilization and leverage during the maximal tests. Thus arm, shoulder, and trunk muscular contractions were added to the lower leg exercise metabolism in determining  $\dot{V}O_{2max}$ . A preliminary continuous HPC  $\dot{V}O_{2max}$  test (Part 1) was performed 3 to 4 weeks before the main experiments in order to estimate the maximal workload (70 rpm with 300 kg-m•min<sup>-1</sup> increments until heart rate reached 180 b•min<sup>-1</sup> or volitional fatigue occurred); this was followed by a 5-min cooling-down period (fig. 8, left panel). After at least 3 days of recovery, the maximal exercise protocol (Part 2) was performed again (fig. 8, right panel) where the subjects warmed up at 70 rpm at about 40% of their maximal workload (O<sub>2</sub> uptake) determined in Part 1 above. They then exercised for 2-min intervals at progressively greater workloads starting at about 400 kg-m•min<sup>-1</sup> below the Part 1 estimated maximal load and continuing to 200 kg-m•min<sup>-1</sup> below maximal; then the maximal load was undertaken. If the subjects completed 2 min at this “maximal” load, it was increased by 200 kg m•min<sup>-1</sup> each min until a heart rate of 180 b•min<sup>-1</sup> or volitional fatigue occurred. A 5-min cooling-down period followed. This abbreviated maximal testing protocol was used to minimize training effects. Heart rate, integrated from the ECG, and ratings of perceived exertion (RPE) were recorded during all maximal tests.




Figure 7.




Figure 8

Maximal human-powered centrifuge protocol. Pre-training familiarization tests and tolerances were determined with the HPC (Greenleaf et al., 1999) 3 to 4 weeks before the experiments began (fig. 3). The subjects were secured in the couch with the 4-point restraint harness and to the pedals with toe-clips and Velcro® straps; a blindfold was worn for all acceleration runs and radio headsets (model H3391, David Clark Co., Inc., Worcester, Mass.) were used to maintain communications among the subject, research team, and medical monitor (fig. 9). The preliminary HPC test (Part 1) consisted of a 9-min warm-up period at +0.5 Gz followed by 4 min at +1.0 Gz, 2-min rest, 4 min at +2.0 Gz, 2-min




Figure 9.

rest, and then the maximal run (rpm) to volitional fatigue – followed by a cooling-down period (fig. 10, left panel). Then, after at least 3 days of recovery, the final maximal protocol (Part 2) was performed; it began with a 5-min warm-up period at about 40% of the maximal rpm (fig. 10, right panel). This was followed by 2 min at 60%, 2 min at 80%, and then increased by 20% of  $\text{rpm-max} \cdot \text{min}^{-1}$  until volitional fatigue – followed by a 5-min cooling-down period. Because of the physical constraints of the couch on the 1.9-m centrifuge radius, all of the taller subjects commented about being cramped during exercise, and felt that it attenuated their performance.

Platform rpm (angular acceleration); that is, pedal cadence, was maintained with a digital tuner metronome (model DTM-12, Korg, Tokyo, Japan) through the subject's headset. Oxygen uptake, heart rate, ECG, RPE (modified Borg scale), and Gz level and platform rpm were recorded during all maximal acceleration protocols.

The CPX Express metabolic analyzer was calibrated by John Hoppe of Vacumetrics Inc./Vacu•Med Division. The difference between the CPX and Vacu•Med data are presented in appendix E. The mean respective  $\text{O}_2$  and  $\text{CO}_2$  differences were -1.13 and +0.09% (medium range) and -1.68 and +5.10% (high range).

Tilt-table (orthostasis) protocol. After one practice tilting without blood sampling, the experimental tilt test was conducted before and after each phase (fig. 3) with the subjects initially in the horizontal, supine body position (fig. 11) on the tilt table (Physical Therapy Treatment Table, Laberne Mfg. Co.,




Figure 10.




Figure 11.

Columbia, S.C.). After 35 min of supine rest (fig. 11), an 18-gauge Teflon catheter (Quick-Cath, Baxter Healthcare Corporation, Deerfield, Ill.) was inserted into the subject's right antecubital vein and 27 ml of blood collected for control measurements. Evans blue dye (T-1824: 2.5 ml, 25 mg•5 ml<sup>-1</sup>, New World Trading Corporation, DeBary, Fla.) was then injected through the catheter for determination of resting plasma volume (PV); the catheter was subsequently flushed with saline (0.9% NaCl, McGaw, Inc., Irvine, Calif.). A venous blood sample (5 ml) was withdrawn at 10 min from the left arm and added carefully into a tube containing lithium-heparin. The table then was tilted to 70° head-up until the onset of pre-syncopal signs or symptoms such as a sudden decrease in systolic blood pressure > 25 mmHg•min<sup>-1</sup>, decrease in diastolic blood pressure > 15 mmHg•min<sup>-1</sup>, sudden drop in heart rate > 15 b•min<sup>-1</sup>, accompanying nausea, clammy skin, profuse sweating, or pallor. If none of these signs or symptoms was severe enough for termination, the test was stopped at 60 min. No subject reached unconsciousness. After only one +Gz<sub>max</sub> run subject RAY reported nausea, and subject SCH was dizzy, nauseated, and vomited after his run. Thus, both passive and active (exercise) acceleration resulted in far fewer adverse signs and symptoms than occurred after tilting (table 4).

Blood (23 ml) was collected from the catheter (after waste withdrawal) following return of the subjects to the supine position (fig. 11). About 480 ml of blood were taken over the 20-week study period. A 24-hr urine sample was collected just prior to each tilt test (fig. 3).



**Table 4. Individual baseline (BL) and pre- and post-training pre-syncope signs-symptoms (PS-S) and tilt time for the three phases during tilt test.**

| <b>Phase I. Pre-Training (16 July) and Passive Training</b> |         |         |                                   |                |      |
|-------------------------------------------------------------|---------|---------|-----------------------------------|----------------|------|
|                                                             |         |         | Signs and symptoms                | Tilt time, min | PS-S |
| FLE                                                         | No Tilt |         |                                   |                |      |
| FRE                                                         | BL      | 16 July | OK                                | 60             | No   |
|                                                             | Pre     | 10 Sept | OK                                | 60             | No   |
|                                                             | Post    | 1 Oct   | OK                                | 60             | No   |
| HUN                                                         | BL      | 16 July | Tachycardia                       | 60             | No   |
|                                                             | Pre     | 10 Sept | OK                                | 60             | No   |
|                                                             | Post    | 1 Oct   | OK                                | 60             | No   |
| JAG                                                         | BL      | 16 July | Dizziness                         | 23             | Yes  |
|                                                             | Pre     | 23 July | Light-headed                      | 22             | Yes  |
|                                                             | Post    | 13 Aug  | Strange feeling in head           | 27             | Yes  |
| RAY                                                         | BL      | 16 July | Cold sweat, dizzy, nausea         | 43             | Yes  |
|                                                             | Pre     | 23 July | Cold sweat, pale, nausea          | 32             | Yes  |
|                                                             | Post    | 13 Aug  | OK                                | 60             | No   |
| RUI                                                         | BL      | 16 July | Sweating, warm                    | 60             | Yes  |
|                                                             | Pre     | 23 July | Pale, nervous                     | 30             | Yes  |
|                                                             | Post    | 13 Aug  | OK                                | 60             | No   |
| SCH                                                         | BL      | 16 July | Can't relax, feet tingling        | 54             | Yes  |
|                                                             | Pre     | 10 Sept | Feet tingling, eye light flashing | 59             | Yes  |
|                                                             | Post    | 1 Oct   | OK                                | 60             | No   |

| <b>Phase II. Exercise Training</b> |         |         |                                    |    |     |
|------------------------------------|---------|---------|------------------------------------|----|-----|
|                                    |         |         |                                    |    |     |
| FLE                                | No Tilt |         |                                    |    |     |
| FRE                                | Pre     | 23 July | OK                                 | 60 | No  |
|                                    | Post    | 13 Aug  | OK                                 | 60 | No  |
| HUN                                | Pre     | 23 July | Tachycardia                        | 60 | No  |
|                                    | Post    | 13 Aug  | OK                                 | 60 | No  |
| JAG                                | Pre     | 10 Sept | Very weak                          | 46 | Yes |
|                                    | Post    | 1 Oct   | Loss of vision                     | 27 | Yes |
| RAY                                | Pre     | 10 Sept | Nausea                             | 48 | Yes |
|                                    | Post    | 1 Oct   | OK                                 | 60 | No  |
| RUI                                | Pre     | 10 Sept | Can't relax                        | 60 | No  |
|                                    | Post    | 1 Oct   | OK                                 | 60 | No  |
| SCH                                | Pre     | 23 July | Light-headed                       | 51 | Yes |
|                                    | Post    | 13 Aug  | Light-headed, facial muscle spasms | 35 | Yes |

Table 4. Concluded.

| Phase III. Combined Training |         |  |                        |    |     |
|------------------------------|---------|--|------------------------|----|-----|
| FLE                          | 29 Oct  |  | OK                     | 60 | No  |
|                              | 19 Nov  |  | OK                     | 60 | No  |
|                              | 17 Dec  |  | OK                     | 60 | No  |
| FRE                          | 29 Oct  |  | OK                     | 60 | No  |
|                              | 19 Nov  |  | OK                     | 60 | No  |
|                              | 17 Dec  |  | OK                     | 60 | No  |
| HUN                          | 29 Oct  |  | BP fell dramatically   | 59 | Yes |
|                              | 19 Nov  |  | BP fell dramatically   | 36 | Yes |
|                              | 17 Dec  |  | OK                     | 58 | No  |
| JAG                          | 29 Oct  |  | Light-headed, headache | 27 | Yes |
|                              | 19 Nov  |  | Nausea                 | 19 | Yes |
|                              | 17 Dec  |  | OK                     | 28 | No  |
| RAY                          | 29 Oct  |  | Sweating, nausea       | 15 | Yes |
|                              | 19 Nov  |  | Sweating, nausea       | 41 | Yes |
|                              | 17 Dec  |  | OK                     | 60 | No  |
| RUI                          | 29 Oct  |  | OK                     | 60 | No  |
|                              | 19 Nov  |  | OK                     | 60 | No  |
|                              | 17 Dec  |  | OK                     | 60 | No  |
| SCH                          | No Tilt |  |                        |    |     |

Physiological variables measured or calculated during the tilt test were heart rate from the ECG, thoracic impedance index, stroke volume (SV), and cardiac output ( $\dot{Q}$ , Cardiodynamic monitor, BoMed Medical Mfg., Ltd., Irvine, Calif.). Additional measurements that were monitored continuously included noninvasive arterial pressure (model 2300, Finapres Ohmeda, Englewood, Colo.; calibrated with a W.A. Baum Co., Inc. Baumanometer sphygmomanometer, Copiague, N.Y.), radial arterial blood flow (model 909, Parks Medical Electronic, Inc., Aloha, Oreg.), forearm skin perfusion (model PF4001, Perimed Periflux Smithtown, N.Y.), temple skin perfusion (model BPM 403A, LaserFlo Skin Perfusion Monitor, TSI, St. Paul, Minn.), and calf circumference with a sylvastic strain gauge (model EC-4, D.E. Hokansen, Issaquah, Wash.); peripheral vascular resistance was calculated off-line from arterial pressure/cardiac output. All data were presented on a chart recorder (model MT 8800, Astro-Med, Inc., West Warwick, R.I.) and stored on an analog tape recorder (model MR-40, Teac, Japan) and in a computer (PC, Comteq Computer Co., Rockville, Md.; Windaq model DI-220, Dataq Instruments, Inc., Akron, Ohio) with a 250-Hz sampling rate.

Blood sampling and analyses. After whole blood was allocated for hematocrit (Hct) and hemoglobin concentration [Hb] analyses, the remaining blood was placed into chilled and treated tubes and centrifuged for 15 min at 1,500 G at 4°C – except growth hormone samples which were centrifuged for 20 min at 1,000 G (model RC2-B, Ivan Sorvall, Inc., Newton, Conn.). Hematocrit was measured immediately, while plasma for osmotic and Evans blue dye analyses was refrigerated at 4°C for analysis later that day. All other samples were stored at -70°C for analysis after the study. Tube treatment was potassium-EDTA plus 400 IU•mL<sup>-1</sup> aprotinin for [Hb] and (PRA), [PVP], and [PA]; lithium-heparin for Hct, growth hormone (I-GH, B-GH, I-GF-1), and plasma osmotic, sodium,

potassium, albumin, and total protein concentrations; and EDTA plus reduced glutathione-saline-NaOH for [PE] and norepinephrine [PNE] concentrations.

Plasma volume was measured in supine subjects with the Evans blue dye technique from one 10-min post-injection blood sample (Campbell et al., 1958; Greenleaf et al., 1979). Syringes (Beckton-Dickenson, Franklin Lakes, N.J.) used for dye injection were weighed pre- and post- injection (model B6 balance, Mettler Instrument Corp., Highstown, N.J.) to determine the amount of dye injected. Blood was centrifuged for 15 min at 1,500 G (4°C) and the plasma eluate was analyzed from Sephadex columns (No. PD-10, Pharmacia LKB Biotechnology, Uppsala, Sweden) and measured at an absorbency of 615 nm (model 35 spectrophotometer, Beckman Instruments Inc., Irvine, Calif.).

Percent change in post-tilt PV was calculated from the Hct and [Hb] transformation equation (Greenleaf et al., 1979):

$$\% \Delta PV = \frac{100 [\text{Hb}_B \times (1 - \text{Hct}_A \times 10^{-2})]}{[\text{Hb}_A \times (1 - \text{Hct}_B \times 10^{-2})]} - 100$$

where: B is before and A is after tilt.

This equation has been validated for a period of only 2 hr, but F-cell ratio errors should have been minimized (more equal distribution of red blood cells) with the subjects in the supine position before tilt testing. The post-tilt PV data were back-calculated to milliliters from percent changes calculated from the Hb-Hct transformation equation.

Hematocrit was measured in quadruplicate from microcapillary tubes, run for 10 min at 11,500 rpm on an International Equipment Company Micro-MB centrifuge (Needham Heights, Mass.), and read on a modified microcapillary tube reader (model CR Micro-capillary reader, International Equipment Company, Needham Heights, Mass.) sensitive to  $\pm 0.1$  Hct units. Raw Hct values were corrected for trapped plasma (0.96) and for whole body Hct (0.91). Blood [Hb] was measured in duplicate with a cyanomethemoglobin method (Sigma Diagnostics®, St. Louis, Mo.).

Plasma osmolality ( $\pm 1$  mosmol $\cdot$ kgH<sub>2</sub>O<sup>-1</sup>) was determined by freezing-point depression (model 3R osmometer, Advanced Instruments, Needham Heights, Mass.). Plasma total protein, albumin, sodium, and potassium analyses variability ( $\pm$  SD, cv) of 10 runs on one sample were determined with a Beckman Coulter Synchron analyzer (model LX20, Beckman Coulter, Inc., Brea, Calif.): total protein ( $\pm 0.05$  g $\cdot$ dL<sup>-1</sup>, 1.0) with the biuret reaction (Kingsley, 1942), albumin ( $\pm 0.05$  g $\cdot$ dL<sup>-1</sup>, 1.5) with Bromecresol purple binding (Pinnell and Northam, 1978), and indirect potentiometry (electrodes) for both sodium ( $\pm 0.05$  mmol $\cdot$ l<sup>-1</sup>, 0.4) and potassium ( $\pm 0.04$  mmol $\cdot$ l<sup>-1</sup>, 0.8) (Eisenman, 1967; Pioda et al., 1969).

Plasma hormone concentrations were analyzed by Helmut Hinghofer-Szalkay with radioimmunoassay, except the catecholamines which were analyzed by Michael Ziegler with the radioenzymatic method of Kennedy and Ziegler (1990). This modification of the catechol-o-methyltransferase (COMT)-based radioenzymatic assay for [PNE] and [PE] improves sensitivity and

selectivity, and eliminates many inhibitors of COMT. Prior to assay, the samples were extracted into heptane with diphenylborate and then put into dilute acetic acid. This extraction procedure has an efficiency of 78% for NE, but less than 2% for S-adenosylmethionine. The extraction procedure also excludes calcium and other COMT inhibitors present in urine, plasma, and every other tissue tested. This eliminated the requirement for individual standardization of tissue and urine samples. Sensitivity of the assay for PNE and PE in 1 ml of plasma was 10 and 6 pg•ml<sup>-1</sup>, respectively. The respective intra-assay cv for PNE and PE were 4% and 13%, and the inter-assay cv for PNE and PE were 10% and 16%, respectively, in human plasma containing low catecholamine levels. The assay permits quantitation of PE levels that were undetectable in prior assays.

Plasma aldosterone was measured with a modified radioimmunoassay (AldoCTK-2, Sorin Biomedica, Italy); its sensitivity, defined as the apparent concentration of analyte that can be distinguished from the zero standard, was below 20 pg•ml<sup>-1</sup> at the 95% confidence limit. The cv for the within- and between-assay variability was 9.7% and 11.5%, respectively. Plasma vasopressin was determined on ethanol-extracted plasma with a radioimmunoassay kit (Nichols Institute, Diagnostics BV, The Netherlands) with <sup>125</sup>I-AVP as the labeled compound. The anti-AVP antiserum did not cross-react with Lys<sup>8</sup>-vasopressin, oxytocin, or vasotocin. Sensitivity at the 99% confidence limit was 1.3 pg•ml<sup>-1</sup>. Plasma renin activity was determined by measurement of ANG-I (RENCTK, Sorin Biomedica, Italy) based on competition between labeled and native ANG-I to be assayed for a fixed number of antibody binding sites; PRA was expressed as nanograms of ANG-II formed per milliliter of plasma after 1 hr of incubation. Sensitivity was < 0.20 ng•ml<sup>-1</sup> at the 95% confidence limit.

Plasma growth hormone [PGH] was measured by radioimmunoassay (RIA) and bioassay by Richard Grindeland. Eight-milliliter aliquots of blood, obtained before and after each experimental treatment, were put into 15-ml plastic centrifuge tubes containing 100 ul (100 units) of heparin, mixed gently, and placed in an ice bath. These tubes were then centrifuged at 1,000 G for 20 min at 5°C and the plasma transferred to cryovials for storage at -70°C. The frozen samples were thawed in a 37°C water bath, care being taken to not unduly heat the plasma. An aliquot of 0.5 – 1.0 ml of plasma was left in each tube for the immunoassay; the balance was pooled by experimental treatment, the volume measured, and the pooled samples put into siliconized vials until injected into bioassay rats. The plasma was bioassayed according to the procedure of Greenspan et al., (1949) in 40-day-old female albino rats which had been hypophysectomized at 26 days of age by the vendor (Hilltop Laboratories, Scottsdale, Pa.). After surgery the rats were allowed to recover for 3 days and were then shipped to Ames Research Center where they were inspected and allowed to acclimate to the new environment and diet and to regain their body water balance. The assay rats were weighed 7 and 14 days after surgery; animals that gained more than 1 g of body weight per day over that interval were considered incompletely hypophysectomized and were not used in the assay. Rats that lost more than 2 g total weight were also removed from the assay.

Five rats were put into each treatment group or dose level of standard hormone. Three dose levels (1, 5, and 15 µg total dose) of standard bovine GH (1.5 IU•mg<sup>-1</sup>) were employed. Pooled plasma, standard, or saline was injected intraperitoneally (0.5 ml•day<sup>-1</sup>) for 4 days; on the fifth day the rats were euthanized with an overdose of CO<sub>2</sub> and a tibia dissected out, split longitudinally, and stained with silver nitrate. The stained tibias were put into individual vials containing 70% ethanol until the proximal growth plates were read microscopically (Carl Zeiss, Germany) to the nearest micrometer. Ten readings of the epiphyseal plate thickness were made, averaged for each rat, then averaged by group, and then

compared to a standard curve derived from bovine growth hormone standards. Because of the limited volume of plasma, only a single dose level of plasma was used. The mean [PGH] and 95% confidence limits were calculated using a "bracketed three point assay" (Bliss, 1952). The [PGH] per milliliter of plasma was calculated as a function of the epiphyseal width. The bovine GH had a specific biological activity one half that of highly purified human GH, so values ( $\text{ng}\cdot\text{ml}^{-1}$ ) obtained from the standard curve were divided by 2 to yield human GH values. The commercial kits used for radioimmunoassays (Diagnostic Products, Inc. Los Angeles, Calif.) employed a double antibody system similar to the procedure of Utiger et al. (1962). Human growth hormone (hGH) standards ( $1\text{-}30\text{ ng}\cdot\text{ml}^{-1}$ ) or sample ( $100\text{ }\mu\text{l}$ ) were pipetted into duplicate assay tubes. The standard hormone was NIH preparation NIAMDD hGH RP-1. Next,  $100\text{-}\mu\text{l}$  aliquots of rabbit anti-human GH serum were added. The reagents were mixed on a vortex mixer and incubated for 1 hr at room temperature followed by addition of the radio-iodinated hGH, mixed again and incubated for another hour at room temperature. Then the cold second antibody (goat anti-rabbit gamma globulin) mixed with polyethylene glycol was added, the contents mixed, and the tubes centrifuged at  $1,500\text{ G}$  for 30 min at  $5^{\circ}\text{C}$ . Except for one pair of tubes used to determine total counts, the supernatant was aspirated and the tubes were inverted to drain. The tubes were counted in a Packard Cobra model crystal scintillation counter for 1 min. After automatic subtraction of nonspecific binding, the results were plotted automatically as a log dose-logit curve and the results expressed as  $\text{ng hGH}\cdot\text{ml}^{-1}$ . The interassay coefficient of variation was 5% and the intraassay coefficient was 4%; outlier samples were reassayed.

Urine collection and analyses. A 24-hr urine volume, collected just before each tilt test, was measured to  $\pm 5\text{ ml}$  in a graduated cylinder and aliquots were frozen for subsequent analysis by Scott Smith. Urinary collagen cross-links (pyridinium, sensitivity  $7.5\text{ nmoles}\cdot\text{l}^{-1}$ ) and deoxypyridinoline (sensitivity  $1.1\text{ nmoles}\cdot\text{l}^{-1}$ ) were determined with the Pylinks™ and Pylinks-D™ kits, respectively (Metra Biosystems, Palo Alto, Calif.). Concentrations of n-telopeptide were measured with the Osteomark® ELISA kit (Ostex International, Seattle, Wash.) which detects the n-telopeptide region of bone collagen specifically in human urine (Smith et al., 1998, 1999) with a sensitivity of  $20\text{ nmoles}\cdot\text{l}^{-1}$ . Hydroxyproline analysis was performed on an amino acid analyzer (model L-8800, Hitachi Corp., San Jose, Calif.) using methods adapted from Paroni et al. (1992) and Slocum and Cummings (1990). Urine samples and internal standard (glucosaminic acid, TCI America, Portland, Oreg.) were hydrolyzed in  $6\text{N HCl}$  for 16 hr at  $110^{\circ}\text{C}$ . The hydrolysate was adjusted to a pH of 2.2 and concentrated using a Speedvac (Savant Corp., Holbrook, N.Y.). The solution was filtered through a  $0.2\text{-}\mu\text{m}$  filter and brought to volume with lithium buffer solution (pH 2.2). The protein-free filtrate was refrigerated until injected onto the analyzer. Post-column ninhydrin derivatization produces a chromagen detectable at  $440\text{ nm}$  with a sensitivity of  $0.021\text{ }\mu\text{moles}\cdot\text{ml}^{-1}$ . Urinary calcium was determined using an atomic absorption spectrophotometer (model 4000, Perkin-Elmer, Norwalk, Mass., with a sensitivity of  $5\text{ mg}\cdot\text{dl}^{-1}$ ). Creatinine was also determined spectrophotometrically (Owen et al., 1954) with a sensitivity of  $0.1\text{ mg}\cdot\text{dl}^{-1}$ .

Magnetic Resonance Imaging (MRI). The lower body of each subject was imaged 2 days before and 1 day after completion of each of the 3-week training regimens on a 1.5 Tesla whole-body MR imager (Siemens Vision™, Iselin, N.J.) at Los Gatos MRI (Los Gatos, Calif.) by Peter Hardy. The subjects were supine with their feet at the center of the imaging magnet; their thighs were imaged by the body coil of the imager. Three types of images were obtained during each session: (1) 1-cm-thick T1-weighted images of the pelvis and thighs to estimate the volume of the rectus femoris, vastus lateralis, vastus intermedius, and vastus medialis; (2) a flow-alternating inversion recovery (FAIR) image

estimated perfusion of the quadriceps muscle; and (3) a multi-echo technique measured the spin-spin relaxation time (T2); that is, values proportional to the amount of work a muscle performs. The multi-echo spin images (T2) were calculated pixel-by-pixel for each of the five slices acquired through the thighs. The IDL software allowed the user to select an arbitrarily defined region in the muscle from which to obtain basic statistics such as average and standard deviation over the region. The average T2 in each head of the quadriceps muscle was extracted from the images taken before and after the subjects did a maximal number of deep knee bends. The change in T2 ( $T2_{\text{post}} - T2_{\text{pre}}$ ) was calculated for a given muscle.

Total imaging time was approximately 45 min. The MR images were transferred onto magneto-optical disks for analysis at the University of Kentucky. The volumes of the four heads of the quadriceps muscle in each leg were determined by manually outlining the muscles using the commercially available image analysis software 3DVIEWNIX. Total muscle volume was the sum of the areas outlined on each image multiplied by the slice-to-slice separation.

The T2 data were determined from the multi-echo images, using user-derived software, and perfusion data were derived from the FAIR images, using custom software written in IDL.

## DATA ANALYSIS

Hemodynamic. A data analysis system devised by David Brown was developed for the Intel Windows platform using Microsoft Visual C. Analysis routines were designed to organize, group, scan, and average a large number of data sets to provide quick and easy access to signal processing, and to statistical and data base functions. A data collection system was developed using the Intel Windows platform and Microsoft Visual C++. Data acquisition used National Instruments E-series analog-to-digital converters and a full-screen high-performance mode to sample and display continuously up to 16 channels of data at sample rates from 1 to 10,000 Hz (or higher depending on the AD board).

All digitally sampled (250-Hz) wave forms (arterial pressure, peripheral flows, respiration, etc.) were analyzed to give both integrated mean values as well as spectral power results using the data analysis system described above. After R-wave detection, instantaneous R-R interval time series were constructed from the ECG. The resulting piecewise constant time series was low-pass filtered and sampled at 5 Hz. Auto- and cross-spectral estimates were computed using averaged periodograms. A Hanning window was used to reduce side-lobe error. Estimates of coherence were computed as a ratio of the squared magnitude of the cross spectrum divided by the product of the two autospectra. For each variable, the 10 min of resting control and 60 min of tilt data were divided into 200-sec segments for analysis. Mean values and spectral power from each rest and control period were averaged within subjects to enhance robustness. Results were then averaged across subjects for plotting and statistical analysis.

Statistical. Means and standard deviations ( $\pm$  SD) were calculated for subject characteristics, and means and standard errors ( $\pm$  SE) were calculated for all other variables. The exercise, tilt, and blood variables were analyzed with a three-factor analysis of variance (SPSS 7.5 for Windows; SPSS, Inc., 1996, Chicago, Ill.) with significant differences between the means determined with the Dunnett post-

hoc procedure. In addition, mean values and spectral power of the hemodynamic data were analyzed using a three-factor ANOVA (PC SAS version 6.12, SAS Institute, Inc., Cary, N.C.) to assess the significance of main effects: training protocols and tilting before and after training. When post-hoc testing was warranted (significant F-ratio for the effect of interest), t-statistics with degrees of freedom determined by Satherwaite's approximation, were constructed to compare mean responses. In all cases statistical significance was determined at  $P < 0.05$  and nonsignificant changes were NS.

## RESULTS

### Maximal Exercise (Passive, Exercise, Combined) Data

Exercise load. The maximal exercise loads were increased ( $P < 0.05$ ) for all three Phases after training (fig. 12a) in stepwise order from Passive (by  $8.3 \pm 4.3\%$ ), to Exercise (by  $12.6 \pm 5.7\%$ ), to Combined (by  $15.4 \pm 2.6\%$ ) (fig. 12b).

Oxygen uptake. However, maximal oxygen uptake tended to decrease (NS) with Passive training and to increase (NS) with Exercise and Combined training (figs. 13a and 13b).

Heart rate. Maximal exercise heart rates ( $HR_{max}$ ) were decreased by  $3 \pm 2\%$  (NS) with Passive, unchanged ( $0 \pm 2\%$ ) with Exercise, and tended to increase by  $2 \pm 2\%$  (NS) with Combined training (figs. 14a and 14b).

Exercise tolerance. Maximal exercise time to fatigue was unchanged ( $+2.6 \pm 3.4\%$ , NS) with Passive, but was increased by  $6.0 \pm 4.9\%$  ( $P < 0.05$ ) with Exercise, and by  $17.9 \pm 3.2\%$  ( $P < 0.05$ ) with Combined training (figs. 15a and 15b).

### Orthostatic (Tilt-Table) Cardiovascular Data

Resting pre-tilt heart rate. Resting HR (mean of 2 min before tilting) was unchanged with the three training Phases (fig. 16a), but was elevated by  $12.9 \pm 5.2\%$  ( $P < 0.05$ ) after Passive training (fig. 16b).

Resting pre-tilt systolic blood pressure. Mean resting SBP was unchanged after training in each of the respective Phases (fig. 17a). However, it tended to increase (NS) with Passive and decrease (NS) with Exercise and Combined Phases (fig. 17b).

Resting pre-tilt diastolic blood pressure. Mean resting DBP was also unchanged after training in each of the respective Phases (fig. 18a), and it also tended to be lower (NS) with Exercise and Combined Phases (fig. 18b).

Resting pre-tilt mean arterial pressure. Average MAP followed the SBP and DBP with a tendency to increase (NS) with Exercise and Combined Phases (figs. 19a and 19b).

Tilt-tolerance time. Mean tilt-tolerance time pre- to post-training was increased only with Passive training from  $43.5 \pm 7.2$  to  $54.7 \pm 5.3$  min, respectively (fig. 20a); that is, by  $37.8 \pm 19.6\%$  ( $P < 0.05$ ) (fig. 20b); tolerances were unchanged with the Exercise and Combined Phases.

Heart rate at tolerance. Mean HR at tolerance pre- to post-training was increased from  $71 \pm 8$  to  $89 \pm 7$  bpm, respectively (fig. 21a); that is, by  $29.1 \pm 7.7\%$  ( $P < 0.05$ ) with Passive training; HR was unchanged with the Exercise and Combined Phases.

Mean arterial pressure at tolerance. Average MAP at tolerance was increased from  $88 \pm 13$  to  $108 \pm 10$  mmHg (by  $33.4 \pm 18.8\%$ ,  $P < 0.05$ ) with Passive, and from  $78 \pm 9$  to  $96 \pm 6$  mmHg (by  $28.8 \pm 14.7\%$ ,  $P < 0.05$ ) with Combined; Exercise MAP was unchanged (figs. 22a and 22b).

Cardiac R-R interval with training. R-R intervals were not different at rest for the six treatments pre- and post-tilt for the three Phases (fig. 23). The greater the R-R interval, the slower the HR. But the intervals were uniformly lower ( $P < 0.05$ ) for the six post-tilt treatments with no differences between or among Phases.

Stroke volume with training. Stroke volume, similar to the R-R interval, was not different at rest for the six treatments for the three Phases (fig. 24), but it was uniformly lower ( $P < 0.05$ ) at about  $60 \text{ ml} \cdot \text{beat}^{-1}$  for the six post-tilt treatments with no differences between or among Phases.

End-diastolic volume with training. E-D volume was not different at rest for the six treatments for the three Phases (fig. 25) but it was also uniformly decreased ( $P < 0.05$ ) for the six post-tilt treatments with no differences between or among Phases.

Cardiac output with training. Cardiac output was not different at rest for the six treatments for the three Phases (fig. 26), and  $\dot{Q}$  was uniformly decreased ( $P < 0.05$ ) for the six post-tilt treatments with no differences between or among Phases.

Cuff arterial pressure with training. Arterial pressure was not different at rest or post-tilt for the six respective treatments for the three Phases (fig. 27). But all post-tilt values tended (NS) to be higher than their corresponding rest values.

Total peripheral resistance with training. Total PR was not different at rest for the six treatments for the three Phases (fig. 28). But PR was uniformly increased ( $P < 0.05$ ) for the six post-tilt treatments with no differences between or among Phases.

Cardiovascular variables. (figs. 29a and 29b).

Blood hemoglobin [Hb]. Hemoglobin concentration was not different at rest for the six treatments or at post-tilt for the six treatments in the three Phases (fig. 30a). But Hb post-tilt was increased ( $P < 0.05$ ) for five treatments (except for post-Combined) reflecting decreases in plasma volume post-tilt. This is also indicated in the percent changes in Hb at rest and post-tilt (fig. 30b). Percent changes in Hb at tilt tolerance pre- and post-training (fig. 30c) indicate significant ( $P < 0.05$ ) increases in five treatments—except for post-training in the Combined Phase.



Raw hematocrit (Hct). Hematocrit was not different at rest for the six treatments or at post-tilt for the six treatments in the three Phases (fig. 31a). But Hct was uniformly increased ( $P < 0.05$ ) for the six post-tilt treatments (reflecting decreases in plasma volume) with no differences between or among Phases. This also reflected decreases in plasma volume post-tilt that are reflected in the percent changes in Hct at rest and post-tilt (fig. 31b). Percent changes in Hct at tilt tolerance pre-and post-training indicated significant ( $P < 0.05$ ) increases in all six treatments (fig. 31c).

Plasma volume (PV). Measured (T-1824) PV was not different at rest or post-tilt for the six treatments in the three Phases (fig. 32a). But PV was uniformly decreased ( $P < 0.05$ ) for five treatments—except for post-training in the Combined Phase. These results are also indicated in the percent changes in Hct at rest and post-tilt (fig. 32b). Percent changes in PV at tilt-tolerance pre-and post-training (fig. 32c) indicate significant ( $P < 0.05$ ) decreases for five treatments—except for post-training in the Combined Phase.

Plasma sodium [PNa]. Plasma sodium concentration varied between 136 and 138  $\text{mmol}\cdot\text{l}^{-1}$  with post-tilt levels generally lower than comparable rest (pre-tilt) values, except for Exercise post-training (fig. 33a). Percent changes in [PNa] at rest and post-tilt (fig. 33b) were uniformly higher with Passive and Exercise Phases, when compared with essentially unchanged Combined data. Conversely, all percent changes in [PNa] at tolerance pre-and post-training were negative and varied from 0.4 to 0.8 (all NS). Sodium accompanied the vascular-to-interstitial-fluid space shift of PV, so [PNa] remained unchanged.

Plasma potassium [PK]. Plasma potassium concentration varied between 3.9 and 4.3  $\text{mmol}\cdot\text{l}^{-1}$  at rest and post-tilt (fig. 34a), with reduced percent change at rest and post-tilt for the three Phases except for Exercise post-tilt (fig. 34b). Percent changes at tolerance with training were positive (NS) with Passive and Exercise and negative (NS) with Combined Phases; the greatest post-training increase in [PK] was with Exercise and the greatest decrease with Combine (fig. 34c).

Plasma osmolality [POsm]. Plasma osmotic concentration varied between 286 and 289  $\text{mOsm}\cdot\text{kg}^{-1}$  at rest and post-tilt with somewhat lower (NS) values post-tilt, except for Combined (fig. 35a). There were greater percent changes with training in Passive and Exercise, and the lower changes with Combined (fig. 35b) were qualitatively similar with comparable [PNa] responses. There was a minimal percent change in [POsm] with training at tilt-tolerance (fig. 35c) which was similar to comparable [PNa] changes in response to outward shifts of PV. Percent changes in [POsm] were in the same negative direction but were much smaller than comparable [PNa] shifts, and [PK] tended to increase.

Plasma albumin [PAIb]. Plasma albumin concentration was increased (most  $P < 0.05$ ) post-tilt pre-and post-training (fig. 36a). It tended to decrease (NS) at rest and post-tilt with training with Passive and Exercise, but to increase (NS) with Combined (fig. 36b). All percent changes at tolerance pre-and post-training were uniformly positive (range 5.1% to 9.1%), some significantly increased from zero, with no difference among them (fig. 36c). These increases in [PAIb] reflect losses (shifts) in PV.

Plasma total protein [PTP]. Plasma total protein concentration responses were qualitatively similar to those of [PAIb]: compare figures 36b and 37b, and 36c and 37c.

**Plasma renin activity [PRA].** Renin activity was not different at rest pre-and post-training and varied within the normal range; our data were 0.45 to 0.77 ngAngI•ml<sup>-1</sup>•hr<sup>-1</sup> but all PRA were elevated (P < 0.05) post-tilt within the range of 1.51 to 2.47 ngAngI•ml<sup>-1</sup>•hr<sup>-1</sup> (fig. 38a). The percent change in both PRA's decreased (NS) with Passive, increased (NS) with Exercise, and were unchanged with Combined (fig. 38b). All percent changes in PRA at tolerance pre-and post-training were increased (P < 0.05) by 181% to 351% (fig. 38c).

**Plasma aldosterone [PA].** In general, the [PA] responses were similar, as expected, to those of PRA. The rest [PA] were not different and were within the normal range; our data were 72 to 115 pg•ml<sup>-1</sup> (fig. 39a); but all [PA] values were elevated (P < 0.05) post-tilt within the range of 238 to 276 pg•ml<sup>-1</sup>. Percent changes in rest [PA] decreased by 29% (NS) with Passive but increased by 95% (NS) with Exercise; post-tilt [PA] were unremarkable (+0.7% and +20.5%, respectively) (fig. 39b). All percent changes in [PA] at tolerance pre-and post-training were increased by 67% to 451% in the Passive and Exercise Phases (fig. 39c) similar to comparable PRA data in fig. 38c.

**Plasma vasopressin [PVP].** Plasma vasopressin concentration at rest was at the high end of normal or higher with a range of 1.4 to 6.7 pg•ml<sup>-1</sup> (fig. 40a). The resting Combined [PVP] were especially elevated at 6.7 and 4.0 pg•ml<sup>-1</sup>, respectively, post-tilt levels were greatly elevated with Passive (72.7 pre vs. 34.2 pg•ml<sup>-1</sup> post) and Exercise (35.0 pre vs. 28.1 pg•ml<sup>-1</sup> post). Percent changes in [PVP] at rest and post-tilt with training varied considerably from -13.2% Combined post-tilt to 259.9% Exercise rest (fig. 40b). However, the percent changes in [PVP] at tilt-tolerance pre-and post-training were most diverse: the increases were about 10,000% with pre-and post-training Passive and pre-training Exercise; the other three varied from 83% to 957% (fig. 40c).

**Plasma epinephrine [PE].** Plasma epinephrine concentration at rest was essentially unchanged among the six treatments; they varied from 25 to 31 pg•ml<sup>-1</sup> (fig. 41a). Post-tilt levels (except for Combined post-training) tended to be elevated (NS) from 47 to 106 pg•ml<sup>-1</sup>. At rest the percent changes in [PE] pre- versus post-training were all positive for the three Phases, but all were negative post-tilt (fig. 41b). All percent changes in [PE] at tilt-tolerance pre-training and post-training (except Combined) were positive (range 80% to 268%) (fig. 41c).

**Plasma norepinephrine [PNE].** Plasma norepinephrine concentration at rest was also essentially unchanged among those six treatments; they varied from 199 to 258 pg•ml<sup>-1</sup> (fig. 42a). All post-tilt levels were elevated (P < 0.05) from rest levels (range 381 to 581 pg•ml<sup>-1</sup>). Percent changes pre-training versus post-training at rest and post-tilt varied within ± 20% (fig. 42b). All percent changes in [PNE] at tilt-tolerance for the three Phases were increased (P < 0.05) by 90% to 132%; there were no differences among those six treatments (fig. 42c).

**Plasma dopamine [PD].** Plasma dopamine concentrations at rest and post-tilt were variable but not significantly different among the six treatments (fig. 43a). Percent changes in [PD] post-tilt were attenuated compared to rest pre-training versus post-training for the three Phases (fig. 43b), and there were similar responses at tilt-tolerance pre- and post-training (fig. 43c).

**Plasma growth hormone [PGH].** Plasma growth hormone concentration as measured by radioimmunoassay (RIA), with two exceptions, was essentially undetectable (value = 1) in the pre-

Phase resting periods (fig. 44a). At post-tilt it varied from  $2.2 \pm \text{SE } 1.1 \text{ ng}\cdot\text{ml}^{-1}$  (pre-Combined) to  $10.0 \pm 4.1 \text{ ng}\cdot\text{ml}^{-1}$  with pre-Passive. Summed post-tilt levels were highest with Passive ( $10.0 + 4.5$ ), somewhat lower with Exercise ( $3.7 + 5.4$ ), and lowest with Combined ( $3.3 + 2.2$ ) suggesting progressively less respective stress during tilting. Of the 16 cases (excluding baseline data) in Table 4 who experienced post-training pre-syncopal signs and symptoms (PS-S), 13 had increased (greater than 1) [PGH]; but there were 10 cases with increased [PGH] that did not have PS-S. Thus, onset of PS-S was minimally associated with increased [PGH].

From bioassay data, the resting levels (with two exceptions of 1375 and 1205) ranged from 720 to  $775 \text{ ng HGH}\cdot\text{ml}^{-1}$  (fig. 44b). However, unlike the RIA findings, the summed post-tilt data were highest with Combined ( $2075 + 2000 = 4075$ ), somewhat lower but similar with the Exercise ( $775 + 1850 = 2625$ ) and Passive ( $2125 + 575 = 2700$ ) Phases. Most individual Phase resting versus post-tilt values were significantly different except for post-Passive and pre-Exercise training. Clearly there are great individual differences in [PGH] after tilt that do not appear to be associated with pre-syncopal signs or symptoms.

In like manner, about half of the post-exercise [PGH] were greater than 1 before and after Combined training (see PGH table in appendix F).

### **Urine Data (24 hr)**

Urinary volume and rate (24 hr). Urinary 24-hr volumes were not different between and among pre- and post-training samples for the three Phases (fig. 45). All urinary excretion rates (milliliters per minute) were within normal limits for uncontrolled food and fluid intakes.

Urinary variables: The urinary variables were creatinine (fig. 46), deoxypyridinoline (fig. 47), deoxypyridinoline / creatinine ratio (fig. 48), n-telopeptide (fig. 49), n-telopeptide / creatinine ratio (fig. 50), pyridinium cross-links (fig. 51), pyridinium cross-links / creatinine ratio (fig. 52), hydroxyproline (fig. 53), hydroxyproline / creatinine ratio (fig. 54), and calcium (fig. 55). There were no significant differences between or among these 24-hr urinary variables pre- or post-training for the three Phases; it is noted that the 24-hr sampling time may have been too short.

### **Magnetic Resonance Imaging**

Volume. There was a significant ( $P < 0.05$ ) increase of 4% to 6% in all four quadriceps muscle volumes (T1), both right and left, post-Combined training (fig. 56).

Excitation. Change in the multi-echo spin images (T2, excitation) of their muscles during pre-Combined training was correlated 0.79 ( $r^2 = 0.62$ ) with the number of knee bends, that is, the amount of muscular work (fig. 57). Each subject performed different numbers of knee bends according to his maximal ability.

## SUMMARY OF RESULTS

### Maximal Exercise (Passive, Exercise, Combined) Data

1. Maximal supine exercise loads increased significantly ( $P < 0.05$ ) by 8.3% (Passive), by 12.6% (Exercise), and by 15.4% (Combined) after training, but their post-training maximal oxygen uptakes and maximal heart rates were unchanged. Maximal time to fatigue (endurance) was unchanged with Passive, but it too increased ( $P < 0.05$ ) with Exercise and Combined training. Thus, the exercise in the Exercise and Combined training Phases resulted in greater maximal loads and endurance without effect on maximal oxygen uptake or heart rate.

### Orthostatic (Tilt-Table) Cardiovascular Data

1. Resting pre-tilt heart rate was elevated by 12.9% ( $P < 0.05$ ) only after Passive training, suggesting that the exercise training attenuated the HR response. Resting pre-tilt blood pressures (SBR, DBP, MAP) were not different pre- or post-training in any Phase. Post-training tilt-tolerance time and heart rate were increased ( $P < 0.05$ ) only with Passive training by 37.8% and by 29.1%, respectively. Thus, addition of exercise training appeared to attenuate the increased Passive tolerance.

2. Resting (pre-tilt) and post-tilt cardiac R-R interval, stroke volume, end-diastolic volume, and cardiac output were all uniformly reduced ( $P < 0.05$ ), and peripheral resistances were uniformly increased ( $P < 0.05$ ) pre- and post-training for the three Phases, indicating there was no effect of the exercise training on these cardiovascular variables.

### Orthostatic (Tilt-Table) Biochemical Data

1. Plasma volume (percent change) was uniformly decreased by 8% to 14% ( $P < 0.05$ ) at tilt-tolerance pre-training versus post-training, indicating essentially no effect of training on the level of hypovolemia. The latter was reflected in the 6% to 12% ( $P < 0.05$ ) increase in [PAIb] and [PTP].

2. Percent changes in [PNa] pre-training versus post-training were minimal (less than -0.8%) as was [POsm] (less than -0.4%) indicating that plasma shifts during tilting were essentially isotonic.

3. The percent changes in [PRA], [PA], [PE], and [PNa] exhibited similar characteristic increases at tolerance both pre- and post-training, whereas the usual increase in [PVP] was greatly attenuated post-training with Exercise, and pre- and post-training with Combined. The explanation for the latter is not obvious but is not likely a result of technical errors.

### Urine Data (24 hr)

1. Urinary volumes were within normal limits ( $1.2$  to  $1.5 \text{ ml} \cdot \text{min}^{-1}$ ) between and among pre- and post-training samples for the three Phases.

2. There were no significant differences between or among the 10 urinary variables pre- and post-training for the three Phases.

### **Magnetic Resonance Imaging**

1. There was a 4% to 6% increase ( $P < 0.05$ ) in all four quadriceps muscle volumes (right and left) after post-Combined training.

## REFERENCES

- Bliss, C.I. The Statistics of Bioassay, with Special Reference to the Vitamins. New York: Academic Press, 1952. p. 480-525.
- Borg, G.A.V. Psychophysical bases of perceived exertion. Med Sci Sports Exerc 14:377-381, 1982.
- Buckey, J.C., Jr, L.D. Lane, B.D. Levine, D.E. Watenpaugh, S.J. Wright, W.E. Moore, F.A. Gaffney, and C.G. Blomqvist. Orthostatic intolerance after spaceflight. J Appl Physiol 81:7-18, 1996.
- Burton, R.R. A human-use centrifuge for space stations: proposed ground-based studies. Aviat Space Environ Med 59:579-582, 1988.
- Campbell, T.J., B. Frohman, and E.B. Reeve. A simple, rapid, and accurate method of extracting T-1824 from plasma, adapted to the routine measurement of blood volume. J Lab Clin Med 52:768-777, 1958.
- Chou, J.L. Human Physiological Responses during +Gz Acceleration with Cycle Ergometer Leg Exercise. San Francisco State University; M.A. thesis in Biology: Physiology and Behavioral Biology, 1997. 43p.
- Chou, J.L., G.P.N. Leftheriotis, N.J. Stad, N.F. Arndt, C.G.R. Jackson, S. Simonson, P.R. Barnes, and J.E. Greenleaf. Human physiological responses to cycle ergometer leg exercise during +Gz acceleration. Ames Research Center, Moffett Field, CA: NASA Tech Memo -1998 - 112237. 22p.
- Chou, J.L., M.A. McKenzie, N.J. Stad, P.R. Barnes, C.G.R. Jackson, F. Ghiasvand, and J.E. Greenleaf. Acceleration tolerance: Effect of exercise, acceleration training; bed rest and weightlessness deconditioning. A compendium of research (1950-1996). Ames Research Center, Moffett Field, CA: NASA Tech Memo 112214, 1997. 52p.
- Collins, J.J. , C.J. DeLuca, A.E. Pavlik, S.H. Roy, and M.S. Emley. The effects of spaceflight on open-loop and closed-loop postural control mechanisms: human neurovestibular studies on SLS-2. Exp Brain Res 107:145-150, 1995.
- Convertino, V.A. Physiological adaptations to weightlessness: Effects on exercise and work performance. Exerc Sport Sci Rev 18:119-166, 1990.
- Eisenman, G. Glass Electrodes for Hydrogen and Other Cations: Principles and Practice. New York: Marcel Dekker, Inc, 1967. 582p.
- Ellis, S., L.C. Kirby, and J.E. Greenleaf. Lower extremity muscle thickness during 30-day 6° head-down bed rest with isotonic and isokinetic exercise training. Aviat Space Environ Med 64:1011-1015, 1993.

- Greenleaf, J.E., E.M. Bernauer, A.C. Ertl, R. Bulbulian, and M. Bond. Isokinetic strength and endurance during 30-day 6° head-down bed rest with isotonic and isokinetic exercise training. Aviat Space Environ Med 65:45-50, 1994.
- Greenleaf, J.E., E.M. Bernauer, A.C. Ertl, T.S. Trowbridge, and C.E. Wade. Work capacity during 30 days of bed rest with isotonic and isokinetic exercise training. J Appl Physiol 67:1820-1826, 1989.
- Greenleaf, J.E., J.L. Chou, N.J. Stad, G.P.N. Leftheriotis, N.F. Arndt, C.G.R. Jackson, S.R. Simonson, and P.R. Barnes. Short-arm (1.9 m) +2.2 Gz acceleration: isotonic exercise load-O<sub>2</sub> uptake relationship. Aviat Space Environ Med 70:1173-1182, 1999.
- Greenleaf, J.E., V.A. Convertino, and G.R. Mangseth. Plasma volume during stress in man: osmolality and red cell volume. J Appl Physiol 47:1031-1038, 1979.
- Greenleaf, J.E., D.P. Gundo, D.E. Watenpugh, G.M. Mulenburg, M.A. McKenzie, R. Looft-Wilson, and A.R. Hargens. Cycle-powered short radius (1.9 m) centrifuge: Effect of exercise versus passive acceleration on heart rate in humans. Ames Research Center, Moffett Field, CA: NASA Tech Memo 110433, 1997. 10p.
- Greenspan, F.X., C.H. Li, M.E. Simpson, and H.M. Evans. Bioassay of hypophyseal growth hormone: tibia test. Endocrinology 45:455-463, 1949.
- Kakurin, L.I., B.S. Katkovskiy, V.A. Tishler, G.I. Kozyrevskaya, V.S. Shashkov, V.S. Georgiyevskiy, A.I. Grigoryev, V.M. Mikhaylov, O.D. Anashkin, G.V. Machinskiy, A.A. Savilov, and Ye.P. Tikhomirov. Substantiation of a set of preventive measures referable to the objectives of missions in the Salyut orbital station. Kosm Biol Aviakosm Med 12:20-27, 1978.
- Kennedy, B., and M.G. Ziegler. A more sensitive and specific radioenzymatic assay for catecholamines. Life Sci 47:2143-2153, 1990.
- Kingsley, G.R. Direct biuret method for determination of serum proteins as applied to photoelectric and visual colorimetry. J Lab Clin Med 27:840-845, 1942.
- Kotovskaya, A.R., R.R. Galle, and A.A. Shipov. Biomedical research on the problem of artificial gravity. Kosm Biol Aviakosm Med 11:12-19, 1977.
- LeBlanc, A.D., V.S. Schneider, H.J. Evans, D.A. Engelbretson, and J.M. Krebs. Bone mineral loss and recovery after 17 weeks of bed rest. J Bone Miner Res 5:843-850, 1990.
- LeBlanc, A., V. Schneider, L. Shackelford, S. West, V. Oganov, A. Bakulin, and L. Veronin. Bone mineral and lean tissue loss after long duration spaceflight. J Bone Miner Res 11:S323, 1996.
- Mulenburg, G.M., and J. Vernikos. Human powered centrifuge. US Patent 5,616,104, April, 1997.

- Nicogossian, A.E. Overall physiological response to space flight. In: Space Physiology and Medicine (2<sup>nd</sup> ed.), edited by A.E. Nicogossian, C.L. Huntoon, and S.L. Pool. Philadelphia: Lea and Febiger, 1994. p. 139-153.
- Owen, J.A., B. Iggo, F.J. Scandrett, and C.P. Stewart. The determination of creatinine in plasma or serum, and in urine; a critical examination. Biochem J 58:426-437, 1954.
- Paloski, W.H., M.F. Reschke, F.O. Black, D.D. Doxey, and D.L. Harm. Recovery of postural equilibrium control following spaceflight. Ann NY Acad Sci 656:747-754, 1992.
- Paroni, R., E. De Vecchi, I. Fermo, C. Arcelloni, L. Diomede, F. Magni, and P.A. Bonini. Total urinary hydroxyproline determined with rapid and simple high-performance liquid chromatography. Clin Chem 38:407-411, 1992.
- Pinnell, A.E., and B.E. Northam. New automated dye-binding method for serum albumin determination with bromocresol purple. Clin Chem 24:80-86, 1978.
- Pioda, L.A.R., V. Stankova, and W. Simon. Highly selective potassium ion responsive liquid-membrane electrode. Anal Lett 2:665-674, 1969.
- Shulzhenko, E.B., I.F. Vil-Vilyams, E.A. Aleksandrova, and K.I. Gogolev. Prophylactic effects of intermittent acceleration against physiological deconditioning in simulated weightlessness. Life Sci Space Res 17:187-192, 1979.
- Shulzhenko, E.B., I.F. Vil-Vilyams, M.A. Khudyakova, and A.I. Grigoryev. Deconditioning during prolonged immersion and possible countermeasures. Life Sci Space Res 14:289-294, 1976.
- Slocum, R.H., and J.G. Cummings. Amino acid analysis of physiological samples. In : Techniques in Diagnostic Human Biochemical Genetics, edited by F.A. Hommes. New York: Wiley-Liss, 1990. p. 87-126.
- Smith S.M., J.L. Nillen, A. LeBlanc, A. Lipton, L.M. Demers, H.W. Lane, and C.S. Leach. Collagen cross-link excretion during space flight and bed rest. J Clin Endocrinol Metab 83:3584-3591, 1998.
- Smith, S.M., M.E. Wastney, B.V. Morukov, I.M. Larina, L.E. Nyquist, S.A. Abrams, E.N. Taran, C.-Y. Shih, J.L. Nillen, J.E. Davis-Street, B.L. Rice, and H.W. Lane. Calcium metabolism before, during, and after a 3-mo spaceflight: kinetic and biochemical changes. Am J Physiol Regulatory Integrative Comp Physiol 277:R1-R10, 1999.
- Sonnenfeld, G. Immune responses in space flight. Int J Sports Med 19 (Suppl 3):S195-S204, 1998.
- Stad, N.J. Exercise-Induced +Gz Acceleration versus Exercise and Acceleration: Effect on Aerobic Work Metabolism. San Francisco State University; M.A. thesis in Biology: Physiology and Behavioral Biology, 1998. 56p.



- Stone, R.W., Jr., W. Letko, and W.R. Hook. Examination of a possible flight experiment to evaluate an onboard centrifuge as a therapeutic device. In: The Role of the Vestibular Organs in Space Exploration, edited by J. Huertas, and A. Graybiel. Washington, DC: NASA Special Publ 115, 1966. p. 245-256.
- Utiger, R.D., M.L. Parker, and W.H. Daughaday. Studies on human growth hormone. I. A radio-immunoassay for human growth hormone. J Clin Invest 41:254-261, 1962.
- Vener, J.M. Cardiopulmonary Responses to Incremental Supine Cycle Ergometry with Concomitant +Gz Acceleration. California State University, Fresno; M.A. thesis in Kinesiology, 2000. 92p.
- Vernikos, J. Artificial gravity intermittent centrifugation as a space flight countermeasure. J Gravit Physiol 4:P13-P16, 1997.
- Vil-Viliams, I.F. Principle approaches to selection of the short-arm centrifuge regimens for extended space flight. Acta Astronautica 33:221-229, 1994.
- Vil-Vilyams, I.F., and Ye.B. Shulzhenko. Functional state of the cardiovascular system under the combined effect of 28-day immersion, rotation on a short-arm centrifuge and exercise on a bicycle ergometer. Kosm Biol Aviakosm Med 14:42-45, 1980.
- White, P.D., J.W. Nyberg, L.M. Finney, and W.J. White. Influence of periodic centrifugation on cardiovascular functions of man during bed rest. Santa Monica, CA: Douglas Aircraft Co., Inc., Rept No DAC-59286, 1966. 47p.
- Zerath, E. Effects of microgravity on bone and calcium homeostasis. Adv Space Res 21:1049-1058, 1998.



Figure 12. The Ames HPC-99 Research Team. (left to right) John Greenleaf, Jon Griffith, Jamie Vener, Heather Wilson, James Klem, Kendra Bailey-Pemberton, Jodie Stocks, Stephenie Cowell, Simon Evetts, Paul Barnes, Shawn Simonson, Sunitra Shastry.

### **Present Addresses:**

S.A. Cowell, BS: California Polytechnic State University, Biological Sciences Department, Bldg. 33, 1 Grand Avenue, San Luis Obispo, CA 93407.

S.N. Evetts, PhD: Captain, 23 Pioneer Regiment, St. David's Barracks, Bicester, Oxon, United Kingdom OX26 6HF.

F.B. Moore, PhD: Lockheed Martin Space Operations Corporation, NASA, Ames Research Center, Moffett Field, CA 94035-1000.

K.N. Pemberton, MA: Miltenyi Biotec, 12740 Earhart Drive, Auburn, CA 95602.

S.R. Simonson, EdD: Department of Wellness and Movement Sciences, Box 680, Western New Mexico University, Silver City, NM 88062.

J.M. Stocks, PhD: Eli Lilly Australia Pty Ltd., Asia Pacific Regional Data Management Center, Level 1, 16 Giffnock Avenue, Macquarie Park, NSW, Australia 2113.

J.M. Vener, MA: Department of Exercise and Sports Science, Room 200, University of Utah, 250 South 1850 East, Salt Lake City, UT 84112-0920.

H.W. Wilson, BS: Lockheed Martin Space Operations Corporation, NASA, Ames Research Center, Moffett Field, CA 94035-1000.

Supported by NASA: Task UPN 111-10-20; KSGC: WKU 522775-00-13; NASA EPSCoR: WKU 522635-00-04; NIH: UKGCRC MOIRR02602; GRRC MOI-RR00827

APPENDIX A1



San Francisco State University  
1600 Holloway Avenue  
San Francisco, California 94132

Office of Research and  
Sponsored Programs  
415/338-2231

TO:

JOHN E. GREENLEAF, PH.D.  
NASA AMES RESEARCH CENTER  
MOFFETT FIELD, CA 94035-1000

DATE:

MAY 08, 1998

FROM:

Associate Vice President for Research and Sponsored Programs

SUBJECT:

Protection of Human or Animal Subject Review of Research:  
EXERCISE TRAINING ON SHORT-ARM CENTRIFUGE, H.R. NO. 158

The Committee for the Protection of Human Subjects (CPHS) or the University Animal Care and Use Committee (UACUC) have requested that you be advised of the current status of your planned research in terms of the protection of the rights of human or animal subjects. Please direct your attention only to the pertinent section(s) marked below. If you have any questions, please contact the Office of Research and Sponsored Programs. (338-2231.)

I.  PENDING STATUS:

The CPHS/UACUC is reviewing your research project with respect to the protections planned for the rights of the human or animal subjects involved. To proceed with the review, it needs, as soon as possible, the material(s) checked below:

PROTECTION OF HUMAN SUBJECTS — Materials Needed:

- 1.  Completion of the attached Protocol Approved Form and all necessary signatures.
- 2.  An abstract of 150-200 words that describes your planned research with specific emphasis on the provisions for the protection of the rights of the human subjects involved. Be as specific as possible.
- 3.  A copy of the informed Consent Form you intend to use, as appropriate, in this research effort.
- 4.  Other: \_\_\_\_\_

PROTECTION OF ANIMAL SUBJECTS — Materials Needed:

- 1.  Completion of the attached Protocol Approval Form and all necessary signatures.
- 2.  An abstract of 150-200 words that describes your planned research protocol with specific emphasis for the humane use of animals. Be as specific as possible.
- 3.  Other: \_\_\_\_\_

II.  RECORDED OR APPROVED:

A. Your Protocol Approval Form and supporting materials appear to be in order and have been recorded in the proceedings of the CPHS as noted below:

- 1.  Exempt. Research may proceed.

B. The CPHS/UACUC has reviewed your research protocol and has taken the following action:

- 1.  Approved, expedited review. Research may proceed.
- 2.  Approved, reasonable risk. Research may proceed.
- 3.  Deferred, see attached memo.
- 4.  Not approved, see attached memo.

## APPENDIX A2

National Aeronautics and  
Space Administration  
**Ames Research Center**  
Moffett Field, CA 94035-1000



Reply to Attn of: QH: 243-2

January 4, 1999

**TO:** John Greenleaf, Principal Investigator  
**FROM:** Chair, Human Research Institutional Review Board  
**SUBJECT: Certification of Approval - HR I Protocol**

Your protocol, HR # 191, "Exercise Training on the Short-arm Centrifuge", has been reviewed and approved by the Human Research Institutional Review Board. You are authorized to begin your research, subject to requirements as outlined in AMI 7170.1 (Human Research Planning and Approval). Your approval date is December 7, 1998. Approval expires on December 6, 1999.

The following conditions must be satisfied for your certification to remain in effect:

- 1) Modifications/changes in the project must be received and approved by the Human Research Institutional Review Board and/or by the Chairman, HRIRB before they are initiated, except when necessary, to eliminate immediate or apparent hazard to the subject.
- 2) The office of the Chairman, HRIRB should be notified immediately of any injuries to human subjects and/or any unanticipated problems that involve risks to human subjects or others.
- 3) Copies of the approved consent form(s) must be used for all investigational studies involving human subjects. In addition, all subjects must be given a copy of the consent form(s) to keep for their own records.
- 4) The Department of Health and Human Services and the FDA requires that the HRIRB conduct continuing review of ongoing research at intervals appropriate to the degree of risk, but not less than once per year. To meet this requirement, you must either submit a Protocol Status Report (Attachment A) by June 7, 1999 or request to address the Board during the regularly scheduled meeting preceeding this date .
- 5) In addition, you are also required to complete a Protocol Renewal Request Form (see Attachment B) 6 weeks prior to your protocol expiration date if you wish to conduct research beyond the certification of approval expiration date.

APPENDIX A3

National Aeronautics and  
Space Administration  
**Ames Research Center**  
Moffett Field, CA 94035-1000



Reply to Attn of:

QH:243-2

February 16, 1999

TO: John Greenleaf, Principal Investigator

FROM: Ralph Pelligra, Chair, Human Research Institutional  
Review Board (HRIRB)

SUBJECT: Modification to HR I #191 entitled, "Human-Powered  
Centrifuge:Hormone Studies", Co-Investigator, Richard E.  
Grindeland

The proposed Subject modification was reviewed and approved by  
the HRIRB on Monday, February 8, 1999.

A handwritten signature in black ink, appearing to be "R. Pelligra", written in a cursive style.

Ralph Pelligra, M.D.

cc: R. E. Grindeland

## APPENDIX A4

National Aeronautics and  
Space Administration  
**Ames Research Center**  
Moffett Field, CA 94035-1000



Reply to Attn of: QH:243-2

November 8, 1999

To: Dr. John Greenleaf, Principal Investigator

From: Ralph Pelligra, Chairman, Human Research Institutional  
Review Board (HRIRB)

Subject: HRI-191, "Exercise Training on the Short-Arm  
Centrifuge"

The HRIRB met on 11/1/99 and approved the following:


- 1) A one-month extension of the Subject protocol
- 2) The addition of two minutes of 1Gz passive warm-up; two minutes passive at 50% max; and two minutes of 1Gz passive exposure prior to HPC exercise and acceleration runs on Nov 2 and Nov 17.
- 3) Your responses, dated 10/14/99, to the HRIRB requests of 10/4/99.

Please contact me if you have further questions. Thank you.

A handwritten signature in black ink, appearing to be "R. Pelligra", written in a cursive style.


Ralph Pelligra, M.D.

# APPENDIX B1





APPENDIX B2



APPENDIX C1

42 Instruments and Equipment Inventory — OXYGEN UPTAKE

| equipment                        | model name/no.                         | manufacturer                                                     | manufacturer location | used for                                                                |
|----------------------------------|----------------------------------------|------------------------------------------------------------------|-----------------------|-------------------------------------------------------------------------|
| Oxygen analyzer                  | CPX Express                            | Medical Graphics Corporation                                     | St. Paul, MN          | VO <sub>2</sub> , HR                                                    |
| Laptop computer                  | H-1330                                 | Quantex Microsystems, Inc.                                       | Somerset, New Jersey  | Download and store VO <sub>2</sub> data                                 |
| ECG                              | 78202 (A, B, C)                        | Hewlett-Packard                                                  | Palo Alto, CA         | ECG, HR during exercise                                                 |
| Directional Ultrasonic Flowmeter | Model 1012                             | LM Electronics Inc.                                              |                       | Doppler                                                                 |
| Blood Pressure Monitor           | Pilot                                  | Colin Medical Instruments                                        | San Antonio, TX       | BP during exercise + acceleration                                       |
| Headset                          | H3391                                  | David Clark Company, Inc.                                        | Worcester, MA         | Audio contact w/subjects                                                |
| Wireless backpack                | WTR-2                                  | Clear-Com                                                        | Emeryville, CA        | Audio contact w/subjects                                                |
| Stopwatches                      |                                        | TAG Heuer                                                        | Springfield, NJ       | Timing testing + training runs                                          |
| Medical monitoring computer      | Micron Client Pro                      | Micron PC, Inc.                                                  | Meridian, ID          | Record ECG, Gz, BP                                                      |
| Biobench computer program        | BioBench version 1.0 for Windows 95/NT | National Instruments Corporation                                 | Austin, Texas         | Record ECG, Gz, BP                                                      |
| Connector Block                  | SCB-100                                | National Instruments Corporation                                 | Austin, Texas         | Collect and channel data off of HPC platform to acquisition computer    |
| Data acquisition card            | PCI-6031E Multifunction I/O board      | National Instruments Corporation                                 | Austin, Texas         | Data acquisition card in HPC computer; collect data off of HPC platform |
| Scale                            | Model 61-1320                          | Fairbanks Morse Weighing Systems Division, Colt Industries, Inc. |                       | Subject height                                                          |
| Digital Weight Display           | Model 5780                             | NCI                                                              | San Carlos, CA        | Subject weight                                                          |
| Uniworl Ergometer                | Model 845                              | Quinton                                                          | Seattle, Washington   | Supine cycle testing + training                                         |
| Human Powered Centrifuge         | Human Powered Centrifuge               | NASA Ames Research Center                                        | Moffett Field, CA     | Acceleration and exercise testing and training                          |
| Digital Tuner Metronome          | DTM-12                                 | Korg                                                             | Tokyo, Japan          | Provide pedalling cadence                                               |
| Printer                          | BJC                                    | Canon Hi-Tech                                                    | Thailand              | O <sub>2</sub> Data                                                     |
| Video Camera                     |                                        |                                                                  |                       | Monitor subjects                                                        |

APPENDIX C2

Instruments and Equipment Inventory — TILT TEST

| equipment                                      | model name/no.                                  | manufacturer                                   | manufacturer location    | used for                              |
|------------------------------------------------|-------------------------------------------------|------------------------------------------------|--------------------------|---------------------------------------|
| Tilt table                                     | Physical Therapy Treatment Table serial no. 499 | Laberne Mfg. Co. Physical Therapy Equipment    | Columbia, South Carolina | Tilting subjects                      |
| Cardiodynamic Monitor                          | Cardiodynamic Monitor (CDM)                     | BoMed Medical Mfg. Ltd. (Cardiodynamics, Inc.) | Irvine, CA               | CO, SV, HR, end diastolic volume, ECG |
| 2 Channel Skin Perfusion Meter (Laser Doppler) | PeriFlux Model PF4001                           | Perimed                                        | Smithtown, NY            | Skin perfusion of forearm and palm    |
| 1 Channel Skin Perfusion Meter (Laser Doppler) | LaserFlo Skin Perfusion Monitor, BPM 403A       | TSI                                            | St. Paul, MN             | Skin Perfusion on head                |
| Finapres                                       | Ohmeda 2300 Finapres BP Monitor                 | Ohmeda, a division of BOC Health Care          | Englewood, CO            | BP                                    |
| Dual Frequency Directional Doppler             | Model 909                                       | Parks Medical Electronics, Inc.                | Aloha, OR                | Doppler for radial artery             |
| Plethysmograph                                 | EC-4                                            | D.E. Hokanson, Inc.                            | Issaquah, WA             | Calf circumference                    |
| Chart recorder                                 | MT 8800                                         | Astro-Med, Inc.                                | West Warwick, RI         | Tilt data acquisition                 |
| Analog Tape Recorder                           | TEAC Model (add 11/17/99)                       |                                                |                          |                                       |
| 16 channel data acquisition system             | Model DI-220                                    | Dataq Instruments, Inc.                        | Akron, OH                | Tilt data acquisition                 |
| Data acquisition computer                      | Pentium II                                      | Comteq Computer Company                        | Rockville, MD            | Tilt data acquisition                 |
| 16 channel data acquisition card               | PCI-6023E Multifunction I/O board               | National Instruments Corporation               | Austin, Texas            | Tilt data acquisition                 |
| 16 channel data acquisition BNC box            | BNC 2090 and SH68-68-EP                         | National Instruments Corporation               | Austin, Texas            | Tilt data acquisition                 |
| Data acquisition software                      | Brown DD EX1, Biomedical Engineering            | University of Kentucky                         | Lexington, KY            | Tilt data acquisition                 |
| Sphygmomanometer                               | Baumanometer, Standby model                     | W.A. Baum Co. Inc.                             | Copliague, New York      | Supine resting BP (pre-tilt)          |
| Accelerometer                                  | ADXLD EM-3                                      | Analog Service                                 | Roewood, MS              | Tilt angle +/- 5%                     |
| Tape recorder                                  | SR 510                                          | TEAC                                           | Japan                    | Analog record of tilt test            |

APPENDIX C3

Instruments and Equipment Inventory — BLOOD HANDLING

| equipment                             | model name/no.                        | manufacturer                      | manufacturer location  | used for                               |
|---------------------------------------|---------------------------------------|-----------------------------------|------------------------|----------------------------------------|
| Catheters                             | Quick-Cath                            | Baxter Healthcare Corporation     | Deerfield, IL          | Blood withdrawal, EB injection         |
| Injection sites                       | Injection sites                       | Baxter Healthcare Corporation     | Deerfield, IL          | Blood withdrawal, EB injection         |
| 30 + 5 ml syringes                    | LuerLok                               | Becton Dickinson + Co.            | Franklin Lakes, NJ     | Blood withdrawal, EB injection         |
| EB dye                                | 25 mg in 5 ml                         | New World Trading Corp.           | Florida                | Plasma volume determination            |
| Balance                               | B6 (43289)                            | Mettler Instrument Corp.          | Highstown, NJ          | Weight of Evans Blue syringes          |
| Heparin Lock Flush Solution           | Hep-Lock                              | Elkins-Sinn, Inc.                 | Cherry Hill, NJ        | Hep saline                             |
| NaCl Injection                        | 0.9% NaCl Injection, 500ml            | McGaw, Inc.                       | Irvine, CA             | Saline                                 |
| Heparinized Microcapillary tubes      | Red Tip                               | Sherwood Medical Industries       | St. Louis, MO          | Collecting Hct samples                 |
| Sealing cork                          | Seal-ease                             | Clay Adams                        | Parsippany, NJ         | Sealing Hct tubes                      |
| Skin Perfusion                        | Periflux-2 channel                    | Perines                           | Smithtown, NY          | Skin Perfusion (forearm, palm)         |
| Microcentrifuge                       | Micro-MB centrifuge                   | International Equipment Company   | Needham Heights, Mass. | Hct tube centrifugation                |
| Micro-Capillary reader                | Model CR                              | International Equipment Company   | Needham Heights, Mass. | Hct tube reading                       |
| Blood collection tubes (EDTA + LiHep) | Vacutainer                            | Becton Dickinson and Company      | Franklin Lakes, NJ     | Initial blood collection and treatment |
| Refrigerated Centrifuge               | RC2-B                                 | Ivan Sorvall Inc.                 | Newtown, Connecticut   | Blood tube centrifugation              |
| Pipette                               | Pipetman                              | Gilson, Inc.                      | Middleton, Wisconsin   | Blood separation                       |
| Advanced Digimatic Osmometer          | Model 3DII                            | Advanced Instruments, Inc.        | Needham Heights, Mass. | Plasma osmolality                      |
| Spectrophotometer                     | Model 35                              | Beckman                           | Irvine, CA             | Hb concentration/PV                    |
| Hemoglobin Kit                        | Cyanmethoemoglobin method, Kit # 525A | Sigma Diagnostics                 | St. Louis, MO          | Hb concentration                       |
| Evans Blue                            | T-1824                                | The New World Trading Corp.       | De Bary, FL            | Plasma volume determination            |
| Total protein analysis                | LX2D                                  | Beckman Coulter Synchron Analyzer | Brea, CA               |                                        |
| Na analysis                           | LX2D                                  | Beckman Coulter Synchron Analyzer | Brea, CA               |                                        |
| K analysis                            | LX2D                                  | Beckman Coulter Synchron Analyzer | Brea, CA               |                                        |
| Albumin analysis                      | LX2D                                  | Beckman Coulter Synchron Analyzer | Brea, CA               |                                        |
| AVP analysis                          | See text                              |                                   |                        |                                        |
| Renin analysis                        | See text                              |                                   |                        |                                        |
| Catecholamines                        | See text                              |                                   |                        |                                        |

APPENDIX C4

Instruments and Equipment Inventory — MISCELLANEOUS


| equipment                         | model name/no.                       | manufacturer            | manufacturer location | used for                  |
|-----------------------------------|--------------------------------------|-------------------------|-----------------------|---------------------------|
| Data analysis                     | Pentium II                           | Comteq Computer Company | Rockville, MD         | Data analysis             |
| Data analysis                     | SPSS 7.5 for Windows                 | SPSS, Inc.              | Chicago, IL           | Data analysis             |
| Data analysis                     | Microsoft Excel 97                   | Microsoft Corporation   | Redmond, WA           | Data analysis             |
| Data analysis                     | Brown DD EX1, Biomedical Engineering | University of Kentucky  | Lexington, KY         | Data analysis             |
| Refractometer                     | Protometer                           | National Instr.         |                       | Total protein and density |
| MRI                               | Vision (-5 Tesla)                    | Seimans                 | Iselin, NJ            | Leg muscles               |
| Data acquisition and analysis     |                                      | David Brown             | Lexington, KY         | Data analysis             |
| Overhead camera                   | BL 600                               | Panasonic               |                       |                           |
| (2) Lip stick camera              |                                      | Panasonic               |                       |                           |
| (2) Power supply                  | GP-KS102                             | Panasonic               |                       |                           |
| (2) 120V AC to 12V DC transformer | CAT No. 273-1653A                    | Archer                  |                       |                           |

APPENDIX D

**INTENSITY OF PERCEIVED EXERTION**

Name \_\_\_\_\_ Date \_\_\_\_\_

Test \_\_\_\_\_



|      |                          |                          |
|------|--------------------------|--------------------------|
| 10++ | <b>Supra-maximal</b>     |                          |
| 10+  | <b>Maximal</b>           |                          |
| 10   | <b>Very, very strong</b> | <b>(almost max)</b>      |
| 9    |                          |                          |
| 8    | <b>Very strong</b>       |                          |
| 7    |                          |                          |
| 6    |                          |                          |
| 5    | <b>Strong</b>            | <b>(heavy)</b>           |
| 4    | <b>Somewhat strong</b>   |                          |
| 3    | <b>Moderate</b>          |                          |
| 2    | <b>Weak</b>              | <b>(light)</b>           |
| 1    | <b>Very weak</b>         |                          |
| 0.5  | <b>Very, very weak</b>   | <b>(just noticeable)</b> |
| 0    | <b>Nothing at all</b>    |                          |

# APPENDIX E

## VacuMed



Enter data into blue fields only

### Precision O2 Calibration & Metabolic Rate Calculation

| Line                                                 | Data Description                                                                   | LOW RANGE | MEDIUM RANGE | HIGH RANGE |
|------------------------------------------------------|------------------------------------------------------------------------------------|-----------|--------------|------------|
| 1                                                    | Enter Room Temperature (C):<br>(10 - 39.2 Degree in 0.2 degree intervals)          | 22.0      |              |            |
| 2                                                    | Enter Baro Pressure (mmHg):                                                        | 770       |              |            |
| 3                                                    | Enter Ambient Humidity (%):                                                        | 43        |              |            |
| 4                                                    | Vapor Pressure from Table*:                                                        | 19.83     |              |            |
| 5                                                    | Read True O2 Concentration:                                                        | 20.71     |              |            |
| 6                                                    | Enter CO2% of Metabolic CAL GAS:                                                   | 20.99     |              |            |
| 7                                                    | Gas Flow Liters per Minute                                                         | 2.38      | 6.97         | 13.56      |
| 8                                                    | Metabolic Flow (ATPs*) in ml                                                       | 500       | 1463         | 2846       |
| 9                                                    | Metabolic Flow (STPD) in ml                                                        | 464       | 1360         | 2645       |
| <b>Correct CPX VO2 (computed) to VO2 (simulated)</b> |                                                                                    |           |              |            |
| 10                                                   | Enter system's exhaled Temp Assumption (C):<br>(10 - 39.2 in 0.2 degree intervals) | 34.0      |              |            |
| 11                                                   | Vapor Pressure from Table*:                                                        | 39.90     |              |            |
| 12                                                   | Enter system's VO2 (computed stpd):                                                | 394       | 1230         | 2380       |
| 13                                                   | Enter system's VCO2 (computed stpd):                                               | 399       | 1245         | 2550       |
| 14                                                   | Corrected VO2 (simulated STPD):                                                    | 431       | 1344         | 2601       |
|                                                      | ERROR VO2                                                                          | -7.81%    | -1.13%       | -1.68%     |
| 15                                                   | Corrected VCO2 (simulated STPD):                                                   | 436       | 1361         | 2787       |
|                                                      | ERROR VCO2                                                                         | -6.46%    | 0.09%        | 5.10%      |

# ATPs = mixture of dry cal gas and room air is somewhat humid  
\* Corrected for % humidity

**CPX Make & Model:**

Copyright 1998 Vacumetrics Inc  
R/O2CAL.xls

10 Oct 1990

**Serial No:**

**Date:**

**John Hoppe**  
President

4483 McGrath St. # 102 Ventura CA 93003  
Tel (805) 644-7461 • (800) 235-3333  
Fax (805) 654-8759 • mobile (805) 320-0654  
E-mail: info@vacumed.com  
Internet: www.vacumed.com

**Vacumetrics Inc. / VacuMed Division**

Serving customers around the world  
in the exercise physiology and cardiopulmonary  
sciences since 1968

## APPENDIX F

### Supine Heart Rate Max (beats • min<sup>-1</sup>)

| Subject | PRE Training |          |          | POST Training |          |          | POST study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 162      |               |          | 171      | 166        |
| FRE     | 169          | 170      | 169      | 160           | 169      | 165      | 181        |
| HUN     | 173          | 170      | 178      | 172           | 182      | 174      | 183        |
| JAG     | 186          | 182      | 171      | 170           | 173      | 173      | 170        |
| RAY     | 188          | 177      | 164      | 180           | 177      | 178      | 179        |
| RUI     | 159          | 166      | 158      | 160           | 159      | 156      | 166        |
| SCH     | 171          | 184      |          | 176           | 185      |          |            |
| Mean    | 174          | 175      | 167      | 170           | 174      | 170      | 174        |
| SD      | 11           | 7        | 7        | 8             | 9        | 8        | 8          |
| SE      | 4            | 3        | 3        | 3             | 4        | 3        | 3          |

### Supine Heart Rate Max (% change)

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 6        |
| FRE     | -5                            | -1       | -2       |
| HUN     | -1                            | 7        | -2       |
| JAG     | -9                            | -5       | 1        |
| RAY     | -4                            | 0        | 9        |
| RUI     | 1                             | -4       | -1       |
| SCH     | 3                             | 1        |          |
| Mean    | -3                            | 0        | 2        |
| SD      | 4                             | 4        | 5        |
| SE      | 2                             | 2        | 2        |

### Supine VO<sub>2</sub> Max (ml • min<sup>-1</sup> • kg<sup>-1</sup>)

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 37.0     |               |          | 38.0     | 38.5       |
| FRE     | 35.2         | 38.3     | 39.1     | 34.4          | 37.2     | 40.5     | 35.2       |
| HUN     | 28.1         | 27.9     | 30.9     | 31.2          | 28.1     | 31.8     | 31.3       |
| JAG     | 37.9         | 38.2     | 37.6     | 36.3          | 33.1     | 36.9     | 32.9       |
| RAY     | 39.6         | 31.6     | 34.8     | 37.1          | 36.8     | 34.7     | shut down  |
| RUI     | 36.7         | 33.5     | 40.3     | 38.3          | 36.1     | 38.7     | 39.1       |
| SCH     | 37.0         | 39.8     |          | 33.9          | 39.3     |          |            |
| Mean    | 35.8         | 34.9     | 36.6     | 35.2          | 35.1     | 36.8     | 35.4       |
| SD      | 4.0          | 4.7      | 3.4      | 2.6           | 4.0      | 3.1      | 3.4        |
| SE      | 1.6          | 1.9      | 1.4      | 1.0           | 1.6      | 1.3      | 1.4        |

### Supine VO<sub>2</sub> Max (% change)

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 2.7      |
| FRE     | -2.3                          | -2.9     | 3.6      |
| HUN     | 11.0                          | 0.7      | 2.9      |
| JAG     | -4.2                          | -13.4    | -1.9     |
| RAY     | -6.3                          | 16.5     | -0.3     |
| RUI     | 4.4                           | 7.8      | -4.0     |
| SCH     | -8.4                          | -1.3     |          |
| Mean    | -1.0                          | 1.2      | 0.5      |
| SD      | 7.3                           | 10.1     | 3.0      |
| SE      | 3.0                           | 4.1      | 1.2      |



**Supine Heart Rate Max (beats • min<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 162      |               |          | 171      | 166        |
| FRE     | 169          | 170      | 169      | 160           | 169      | 165      | 181        |
| HUN     | 173          | 170      | 178      | 172           | 182      | 174      | 183        |
| JAG     | 186          | 182      | 171      | 170           | 173      | 173      | 170        |
| RAY     | 188          | 177      | 164      | 180           | 177      | 178      | 179        |
| RUI     | 159          | 166      | 158      | 160           | 159      | 156      | 166        |
| SCH     | 171          | 184      |          | 176           | 185      |          |            |
| Mean    | 174          | 175      | 167      | 170           | 174      | 170      | 174        |
| SD      | 11           | 7        | 7        | 8             | 9        | 8        | 8          |
| SE      | 4            | 3        | 3        | 3             | 4        | 3        | 3          |

**Supine Heart Rate Max (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 6        |
| FRE     | -5                            | -1       | -2       |
| HUN     | -1                            | 7        | -2       |
| JAG     | -9                            | -5       | 1        |
| RAY     | -4                            | 0        | 9        |
| RUI     | 1                             | -4       | -1       |
| SCH     | 3                             | 1        |          |
| Mean    | -3                            | 0        | 2        |
| SD      | 4                             | 4        | 5        |
| SE      | 2                             | 2        | 2        |

**Supine Time to Fatigue (min)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 725      |               |          | 852      | 810        |
| FRE     | 780          | 815      | 780      | 750           | 900      | 865      | 850        |
| HUN     | 840          | 750      | 715      | 745           | 900      | 830      | 840        |
| JAG     | 760          | 807      | 660      | 810           | 700      | 840      | 722        |
| RAY     | 725          | 710      | 640      | 790           | 710      | 812      | 800        |
| RUI     | 709          | 700      | 730      | 750           | 725      | 795      | 807        |
| SCH     | 780          | 780      |          | 855           | 900      |          |            |
| Mean    | 766          | 760      | 708      | 783           | 806      | 832      | 805        |
| SD      | 47           | 49       | 51       | 44            | 103      | 26       | 45         |
| SE      | 19           | 20       | 21       | 18            | 42       | 11       | 18         |

**Supine Time to Fatigue (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 17.5     |
| FRE     | -3.8                          | 10.4     | 10.9     |
| HUN     | -11.3                         | 20.0     | 16.1     |
| JAG     | 6.6                           | -13.3    | 27.3     |
| RAY     | 9.0                           | 0.0      | 26.9     |
| RUI     | 5.8                           | 3.6      | 8.9      |
| SCH     | 9.6                           | 15.4     |          |
| Mean    | 2.6                           | 6.0      | 17.9     |
| SD      | 8.4                           | 12.0     | 7.8      |
| SE      | 3.4                           | 4.9      | 3.2      |

**Supine Resting Heart Rate (average of tilt -5min to -2min) (beats • min<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 52.7     |               |          | 53.9     |            |
| FRE     | 76.0         | 74.9     | 75.3     | 86.4          | 75.3     | 75.3     |            |
| HUN     | 58.3         | 67.2     | 57.2     | 65.6          | 77.2     | 67.5     |            |
| JAG     | 71.0         | 65.9     | 68.8     | 76.4          | 67.8     | 80.6     |            |
| RAY     | 50.6         | 46.3     | 55.4     | 49.0          | 45.2     | 47.5     |            |
| RUI     | 60.9         | 64.8     | 60.7     | 67.6          | 58.6     | 59.3     |            |
| SCH     | 58.2         | 68.5     |          | 79.1          | 55.4     |          |            |
| Mean    | 62.5         | 64.6     | 61.7     | 70.7          | 63.3     | 64.0     |            |
| SD      | 9.3          | 9.6      | 8.7      | 13.1          | 12.4     | 12.7     |            |
| SE      | 3.8          | 3.9      | 3.5      | 5.3           | 5.1      | 5.2      |            |

**Supine Resting Heart Rate (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 2.3      |
| FRE     | 13.7                          | 0.5      | 0.0      |
| HUN     | 12.5                          | 14.9     | 18.0     |
| JAG     | 7.6                           | 2.9      | 17.2     |
| RAY     | -3.2                          | -2.4     | -14.3    |
| RUI     | 11.0                          | -9.6     | -2.3     |
| SCH     | 35.9                          | -19.1    |          |
| Mean    | 12.9                          | -2.1     | 3.5      |
| SD      | 12.8                          | 11.5     | 12.3     |
| SE      | 5.2                           | 4.7      | 5.0      |

**Resting Systolic BP (mmHg)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 115.8    |               |          | 121.1    |            |
| FRE     | 131.6        | 130.2    | 137.5    | 154.0         | 129.7    | 129.7    |            |
| HUN     | 150.0        | 149.3    | 119.2    | 162.8         | 136.0    | 140.0    |            |
| JAG     | 149.8        | 154.7    | 155.6    | 125.1         | 145.8    | 140.1    |            |
| RAY     | 141.9        | 152.9    | 147.1    | 152.9         | 149.3    | 120.7    |            |
| RUI     | 162.2        | 146.1    | 151.9    | 163.0         | 144.5    | 153.0    |            |
| SCH     | 149.8        | 157.0    |          | 144.9         | 134.8    |          |            |
| Mean    | 147.6        | 148.4    | 137.9    | 150.5         | 140.0    | 134.1    |            |
| SD      | 10.2         | 9.7      | 16.9     | 14.2          | 7.6      | 12.6     |            |
| SE      | 4.2          | 4.0      | 6.9      | 5.8           | 3.1      | 5.2      |            |

**Resting Systolic BP (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 4.6      |
| FRE     | 17.0                          | -0.4     | -5.7     |
| HUN     | 8.5                           | -8.9     | 17.4     |
| JAG     | -16.5                         | -5.8     | -10.0    |
| RAY     | 7.8                           | -2.4     | -17.9    |
| RUI     | 0.5                           | -1.1     | 0.7      |
| SCH     | -3.3                          | -14.1    |          |
| Mean    | 2.3                           | -5.4     | -1.8     |
| SD      | 11.6                          | 5.3      | 12.3     |
| SE      | 4.7                           | 2.2      | 5.0      |

**Resting Diastolic BP (mmHg)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 72.5     |               |          | 71.8     |            |
| FRE     | 76.1         | 90.6     | 81.7     | 90.1          | 70.3     | 78.2     |            |
| HUN     | 88.7         | 76.5     | 65.9     | 102.2         | 82.0     | 79.5     |            |
| JAG     | 82.0         | 84.5     | 93.3     | 73.8          | 81.9     | 70.4     |            |
| RAY     | 95.6         | 85.2     | 78.8     | 78.6          | 82.5     | 78.3     |            |
| RUI     | 90.0         | 76.9     | 88.5     | 99.9          | 78.9     | 83.3     |            |
| SCH     | 81.8         | 88.3     |          | 70.9          | 80.0     |          |            |
| Mean    | 85.7         | 83.7     | 80.1     | 85.9          | 79.3     | 76.9     |            |
| SD      | 7.0          | 5.8      | 10.1     | 13.4          | 4.6      | 4.9      |            |
| SE      | 2.9          | 2.4      | 4.1      | 5.5           | 1.9      | 2.0      |            |

**Resting Diastolic BP (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -1.0     |
| FRE     | 18.4                          | -22.4    | -4.3     |
| HUN     | 15.2                          | 7.2      | 20.6     |
| JAG     | -10.0                         | -3.1     | -24.5    |
| RAY     | -17.8                         | -3.2     | -0.6     |
| RUI     | 11.0                          | 2.6      | -5.9     |
| SCH     | -13.3                         | -9.4     |          |
| Mean    | 0.6                           | -4.7     | -2.6     |
| SD      | 16.0                          | 10.3     | 14.4     |
| SE      | 6.5                           | 4.2      | 5.9      |

**Resting MAP (mmHg)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 88.2     |               |          | 89.3     |            |
| FRE     | 91.0         | 104.1    | 98.5     | 106.1         | 87.0     | 94.5     |            |
| HUN     | 105.1        | 95.7     | 81.7     | 120.8         | 97.4     | 97.8     |            |
| JAG     | 101.6        | 102.6    | 111.5    | 90.1          | 98.6     | 88.5     |            |
| RAY     | 107.8        | 104.5    | 96.8     | 98.3          | 100.2    | 91.6     |            |
| RUI     | 110.1        | 94.8     | 107.7    | 119.4         | 97.1     | 102.9    |            |
| SCH     | 100.9        | 106.5    |          | 90.3          | 96.5     |          |            |
| Mean    | 102.8        | 101.4    | 97.4     | 104.2         | 96.1     | 94.1     |            |
| SD      | 6.8          | 4.9      | 11.3     | 13.7          | 4.7      | 5.5      |            |
| SE      | 2.8          | 2.0      | 4.6      | 5.6           | 1.9      | 2.3      |            |

**Resting MAP (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 1.2      |
| FRE     | 16.6                          | -16.4    | -4.1     |
| HUN     | 14.9                          | 1.8      | 19.7     |
| JAG     | -11.3                         | -3.9     | -20.6    |
| RAY     | -8.8                          | -4.1     | -5.4     |
| RUI     | 8.4                           | 2.4      | -4.5     |
| SCH     | -10.5                         | -9.4     |          |
| Mean    | 1.6                           | -4.9     | -2.3     |
| SD      | 13.2                          | 7.1      | 13.0     |
| SE      | 5.4                           | 2.9      | 5.3      |

**Resting Pulse Pressure (mmHg)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 43.3     |               |          | 49.3     |            |
| FRE     | 55.5         | 39.6     | 55.8     | 63.9          | 59.4     | 51.5     |            |
| HUN     | 61.3         | 72.8     | 53.3     | 60.6          | 54.0     | 60.5     |            |
| JAG     | 67.8         | 70.2     | 62.3     | 51.3          | 63.9     | 69.7     |            |
| RAY     | 46.3         | 67.7     | 68.3     | 74.3          | 66.8     | 42.4     |            |
| RUI     | 72.2         | 69.2     | 63.4     | 63.1          | 65.6     | 69.7     |            |
| SCH     | 68.0         | 68.7     |          | 74.0          | 54.8     |          |            |
| Mean    | 61.9         | 64.7     | 57.7     | 64.5          | 60.8     | 57.2     |            |
| SD      | 9.6          | 12.4     | 8.9      | 8.7           | 5.5      | 11.3     |            |
| SE      | 3.9          | 5.1      | 3.6      | 3.5           | 2.3      | 4.6      |            |

**Resting Pulse Pressure (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 13.9     |
| FRE     | 15.1                          | 50.0     | -7.7     |
| HUN     | -1.1                          | -25.8    | 13.5     |
| JAG     | -24.3                         | -9.0     | 11.9     |
| RAY     | 60.5                          | -1.3     | -37.9    |
| RUI     | -12.6                         | -5.2     | 9.9      |
| SCH     | 8.8                           | -20.2    |          |
| MEAN    | 7.7                           | -1.9     | 0.6      |
| SD      | 29.5                          | 27.1     | 20.5     |
| SE      | 12.1                          | 11.1     | 8.4      |

**Tilt Table Tolerance Time (min)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 60.00    |               |          | 60.00    | 60.00      |
| FRE     | 60.00        | 60.00    | 60.00    | 60.00         | 60.00    | 60.00    | 60.00      |
| HUN     | 60.00        | 60.00    | 59.00    | 60.00         | 60.00    | 35.67    | 58.00      |
| JAG     | 22.20        | 45.42    | 25.75    | 28.15         | 26.68    | 17.92    | 28.00      |
| RAY     | 31.32        | 45.97    | 15.58    | 60.00         | 60.00    | 41.25    | 60.00      |
| RUI     | 29.30        | 60.00    | 60.00    | 60.00         | 60.00    | 60.00    | 60.00      |
| SCH     | 58.00        | 49.92    |          | 60.00         | 34.05    |          |            |
| Mean    | 43.47        | 53.55    | 46.72    | 54.69         | 50.12    | 45.81    | 54.33      |
| SD      | 17.65        | 7.23     | 20.44    | 13.00         | 15.48    | 17.35    | 12.93      |
| SE      | 7.21         | 2.95     | 8.35     | 5.31          | 6.32     | 7.08     | 5.28       |

**Tilt Table Tolerance (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 0.00     |
| FRE     | 0.00                          | 0.00     | 0.00     |
| HUN     | 0.00                          | 0.00     | -39.54   |
| JAG     | 26.80                         | -41.26   | -30.41   |
| RAY     | 91.57                         | 30.52    | 164.76   |
| RUI     | 104.78                        | 0.00     | 0.00     |
| SCH     | 3.45                          | -31.79   |          |
| Mean    | 37.77                         | -7.09    | 15.80    |
| SD      | 48.03                         | 25.86    | 75.02    |
| SE      | 19.61                         | 10.56    | 30.62    |

**Heart Rate @ Tilt Table Tolerance Time (beats • min<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 84       |               |          | 78       | 90         |
| FRE     | 90           | 104      | 96       | 96            | 96       | 96       |            |
| HUN     | 90           | 108      | 96       | 96            | 108      | 90       | 120        |
| JAG     | 66           | 78       | 108      | 96            | 120      | 126      | 108        |
| RAY     | 48           | 66       | 54       | 66            | 60       | 72       | 72         |
| RUI     | 48           | 72       | 72       | 72            | 60       | 72       | 72         |
| SCH     | 84           | 102      |          | 108           | 90       |          |            |
| Mean    | 71           | 88       | 85       | 89            | 89       | 89       | 92         |
| SD      | 20           | 18       | 20       | 16            | 25       | 21       | 21         |
| SE      | 8            | 8        | 8        | 7             | 10       | 8        | 9          |

**Heart Rate @ Tilt Table Tolerance Time (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -7.1     |
| FRE     | 6.7                           | -7.7     | 0.0      |
| HUN     | 6.7                           | 0.0      | -6.3     |
| JAG     | 45.5                          | 53.8     | 16.7     |
| RAY     | 37.5                          | -9.1     | 33.3     |
| RUI     | 50.0                          | -16.7    | 0.0      |
| SCH     | 28.6                          | -11.8    |          |
| Mean    | 29.1                          | 1.4      | 6.1      |
| SD      | 18.9                          | 26.2     | 15.8     |
| SE      | 7.7                           | 10.7     | 6.5      |

**MAP @ Tilt Table Tolerance (mmHg)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 90       |               |          | 100      | 95         |
| FRE     | 110          | 115      | 70       | 110           | 100      | 105      | 115        |
| HUN     | 110          | 90       | 60       | 115           | 100      | 70       | 100        |
| JAG     | 40           | 85       | 115      | 60            | 75       | 90       | 80         |
| RAY     | 90           | 110      | 75       | 120           | 115      | 100      | 120        |
| RUI     | 60           | 110      | 60       | 130           | 120      | 110      | 120        |
| SCH     | 115          | 100      |          | 110           | 120      |          |            |
| Mean    | 88           | 102      | 78       | 108           | 105      | 96       | 105        |
| SD      | 31           | 12       | 21       | 24            | 17       | 14       | 16         |
| SE      | 13           | 5        | 9        | 10            | 7        | 6        | 7          |

**MAP @ Tilt Table Tolerance (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | 11.1     |
| FRE     | 0.0                           | -13.0    | 50.0     |
| HUN     | 4.5                           | 11.1     | 16.7     |
| JAG     | 50.0                          | -11.8    | -21.7    |
| RAY     | 33.3                          | 4.5      | 33.3     |
| RUI     | 116.7                         | 9.1      | 83.3     |
| SCH     | -4.3                          | 20.0     |          |
| Mean    | 33.4                          | 3.3      | 28.8     |
| SD      | 46.0                          | 13.2     | 35.9     |
| SE      | 18.8                          | 5.4      | 14.7     |

**R-R Interval (msec)**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt |
| RUI     | 984                  | 841       | 879                   | 792       | 894                   | 855       | 1010                   | 929       | 980                   | 850       | 1020                   | 890       |
| RAY     | 1190                 | 858       | 1220                  | 941       | 1260                  | 932       | 1300                   | 999       | 1100                  | 880       | 1250                   | 940       |
| HUN     | 1020                 | 696       | 912                   | 662       | 917                   | 619       | 772                    | 560       | 1020                  | 700       | 900                    | 680       |
| JAG     | 844                  | 615       | 776                   | 590       | 905                   | 634       | 886                    | 607       | 890                   | 620       | 750                    | 540       |
| SCH     | 1020                 | 738       | 761                   | 599       | 889                   | 671       | 1060                   | 795       |                       |           |                        |           |
| FLE     |                      |           |                       |           |                       |           |                        |           |                       |           |                        |           |
| FRE     | 776                  | 634       | 701                   | 555       | 813                   | 679       | 788                    | 642       | 790                   | 650       | 810                    | 640       |
| Mean    | 972                  | 730       | 875                   | 690       | 946                   | 732       | 969                    | 755       | 987                   | 738       | 975                    | 735       |
| SD      | 146                  | 102       | 186                   | 149       | 158                   | 130       | 199                    | 181       | 131                   | 106       | 191                    | 153       |
| SE      | 60                   | 42        | 76                    | 61        | 64                    | 53        | 81                     | 74        | 53                    | 43        | 78                     | 62        |

**R-R Interval (% change)**

| Subject | % Change Resting to Tilt |          |          |               |          |          | % Change PRE to POST Training, Resting |          |               |          |                       |         |                        |          |
|---------|--------------------------|----------|----------|---------------|----------|----------|----------------------------------------|----------|---------------|----------|-----------------------|---------|------------------------|----------|
|         | PRE Training             |          |          | POST Training |          |          | PRE Training                           |          | POST Training |          | PRE Training, Resting |         | POST Training, Resting |          |
|         | passive                  | exercise | combined | passive       | exercise | combined | passive                                | exercise | passive       | exercise | combined              | passive | exercise               | combined |
| RUI     | -14.53                   | -4.36    | -13.27   | -9.90         | -8.02    | -12.75   | -10.67                                 | 12.98    | -10.67        | 12.98    | 4.08                  | -10.67  | 12.98                  | 4.08     |
| RAY     | -27.90                   | -26.03   | -20.00   | -22.87        | -23.15   | -24.80   | 2.52                                   | 3.17     | 2.52          | 3.17     | 13.64                 | 2.52    | 3.17                   | 13.64    |
| HUN     | -31.76                   | -32.50   | -31.37   | -27.41        | -27.46   | -24.44   | -10.59                                 | -15.81   | -10.59        | -15.81   | -11.76                | -10.59  | -15.81                 | -11.76   |
| JAG     | -27.13                   | -29.94   | -30.34   | -23.97        | -31.49   | -28.00   | -8.06                                  | -2.10    | -8.06         | -2.10    | -15.73                | -8.06   | -2.10                  | -15.73   |
| SCH     | -27.65                   | -24.52   |          | -21.29        | -25.00   |          | -25.39                                 | 19.24    | -25.39        | 19.24    |                       | -25.39  | 19.24                  |          |
| FLE     |                          |          | -35.96   |               |          | -35.71   |                                        |          |               |          | -1.75                 |         |                        | -1.75    |
| FRE     | -18.30                   | -16.48   | -17.72   | -20.83        | -18.53   | -20.99   | -9.66                                  | -3.08    | -9.66         | -3.08    | 2.53                  | -9.66   | -3.08                  | 2.53     |
| Mean    | -24.55                   | -22.31   | -24.78   | -21.04        | -22.28   | -24.45   | -10.31                                 | 2.40     | -10.31        | 2.40     | -1.50                 | -10.31  | 2.40                   | -1.50    |
| SD      | 6.62                     | 10.36    | 9.00     | 5.95          | 8.21     | 7.60     | 8.92                                   | 12.47    | 8.92          | 12.47    | 10.81                 | 8.92    | 12.47                  | 10.81    |
| SE      | 2.70                     | 4.23     | 3.67     | 2.43          | 3.35     | 3.10     | 3.64                                   | 5.09     | 3.64          | 5.09     | 4.41                  | 3.64    | 5.09                   | 4.41     |

**Stroke Volume (ml • beat<sup>-1</sup>)**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 115                  | 71        | 107                   | 71        | 84                    | 59        | 112                    | 80        | 109                   | 75        | 111                    | 73        |            |           |
| RAY     | 119                  | 69        | 131                   | 77        | 107                   | 68        | 119                    | 74        | 120                   | 75        | 129                    | 83        |            |           |
| HUN     | 135                  | 74        | 117                   | 69        | 120                   | 65        | 92                     | 62        | 147                   | 76        | 108                    | 69        |            |           |
| JAG     | 96                   | 51        | 101                   | 56        | 117                   | 60        | 100                    | 51        | 112                   | 58        | 115                    | 57        |            |           |
| SCH     | 96                   | 55        | 77                    | 52        | 71                    | 44        | 106                    | 63        |                       |           |                        |           |            |           |
| FLE     |                      |           |                       |           |                       |           |                        |           |                       |           |                        |           |            |           |
| FRE     | 59                   | 47        | 65                    | 48        | 68                    | 53        | 67                     | 46        | 105                   | 55        | 110                    | 49        |            |           |
| Mean    | 103                  | 61        | 100                   | 62        | 95                    | 58        | 99                     | 63        | 110                   | 65        | 106                    | 63        |            |           |
| SD      | 26                   | 12        | 25                    | 12        | 23                    | 9         | 18                     | 13        | 27                    | 12        | 22                     | 15        |            |           |
| SE      | 11                   | 5         | 10                    | 5         | 9                     | 4         | 8                      | 5         | 11                    | 5         | 9                      | 6         |            |           |

**Stroke Volume (% change)**

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |                                        |          |          |  |  |  |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|----------------------------------------|----------|----------|--|--|--|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          | % Change PRE to POST Training, Resting |          |          |  |  |  |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined | passive                                | exercise | combined |  |  |  |
| RUI     | -38                      | -30      | -31      | -34     | -29      | -34      | -7            | 33       | 2        |         |          |          |                                        |          |          |  |  |  |
| RAY     | -42                      | -36      | -38      | -41     | -38      | -36      | 10            | 11       | 8        |         |          |          |                                        |          |          |  |  |  |
| HUN     | -45                      | -46      | -48      | -41     | -33      | -36      | -13           | -23      | -27      |         |          |          |                                        |          |          |  |  |  |
| JAG     | -47                      | -49      | -48      | -45     | -49      | -50      | 5             | -15      | 3        |         |          |          |                                        |          |          |  |  |  |
| SCH     | -43                      | -38      |          | -32     | -41      |          | -20           | 49       | 5        |         |          |          |                                        |          |          |  |  |  |
| FLE     |                          |          | -48      |         |          | -55      |               |          | 5        |         |          |          |                                        |          |          |  |  |  |
| FRE     | -20                      | -22      | -25      | -26     | -31      | -32      | 10            | -1       | 2        |         |          |          |                                        |          |          |  |  |  |
| Mean    | -39                      | -37      | -40      | -37     | -37      | -41      | -2            | 9        | -1       |         |          |          |                                        |          |          |  |  |  |
| SD      | 10                       | 10       | 10       | 7       | 7        | 10       | 13            | 28       | 13       |         |          |          |                                        |          |          |  |  |  |
| SE      | 4                        | 4        | 4        | 3       | 3        | 4        | 5             | 11       | 5        |         |          |          |                                        |          |          |  |  |  |

**End-Diastolic Volume (ml)**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt |
| RUI     | 183                  | 130       | 175                   | 130       | 133                   | 107       | 184                    | 142       | 180                   | 147       | 180                    | 138       |
| RAY     | 191                  | 133       | 220                   | 162       | 174                   | 130       | 198                    | 154       | 195                   | 146       | 218                    | 169       |
| HUN     | 226                  | 158       | 205                   | 163       | 208                   | 153       | 175                    | 149       | 257                   | 177       | 190                    | 160       |
| JAG     | 170                  | 135       | 174                   | 132       | 203                   | 154       | 180                    | 136       | 194                   | 143       | 190                    | 129       |
| SCH     | 152                  | 109       | 119                   | 106       | 120                   | 103       | 168                    | 126       |                       |           |                        |           |
| FLE     |                      |           |                       |           |                       |           |                        |           | 166                   | 109       | 181                    | 142       |
| FRE     | 112                  | 112       | 128                   | 101       | 122                   | 121       | 119                    | 113       | 120                   | 112       | 127                    | 114       |
| Mean    | 172                  | 130       | 170                   | 132       | 160                   | 128       | 171                    | 137       | 185                   | 139       | 181                    | 142       |
| SD      | 38                   | 18        | 40                    | 26        | 40                    | 22        | 27                     | 15        | 45                    | 25        | 30                     | 20        |
| SE      | 16                   | 7         | 16                    | 11        | 16                    | 9         | 11                     | 6         | 18                    | 10        | 12                     | 8         |

**End-Diastolic Volume (% change)**

| Subject | % Change Resting to Tilt |          |          |               |          |          | % Change PRE to POST Training, Resting |          |          |               |          |          |
|---------|--------------------------|----------|----------|---------------|----------|----------|----------------------------------------|----------|----------|---------------|----------|----------|
|         | PRE Training             |          |          | POST Training |          |          | PRE Training                           |          |          | POST Training |          |          |
|         | passive                  | exercise | combined | passive       | exercise | combined | passive                                | exercise | combined | passive       | exercise | combined |
| RUI     | -29                      | -20      | -18      | -26           | -23      | -23      | -4                                     | 38       | 0        |               |          |          |
| RAY     | -30                      | -25      | -25      | -26           | -22      | -22      | 15                                     | 14       | 12       |               |          |          |
| HUN     | -30                      | -26      | -31      | -20           | -15      | -16      | -9                                     | -16      | -26      |               |          |          |
| JAG     | -21                      | -24      | -26      | -24           | -24      | -32      | 2                                      | -11      | -2       |               |          |          |
| SCH     | -28                      | -14      |          | -11           | -25      |          | -22                                    | 40       |          |               |          |          |
| FLE     |                          |          | -34      |               |          | -22      |                                        |          | 9        |               |          |          |
| FRE     | 0                        | -1       | -7       | -21           | -5       | -10      | 14                                     | -2       | 6        |               |          |          |
| Mean    | -23                      | -18      | -24      | -21           | -19      | -21      | -1                                     | 10       | 0        |               |          |          |
| SD      | 12                       | 10       | 10       | 6             | 8        | 7        | 14                                     | 24       | 14       |               |          |          |
| SE      | 5                        | 4        | 4        | 2             | 3        | 3        | 6                                      | 10       | 6        |               |          |          |



**Cardiac Output ( $L \cdot \text{min}^{-1}$ )**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt |
| RUI     | 7.04                 | 5.15      | 7.38                  | 5.46      | 5.75                  | 4.19      | 6.77                   | 5.25      | 6.75                  | 5.29      | 6.54                   | 4.96      |
| RAY     | 6.05                 | 4.90      | 6.53                  | 4.95      | 5.15                  | 4.42      | 5.55                   | 4.48      | 6.60                  | 5.16      | 6.23                   | 5.39      |
| HUN     | 8.06                 | 6.47      | 7.85                  | 6.43      | 8.18                  | 6.44      | 7.22                   | 6.75      | 8.85                  | 6.66      | 7.27                   | 6.23      |
| JAG     | 6.87                 | 5.11      | 7.97                  | 5.82      | 7.81                  | 5.42      | 6.85                   | 5.00      | 7.61                  | 5.67      | 9.26                   | 6.32      |
| SCH     | 5.69                 | 4.55      | 6.13                  | 5.30      | 4.85                  | 4.03      | 6.05                   | 4.76      |                       |           |                        |           |
| FLE     |                      |           |                       |           |                       |           |                        |           |                       |           |                        |           |
| FRE     | 4.59                 | 4.37      | 5.58                  | 5.00      | 5.12                  | 4.69      | 5.18                   | 4.39      | 4.93                  | 4.51      | 4.85                   | 4.11      |
| Mean    | 6.38                 | 5.09      | 6.91                  | 5.49      | 6.14                  | 4.87      | 6.27                   | 5.11      | 6.72                  | 5.31      | 6.67                   | 5.10      |
| SD      | 1.21                 | 0.74      | 0.97                  | 0.56      | 1.47                  | 0.91      | 0.81                   | 0.87      | 1.40                  | 0.80      | 1.50                   | 1.12      |
| SE      | 0.49                 | 0.30      | 0.40                  | 0.23      | 0.60                  | 0.37      | 0.33                   | 0.35      | 0.57                  | 0.33      | 0.61                   | 0.46      |

**Cardiac Output (% change)**

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |                                        |          |          |  |  |  |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|----------------------------------------|----------|----------|--|--|--|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          | % Change PRE to POST Training, Resting |          |          |  |  |  |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined | passive                                | exercise | combined |  |  |  |
| RUI     | -26.85                   | -27.13   | -21.63   | -26.02  | -22.45   | -24.16   | 4.83          | 17.74    | -3.11    | 4.83    | 17.74    | -3.11    | 4.83                                   | 17.74    | -3.11    |  |  |  |
| RAY     | -19.01                   | -14.17   | -21.82   | -24.20  | -19.28   | -13.48   | 7.93          | 7.77     | -5.61    | 7.93    | 7.77     | -5.61    | 7.93                                   | 7.77     | -5.61    |  |  |  |
| HUN     | -19.73                   | -21.27   | -24.75   | -18.09  | -6.51    | -14.31   | -2.61         | -11.74   | -17.85   | -2.61   | -11.74   | -17.85   | -2.61                                  | -11.74   | -17.85   |  |  |  |
| JAG     | -25.62                   | -30.60   | -25.49   | -26.98  | -27.01   | -31.75   | 16.01         | -12.29   | 21.68    | 16.01   | -12.29   | 21.68    | 16.01                                  | -12.29   | 21.68    |  |  |  |
| SCH     | -20.04                   | -16.91   |          | -13.54  | -21.32   |          | 7.73          | 24.74    |          | 7.73    | 24.74    |          | 7.73                                   | 24.74    |          |  |  |  |
| FLE     |                          |          | -18.43   |         |          | -39.56   |               |          |          |         |          |          |                                        |          |          |  |  |  |
| FRE     | -4.79                    | -8.40    | -8.52    | -10.39  | -15.25   | -15.26   | 21.57         | 1.17     | -1.62    | 21.57   | 1.17     | -1.62    | 21.57                                  | 1.17     | -1.62    |  |  |  |
| Mean    | -19.34                   | -19.75   | -20.11   | -19.87  | -18.64   | -23.09   | 9.25          | 4.57     | -0.19    | 9.25    | 4.57     | -0.19    | 9.25                                   | 4.57     | -0.19    |  |  |  |
| SD      | 7.85                     | 8.28     | 6.21     | 6.93    | 7.08     | 10.75    | 8.50          | 15.18    | 13.12    | 8.50    | 15.18    | 13.12    | 8.50                                   | 15.18    | 13.12    |  |  |  |
| SE      | 3.20                     | 3.38     | 2.54     | 2.83    | 2.89     | 4.39     | 3.47          | 6.20     | 5.36     | 3.47    | 6.20     | 5.36     | 3.47                                   | 6.20     | 5.36     |  |  |  |

## Arterial Pressure (mmHg)

| Subject | PRE Passive Training |           |  | POST Passive Training |           |  | PRE Exercise Training |           |  | POST Exercise Training |           |  | PRE Combined Training |           |  | POST Combined Training |           |  |
|---------|----------------------|-----------|--|-----------------------|-----------|--|-----------------------|-----------|--|------------------------|-----------|--|-----------------------|-----------|--|------------------------|-----------|--|
|         | resting              | post-tilt |  | resting               | post-tilt |  | resting               | post-tilt |  | resting                | post-tilt |  | resting               | post-tilt |  | resting                | post-tilt |  |
| RUI     | 111                  | 109       |  | 120                   | 123       |  | 94                    | 101       |  | 98                     | 103       |  | 108                   | 110       |  | 103                    | 111       |  |
| RAY     | 107                  | 104       |  | 101                   | 104       |  | 103                   | 111       |  | 101                    | 106       |  | 97                    | 97        |  | 95                     | 97        |  |
| HUN     | 105                  | 111       |  | 122                   | 123       |  | 96                    | 95        |  | 98                     | 99        |  | 83                    | 78        |  | 98                     | 92        |  |
| JAG     | 100                  | 95        |  | 90                    | 89        |  | 102                   | 114       |  | 100                    | 107       |  | 113                   | 119       |  | 89                     | 97        |  |
| SCH     | 99                   | 113       |  | 90                    | 101       |  | 107                   | 104       |  | 97                     | 99        |  |                       |           |  |                        |           |  |
| FLE     |                      |           |  |                       |           |  |                       |           |  |                        |           |  |                       |           |  |                        |           |  |
| FRE     | 90                   | 108       |  | 107                   | 116       |  | 105                   | 124       |  | 88                     | 98        |  | 100                   | 117       |  | 88                     | 86        |  |
| Mean    | 102                  | 107       |  | 105                   | 109       |  | 101                   | 108       |  | 97                     | 102       |  | 99                    | 102       |  | 95                     | 100       |  |
| SD      | 7                    | 6         |  | 14                    | 14        |  | 5                     | 10        |  | 5                      | 4         |  | 11                    | 17        |  | 6                      | 11        |  |
| SE      | 3                    | 3         |  | 6                     | 6         |  | 2                     | 4         |  | 2                      | 2         |  | 5                     | 7         |  | 2                      | 4         |  |

## Arterial Pressure (% change)

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined |
| RUI     | -2                       | 7        | 2        | 3       | 5        | 8        | 8             | 4        | 8        | 8       | 4        | -5       |
| RAY     | -3                       | 8        | 0        | 3       | 5        | 2        | 2             | -2       | -6       | -2      | -2       | -2       |
| HUN     | 6                        | -1       | -6       | 1       | 1        | -6       | 2             | 2        | 16       | 2       | 18       | 18       |
| JAG     | -5                       | 12       | 5        | -1      | 7        | 9        | 9             | -2       | -10      | -2      | -21      | -21      |
| SCH     | 14                       | -3       |          | 12      | 2        |          |               | -9       | -9       | -9      |          |          |
| FLE     |                          |          | -2       |         |          | -2       |               |          |          |         |          | -2       |
| FRE     | 20                       | 18       | 17       | 8       | 11       | 20       | 19            | -16      | 19       | -16     | -5       | -5       |
| Mean    | 5                        | 7        | 3        | 4       | 5        | 5        | 3             | -4       | 3        | -4      | -3       | -3       |
| SD      | 10                       | 8        | 8        | 5       | 4        | 9        | 13            | 8        | 13       | 8       | 13       | 13       |
| SE      | 4                        | 3        | 3        | 2       | 2        | 4        | 5             | 3        | 5        | 3       | 5        | 5        |

## % Change PRE to POST Training, Resting

**Total Peripheral Resistance (mmHg/(L • min<sup>-1</sup>))**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt |
| RUI     | 15.77                | 21.17     | 16.26                 | 22.53     | 16.31                 | 24.11     | 14.42                  | 19.62     | 16.04                 | 20.87     | 15.82                  | 22.46     |
| RAY     | 17.69                | 21.22     | 15.47                 | 21.01     | 20.00                 | 25.11     | 18.20                  | 23.66     | 14.67                 | 18.76     | 15.20                  | 18.00     |
| HUN     | 13.03                | 17.16     | 15.54                 | 19.13     | 11.67                 | 14.77     | 13.55                  | 14.65     | 9.36                  | 11.94     | 13.47                  | 14.83     |
| JAG     | 14.48                | 18.63     | 11.29                 | 15.34     | 13.06                 | 21.03     | 14.53                  | 21.40     | 14.81                 | 21.01     | 9.57                   | 15.33     |
| SCH     | 17.45                | 24.84     | 14.60                 | 19.06     | 22.06                 | 25.81     | 16.03                  | 20.80     |                       |           |                        |           |
| FLE     | 19.54                | 24.71     | 19.18                 | 23.20     | 20.51                 | 26.44     | 17.03                  | 22.39     | 16.07                 | 19.23     | 14.90                  | 24.19     |
| FRE     | 16.33                | 21.29     | 15.39                 | 20.05     | 17.27                 | 22.88     | 15.63                  | 20.42     | 20.22                 | 25.98     | 19.67                  | 27.80     |
| Mean    | 2.37                 | 3.11      | 2.55                  | 2.86      | 4.26                  | 4.40      | 1.77                   | 3.14      | 3.50                  | 4.66      | 3.29                   | 5.21      |
| SD      | 0.97                 | 1.27      | 1.04                  | 1.17      | 1.74                  | 1.80      | 0.72                   | 1.28      | 1.43                  | 1.90      | 1.34                   | 2.13      |

**Total Peripheral Resistance (% change)**

| Subject | % Change Resting to Tilt |          |          |               |          |          | % Change PRE to POST Training, Resting |          |          |          |          |  |
|---------|--------------------------|----------|----------|---------------|----------|----------|----------------------------------------|----------|----------|----------|----------|--|
|         | PRE Training             |          |          | POST Training |          |          | passive                                |          | exercise |          | combined |  |
|         | passive                  | exercise | combined | passive       | exercise | combined | passive                                | exercise | passive  | exercise | combined |  |
| RUI     | 34.24                    | 47.82    | 30.11    | 38.56         | 36.06    | 41.97    | 3.11                                   | -11.59   | 3.11     | -11.59   | -1.37    |  |
| RAY     | 19.95                    | 25.55    | 27.88    | 35.81         | 30.00    | 18.42    | -12.55                                 | -9.00    | -12.55   | -9.00    | 3.61     |  |
| HUN     | 31.70                    | 26.56    | 24.36    | 23.10         | 8.12     | 10.10    | 19.26                                  | 16.11    | 19.26    | 16.11    | 43.91    |  |
| JAG     | 28.66                    | 61.03    | 41.86    | 35.87         | 47.28    | 60.19    | -22.03                                 | 11.26    | -22.03   | 11.26    | -35.38   |  |
| SCH     | 42.35                    | 17.00    |          | 30.55         | 29.76    |          | -16.33                                 | -27.33   | -16.33   | -27.33   |          |  |
| FLE     |                          |          | 19.66    |               |          | 62.35    |                                        |          |          |          | -7.28    |  |
| FRE     | 26.46                    | 28.91    | 28.49    | 20.96         | 31.47    | 41.33    | -1.84                                  | -16.97   | -1.84    | -16.97   | -2.72    |  |
| Mean    | 30.56                    | 34.48    | 28.73    | 30.81         | 30.45    | 39.06    | -5.06                                  | -6.25    | -5.06    | -6.25    | 0.13     |  |
| SD      | 7.57                     | 16.50    | 7.44     | 7.31          | 12.77    | 21.29    | 15.10                                  | 16.74    | 15.10    | 16.74    | 25.52    |  |
| SE      | 3.09                     | 6.74     | 3.04     | 2.98          | 5.21     | 8.69     | 6.16                                   | 6.84     | 6.16     | 6.84     | 10.42    |  |

Mean Data Summary

|     | MEAN |      | SE   |      | RR INT |     | MEAN |      | SE    |       | MEAN |      | SE    |      |
|-----|------|------|------|------|--------|-----|------|------|-------|-------|------|------|-------|------|
|     | CONT | TILT | CONT | TILT | EXB    | EXA | CONT | TILT | CONT  | TILT  | CONT | TILT | CONT  | TILT |
| AP  | 100  | 107  | 1.79 | 4.43 | EXB    | EXA | 1008 | 758  | 81.02 | 53.53 | 100  | 61   | 8.31  | 2.10 |
| EXB | 94   | 100  | 3.16 | 3.43 | EXA    | PSB | 993  | 749  | 89.39 | 73.55 | 98   | 60   | 7.40  | 5.73 |
| EXA | 102  | 107  | 3.05 | 2.61 | PSB    | PSA | 972  | 730  | 59.74 | 41.78 | 103  | 61   | 10.79 | 4.72 |
| PSB | 105  | 109  | 5.77 | 5.53 | PSA    | COB | 875  | 690  | 76.04 | 60.74 | 100  | 62   | 10.13 | 4.79 |
| PSA | 98   | 101  | 4.55 | 6.88 | COB    | COA | 987  | 738  | 53.40 | 43.16 | 110  | 64   | 10.94 | 5.09 |
| COB | 95   | 100  | 2.40 | 4.48 | COA    |     | 975  | 735  | 77.92 | 62.28 | 106  | 63   | 8.78  | 6.19 |
| COA |      |      |      |      |        |     |      |      |       |       |      |      |       |      |

|     | MEAN |      | SE    |       | CO  |     | MEAN |      | SE   |      | MEAN  |       | SE   |      |
|-----|------|------|-------|-------|-----|-----|------|------|------|------|-------|-------|------|------|
|     | CONT | TILT | CONT  | TILT  | EXB | EXA | CONT | TILT | CONT | TILT | CONT  | TILT  | CONT | TILT |
| EDV | 167  | 130  | 14.39 | 8.11  | EXB | EXA | 6.17 | 4.93 | 0.59 | 0.35 | 16.91 | 22.30 | 1.57 | 1.70 |
| EXB | 170  | 133  | 11.21 | 7.99  | EXA | PSB | 6.11 | 4.96 | 0.39 | 0.41 | 15.62 | 20.57 | 0.72 | 1.30 |
| EXA | 172  | 130  | 15.71 | 7.26  | PSB | PSA | 6.38 | 5.09 | 0.49 | 0.30 | 16.33 | 21.29 | 0.97 | 1.27 |
| PSB | 170  | 132  | 16.46 | 10.80 | PSA | COB | 6.91 | 5.49 | 0.40 | 0.23 | 15.39 | 20.04 | 1.04 | 1.17 |
| PSA | 186  | 139  | 18.28 | 10.36 | COB | COA | 6.72 | 5.31 | 0.57 | 0.33 | 15.19 | 19.68 | 1.43 | 1.90 |
| COB | 181  | 142  | 12.18 | 8.10  | COA |     | 6.67 | 5.10 | 0.61 | 0.46 | 14.77 | 20.43 | 1.34 | 2.13 |
| COA |      |      |       |       |     |     |      |      |      |      |       |       |      |      |

|     | MEAN |      | SE   |      |
|-----|------|------|------|------|
|     | CONT | TILT | CONT | TILT |
| HIR | 65   | 84   | 3.60 | 5.50 |
| EXB | 64   | 83   | 5.00 | 7.90 |
| EXA | 63   | 83   | 3.90 | 4.70 |
| PSB | 71   | 90   | 5.20 | 7.00 |
| PSA | 62   | 88   | 3.60 | 7.30 |
| COB | 66   | 85   | 5.60 | 8.50 |
| COA |      |      |      |      |

Hemoglobin - raw data ( $g \cdot dl^{-1}$ )

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 14.504               | 14.869    | 14.830                | 15.771    | 13.679                | 15.387    | 14.062                 | 14.254    | 14.101                | 14.929    | 13.755                 | 14.987    | 13.427     | 15.318    |
| RAY     | 14.005               | 14.466    | 13.218                | 14.024    | 12.834                | 14.120    | 13.295                 | 13.775    | 13.986                | 14.621    | 14.352                 | 14.486    | 14.662     | 15.086    |
| JAG     | 14.696               | 14.561    | 12.604                | 14.715    | 14.888                | 15.598    | 14.907                 | 15.617    | 15.468                | 15.680    | 15.295                 | 14.660    | 14.720     | 15.723    |
| HUN     | 12.949               | 14.888    | 13.410                | 15.195    | 12.872                | 12.546    | 13.698                 | 15.752    | 13.928                | 14.717    | 14.024                 | 14.506    | 13.678     | 15.067    |
| SCH     | 13.487               | 15.176    | 13.487                | 16.385    | 13.909                | 15.771    | 13.391                 | 15.080    |                       |           |                        |           |            |           |
| FRE     | 15.598               | 17.018    | 15.579                | 16.942    | 15.195                | 16.596    | 14.293                 | 16.078    | 15.545                | 16.161    | 14.929                 | 16.527    | 15.067     | 17.189    |
| FLE     |                      |           |                       |           |                       |           |                        |           | 13.466                | 15.276    | 13.331                 | 14.198    | 13.948     | 14.913    |
| Mean    | 14.206               | 15.163    | 13.854                | 15.505    | 13.896                | 15.003    | 13.941                 | 15.093    | 14.416                | 15.231    | 14.281                 | 14.894    | 14.250     | 15.549    |
| SD      | 0.938                | 0.944     | 1.116                 | 1.079     | 0.989                 | 1.445     | 0.609                  | 0.908     | 0.873                 | 0.599     | 0.735                  | 0.840     | 0.656      | 0.851     |
| SE      | 0.383                | 0.385     | 0.456                 | 0.441     | 0.404                 | 0.590     | 0.248                  | 0.371     | 0.356                 | 0.245     | 0.300                  | 0.343     | 0.268      | 0.348     |

Hemoglobin (% change)

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |                                        |          |                                     |         |          |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|----------------------------------------|----------|-------------------------------------|---------|----------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          | % Change PRE to POST Training, Resting |          | % Change PRE to POST Training, Tilt |         |          |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined | passive                                | exercise | combined                            | passive | exercise |
| RUI     | 2.514                    | 12.489   | 5.370    | 6.342   | 1.365    | 8.957    | 2.250         | 2.806    | -2.457   | 6.067   | -7.360   | 0.387    |                                        |          |                                     |         |          |
| RAY     | 3.289                    | 10.020   | 4.542    | 6.099   | 3.609    | 0.939    | -5.619        | 3.589    | 2.615    | -3.052  | -2.447   | -0.922   |                                        |          |                                     |         |          |
| JAG     | -0.914                   | 4.770    | 1.369    | 16.752  | 4.764    | -4.153   | -14.236       | 0.129    | -1.120   | 1.055   | 0.123    | -6.507   |                                        |          |                                     |         |          |
| HUN     | 14.971                   | -2.535   | 5.667    | 13.312  | 14.993   | 3.432    | 3.557         | 6.412    | 0.691    | 2.063   | 25.549   | -1.439   |                                        |          |                                     |         |          |
| SCH     | 12.524                   | 13.386   |          | 21.490  | 12.614   |          | 0.000         | -3.726   |          | 7.968   | -4.381   |          |                                        |          |                                     |         |          |
| FRE     | 9.106                    | 9.221    | 3.963    | 8.748   | 12.489   | -4.900   | -0.123        | -5.937   | -3.963   | -0.451  | -3.123   | 2.263    |                                        |          |                                     |         |          |
| FLE     |                          |          | 13.438   |         |          | 11.721   |               |          | -1.001   |         |          | -7.057   |                                        |          |                                     |         |          |
| Mean    | 6.915                    | 7.892    | 5.808    | 12.124  | 8.306    | 2.666    | -2.362        | 0.546    | -0.872   | 2.275   | 1.394    | -2.212   |                                        |          |                                     |         |          |
| SD      | 6.244                    | 5.934    | 4.073    | 6.192   | 5.719    | 6.766    | 6.611         | 4.674    | 2.313    | 4.102   | 12.085   | 3.767    |                                        |          |                                     |         |          |
| SE      | 2.549                    | 2.423    | 1.663    | 2.528   | 2.335    | 2.762    | 2.699         | 1.908    | 0.944    | 1.675   | 4.934    | 1.538    |                                        |          |                                     |         |          |

**Hemoglobin - raw data - acute exercise ( $\text{g} \cdot \text{dl}^{-1}$ )**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 15.526                | 15.699        | 14.082                 | 16.026        |
| RAY     | 12.927                | 13.659        | 12.966                 | 17.490        |
| JAG     | 12.600                | 13.177        | 14.429                 | 15.526        |
| HUN     | 13.890                | 14.486        | 14.101                 | 15.102        |
| FRE     |                       |               | 16.931                 | 16.758        |
| FLE     | 15.160                | 15.603        | 14.044                 | 16.566        |
| Mean    | 14.021                | 14.525        | 14.425                 | 16.245        |
| SD      | 1.303                 | 1.130         | 1.325                  | 0.870         |
| SE      | 0.583                 | 0.505         | 0.541                  | 0.355         |

**Hemoglobin - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     | 1.1                      | 13.8          |
| RAY     | 5.7                      | 34.9          |
| JAG     | 4.6                      | 7.6           |
| HUN     | 4.3                      | 7.1           |
| FRE     |                          | -1.0          |
| FLE     | 2.9                      | 18.0          |
| Mean    | 3.7                      | 13.4          |
| SD      | 1.8                      | 12.4          |
| SE      | 0.8                      | 5.0           |

Hematocrit - raw data (%)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 42.9                 | 46.1      | 44.7                  | 46.3      | 45.4                  | 46.9      | 44.5                   | 46.7      | 42.6                  | 45.8      | 42.5                   | 45.0      | 43.6       | 51.3      |
| RAY     | 44.2                 | 45.7      | 41.3                  | 44.1      | 42.7                  | 45.9      | 42.6                   | 44.7      | 44.1                  | 45.2      | 45.5                   | 47.8      | 45.6       | 48.0      |
| JAG     | 46.9                 | 47.6      | 41.6                  | 46.2      | 45.9                  | 47.6      | 45.4                   | 48.6      | 43.3                  | 47.5      | 45.9                   | 45.5      | 42.7       | 47.5      |
| HUN     | 42.9                 | 45.9      | 43.0                  | 48.2      | 41.7                  | 46.4      | 41.2                   | 45.9      | 41.3                  | 46.9      | 43.4                   | 44.9      | 42.0       | 44.9      |
| SCH     | 43.8                 | 47.8      | 41.8                  | 47.1      | 46.4                  | 49.5      | 42.7                   | 46.6      |                       |           |                        |           |            |           |
| FRE     | 47.2                 | 50.6      | 45.8                  | 49.6      | 44.7                  | 51.5      | 43.3                   | 48.5      | 46.5                  | 48.8      | 46.3                   | 49.3      | 44.5       | 49.9      |
| FLE     |                      |           |                       |           |                       |           |                        |           | 41.4                  | 47.5      | 42.5                   | 46.0      | 42.0       | 47.0      |
| Mean    | 44.7                 | 47.3      | 43.0                  | 46.9      | 44.5                  | 48.0      | 43.3                   | 46.8      | 43.2                  | 47.0      | 44.3                   | 46.4      | 43.4       | 48.1      |
| SD      | 1.9                  | 1.8       | 0.8                   | 1.9       | 0.8                   | 0.9       | 0.6                    | 0.6       | 1.9                   | 1.3       | 1.7                    | 1.8       | 1.5        | 2.3       |
| SE      | 0.8                  | 0.8       | 0.8                   | 0.8       | 0.8                   | 0.9       | 0.6                    | 0.6       | 0.8                   | 0.5       | 0.7                    | 0.7       | 0.6        | 0.9       |

Hematocrit (% change)

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined |
| RUI     | 7.3                      | 3.2      | 7.5      | 3.4     | 4.8      | 5.9      | 4.2           | -2.0     | -0.3     | 0.4     | -0.4     | -1.7     |
| RAY     | 3.3                      | 7.6      | 2.6      | 6.7     | 4.9      | 5.1      | -6.6          | -0.1     | 3.3      | -3.5    | -2.6     | 5.7      |
| JAG     | 1.6                      | 3.8      | 9.8      | 11.0    | 7.1      | -0.8     | -11.3         | -1.1     | 6.0      | -3.1    | 2.0      | -4.3     |
| HUN     | 7.0                      | 11.3     | 13.6     | 12.2    | 11.4     | 3.3      | 0.2           | -1.2     | 5.1      | 5.1     | -1.1     | -4.4     |
| SCH     | 9.2                      | 6.7      |          | 12.7    | 9.1      |          | -4.5          | -8.0     |          | -1.4    | -5.9     |          |
| FRE     | 7.0                      | 15.3     | 4.9      | 8.3     | 12.0     | 6.5      | -3.0          | -3.1     | -0.5     | -1.8    | -5.9     | 1.0      |
| FLE     |                          |          | 14.7     |         |          | 8.1      |               |          | 2.7      |         |          | -3.3     |
| Mean    | 5.9                      | 8.0      | 8.9      | 9.1     | 8.2      | 4.7      | -3.5          | -2.6     | 2.7      | -0.7    | -2.3     | -1.2     |
| SD      | 2.9                      | 4.6      | 4.8      | 3.6     | 3.1      | 3.1      | 5.4           | 2.8      | 2.7      | 3.2     | 3.1      | 3.9      |
| SE      | 1.2                      | 1.9      | 1.9      | 1.5     | 1.3      | 1.3      | 2.2           | 1.3      | 1.1      | 1.3     | 1.3      | 1.6      |

**Hematocrit - raw data - acute exercise (%)**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 44.3                  | 49.0          | 43.3                   | 46.6          |
| RAY     | 44.8                  | 46.5          | 45.7                   | 46.7          |
| JAG     | 46.6                  | 47.3          | 43.9                   | 45.5          |
| HUN     | 43.1                  | 44.6          | 43.6                   | 44.6          |
| FRE     |                       |               | 48.0                   | 49.5          |
| FLE     | 46.4                  | 49.1          | 43.3                   | 47.5          |
| Mean    | 45.0                  | 47.3          | 44.6                   | 46.7          |
| SD      | 1.5                   | 1.9           | 1.9                    | 1.7           |
| SE      | 0.7                   | 0.8           | 0.8                    | 0.7           |

**Hematocrit - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     | 10.6                     | 7.6           |
| RAY     | 3.8                      | 2.2           |
| JAG     | 1.5                      | 3.5           |
| HUN     | 3.3                      | 2.4           |
| FRE     |                          | 3.1           |
| FLE     | 5.8                      | 9.7           |
| Mean    | 5.0                      | 4.8           |
| SD      | 3.5                      | 3.1           |
| SE      | 1.6                      | 1.3           |



**Absolute Plasma Volume - raw data (ml)**

| Subject | PRE Passive Training |           |          | POST Passive Training |           |          | PRE Exercise Training |           |          | POST Exercise Training |          |           | PRE Combined Training |           |         | POST Combined Training |         |           | POST Study |           |  |
|---------|----------------------|-----------|----------|-----------------------|-----------|----------|-----------------------|-----------|----------|------------------------|----------|-----------|-----------------------|-----------|---------|------------------------|---------|-----------|------------|-----------|--|
|         | resting              | post-tilt | resting  | resting               | post-tilt | resting  | resting               | post-tilt | resting  | post-tilt              | resting  | post-tilt | resting               | post-tilt | resting | post-tilt              | resting | post-tilt | resting    | post-tilt |  |
| RUI     | 4079.630             | 3805.742  | 3470.890 | 3192.531              | 4486.430  | 3905.061 | 4045.867              | 3868.697  | 3432.603 | 3097.912               | 4241.387 | 3756.186  | 4039.263              | 3157.155  |         |                        |         |           |            |           |  |
| RAY     | 3318.130             | 3148.151  | 3543.230 | 3212.897              | 3456.970  | 2999.932 | 3550.740              | 3326.873  | 3776.389 | 3553.297               | 3508.930 | 3360.318  | 3899.327              | 3656.194  |         |                        |         |           |            |           |  |
| JAG     | 3242.540             | 3236.149  | 3310.230 | 2657.322              | 3493.460  | 3249.333 | 3317.526              | 3020.003  | 3469.301 | 3218.142               | 3793.441 | 3979.466  | 3177.299              | 2772.676  |         |                        |         |           |            |           |  |
| HUN     | 3808.510             | 3173.732  | 3717.951 | 3040.232              | 3616.244  | 3469.462 | 3566.740              | 2902.793  | 4130.601 | 3609.798               | 3559.524 | 3371.187  | 3957.060              | 3450.962  |         |                        |         |           |            |           |  |
| SCH     | 2802.510             | 2348.834  | 3036.712 | 2316.454              | 2863.570  | 2410.745 | 3024.270              | 2539.537  | 3443.819 | 3201.675               | 3592.624 | 3102.580  | 3823.708              | 3094.199  |         |                        |         |           |            |           |  |
| FRE     | 3295.060             | 2870.720  | 3688.616 | 3204.134              | 3780.790  | 3123.085 | 3691.250              | 3041.752  | 3300.000 | 2666.140               | 4024.496 | 3597.724  | 3304.174              | 2877.385  |         |                        |         |           |            |           |  |
| FLE     | 3424.397             | 3096.888  | 3461.272 | 2937.262              | 3616.244  | 3192.936 | 3532.732              | 3116.609  | 3592.119 | 3224.494               | 3786.734 | 3527.910  | 3700.139              | 3168.095  |         |                        |         |           |            |           |  |
| Mean    | 452.705              | 477.689   | 285.908  | 370.602               | 527.496   | 493.141  | 344.895               | 447.998   | 307.172  | 342.027                | 293.177  | 314.465   | 364.996               | 336.066   |         |                        |         |           |            |           |  |
| SD      | 184.816              | 195.016   | 104.474  | 151.298               | 215.349   | 203.365  | 140.803               | 182.895   | 125.403  | 139.632                | 119.689  | 128.380   | 149.009               | 137.199   |         |                        |         |           |            |           |  |

**Plasma Volume (% change)**

| Subject | % Change Resting to Tilt |          |          |               |          |          | % Change PRE to POST Training, Resting |          |          |                        |          |          | % Change PRE to POST Training, Tilt |          |          |                     |          |          |
|---------|--------------------------|----------|----------|---------------|----------|----------|----------------------------------------|----------|----------|------------------------|----------|----------|-------------------------------------|----------|----------|---------------------|----------|----------|
|         | PRE Training             |          |          | POST Training |          |          | PRE Training, Resting                  |          |          | POST Training, Resting |          |          | PRE Training, Tilt                  |          |          | POST Training, Tilt |          |          |
|         | passive                  | exercise | combined | passive       | exercise | combined | passive                                | exercise | combined | passive                | exercise | combined | passive                             | exercise | combined | passive             | exercise | combined |
| RUI     | -6.714                   | -12.958  | -9.750   | -8.020        | -4.379   | -11.440  | -14.921                                | -9.820   | 23.562   | -16.113                | -0.931   | 21.249   |                                     |          |          |                     |          |          |
| RAY     | -5.183                   | -13.221  | -5.908   | -9.323        | -6.305   | -4.235   | 6.784                                  | 2.712    | -7.082   | 2.122                  | 10.898   | -5.431   |                                     |          |          |                     |          |          |
| JAG     | -0.197                   | -6.988   | -7.239   | -19.724       | -8.968   | 4.904    | 2.088                                  | -5.036   | 9.343    | -17.886                | -7.058   | 23.657   |                                     |          |          |                     |          |          |
| HUN     | -16.667                  | -4.059   | -12.608  | -18.228       | -18.615  | -5.291   | -2.378                                 | -1.369   | -13.826  | -4.206                 | -16.333  | -6.610   |                                     |          |          |                     |          |          |
| SCH     | -16.188                  | -15.813  | -17.396  | -23.718       | -16.028  | 8.357    | 5.612                                  | 5.342    | 5.342    | -1.379                 | 5.342    | -3.095   |                                     |          |          |                     |          |          |
| FRE     | -12.878                  | -17.396  | -7.031   | -13.135       | -17.396  | -13.640  | 11.944                                 | -2.368   | 4.321    | 11.614                 | -2.604   | 34.941   |                                     |          |          |                     |          |          |
| FLE     |                          |          | -19.208  |               |          | -10.604  |                                        |          | 21.954   |                        |          |          |                                     |          |          |                     |          |          |
| Mean    | -9.638                   | -11.739  | -10.291  | -15.358       | -11.982  | -6.718   | 1.979                                  | -1.711   | 6.379    | -4.308                 | -1.781   | 10.785   |                                     |          |          |                     |          |          |
| SD      | 6.638                    | 5.174    | 4.986    | 6.203         | 6.181    | 6.761    | 9.671                                  | 5.487    | 15.102   | 11.200                 | 9.513    | 17.983   |                                     |          |          |                     |          |          |
| SE      | 2.710                    | 2.112    | 2.036    | 2.533         | 2.523    | 2.760    | 3.948                                  | 2.240    | 6.165    | 4.572                  | 3.884    | 7.341    |                                     |          |          |                     |          |          |

**Plasma Volume - raw data (ml • kg<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |          |          |
|---------|--------------|----------|----------|---------------|----------|----------|------------|----------|----------|
|         | passive      | exercise | combined | passive       | exercise | combined | passive    | exercise | combined |
| RUI     | 47.438       | 52.289   | 39.455   | 40.359        | 46.719   | 48.584   | 45.385     | 45.385   | 45.385   |
| RAY     | 40.514       | 41.154   | 46.853   | 42.767        | 43.196   | 43.054   | 46.960     | 46.960   | 46.960   |
| JAG     | 37.969       | 41.969   | 41.749   | 38.224        | 40.359   | 44.110   | 38.281     | 38.281   | 38.281   |
| HUN     | 39.385       | 42.938   | 42.938   | 38.568        | 38.352   | 36.963   | 40.092     | 40.092   | 40.092   |
| SCH     | 36.972       | 37.778   | 37.596   | 40.062        | 40.324   | 39.178   | 41.115     | 41.115   | 41.115   |
| FRE     | 36.734       | 42.149   | 40.490   | 40.490        | 41.521   | 41.521   | 45.891     | 45.891   | 45.891   |
| FLE     |              |          | 46.479   |               |          | 56.051   |            |          |          |
| Mean    | 39.835       | 43.068   | 42.512   | 40.078        | 41.745   | 44.657   | 42.957     | 42.957   | 42.957   |
| SD      | 3.994        | 5.448    | 3.711    | 1.625         | 2.910    | 6.890    | 3.582      | 3.582    | 3.582    |
| SE      | 1.631        | 2.436    | 1.515    | 0.663         | 1.188    | 2.813    | 1.462      | 1.462    | 1.462    |

Sodium - raw data (mmol • L<sup>-1</sup>)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 137                  | 135       | 138                   | 137       | 136                   | 136       | 138                    | 136       | 138                   | 137       | 138                    | 136       | 137        | 136       |
| RAY     | 137                  | 136       | 137                   | 137       | 137                   | 136       | 137                    | 136       | 136                   | 136       | 137                    | 137       | 132        | 135       |
| HUN     | 134                  | 135       | 138                   | 137       | 140                   | 135       | 134                    | 135       | 137                   | 136       | 136                    | 135       | 139        | 135       |
| JAG     | 138                  | 135       | 139                   | 137       | 146                   | 137       | 138                    | 137       | 138                   | 136       | 137                    | 136       | 138        | 137       |
| SCH     | 136                  | 135       | 137                   | 137       | 136                   | 134       | 139                    | 137       |                       |           |                        |           |            |           |
| FLE     |                      |           |                       |           |                       |           |                        |           |                       |           |                        |           |            |           |
| FRE     | 138                  | 137       | 137                   | 138       | 136                   | 137       | 138                    | 139       | 139                   | 139       | 139                    | 137       | 138        | 138       |
| Mean    | 137                  | 136       | 138                   | 137       | 137                   | 136       | 137                    | 137       | 138                   | 137       | 138                    | 139       | 138        | 138       |
| SD      | 2                    | 1         | 1                     | 0         | 2                     | 1         | 2                      | 1         | 1                     | 1         | 1                      | 1         | 3          | 1         |
| SE      | 1                    | 0         | 0                     | 0         | 1                     | 0         | 1                      | 1         | 0                     | 0         | 0                      | 1         | 1          | 1         |

## Sodium (% change)

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined |
| RUI     | -1.46                    | 0.00     | -0.72    | -0.72   | -1.45    | -1.45    | 0.73          | 1.47     | 0.00     | 1.48    | 0.00     | -0.73    |
| RAY     | -0.73                    | -0.73    | 0.00     | 0.00    | -0.73    | 0.00     | 0.00          | 0.00     | 0.74     | 0.74    | 0.00     | 0.74     |
| HUN     | 0.75                     | -3.57    | -0.73    | -0.72   | 0.75     | -0.74    | 2.99          | -4.29    | -0.73    | 1.48    | 0.00     | -0.74    |
| JAG     | -2.17                    | 0.74     | -1.45    | -1.44   | -0.72    | -0.73    | 0.72          | 1.47     | -0.72    | 1.48    | 0.00     | 0.00     |
| SCH     | -0.74                    | -1.47    |          | 0.00    | -1.44    |          | 0.74          | 2.21     |          | 1.48    | 2.24     |          |
| FLE     |                          |          | 0.00     |         | -1.44    |          |               |          | 0.00     |         |          | -1.44    |
| FRE     | -0.72                    | 0.74     | 0.00     | 0.73    | 0.72     | 0.72     | -0.72         | 1.47     | 0.73     | 0.73    | 1.46     | 1.46     |
| Mean    | -0.85                    | -0.72    | -0.48    | -0.36   | -0.48    | -0.60    | 0.74          | 0.39     | 0.00     | 1.23    | 0.62     | -0.12    |
| SD      | 0.97                     | 1.64     | 0.59     | 0.76    | 0.99     | 0.85     | 1.24          | 2.40     | 0.65     | 0.39    | 0.99     | 1.07     |
| SE      | 0.40                     | 0.67     | 0.24     | 0.31    | 0.41     | 0.35     | 0.51          | 0.98     | 0.27     | 0.16    | 0.40     | 0.44     |

**Sodium - raw data - acute exercise (mmol • L<sup>-1</sup>)**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 137                   | 139           | 137                    | 138           |
| RAY     | 137                   | 138           | 139                    | 140           |
| HUN     | 139                   | 137           | 139                    | 137           |
| JAG     | 138                   | 138           | 139                    | 139           |
| FRE     |                       |               | 138                    | 138           |
| FLE     | 140                   | 139           | 138                    | 139           |
| Mean    | 138                   | 138           | 138                    | 139           |
| SD      | 1                     | 1             | 1                      | 1             |
| SE      | 1                     | 0             | 0                      | 0             |

**Sodium - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | Pre-training             | post-training |
| RUI     | 1.46                     | 0.73          |
| RAY     | 0.73                     | 0.72          |
| HUN     | -1.44                    | -1.44         |
| JAG     | 0.00                     | 0.00          |
| FRE     |                          | 0.00          |
| FLE     | -0.71                    | 0.72          |
| Mean    | 0.01                     | 0.12          |
| SD      | 1.15                     | 0.84          |
| SE      | 0.51                     | 0.34          |

**Potassium - raw data (mmol • L<sup>-1</sup>)**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 4.0                  | 4.1       | 3.8                   | 4.3       | 4.0                   | 4.1       | 4.1                    | 4.3       | 4.4                   | 4.5       | 4.6                    | 4.3       | 3.7        | 4.4       |
| RAY     | 4.2                  | 4.4       | 3.9                   | 3.9       | 3.7                   | 3.8       | 4.0                    | 4.1       | 4.0                   | 3.8       | 4.0                    | 4.1       | 7.3        | 4.0       |
| HUN     | 4.1                  | 4.2       | 4.2                   | 4.2       | 4.2                   | 4.0       | 4.2                    | 4.4       | 4.1                   | 4.3       | 4.6                    | 4.1       | 4.4        | 4.3       |
| JAG     | 4.1                  | 3.9       | 3.8                   | 4.0       | 4.3                   | 4.0       | 3.8                    | 3.8       | 3.8                   | 3.7       | 3.7                    | 3.5       | 3.6        | 3.8       |
| SCH     | 4.3                  | 4.5       | 3.5                   | 3.8       | 4.0                   | 4.5       | 3.9                    | 4.9       |                       |           |                        |           |            |           |
| FLE     |                      |           |                       |           |                       |           |                        |           |                       |           |                        |           |            |           |
| FRE     | 3.9                  | 4.0       | 3.7                   | 3.7       | 3.8                   | 4.1       | 3.4                    | 3.8       | 5.4                   | 5.6       | 4.3                    | 4.7       | 4.0        | 4.5       |
| Mean    | 4.1                  | 4.2       | 3.8                   | 4.0       | 4.0                   | 4.1       | 3.9                    | 4.2       | 4.3                   | 4.3       | 4.2                    | 4.1       | 4.5        | 4.2       |
| SD      | 0.1                  | 0.2       | 0.2                   | 0.2       | 0.2                   | 0.2       | 0.3                    | 0.4       | 0.6                   | 0.7       | 0.4                    | 0.4       | 1.4        | 0.3       |
| SE      | 0.1                  | 0.1       | 0.1                   | 0.1       | 0.1                   | 0.1       | 0.1                    | 0.2       | 0.2                   | 0.3       | 0.1                    | 0.2       | 0.6        | 0.1       |

**Potassium (% change)**

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined |
| RUI     | 2.5                      | 2.5      | 2.3      | 13.2    | 4.9      | -6.5     | -5.0          | 2.5      | 4.5      | 4.9     | 4.9      | -4.4     |
| RAY     | 4.8                      | 2.7      | -5.0     | 0.0     | 2.5      | 2.5      | -7.1          | 8.1      | 0.0      | -11.4   | 7.9      | 7.9      |
| HUN     | 2.4                      | -4.8     | 4.9      | 0.0     | 7.3      | -10.9    | 2.4           | -2.4     | 12.2     | 0.0     | 10.0     | -4.7     |
| JAG     | -4.9                     | -7.0     | -2.6     | 5.3     | 0.0      | -5.4     | -7.3          | -11.6    | -2.6     | 2.6     | -5.0     | -5.4     |
| SCH     | 4.7                      | 12.5     |          | 8.6     | 25.6     |          | -16.6         | -2.5     | -20.4    | -15.6   | 8.9      | -16.1    |
| FLE     |                          |          | 3.7      |         |          | 9.3      |               |          |          |         |          |          |
| FRE     | 2.6                      | 7.9      | -4.9     | 0.0     | 11.8     | -2.4     | -5.1          | -10.5    | 0.0      | -7.5    | -7.3     | 2.6      |
| Mean    | 2.0                      | 2.3      | -0.3     | 4.5     | 8.7      | -2.2     | -6.8          | -2.7     | -1.0     | -4.5    | 3.2      | -3.4     |
| SD      | 3.5                      | 7.4      | 4.4      | 5.5     | 9.2      | 7.2      | 6.8           | 7.5      | 10.8     | 8.2     | 7.5      | 8.1      |
| SE      | 1.4                      | 3.0      | 1.8      | 2.3     | 3.8      | 2.9      | 2.8           | 3.1      | 4.4      | 3.3     | 3.1      | 3.3      |

**Potassium - raw data - acute exercise (mmol • L<sup>-1</sup>)**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 4.0                   | 3.8           | 3.9                    | 3.8           |
| RAY     | 4.2                   | 3.7           | 4.0                    | 3.6           |
| HUN     | 5.0                   | 4.2           | 4.4                    | 4.1           |
| JAG     | 3.7                   | 3.5           | 3.8                    | 3.6           |
| FRE     |                       |               | 4.3                    | 3.5           |
| FLE     | 4.0                   | 4.1           | 4.1                    | 4.2           |
| Mean    | 4.2                   | 3.9           | 4.1                    | 3.8           |
| SD      | 0.5                   | 0.3           | 0.2                    | 0.3           |
| SE      | 0.2                   | 0.1           | 0.1                    | 0.1           |

**Potassium - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     | -5.000                   | -2.564        |
| RAY     | -11.905                  | -10.000       |
| HUN     | -16.000                  | -6.818        |
| JAG     | -5.405                   | -5.263        |
| FRE     |                          | -18.605       |
| FLE     | 2.500                    | 2.439         |
| Mean    | -7.162                   | -6.802        |
| SD      | 7.101                    | 7.145         |
| SE      | 3.176                    | 2.917         |

Osmolality - raw data (mmol • L<sup>-1</sup>)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 288.281              | 286.269   | 288.302               | 287.800   | 286.963               | 283.952   | 290.301                | 286.281   | 291.977               | 290.972   | 289.623                | 287.610   | 287.822    | 287.822   |
| RAY     | 288.281              | 287.275   | 287.800               | 286.797   | 284.454               | 285.457   | 288.794                | 287.789   | 286.952               | 287.957   | 292.138                | 293.648   | 285.318    | 286.820   |
| JAG     | 288.784              | 288.784   | 288.804               | 287.800   | 287.465               | 289.473   | 287.789                | 291.809   | 286.952               | 288.459   | 290.629                | 289.623   | 289.824    | 290.325   |
| HUN     | 285.457              | 286.461   | 289.296               | 288.794   | 292.809               | 284.759   | 281.778                | 283.785   | 287.957               | 286.449   | 287.610                | 282.578   | 295.332    | 289.324   |
| SCH     | 284.454              | 283.450   | 286.281               | 285.779   | 282.747               | 282.747   | 290.812                | 288.804   |                       |           |                        |           |            |           |
| FRE     | 287.967              | 289.473   | 289.296               | 287.789   | 286.460               | 286.962   | 288.804                | 288.804   | 285.946               | 283.936   | 284.591                | 285.597   | 284.317    | 286.820   |
| FLE     |                      |           |                       |           |                       |           |                        |           | 291.977               | 293.987   | 284.591                | 291.635   | 283.329    | 291.326   |
| Mean    | 287.204              | 286.962   | 288.297               | 287.460   | 286.816               | 285.558   | 288.046                | 287.879   | 288.627               | 288.197   | 288.449                | 288.449   | 289.324    | 288.739   |
| SD      | 1.790                | 2.135     | 1.146                 | 1.038     | 3.424                 | 2.384     | 3.263                  | 2.701     | 2.672                 | 3.506     | 3.157                  | 4.046     | 4.376      | 1.884     |
| SE      | 0.731                | 0.872     | 0.468                 | 0.424     | 1.398                 | 0.973     | 1.332                  | 1.103     | 1.091                 | 1.431     | 1.289                  | 1.652     | 1.787      | 0.769     |

Osmolality (% change)

| Subject | % Change Resting to Tilt |          |          |               |          |          | % Change PRE to POST Training, Tilt |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------------|----------|----------|-------------------------------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          | POST Training |          |          | Resting                             |          |          | Tilt    |          |          |
|         | passive                  | exercise | combined | passive       | exercise | combined | passive                             | exercise | combined | passive | exercise | combined |
| RUI     | -0.698                   | -1.049   | -0.344   | -0.174        | -1.385   | -0.695   | 0.007                               | 1.163    | -0.806   | 0.535   | 0.820    | -1.155   |
| RAY     | -0.349                   | 0.353    | 0.350    | -0.349        | -0.348   | 0.517    | -0.167                              | 1.526    | 1.808    | -0.166  | 0.817    | 1.976    |
| JAG     | 0.000                    | 0.698    | 0.525    | -0.348        | 1.397    | -0.346   | 0.007                               | 0.113    | 1.282    | -0.341  | 0.807    | 0.403    |
| HUN     | 0.352                    | -2.749   | -0.524   | -0.174        | 0.712    | -1.749   | 1.345                               | -3.767   | -0.120   | 0.814   | -0.342   | -1.351   |
| SCH     | -0.353                   | 0.000    |          | -0.176        | -0.690   |          | 0.642                               | 2.852    |          | 0.822   | 2.142    |          |
| FRE     | 0.523                    | 0.175    | -0.703   | -0.521        | 0.000    | 0.354    | 0.462                               | 0.818    | -0.474   | -0.582  | 0.642    | 0.585    |
| FLE     |                          |          | 0.688    |               |          | 2.475    |                                     |          | -2.530   |         |          | -0.800   |
| Mean    | -0.088                   | -0.429   | -0.001   | -0.290        | -0.052   | 0.092    | 0.383                               | 0.451    | -0.140   | 0.180   | 0.814    | -0.057   |
| SD      | 0.466                    | 1.281    | 0.593    | 0.142         | 0.996    | 1.423    | 0.563                               | 2.257    | 1.554    | 0.618   | 0.791    | 1.280    |
| SE      | 0.190                    | 0.523    | 0.242    | 0.058         | 0.407    | 0.581    | 0.230                               | 0.921    | 0.635    | 0.252   | 0.323    | 0.523    |

**Osmolality - raw data - acute exercise (mmol • L<sup>-1</sup>)**

| Subject | PRE Combined Training |               |  | POST Combined Training |               |  |
|---------|-----------------------|---------------|--|------------------------|---------------|--|
|         | resting               | post-exercise |  | resting                | post-exercise |  |
| RUI     | 293.333               | 301.334       |  | 289.306                | 293.823       |  |
| RAY     | 288.333               | 291.333       |  | 293.321                | 298.340       |  |
| JAG     | 288.333               | 295.333       |  | 294.325                | 297.336       |  |
| HUN     | 290.333               | 289.333       |  | 293.321                | 290.310       |  |
| FRE     |                       |               |  | 289.808                | 294.325       |  |
| FLE     | 292.833               | 295.833       |  | 294.325                | 294.325       |  |
| Mean    | 290.633               | 294.633       |  | 292.016                | 294.826       |  |
| SD      | 2.388                 | 4.632         |  | 2.289                  | 3.174         |  |
| SE      | 1.068                 | 2.071         |  | 0.934                  | 1.296         |  |

**Osmolality - acute exercise (% change)**

| Subject | % Change Resting to Tilt |              |
|---------|--------------------------|--------------|
|         | pre-training             | pre-training |
| RUI     | 2.727                    | 1.561        |
| RAY     | 1.041                    | 1.711        |
| JAG     | 2.428                    | 1.023        |
| HUN     | -0.344                   | -1.027       |
| FRE     |                          | 1.559        |
| FLE     | 1.025                    | 0.000        |
| Mean    | 1.375                    | 0.805        |
| SD      | 1.238                    | 1.096        |
| SE      | 0.554                    | 0.448        |

Albumin - raw data (g • dl<sup>-1</sup>)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 4.4                  | 4.4       | 4.1                   | 4.5       | 4.1                   | 4.3       | 4.2                    | 4.3       | 3.9                   | 4.3       | 4.0                    | 4.3       | 4.1        | 5.1       |
| RAY     | 4.3                  | 4.5       | 4.0                   | 4.3       | 4.1                   | 4.4       | 4.1                    | 4.4       | 4.2                   | 4.3       | 4.5                    | 4.7       | 4.5        | 4.8       |
| HUN     | 4.0                  | 4.5       | 4.1                   | 4.0       | 4.4                   | 4.8       | 4.4                    | 4.9       | 4.2                   | 4.6       | 4.6                    | 4.7       | 4.2        | 4.5       |
| JAG     | 4.7                  | 4.6       | 3.8                   | 4.3       | 4.4                   | 4.6       | 4.5                    | 4.8       | 4.3                   | 4.7       | 4.6                    | 4.6       | 4.2        | 4.7       |
| SCH     | 4.3                  | 4.8       | 4.1                   | 4.8       | 4.7                   | 5.0       | 4.2                    | 4.5       |                       |           |                        |           |            |           |
| FLE     |                      |           |                       |           |                       |           |                        |           |                       |           |                        |           |            |           |
| FRE     | 4.1                  | 4.4       | 4.1                   | 4.5       | 3.9                   | 4.6       | 3.8                    | 4.4       | 4.1                   | 5.0       | 4.2                    | 4.6       | 4.1        | 4.7       |
| Mean    | 4.3                  | 4.5       | 4.0                   | 4.4       | 4.3                   | 4.6       | 4.2                    | 4.6       | 4.2                   | 4.5       | 4.2                    | 4.5       | 3.9        | 4.6       |
| SD      | 0.2                  | 0.2       | 0.1                   | 0.3       | 0.3                   | 0.3       | 0.2                    | 0.2       | 0.2                   | 0.3       | 0.3                    | 0.2       | 0.2        | 0.2       |
| SE      | 0.1                  | 0.1       | 0.0                   | 0.1       | 0.1                   | 0.1       | 0.1                    | 0.1       | 0.1                   | 0.1       | 0.1                    | 0.1       | 0.1        | 0.1       |

Albumin (% change)

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined |
| RUI     | 0.0                      | 4.9      | 10.3     | 9.8     | 2.4      | 7.5      | -6.8          | 2.4      | 2.6      | 2.3     | 0.0      | 0.0      |
| RAY     | 4.7                      | 7.3      | 2.4      | 7.5     | 7.3      | 4.4      | -7.0          | 0.0      | 7.1      | -4.4    | 0.0      | 9.3      |
| HUN     | 12.5                     | 9.1      | 9.5      | -2.4    | 11.4     | 2.2      | 2.5           | 0.0      | 9.5      | -11.1   | 2.1      | 2.2      |
| JAG     | -2.1                     | 4.5      | 9.3      | 13.2    | 6.7      | 0.0      | -19.1         | 2.3      | 7.0      | -6.5    | 4.3      | -2.1     |
| SCH     | 11.6                     | 6.4      |          | 17.1    | 7.1      |          | -4.7          | -10.6    |          | 0.0     | -10.0    |          |
| FLE     |                          |          | 22.0     |         |          | 9.5      |               |          | 2.4      |         |          | -8.0     |
| FRE     | 7.3                      | 17.9     | -6.7     | 9.8     | 15.8     | 7.1      | 0.0           | -2.6     | -6.7     | 2.3     | -4.3     | 7.1      |
| Mean    | 5.7                      | 8.4      | 7.8      | 9.1     | 8.4      | 5.1      | -5.8          | -1.4     | 3.7      | -2.9    | -1.3     | 1.4      |
| SD      | 6.0                      | 5.0      | 9.5      | 6.6     | 4.6      | 3.6      | 7.5           | 4.9      | 5.8      | 5.4     | 5.1      | 6.3      |
| SE      | 2.4                      | 2.0      | 3.9      | 2.7     | 1.9      | 1.5      | 3.1           | 2.0      | 2.4      | 2.2     | 2.1      | 2.6      |



**Albumin - raw data - acute exercise (g • dl<sup>-1</sup>)**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 4.1                   | 4.8           | 4.1                    | 4.5           |
| RAY     | 4.2                   | 4.5           | 4.3                    | 4.4           |
| HUN     | 4.3                   | 4.5           | 4.4                    | 4.4           |
| JAG     | 4.4                   | 4.6           | 4.2                    | 4.4           |
| FRE     |                       |               | 4.3                    | 4.4           |
| FLE     | 4.6                   | 4.9           | 4.2                    | 4.7           |
| Mean    | 4.3                   | 4.7           | 4.3                    | 4.5           |
| SD      | 0.2                   | 0.2           | 0.1                    | 0.1           |
| SE      | 0.1                   | 0.1           | 0.0                    | 0.0           |

**Albumin - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     | 17.07                    | 9.76          |
| RAY     | 7.14                     | 2.33          |
| HUN     | 4.65                     | 0.00          |
| JAG     | 4.55                     | 4.76          |
| FRE     |                          | 2.33          |
| FLE     | 6.52                     | 11.90         |
| Mean    | 7.99                     | 5.18          |
| SD      | 5.21                     | 4.68          |
| SE      | 2.33                     | 1.91          |

**Total Protein - raw data ( $g \cdot dl^{-1}$ )**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 7.0                  | 6.8       | 6.9                   | 7.4       | 6.6                   | 6.7       | 7.0                    | 7.3       | 6.8                   | 7.3       | 6.7                    | 7.3       | 6.7        | 8.5       |
| RAY     | 7.3                  | 7.6       | 6.9                   | 7.6       | 7.0                   | 7.8       | 7.0                    | 7.5       | 7.4                   | 7.6       | 7.9                    | 8.2       | 7.8        | 8.4       |
| HUN     | 6.9                  | 7.7       | 6.9                   | 8.3       | 6.8                   | 7.6       | 7.3                    | 8.3       | 7.2                   | 7.7       | 7.4                    | 7.8       | 6.4        | 7.6       |
| JAG     | 7.5                  | 7.6       | 6.4                   | 7.2       | 7.3                   | 7.6       | 7.4                    | 8.0       | 7.0                   | 7.7       | 7.5                    | 7.5       | 6.9        | 7.7       |
| SCH     | 7.0                  | 7.7       | 6.7                   | 7.6       | 7.8                   | 7.9       | 6.9                    | 7.4       |                       |           |                        |           |            |           |
| FLE     |                      |           |                       |           |                       |           |                        |           | 6.1                   | 7.5       | 6.2                    | 7.0       | 6.8        | 7.2       |
| FRE     | 6.5                  | 7.1       | 6.5                   | 7.1       | 6.0                   | 7.0       | 6.1                    | 7.0       | 7.1                   | 6.7       | 6.6                    | 7.3       | 5.8        | 7.0       |
| Mean    | 7.0                  | 7.4       | 6.7                   | 7.5       | 6.9                   | 7.4       | 7.0                    | 7.6       | 6.9                   | 7.4       | 7.1                    | 7.5       | 6.7        | 7.7       |
| SD      | 0.3                  | 0.4       | 0.2                   | 0.4       | 0.6                   | 0.5       | 0.5                    | 0.5       | 0.5                   | 0.4       | 0.6                    | 0.4       | 0.7        | 0.6       |
| SE      | 0.1                  | 0.2       | 0.1                   | 0.2       | 0.3                   | 0.2       | 0.2                    | 0.2       | 0.2                   | 0.2       | 0.3                    | 0.2       | 0.3        | 0.2       |

**Total Protein (% change)**

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined |
| RUI     | -2.9                     | 1.5      | 7.4      | 7.2     | 4.3      | 9.0      | -1.4          | 6.1      | -1.5     | 8.8     | 9.0      | 0.0      |
| RAY     | 4.1                      | 11.4     | 2.7      | 10.1    | 7.1      | 3.8      | -5.5          | 0.0      | 6.8      | 0.0     | -3.8     | 7.9      |
| HUN     | 11.6                     | 11.8     | 6.9      | 20.3    | 13.7     | 5.4      | 0.0           | 7.4      | 2.8      | 7.8     | 9.2      | 1.3      |
| JAG     | 1.3                      | 4.1      | 10.0     | 12.5    | 8.1      | 0.0      | -14.7         | 1.4      | 7.1      | -5.3    | 5.3      | -2.6     |
| SCH     | 10.0                     | 1.3      |          | 13.4    | 7.2      |          | -4.3          | -11.5    |          | -1.3    | -6.3     |          |
| FLE     |                          |          | 23.0     |         |          | 12.9     |               |          | 1.6      |         |          | -6.7     |
| FRE     | 9.2                      | 16.7     | -5.6     | 9.2     | 14.8     | 10.6     | 0.0           | 1.7      | -7.0     | 0.0     | 0.0      | 9.0      |
| Mean    | 5.6                      | 7.8      | 7.4      | 12.1    | 9.2      | 6.9      | -4.3          | 0.8      | 1.6      | 1.7     | 2.2      | 1.5      |
| SD      | 5.7                      | 6.4      | 9.4      | 4.6     | 4.1      | 4.8      | 5.6           | 6.7      | 5.3      | 5.5     | 6.6      | 6.0      |
| SE      | 2.3                      | 2.6      | 3.8      | 1.9     | 1.7      | 1.9      | 2.3           | 2.7      | 2.2      | 2.2     | 2.7      | 2.5      |

**Total Protein - raw data - acute exercise (g • dl<sup>-1</sup>)**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 7.0                   | 8.0           | 6.8                    | 7.4           |
| RAY     | 7.6                   | 8.0           | 7.6                    | 7.8           |
| HUN     | 7.1                   | 7.5           | 7.2                    | 7.6           |
| JAG     | 7.2                   | 7.6           | 6.9                    | 7.3           |
| FRE     |                       |               | 6.9                    | 7.1           |
| FLE     | 6.7                   | 7.2           | 6.3                    | 7.2           |
| Mean    | 7.1                   | 7.7           | 7.0                    | 7.4           |
| SD      | 0.3                   | 0.3           | 0.4                    | 0.3           |
| SE      | 0.1                   | 0.2           | 0.2                    | 0.1           |

**Total Protein - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     | 14.29                    | 8.82          |
| RAY     | 5.26                     | 2.63          |
| HUN     | 5.63                     | 5.56          |
| JAG     | 5.56                     | 5.80          |
| FRE     |                          | 2.90          |
| FLE     | 7.46                     | 14.29         |
| Mean    | 7.64                     | 6.67          |
| SD      | 3.81                     | 4.36          |
| SE      | 1.71                     | 1.78          |

PRA - raw data (ngAng1 • ml<sup>-1</sup> • hr<sup>-1</sup>)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 0.493                | 1.215     | 0.406                 | 0.821     | 0.485                 | 0.939     | 0.435                  | 0.844     | 0.497                 | 1.390     | 0.527                  | 1.114     |            |           |
| RAY     | 0.596                | 1.32      | 0.164                 | 1.342     | 0.186                 | 1.344     | 0.208                  | 1.425     | 0.338                 | 0.861     | 0.201                  | 0.767     |            |           |
| HUN     | 0.602                | 1.646     | 0.63                  | 1.912     | 0.468                 | 1.14      | 0.737                  | 2.277     | 0.550                 | 1.500     | 0.638                  | 1.583     | 0.464      | 1.388     |
| JAG     | 0.667                | 2.471     | 0.435                 | 1.292     | 0.42                  | 2.07      | 0.692                  | 2.633     | 2.135                 | 1.390     | 0.338                  | 1.114     | 0.419      | 2.135     |
| SCH     | 1.39                 | 5.633     | 0.894                 | 4.338     | 0.781                 | 4.539     | 0.578                  | 2.052     |                       |           |                        |           |            |           |
| FLE     | 0.86                 | 2.552     | 0.91                  | 2.912     | 0.363                 | 1.709     | 0.717                  | 2.094     | 0.612                 | 2.497     | 0.819                  | 2.927     | 0.648      | 1.994     |
| Mean    | 0.768                | 2.473     | 0.573                 | 2.103     | 0.451                 | 1.957     | 0.561                  | 1.888     | 0.747                 | 1.511     | 0.529                  | 1.583     | 0.442      | 1.864     |
| SD      | 0.328                | 1.648     | 0.295                 | 1.309     | 0.195                 | 1.328     | 0.207                  | 0.645     | 0.688                 | 0.585     | 0.225                  | 0.785     | 0.032      | 0.358     |
| SE      | 0.134                | 0.673     | 0.120                 | 0.534     | 0.079                 | 0.542     | 0.084                  | 0.263     | 0.281                 | 0.218     | 0.092                  | 0.320     | 0.023      | 0.179     |

PRA (% change)

| Subject | % Change Resting to Tilt |          |          |         |          |          |               |          |          |         |          |          |                                     |
|---------|--------------------------|----------|----------|---------|----------|----------|---------------|----------|----------|---------|----------|----------|-------------------------------------|
|         | PRE Training             |          |          |         |          |          | POST Training |          |          |         |          |          | % Change PRE to POST Training, Tilt |
|         | passive                  | exercise | combined | passive | exercise | combined | passive       | exercise | combined | passive | exercise | combined |                                     |
| RUI     | 146.450                  | 93.608   | 179.678  | 102.217 | 94.023   | 111.385  | -17.647       | -10.309  | 6.036    | -32.428 | -10.117  | -19.856  |                                     |
| RAY     | 121.477                  | 622.581  | 154.734  | 718.293 | 585.096  | 281.592  | -72.483       | 11.828   | -40.533  | 1.667   | 6.027    | -10.918  |                                     |
| HUN     | 173.422                  | 143.590  | 172.727  | 203.492 | 208.955  | 148.119  | 4.651         | 57.479   | 16.000   | 16.160  | 99.737   | 5.533    |                                     |
| JAG     | 270.465                  | 392.857  | -34.895  | 197.011 | 280.491  | 229.586  | -34.783       | 64.762   | -84.169  | -47.713 | 27.198   | -19.856  |                                     |
| SCH     | 305.252                  | 481.178  |          | 385.235 | 255.017  |          | -35.683       | -25.992  |          | -22.990 | -54.792  |          |                                     |
| FLE     | 196.744                  | 370.799  | 304.830  | 220.000 | 192.050  | 207.716  | 5.814         | 97.521   | 33.824   | 14.107  | 22.528   | 39.930   |                                     |
| Mean    | 202.302                  | 350.769  | 180.847  | 304.375 | 269.272  | 205.964  | -25.022       | 32.548   | 2.542    | -11.866 | 15.097   | 2.009    |                                     |
| SD      | 71.780                   | 201.055  | 125.567  | 222.512 | 167.563  | 65.140   | 29.486        | 48.112   | 58.645   | 26.362  | 50.927   | 23.686   |                                     |
| SE      | 29.304                   | 82.080   | 51.262   | 90.840  | 68.407   | 26.593   | 12.038        | 19.642   | 23.942   | 10.762  | 20.791   | 9.670    |                                     |

**Aldosterone - raw data (pg • ml<sup>-1</sup>)**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt |
| RUI     | 149.087              | 357.806   | 102.642               | 250.962   | 125.598               | 177.032   | 70.365                 | 278.832   |
| RAY     | 74.245               | 201.832   | 60.847                | 222.110   | 83.356                | 279.259   | 88.277                 | 397.239   |
| HUN     | 111.086              | 237.211   | 64.978                | 351.807   | 14.914                | 147.741   | 104.705                | 299.916   |
| JAG     | 148.937              | 197.230   | 61.102                | 198.811   | 128.596               | 293.518   | 115.886                | 154.855   |
| SCH     | 142.308              | 332.195   | 55.110                | 187.267   | 103.822               | 217.047   | 69.904                 | 137.984   |
| FLE     |                      |           |                       |           |                       |           |                        |           |
| FRE     | 64.680               | 331.408   | 89.960                | 392.905   | 22.154                | 310.478   | 33.381                 | 323.423   |
| Mean    | 115.057              | 276.280   | 72.440                | 267.310   | 79.740                | 237.513   | 80.420                 | 265.375   |
| SD      | 38.125               | 72.291    | 19.173                | 85.233    | 50.202                | 66.842    | 29.440                 | 100.573   |
| SE      | 15.565               | 29.513    | 7.827                 | 34.796    | 20.495                | 27.288    | 12.019                 | 41.059    |

**Aldosterone (% change)**

| Subject | % Change Resting to Tilt |          |           |               |          |           | % Change PRE to POST Training, Resting |          | % Change PRE to POST Training, Tilt |          |
|---------|--------------------------|----------|-----------|---------------|----------|-----------|----------------------------------------|----------|-------------------------------------|----------|
|         | PRE Training             |          |           | POST Training |          |           | passive                                | exercise | passive                             | exercise |
|         | passive                  | exercise | post-tilt | passive       | exercise | post-tilt |                                        |          |                                     |          |
| RUI     | 139.998                  | 40.951   | 144.502   | 144.502       | 296.265  | 296.265   | -31.153                                | -43.976  | -29.861                             | 57.504   |
| RAY     | 171.846                  | 235.020  | 265.030   | 265.030       | 349.992  | 349.992   | -18.046                                | 5.904    | 10.047                              | 42.248   |
| HUN     | 113.538                  | 890.620  | 441.425   | 441.425       | 186.439  | 186.439   | -41.507                                | 602.058  | 48.310                              | 103.001  |
| JAG     | 32.425                   | 128.248  | 225.376   | 225.376       | 33.627   | 33.627    | -58.975                                | -9.884   | 0.802                               | -47.242  |
| SCH     | 133.434                  | 109.057  | 239.806   | 239.806       | 97.391   | 97.391    | -61.274                                | -32.669  | -43.627                             | -36.427  |
| FLE     |                          |          |           |               |          |           |                                        |          |                                     |          |
| FRE     | 412.381                  | 1301.453 | 336.755   | 336.755       | 868.863  | 868.863   | 39.085                                 | 50.677   | 18.556                              | 4.169    |
| Mean    | 167.270                  | 450.891  | 275.482   | 275.482       | 305.433  | 305.433   | -28.645                                | 95.352   | 0.704                               | 20.542   |
| SD      | 128.887                  | 520.080  | 102.313   | 102.313       | 300.288  | 300.288   | 37.029                                 | 250.449  | 33.382                              | 57.870   |
| SE      | 52.618                   | 212.322  | 41.769    | 41.769        | 122.592  | 122.592   | 15.117                                 | 102.246  | 13.628                              | 23.626   |

AVP - raw data (pg • ml<sup>-1</sup>)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 0.430                | 148.935   | 0.300                 | 1.380     | 1.855                 | 3.612     | 0.836                  | 0.602     | 12.410                | 6.700     | 8.110                  | 7.560     |            |           |
| RAY     | 0.820                | 43.913    | 2.798                 | 4.434     | 1.092                 | 10.451    | 1.956                  | 2.035     | 6.180                 | 64.050    | 2.320                  | 4.830     |            |           |
| HUN     | 3.500                | 5.928     | 2.035                 | 5.316     | 3.500                 | 7.477     | 2.213                  | 5.435     | 2.530                 | 8.460     | 3.070                  | 5.970     | 1.290      | 5.160     |
| JAG     | 0.955                | 230.047   | 0.300                 | 188.264   | 0.300                 | 184.579   | 2.709                  | 154.067   | 5.580                 | 5.780     | 2.460                  | 10.080    | 0.300      | 116.490   |
| SCH     | 3.388                | 4.050     | 1.371                 | 2.297     | 0.300                 | 1.577     | 2.346                  | 1.952     |                       |           |                        |           |            |           |
| FLE     |                      |           |                       |           |                       |           |                        |           | 7.010                 | 3.470     | 2.080                  | 2.200     |            | 10.480    |
| FRE     | 1.248                | 3.353     | 3.458                 | 3.210     | 1.641                 | 2.341     | 3.073                  | 4.528     | 6.742                 | 5.630     | 5.810                  | 5.180     |            | 6.220     |
| Mean    | 1.724                | 72.704    | 1.710                 | 34.150    | 1.448                 | 35.006    | 2.189                  | 28.103    | 6.742                 | 15.682    | 3.975                  | 5.970     | 0.795      | 34.588    |
| SD      | 1.359                | 95.267    | 1.299                 | 75.513    | 1.198                 | 73.353    | 0.770                  | 61.735    | 3.213                 | 23.751    | 2.446                  | 2.667     | 0.700      | 54.650    |
| SE      | 0.555                | 38.893    | 0.530                 | 30.828    | 0.489                 | 29.946    | 0.314                  | 25.203    | 1.312                 | 9.696     | 0.998                  | 1.089     | 0.495      | 27.325    |

AVP (% change)

| Subject | % Change Resting to Tilt |           |          |           |          |          |               |          |          |         |          |          |
|---------|--------------------------|-----------|----------|-----------|----------|----------|---------------|----------|----------|---------|----------|----------|
|         | PRE Training             |           |          |           |          |          | POST Training |          |          |         |          |          |
|         | passive                  | exercise  | combined | passive   | exercise | combined | passive       | exercise | combined | passive | exercise | combined |
| RUI     | 34536.047                | 94.717    | -46.011  | 360.000   | -27.990  | -6.782   | -30.233       | -54.933  | -34.649  | -99.073 | -83.333  | 12.836   |
| RAY     | 5255.244                 | 857.051   | 936.408  | 58.470    | 4.039    | 108.190  | 241.220       | 79.121   | -62.460  | -89.903 | -80.528  | -92.459  |
| HUN     | 69.371                   | 113.629   | 234.387  | 161.229   | 145.594  | 94.463   | 41.111        | -36.771  | 21.344   | -10.324 | -27.310  | -29.433  |
| JAG     | 23988.691                | 61426.333 | 3.584    | 62654.667 | 5587.228 | 309.756  | 303.000       | 803.000  | -55.914  | -18.163 | -16.531  | 74.394   |
| SCH     | 19.540                   | 425.667   |          | 67.542    | -16.795  |          | -59.534       | 682.000  |          | -43.284 | 23.779   |          |
| FLE     |                          |           | -50.499  |           |          | 5.769    |               |          | -70.328  |         |          | -36.599  |
| FRE     | 168.670                  | 42.657    | -16.494  | -7.172    | 47.348   | -10.843  | 177.083       | 87.264   | -13.824  | -4.265  | 93.422   | -7.993   |
| Mean    | 10672.927                | 10493.342 | 176.896  | 10549.123 | 956.571  | 83.425   | 36.349        | 259.947  | -35.972  | -44.169 | -15.084  | -13.209  |
| SD      | 14915.621                | 24953.847 | 387.039  | 25526.719 | 2269.426 | 122.487  | 136.038       | 380.196  | 34.805   | 41.282  | 66.897   | 55.625   |
| SE      | 6089.277                 | 10187.365 | 158.008  | 10421.239 | 926.489  | 50.005   | 55.537        | 155.214  | 14.209   | 16.853  | 27.311   | 22.709   |

Epinephrine - raw data (pg • ml<sup>-1</sup>)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 28.029               | 144.016   | 20.248                | 34.242    | 16.376                | 30.616    | 36.512                 | 15.312    | 15.563                | 25.002    | 21.920                 | 22.313    | 18.430     | 22.880    |
| RAY     | 17.545               | 40.960    | 14.917                | 30.132    | 18.444                | 65.170    | 27.253                 | 34.269    | 17.131                | 84.813    | 17.092                 | 23.246    | 14.471     | 13.906    |
| HUN     | 18.048               | 59.483    | 38.082                | 63.720    | 36.772                | 96.789    | 40.816                 | 86.018    | 25.210                | 83.093    | 58.276                 | 51.451    | 20.651     | 75.989    |
| JAG     | 41.816               | 334.930   | 38.237                | 138.135   | 36.268                | 124.241   | 27.369                 | 163.797   | 33.476                | 34.893    | 30.851                 | 46.949    | 20.543     | missing   |
| SCH     | 13.125               | 33.273    | 20.450                | 16.515    | 47.070                | 52.434    | 18.654                 | 46.616    |                       |           |                        |           |            |           |
| FRE     | 30.916               | 23.385    | 24.923                | 24.689    | 26.457                | 33.069    | 25.007                 | 37.487    | 35.293                | 22.777    | 28.727                 | 18.352    | 20.878     | 20.970    |
| FLE     |                      |           |                       |           |                       |           |                        |           | 48.126                | 29.167    | 30.435                 | 22.222    | 10.080     | 44.092    |
| Mean    | 24.913               | 106.008   | 26.143                | 51.239    | 30.231                | 67.053    | 29.269                 | 63.917    | 29.133                | 46.624    | 31.217                 | 30.756    | 17.509     | 35.567    |
| SD      | 10.694               | 120.302   | 9.833                 | 45.501    | 11.899                | 37.026    | 8.054                  | 54.229    | 12.338                | 29.213    | 14.318                 | 14.456    | 4.375      | 25.248    |
| SE      | 4.366                | 49.113    | 4.014                 | 18.576    | 4.858                 | 15.116    | 3.288                  | 22.139    | 5.037                 | 11.926    | 5.845                  | 5.902     | 1.786      | 11.291    |

Epinephrine (% change)

| Subject | % Change Resting to Tilt |          |          |               |          |          | % Change PRE to POST Training, Resting |          |          |         |          |          | % Change PRE to POST Training, Tilt |          |          |         |          |          |
|---------|--------------------------|----------|----------|---------------|----------|----------|----------------------------------------|----------|----------|---------|----------|----------|-------------------------------------|----------|----------|---------|----------|----------|
|         | PRE Training             |          |          | POST Training |          |          | Resting                                |          |          | Tilt    |          |          | Resting                             |          |          | Tilt    |          |          |
|         | passive                  | exercise | combined | passive       | exercise | combined | passive                                | exercise | combined | passive | exercise | combined | passive                             | exercise | combined | passive | exercise | combined |
| RUI     | 413.811                  | 86.957   | 60.650   | 69.113        | -58.063  | 1.793    | -27.761                                | 122.960  | 40.847   | -76.223 | -49.987  | -10.755  |                                     |          |          |         |          |          |
| RAY     | 133.457                  | 253.340  | 395.085  | 101.998       | 25.744   | 36.005   | -14.979                                | 47.761   | -0.228   | -26.436 | -47.416  | -72.591  |                                     |          |          |         |          |          |
| HUN     | 229.582                  | 163.214  | 229.603  | 67.323        | 110.746  | -11.712  | 111.004                                | 10.997   | 131.162  | 7.123   | 111.128  | -38.080  |                                     |          |          |         |          |          |
| JAG     | 700.961                  | 242.564  | 4.233    | 261.260       | 498.476  | 52.180   | -8.559                                 | -24.537  | -7.841   | -58.757 | 31.838   | 34.551   |                                     |          |          |         |          |          |
| SCH     | 153.509                  | 11.396   |          | -19.242       | 149.896  |          | 55.810                                 | -60.370  |          |         |          |          |                                     |          |          |         |          |          |
| FRE     | -24.360                  | 24.991   | -35.463  | -0.939        | 49.906   | -36.116  | -19.385                                | -5.481   | -18.604  | 5.576   | 13.360   | -19.427  |                                     |          |          |         |          |          |
| FLE     |                          |          | -39.395  |               |          | -26.985  |                                        |          | -36.760  |         |          |          |                                     |          |          |         |          |          |
| Mean    | 267.827                  | 130.410  | 102.452  | 79.919        | 129.451  | 2.528    | 16.022                                 | 15.222   | 18.096   | -33.180 | -12.405  | -21.686  |                                     |          |          |         |          |          |
| SD      | 255.686                  | 105.766  | 174.549  | 100.018       | 194.521  | 35.082   | 55.390                                 | 63.887   | 61.100   | 34.566  | 32.443   | 35.077   |                                     |          |          |         |          |          |
| SE      | 104.383                  | 43.179   | 71.259   | 40.832        | 79.413   | 14.322   | 22.613                                 | 26.082   | 24.944   | 14.112  | 13.245   | 14.320   |                                     |          |          |         |          |          |

**Epinephrine - raw data - acute exercise (pg • ml<sup>-1</sup>)**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 54.082                | missing       | missing                | 54.266        |
| RAY     | 31.500                | 64.149        | 18.761                 | 66.705        |
| HUN     | 29.036                | 48.546        | 58.940                 | 43.083        |
| JAG     | 43.424                | 56.128        | 23.978                 | 55.807        |
| FRE     |                       |               | 19.834                 | 102.327       |
| FLE     | 22.554                | 69.892        | 23.978                 | 67.166        |
| Mean    | 36.119                | 59.679        | 29.098                 | 64.892        |
| SD      | 12.565                | 9.325         | 16.850                 | 20.401        |
| SE      | 5.619                 | 4.662         | 7.535                  | 8.329         |

**Epinephrine - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     |                          |               |
| RAY     | 103.648                  | 255.551       |
| HUN     | 67.192                   | -26.904       |
| JAG     | 29.256                   | 132.743       |
| FRE     |                          | 415.917       |
| FLE     | 209.887                  | 180.115       |
| Mean    | 102.496                  | 191.485       |
| SD      | 77.770                   | 162.594       |
| SE      | 38.885                   | 72.714        |



Norepinephrine - raw data (pg • ml<sup>-1</sup>)

| Subject | PRE Passive Training |           |         | POST Passive Training |         |           | PRE Exercise Training |           |         | POST Exercise Training |         |           | PRE Combined Training |           |         | POST Combined Training |         |           | POST Study |           |  |
|---------|----------------------|-----------|---------|-----------------------|---------|-----------|-----------------------|-----------|---------|------------------------|---------|-----------|-----------------------|-----------|---------|------------------------|---------|-----------|------------|-----------|--|
|         | resting              | post-tilt | resting | post-tilt             | resting | post-tilt | resting               | post-tilt | resting | post-tilt              | resting | post-tilt | resting               | post-tilt | resting | post-tilt              | resting | post-tilt | resting    | post-tilt |  |
| RUI     | 169.172              | 212.949   | 175.499 | 314.789               | 162.432 | 257.912   | 215.705               | 245.747   | 223.676 | 399.289                | 171.708 | 267.670   | 189.042               | 375.817   |         |                        |         |           |            |           |  |
| RAY     | 276.897              | 326.257   | 208.384 | 362.776               | 453.670 | 544.551   | 372.298               | 473.356   | 267.709 | 446.010                | 320.078 | 385.135   | 439.785               | 375.905   |         |                        |         |           |            |           |  |
| HUN     | 173.142              | 489.283   | 145.396 | 525.802               | 170.992 | 470.814   | 264.716               | 767.811   | 166.873 | 407.244                | 190.477 | 387.404   | 194.068               | 551.196   |         |                        |         |           |            |           |  |
| JAG     | 323.645              | 671.999   | 185.571 | 395.282               | 192.055 | 391.816   | 211.841               | 617.460   | 140.300 | 335.193                | 192.742 | 221.498   | 144.963               | missing   |         |                        |         |           |            |           |  |
| SCH     | 326.938              | 614.880   | 273.698 | 575.111               | 263.230 | 657.913   | 244.534               | 663.757   |         |                        |         |           |                       |           |         |                        |         |           |            |           |  |
| FRE     | 194.895              | 503.182   | 206.904 | 531.035               | 179.355 | 563.161   | 241.860               | 718.770   | 277.457 | 385.217                | 288.196 | 664.631   | 241.349               | 414.663   |         |                        |         |           |            |           |  |
| FLE     |                      |           |         |                       |         |           |                       |           | 178.433 | 314.192                | 143.585 | 390.813   | 151.927               | 381.555   |         |                        |         |           |            |           |  |
| Mean    | 244.115              | 469.758   | 199.242 | 450.799               | 236.956 | 481.028   | 258.492               | 581.150   | 209.075 | 381.191                | 217.798 | 386.192   | 226.856               | 419.827   |         |                        |         |           |            |           |  |
| SD      | 73.940               | 173.207   | 43.211  | 106.626               | 112.156 | 141.360   | 59.108                | 192.893   | 56.173  | 48.639                 | 69.890  | 154.093   | 109.886               | 75.192    |         |                        |         |           |            |           |  |
| SE      | 30.186               | 70.712    | 17.641  | 43.530                | 45.787  | 57.710    | 24.131                | 78.748    | 22.933  | 19.857                 | 28.533  | 62.908    | 44.861                | 33.627    |         |                        |         |           |            |           |  |

Norepinephrine (% change)

| Subject | % Change Resting to Tilt |          |          |               |          |          | % Change PRE to POST Training, Resting |          |          |          |          |          | % Change PRE to POST Training, Tilt |          |          |          |          |          |          |  |  |
|---------|--------------------------|----------|----------|---------------|----------|----------|----------------------------------------|----------|----------|----------|----------|----------|-------------------------------------|----------|----------|----------|----------|----------|----------|--|--|
|         | PRE Training             |          |          | POST Training |          |          | passive                                |          |          | exercise |          |          | passive                             |          |          | exercise |          |          | combined |  |  |
|         | passive                  | exercise | combined | passive       | exercise | combined | passive                                | exercise | combined | passive  | exercise | combined | passive                             | exercise | combined | passive  | exercise | combined |          |  |  |
| RUI     | 25.877                   | 58.782   | 78.512   | 79.368        | 13.927   | 55.887   | 3.740                                  | 32.797   | -23.234  | 47.824   | -4.717   | -32.963  |                                     |          |          |          |          |          |          |  |  |
| RAY     | 17.826                   | 20.032   | 66.603   | 74.090        | 27.144   | 20.325   | -24.743                                | -17.936  | 19.562   | 11.193   | -13.074  | -13.649  |                                     |          |          |          |          |          |          |  |  |
| HUN     | 162.591                  | 175.343  | 144.044  | 261.634       | 190.051  | 103.386  | -16.025                                | 54.812   | 14.145   | 7.464    | 63.082   | -4.872   |                                     |          |          |          |          |          |          |  |  |
| JAG     | 107.635                  | 104.012  | 138.912  | 113.008       | 191.473  | 14.919   | -42.662                                | 10.302   | 37.378   | -41.178  | 57.589   | -33.919  |                                     |          |          |          |          |          |          |  |  |
| SCH     | 88.072                   | 149.938  |          | 110.126       | 171.438  |          | -16.284                                | -7.103   |          | -6.468   | 0.888    |          |                                     |          |          |          |          |          |          |  |  |
| FRE     | 158.181                  | 213.992  | 38.838   | 156.658       | 197.184  | 130.618  | 6.162                                  | 34.850   | 3.871    | 5.535    | 27.631   | 72.534   |                                     |          |          |          |          |          |          |  |  |
| FLE     |                          |          | 76.084   |               |          | 172.182  |                                        |          | -19.530  |          |          | 24.387   |                                     |          |          |          |          |          |          |  |  |
| Mean    | 96.697                   | 120.350  | 90.499   | 132.481       | 131.870  | 82.886   | -14.969                                | 17.954   | 5.365    | 4.062    | 21.900   | 1.920    |                                     |          |          |          |          |          |          |  |  |
| SD      | 67.213                   | 73.198   | 41.958   | 69.821        | 86.772   | 63.142   | 18.231                                 | 27.708   | 23.424   | 28.765   | 32.789   | 40.696   |                                     |          |          |          |          |          |          |  |  |
| SE      | 27.440                   | 29.883   | 17.129   | 28.504        | 35.425   | 25.778   | 7.443                                  | 11.312   | 9.563    | 11.743   | 13.386   | 16.614   |                                     |          |          |          |          |          |          |  |  |

**Norepinephrine - raw data - acute exercise ( $\text{pg} \cdot \text{ml}^{-1}$ )**

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 237.392               | missing       | missing                | 235.102       |
| RAY     | 266.089               | 781.450       | 457.296                | 1079.190      |
| HUN     | 173.154               | 437.889       | 240.208                | 556.476       |
| JAG     | 139.143               | 663.536       | 110.318                | 400.582       |
| FRE     |                       |               | 354.625                | 1292.650      |
| FLE     | 208.973               | 735.650       | 181.977                | 467.804       |
| Mean    | 204.950               | 654.631       | 268.885                | 671.967       |
| SD      | 50.369                | 152.429       | 138.175                | 417.277       |
| SE      | 22.526                | 76.214        | 61.794                 | 170.353       |

**Norepinephrine - acute exercise (% change)**

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     | 193.680                  | 135.994       |
| RAY     | 152.890                  | 131.664       |
| JAG     | 376.873                  | 263.116       |
| FRE     |                          | 264.512       |
| FLE     | 252.031                  | 157.068       |
| Mean    | 243.869                  | 190.471       |
| SD      | 97.558                   | 67.641        |
| SE      | 48.779                   | 30.250        |

Dopamine - raw data (pg • ml<sup>-1</sup>)

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 4.193                | 9.119     | 10.500                | 11.567    | 18.684                | 8.971     | 13.661                 | 0.497     | 7.468                 | 8.899     | 29.816                 | 22.133    | 4.881      | 15.548    |
| RAY     | 18.871               | 16.748    | 10.219                | 7.704     | 10.475                | 8.514     | 9.033                  | 7.042     | 4.111                 | 6.655     | 13.452                 | 8.164     | 3.380      | 3.764     |
| HUN     | 17.376               | 50.139    | 202.880               | 14.158    | 1.611                 | 14.401    | 20.237                 | 35.581    | 3.501                 | 12.495    | 10.893                 | 9.545     | 8.397      | 13.340    |
| JAG     | 24.251               | 23.293    | 24.206                | 9.646     | 12.482                | 33.404    | 24.811                 | 16.466    | 2.411                 | 5.992     | 10.893                 | 10.001    | 1.736      | missing   |
| SCH     | 38.563               | 24.781    | 25.591                | 12.282    | 32.459                | 32.878    | 8.341                  | 20.785    |                       |           |                        |           |            |           |
| FRE     | 14.231               | 22.141    | 7.004                 | 16.473    | 11.954                | 31.145    | 192.340                | 17.557    | 10.530                | -4.145    | 4.653                  | 11.189    | 15.204     | 10.004    |
| FLE     |                      |           |                       |           |                       |           |                        |           | 24.049                | 8.685     | 12.671                 | 6.030     | 2.886      | 11.001    |
| Mean    | 19.581               | 24.370    | 46.733                | 11.972    | 14.606                | 21.552    | 44.737                 | 16.321    | 8.682                 | 6.430     | 13.439                 | 11.177    | 6.081      | 10.731    |
| SD      | 11.427               | 13.864    | 76.891                | 3.128     | 10.324                | 12.167    | 72.594                 | 12.074    | 8.100                 | 5.658     | 8.614                  | 5.651     | 5.028      | 4.448     |
| SE      | 4.665                | 5.660     | 31.390                | 1.277     | 4.215                 | 4.967     | 29.636                 | 4.929     | 3.307                 | 2.310     | 3.517                  | 2.307     | 2.053      | 1.989     |

Dopamine (% change)

| Subject | % Change Resting to Tilt |         |               |         |               |         | % Change PRE to POST Training, Resting |          |          |         |          |          | % Change PRE to POST Training, Tilt |         |          |         |          |          |
|---------|--------------------------|---------|---------------|---------|---------------|---------|----------------------------------------|----------|----------|---------|----------|----------|-------------------------------------|---------|----------|---------|----------|----------|
|         | PRE Training             |         | POST Training |         | POST Training |         | passive                                |          | exercise |         | combined |          | passive                             |         | exercise |         | combined |          |
| RUI     | 117.482                  | -51.986 | 18.843        | 10.162  | -96.382       | -25.773 | 150.417                                | -26.884  | 298.210  | 26.845  | -94.460  | 148.713  | 28.845                              | -84.460 | 148.713  | 28.845  | -84.460  | 148.713  |
| RAY     | -11.250                  | -18.721 | 61.883        | -24.611 | -22.041       | -39.310 | -45.848                                | -13.766  | 227.220  | -4.000  | -17.289  | 22.675   | -4.000                              | -17.289 | 22.675   | -4.000  | -17.289  | 22.675   |
| HUN     | 188.583                  | 793.917 | 256.898       | -93.021 | 75.822        | -12.375 | 1067.587                               | 1156.176 | 211.140  | -71.763 | 147.073  | -23.609  | -71.763                             | 147.073 | -23.609  | -71.763 | 147.073  | -23.609  |
| JAG     | -3.950                   | 168.282 | 148.528       | -60.150 | -33.694       | 9.372   | -0.186                                 | 99.283   | 279.282  | -58.588 | -50.707  | 66.906   | -58.588                             | -50.707 | 66.906   | -58.588 | -50.707  | 66.906   |
| SCH     | -35.739                  | 1.291   |               | -52.007 | 149.191       |         | -33.638                                | -74.303  |          | -50.438 |          |          | -50.438                             |         |          | -50.438 |          |          |
| FRE     | 55.683                   | 160.540 | -139.364      | 135.194 | -90.872       | 140.469 | -50.784                                | 1509.001 | -55.812  | -25.600 | -43.628  | -369.940 | -25.600                             | -43.628 | -369.940 | -25.600 | -43.628  | -369.940 |
| FLE     |                          |         | -63.866       |         |               | -52.411 |                                        |          | -47.312  |         |          |          |                                     |         |          |         |          |          |
| Mean    | 51.780                   | 175.551 | 47.150        | -14.072 | -2.983        | 3.329   | 181.258                                | 441.580  | 162.118  | -38.924 | -15.965  | -30.971  | -38.924                             | -15.965 | -30.971  | -38.924 | -15.965  | -30.971  |
| SD      | 86.921                   | 317.029 | 143.113       | 80.962  | 97.071        | 70.516  | 440.693                                | 701.450  | 161.021  | 35.575  | 83.845   | 178.717  | 35.575                              | 83.845  | 178.717  | 35.575  | 83.845   | 178.717  |
| SE      | 35.485                   | 129.426 | 58.426        | 33.052  | 39.629        | 28.788  | 179.912                                | 286.366  | 65.737   | 14.524  | 34.230   | 72.981   | 14.524                              | 34.230  | 72.981   | 14.524  | 34.230   | 72.981   |

Dopamine - raw data - acute exercise ( $\text{pg} \cdot \text{ml}^{-1}$ )

| Subject | PRE Combined Training |               | POST Combined Training |               |
|---------|-----------------------|---------------|------------------------|---------------|
|         | resting               | post-exercise | resting                | post-exercise |
| RUI     | 3.779                 | missing       | missing                | 52.459        |
| RAY     | 8.098                 | 28.476        | 11.667                 | 27.252        |
| HUN     | 4.037                 | 9.704         | 32.832                 | 15.210        |
| JAG     | 6.271                 | 22.096        | -1.084                 | 11.547        |
| FRE     |                       |               | 9.247                  | 53.940        |
| FLE     | 4.602                 | 25.677        | 8.913                  | 20.596        |
| Mean    | 5.357                 | 21.488        | 12.315                 | 30.167        |
| SD      | 1.813                 | 8.279         | 12.469                 | 18.615        |
| SE      | 0.811                 | 4.139         | 5.576                  | 7.600         |

## Dopamine - acute exercise (% change)

| Subject | % Change Resting to Tilt |               |
|---------|--------------------------|---------------|
|         | pre-training             | post-training |
| RUI     |                          |               |
| RAY     | 251.642                  | 133.582       |
| HUN     | 140.377                  | -53.673       |
| JAG     | 252.352                  | -1165.221     |
| FRE     |                          | 483.324       |
| FLE     | 457.953                  | 131.078       |
| Mean    | 275.581                  | -94.182       |
| SD      | 132.480                  | 629.464       |
| SE      | 66.240                   | 281.505       |

**Plasma Human Growth Hormone (ng • ml<sup>-1</sup>)**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           | POST Study |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt | resting    | post-tilt |
| RUI     | 1                    | 1 #       | 1                     | 1         | 1                     | 1         | 1                      | 1         | 1                     | 1         | 1                      | 1         | 1          | 1         |
| RAY     | 1                    | 16.0 #    | 1                     | 17.1      | 1                     | 7.7 #     | 1                      | 6.4       | 1                     | 6.8 #     | 1                      | 7.4 #     | 1          | 9.3       |
| JAG     | 1                    | 22.4 #    | 1                     | 3.7 #     | 6.2                   | 4.7 #     | 1                      | 16.0 #    | 6.6                   | 7.1 #     | 1                      | 1 #       | 1          | 1.6       |
| HUN     | 1                    | 18.5      | 1                     | 3.0       | 1                     | 6.0       | 1                      | 7.1       | 1                     | 2.9 #     | 1                      | 1.5 #     | 1          | 2.4       |
| SCH     | 1                    | 1 #       | 1                     | 1         | 1                     | 2.0 #     | 1                      | 1 #       | 1                     | 1         | 1                      | 1         | 1          | 1         |
| FRE     | 1                    | 1         | 1                     | 1         | 1                     | 1         | 1                      | 1         | 1                     | 1         | 1                      | 1         | 1          | 1         |
| Mean    | 1                    | 10.0      | 1                     | 4.5       | 1.9                   | 3.7       | 1                      | 5.4       | 1.9                   | 3.3       | 1                      | 2.2       | 1          | 2.7       |
| SD      | 0                    | 10.0      | 0                     | 6.3       | 2.1                   | 2.8       | 0                      | 5.9       | 2.3                   | 2.9       | 0                      | 2.6       | 0          | 3.3       |
| SE      | 0                    | 4.1       | 0                     | 2.6       | 0.9                   | 1.2       | 0                      | 2.4       | 0.9                   | 1.2       | 0                      | 1.1       | 0          | 1.3       |

Individual radioimmunoassay data. # Pre-syncope signs or symptoms.

**Plasma Human Growth Hormone (ng HGH • ml<sup>-1</sup>)**

| Subject | PRE Passive Training |           | POST Passive Training |           | PRE Exercise Training |           | POST Exercise Training |           | PRE Combined Training |           | POST Combined Training |           |
|---------|----------------------|-----------|-----------------------|-----------|-----------------------|-----------|------------------------|-----------|-----------------------|-----------|------------------------|-----------|
|         | resting              | post-tilt | resting               | post-tilt | resting               | post-tilt | resting                | post-tilt | resting               | post-tilt | resting                | post-tilt |
| Mean    | 725                  | 2125      | 775                   | 575       | 720                   | 775       | 730                    | 1850      | 1375                  | 2075      | 1205                   | 2000      |
| SE      | 84                   | 99        | 85                    | 81        | 84                    | 85        | 84                     | 97        | 93                    | 99        | 91                     | 98        |
| CL      | 693-763              | 1859-2428 | 660-910               | 498-703   | 643-818               | 730-823   | 643-838                | 1688-2033 | 1180-1595             | 1820-2388 | 105-1315               | 1893-2103 |
|         | P < 0.001            |           | NS                    |           | NS                    |           | P < 0.001              |           | P < 0.001             |           | P < 0.001              |           |

Mean summed bioassay data. CL = 95% confidence limits. NS = not significant.

**Plasma Human Growth Hormone (ng • ml<sup>-1</sup>)**

| Subject | BEFORE Combined Training |               | AFTER Combined Training |               |
|---------|--------------------------|---------------|-------------------------|---------------|
|         | pre exercise             | post exercise | pre exercise            | post exercise |
| RUI     | 1                        | 1             | 1                       | 1             |
| RAY     | 1                        | 10.9          | 1                       | 1.5           |
| JAG     | 1                        | 6.2           | 1                       | 13.5          |
| HUN     | 1                        | 1             | 1                       | 1             |
| SCH     |                          |               |                         |               |
| FRE     | 1                        | 1             | 1                       | 26.6          |
| Mean    | 1                        | 4.0           | 1                       | 7.4           |
| SD      | 0                        | 4.5           | 0                       | 10.6          |
| SE      | 0                        | 2.0           | 0                       | 4.3           |

Individual radioimmunoassay data.

**Urine Volume (ml)**

| Subject                | PRE Training |          |          | POST Training |          |          | POST Study |
|------------------------|--------------|----------|----------|---------------|----------|----------|------------|
|                        | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE                    |              |          | 2295     |               |          | 1935     | 1085       |
| FRE                    | 3945         | 3065     | 2975     | 3000          | 3350     | 3395     | 3355       |
| HUN                    | 2730         | 2990     | 1435     | 2390          | 2020     | 2645     | 2015       |
| JAG                    | 3162         | 1640     | 1365     | 2570          | 1710     | 1485     | 2065       |
| RAY                    | 710          | 965      | 1025     | 1030          | 780      | 1045     | 926        |
| RUI                    | 1760         | 1805     | 1575     | 1270          | 1760     | 2185     | 1815       |
| SCH                    | 750          | 1082     |          | 810           | 1140     |          |            |
| Mean                   | 2176         | 1925     | 1778     | 1845          | 1793     | 2115     | 1877       |
| SD                     | 1324         | 912      | 721      | 919           | 887      | 836      | 869        |
| SE                     | 540          | 372      | 294      | 375           | 362      | 341      | 355        |
| ml / min <sup>-1</sup> | 1.5          | 1.3      | 1.2      | 1.3           | 1.2      | 1.5      | 1.3        |

**Urine Volume (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -16      |
| FRE     | -24                           | 9        | 14       |
| HUN     | -12                           | -32      | 84       |
| JAG     | -19                           | 4        | 9        |
| RAY     | 45                            | -19      | 2        |
| RUI     | -28                           | -2       | 39       |
| SCH     | 8                             | 5        |          |
| Mean    | -5                            | -6       | 22       |
| SD      | 28                            | 16       | 35       |
| SE      | 11                            | 7        | 14       |

**Creatinine (mg • 24 hr<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 6265     |               |          | 2187     | 1172       |
| FRE     | 986          | 1624     | 1190     | 1080          | 1173     | 985      | 772        |
| HUN     | 956          | 2063     | 1521     | 1745          | 1252     | 2566     | 3949       |
| JAG     | 3826         | 2214     | 1119     | 1491          | 770      | 3104     | 2003       |
| RAY     | 2109         | 1891     | 2081     | 2410          | 1724     | 2174     | 1889       |
| RUI     | 1549         | 2274     | 2741     | 2337          | 2534     | 1901     | 1289       |
| SCH     | 705          | 822      |          | 1361          | 1516     |          |            |
| Mean    | 1689         | 1815     | 2486     | 1737          | 1495     | 2153     | 1846       |
| SD      | 1163         | 540      | 1949     | 538           | 603      | 708      | 1128       |
| SE      | 475          | 221      | 796      | 220           | 246      | 289      | 461        |

**Creatinine (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -65.1    |
| FRE     | 9.5                           | -27.8    | -17.2    |
| HUN     | 82.5                          | -39.3    | 68.7     |
| JAG     | -61.0                         | -65.2    | 177.4    |
| RAY     | 14.3                          | -8.8     | 4.5      |
| RUI     | 50.9                          | 11.4     | -30.6    |
| SCH     | 93.0                          | 84.4     |          |
| Mean    | 31.5                          | -7.5     | 22.9     |
| SD      | 56.8                          | 52.1     | 87.8     |
| SE      | 23.2                          | 21.3     | 35.8     |

**Deoxyypyridinoline (DPD) (nmol • L<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 112.8    |               |          | 40.9     | 37.3       |
| FRE     | 13.0         | 8.4      | 9.2      | 12.3          | 7.3      | 10.1     | 9.6        |
| HUN     | 17.0         | 25.2     | 53.4     | 27.5          | 32.3     | 50.2     | 127.8      |
| JAG     | 28.3         | 33.9     | 30.5     | 15.5          | 14.3     | 60.1     | 26.3       |
| RAY     | 63.1         | 49.3     | 38.3     | 66.3          | 93.2     | 50.3     | 63.6       |
| RUI     | 42.5         | 61.5     | 97.8     | 83.7          | 68.9     | 33.2     | 26.7       |
| SCH     | 31.0         | 16.1     |          | 60.6          | 43.1     |          |            |
| Mean    | 32.5         | 32.4     | 57.0     | 44.3          | 43.2     | 40.8     | 48.6       |
| SD      | 18.3         | 20.1     | 40.3     | 29.8          | 32.9     | 17.6     | 42.7       |
| SE      | 7.5          | 8.2      | 16.5     | 12.2          | 13.4     | 7.2      | 17.4       |

**DPD (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -63.7    |
| FRE     | -5.4                          | -13.1    | 9.8      |
| HUN     | 61.8                          | 28.2     | -6.0     |
| JAG     | -45.2                         | -57.8    | 97.0     |
| RAY     | 5.1                           | 89.0     | 31.3     |
| RUI     | 96.9                          | 12.0     | -66.1    |
| SCH     | 95.5                          | 167.7    |          |
| Mean    | 34.8                          | 37.7     | 0.4      |
| SD      | 58.6                          | 80.0     | 61.6     |
| SE      | 23.9                          | 32.7     | 25.1     |

**DPD/Creatinine (nmol/nmol creat)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 4.7      |               |          | 4.1      | 3.9        |
| FRE     | 5.9          | 1.8      | 2.6      | 3.9           | 2.4      | 3.9      | 4.7        |
| HUN     | 5.5          | 4.1      | 5.7      | 4.3           | 5.9      | 5.8      | 7.4        |
| JAG     | 2.6          | 2.8      | 4.2      | 3.0           | 3.6      | 3.2      | 3.1        |
| RAY     | 2.4          | 2.8      | 2.1      | 3.2           | 4.8      | 2.7      | 3.5        |
| RUI     | 5.5          | 5.5      | 6.4      | 5.1           | 5.4      | 4.3      | 4.2        |
| SCH     | 3.7          | 2.4      |          | 4.1           | 3.7      |          |            |
| Mean    | 4.3          | 3.2      | 4.3      | 3.9           | 4.3      | 4.0      | 4.5        |
| SD      | 1.6          | 1.3      | 1.7      | 0.8           | 1.3      | 1.1      | 1.5        |
| SE      | 0.6          | 0.5      | 0.7      | 0.3           | 0.5      | 0.4      | 0.6        |

**DPD/Creatinine (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -12.8    |
| FRE     | -33.9                         | 33.3     | 50.0     |
| HUN     | -21.8                         | 43.9     | 1.8      |
| JAG     | 15.4                          | 28.6     | -23.8    |
| RAY     | 33.3                          | 71.4     | 28.6     |
| RUI     | -7.3                          | -1.8     | -32.8    |
| SCH     | 10.8                          | 54.2     |          |
| Mean    | -0.6                          | 38.3     | 1.8      |
| SD      | 25.1                          | 24.9     | 32.0     |
| SE      | 10.2                          | 10.2     | 13.0     |

**NTx (nmol • 24 hr<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 1643.7   |               |          | 566.0    | 295.4      |
| FRE     | 199.6        | 345.4    | 298.7    | 320.1         | 242.5    | 247.8    | 227.5      |
| HUN     | 230.1        | 655.7    | 322.3    | 408.9         | 235.5    | 610.2    | 1000.4     |
| JAG     | 1215.2       | 804.7    | 640.9    | 507.1         | 465.6    | 1125.5   | 667.8      |
| RAY     | 496.5        | 332.0    | 375.5    | 589.1         | 566.5    | 514.8    | 479.8      |
| RUI     | 459.2        | 924.9    | 1135.9   | 647.1         | 804.0    | 531.8    | 359.6      |
| SCH     | 148.9        | 195.5    |          | 333.6         | 289.8    |          |            |
| Mean    | 458.3        | 543.0    | 736.2    | 467.7         | 434.0    | 599.4    | 505.1      |
| SD      | 397.3        | 293.7    | 545.0    | 135.4         | 224.6    | 287.5    | 288.0      |
| SE      | 162.2        | 119.9    | 222.5    | 55.3          | 91.7     | 117.4    | 117.6      |

**NTx (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -65.6    |
| FRE     | 60.4                          | -29.8    | -17.0    |
| HUN     | 77.7                          | -64.1    | 89.3     |
| JAG     | -58.3                         | -42.1    | 75.6     |
| RAY     | 18.7                          | 70.6     | 37.1     |
| RUI     | 40.9                          | -13.1    | -53.2    |
| SCH     | 124.0                         | 48.2     |          |
| Mean    | 43.9                          | -5.0     | 11.0     |
| SD      | 61.5                          | 53.1     | 66.0     |
| SE      | 25.1                          | 21.7     | 26.9     |



**NTx/Creatinine (nmol/nmol)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 29.6     |               |          | 29.3     | 28.5       |
| FRE     | 22.9         | 24.0     | 28.4     | 33.5          | 23.4     | 28.4     | 33.3       |
| HUN     | 27.2         | 35.9     | 23.9     | 26.5          | 21.3     | 26.9     | 28.6       |
| JAG     | 35.9         | 41.1     | 64.7     | 38.4          | 68.4     | 41.0     | 37.7       |
| RAY     | 26.6         | 19.8     | 20.4     | 27.6          | 37.1     | 26.8     | 28.7       |
| RUI     | 33.5         | 46.0     | 46.8     | 31.3          | 35.8     | 31.6     | 31.5       |
| SCH     | 23.9         | 26.9     |          | 27.7          | 22.3     |          |            |
| Mean    | 28.3         | 32.3     | 35.6     | 30.8          | 34.7     | 30.7     | 31.4       |
| SD      | 5.2          | 10.3     | 16.9     | 4.6           | 17.9     | 5.4      | 3.7        |
| SE      | 2.1          | 4.2      | 6.9      | 1.9           | 7.3      | 2.2      | 1.5        |

**NTx/Creatinine (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -1.0     |
| FRE     | 46.3                          | -2.5     | 0.0      |
| HUN     | -2.6                          | -40.7    | 12.6     |
| JAG     | 7.0                           | 66.4     | -36.6    |
| RAY     | 3.8                           | 87.4     | 31.4     |
| RUI     | -6.6                          | -22.2    | -32.5    |
| SCH     | 15.9                          | -17.1    |          |
| Mean    | 10.6                          | 11.9     | -4.4     |
| SD      | 19.1                          | 52.2     | 26.2     |
| SE      | 7.8                           | 21.3     | 10.7     |

**Pyridinium Crosslinks (nmol • 24 hr<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 716.5    |               |          | 222.3    | 186.4      |
| FRE     | 185.0        | 236.6    | 205.9    | 247.5         | 157.8    | 116.8    | 99.0       |
| HUN     | 167.1        | 331.0    | 164.0    | 214.6         | 148.5    | 310.5    | 653.9      |
| JAG     | 496.8        | 286.0    | 172.0    | 194.0         | 123.6    | 361.7    | 207.7      |
| RAY     | 140.2        | 209.0    | 206.7    | 252.4         | 294.1    | 239.1    | 211.5      |
| RUI     | 222.5        | 277.8    | 405.2    | 317.8         | 295.9    | 314.9    | 155.7      |
| SCH     | 98.2         | 86.0     |          | 166.1         | 195.7    |          |            |
| Mean    | 218.3        | 237.7    | 311.7    | 232.1         | 202.6    | 260.9    | 252.4      |
| SD      | 142.7        | 85.4     | 217.3    | 53.1          | 75.2     | 87.5     | 201.1      |
| SE      | 58.3         | 34.9     | 88.7     | 21.7          | 30.7     | 35.7     | 82.1       |

**Pyridinium Crosslinks (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -69.0    |
| FRE     | 33.8                          | -33.3    | -43.3    |
| HUN     | 28.4                          | -55.1    | 89.3     |
| JAG     | -61.0                         | -56.8    | 110.3    |
| RAY     | 80.0                          | 40.7     | 15.7     |
| RUI     | 42.8                          | 6.5      | -22.3    |
| SCH     | 69.1                          | 127.6    |          |
| Mean    | 32.2                          | 4.9      | 13.5     |
| SD      | 49.9                          | 71.0     | 72.7     |
| SE      | 20.4                          | 29.0     | 29.7     |

**Pyridinium Crosslinks/Creatinine (nmol/nmol)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 12.9     |               |          | 11.5     | 18.0       |
| FRE     | 21.2         | 16.5     | 19.5     | 25.9          | 15.2     | 13.4     | 14.5       |
| HUN     | 19.8         | 18.1     | 12.2     | 13.9          | 13.4     | 13.7     | 18.7       |
| JAG     | 14.7         | 14.6     | 17.4     | 14.7          | 18.2     | 13.2     | 11.7       |
| RAY     | 7.5          | 12.5     | 11.2     | 11.8          | 19.3     | 12.4     | 12.7       |
| RUI     | 16.2         | 13.8     | 16.7     | 15.4          | 13.2     | 18.7     | 13.7       |
| SCH     | 15.7         | 11.8     |          | 13.8          | 14.6     |          |            |
| Mean    | 15.9         | 14.6     | 15.0     | 15.9          | 15.7     | 13.8     | 14.9       |
| SD      | 4.8          | 2.4      | 3.3      | 5.0           | 2.5      | 2.5      | 2.9        |
| SE      | 2.0          | 1.0      | 1.4      | 2.1           | 1.0      | 1.0      | 1.2        |

**Pyridinium Crosslinks/Creatinine (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -10.9    |
| FRE     | 22.2                          | -7.9     | -31.3    |
| HUN     | -29.8                         | -26.0    | 12.3     |
| JAG     | 0.0                           | 24.7     | -24.1    |
| RAY     | 57.3                          | 54.4     | 10.7     |
| RUI     | -4.9                          | -4.3     | 12.0     |
| SCH     | -12.1                         | 23.7     |          |
| Mean    | 5.4                           | 10.8     | -5.2     |
| SD      | 30.5                          | 28.9     | 19.6     |
| SE      | 12.5                          | 11.8     | 8.0      |

**Hydroxyproline ( mol • 24 hr<sup>-1</sup>)**

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 257.0    |               |          | 74.3     | 40.8       |
| FRE     | 15.8         | 55.2     | 54.7     | 44.4          | 52.3     | 44.8     | 40.3       |
| HUN     | 47.0         | 80.1     | 40.2     | 51.6          | 50.9     | 105.8    | 141.9      |
| JAG     | 297.2        | 146.9    | 68.3     | 99.7          | 48.6     | 137.2    | 127.2      |
| RAY     | 96.8         | 55.2     | 54.5     | 122.8         | 94.2     | 74.0     | 80.0       |
| RUI     | 83.8         | 96.0     | 98.3     | 73.2          | 96.4     | 88.3     | 89.3       |
| SCH     | 18.3         | 54.1     |          | 69.7          | 58.4     |          |            |
| Mean    | 93.2         | 81.3     | 95.5     | 76.9          | 66.8     | 87.4     | 86.6       |
| SD      | 105.3        | 36.4     | 81.5     | 29.6          | 22.3     | 31.6     | 42.4       |
| SE      | 43.0         | 14.9     | 33.3     | 12.1          | 9.1      | 12.9     | 17.3       |

**Hydroxyproline (% change)**

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -71.1    |
| FRE     | 181.0                         | -5.3     | -18.1    |
| HUN     | 9.8                           | -36.5    | 163.2    |
| JAG     | -66.5                         | -66.9    | 100.9    |
| RAY     | 26.9                          | 70.7     | 35.8     |
| RUI     | -12.6                         | 0.4      | -10.2    |
| SCH     | 280.9                         | 7.9      |          |
| Mean    | 69.9                          | -4.9     | 33.4     |
| SD      | 132.5                         | 46.4     | 85.9     |
| SE      | 54.1                          | 18.9     | 35.1     |

### Hydroxyproline/Creatinine ( mol • g<sup>-1</sup>)

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 41.0     |               |          | 34.0     | 34.8       |
| FRE     | 16.0         | 34.0     | 46.0     | 41.1          | 44.6     | 45.5     | 52.2       |
| HUN     | 49.2         | 38.8     | 26.4     | 29.6          | 40.7     | 41.2     | 35.9       |
| JAG     | 77.7         | 66.4     | 61.0     | 66.9          | 63.1     | 44.2     | 63.5       |
| RAY     | 45.9         | 29.2     | 26.2     | 51.0          | 54.6     | 34.0     | 42.2       |
| RUI     | 54.1         | 42.2     | 35.9     | 31.3          | 38.0     | 46.4     | 69.3       |
| SCH     | 26.0         | 65.8     |          | 51.2          | 38.5     |          |            |
| Mean    | 44.8         | 46.1     | 39.4     | 45.2          | 46.6     | 40.9     | 49.7       |
| SD      | 21.8         | 16.1     | 13.2     | 14.1          | 10.1     | 5.6      | 14.5       |
| SE      | 8.9          | 6.6      | 5.4      | 5.8           | 4.1      | 2.3      | 5.9        |

### Hydroxyproline/Creatinine (% change)

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -17.1    |
| FRE     | 156.9                         | 31.2     | -1.1     |
| HUN     | -39.8                         | 4.9      | 56.1     |
| JAG     | -13.9                         | -5.0     | -27.5    |
| RAY     | 11.1                          | 87.0     | 29.8     |
| RUI     | -42.1                         | -10.0    | 29.2     |
| SCH     | 96.9                          | -41.5    |          |
| Mean    | 28.2                          | 11.1     | 11.6     |
| SD      | 81.1                          | 44.0     | 32.0     |
| SE      | 33.1                          | 18.0     | 13.1     |

### Calcium (mg • 24 hr<sup>-1</sup>)

| Subject | PRE Training |          |          | POST Training |          |          | POST Study |
|---------|--------------|----------|----------|---------------|----------|----------|------------|
|         | passive      | exercise | combined | passive       | exercise | combined |            |
| FLE     |              |          | 301      |               |          | 145      | 105        |
| FRE     | 114          | 656      | 95       | 111           | 144      | 85       | 84         |
| HUN     | 306          | 239      | 172      | 50            | 148      | 270      | 248        |
| JAG     | 389          | 57       | 46       | 242           | 70       | 113      | 233        |
| RAY     | 92           | 98       | 113      | 202           | 195      | 237      | 120        |
| RUI     | 92           | 444      | 462      | 66            | 429      | 575      | 236        |
| SCH     | 79           | 104      |          | 59            | 56       |          |            |
| Mean    | 179          | 266      | 198      | 122           | 174      | 238      | 171        |
| SD      | 134          | 238      | 156      | 81            | 135      | 180      | 76         |
| SE      | 55           | 97       | 64       | 33            | 55       | 74       | 31         |

### Calcium (% change)

| Subject | % Change PRE to POST Training |          |          |
|---------|-------------------------------|----------|----------|
|         | passive                       | exercise | combined |
| FLE     |                               |          | -52      |
| FRE     | -3                            | -78      | -11      |
| HUN     | -84                           | -38      | 57       |
| JAG     | -38                           | 23       | 146      |
| RAY     | 120                           | 99       | 110      |
| RUI     | -28                           | -3       | 24       |
| SCH     | -25                           | -46      |          |
| Mean    | -10                           | -7       | 46       |
| SD      | 69                            | 63       | 74       |
| SE      | 28                            | 26       | 30       |

**MRI Muscle Volume (cm<sup>3</sup>)**

**PRE Exercise Training**

| Subject | R. Femoris |       | V. Lateralis |       | V. Intermedius |       | V. Medialis |       |
|---------|------------|-------|--------------|-------|----------------|-------|-------------|-------|
|         | Left       | Right | Left         | Right | Left           | Right | Left        | Right |
| RUI     | 297        | 294   | 919          | 895   | 542            | 619   | 609         | 646   |
| RAY     | 347        | 358   | 1061         | 993   | 705            | 657   | 608         | 593   |
| FRE     | 242        | 242   | 779          | 718   | 510            | 544   | 511         | 535   |
| HUN     | 333        | 333   | 876          | 876   | 565            | 565   | 585         | 585   |
| JAG     | 375        | 382   | 784          | 733   | 679            | 642   | 540         | 532   |
| FLE     | 229        | 229   | 663          | 663   | 504            | 504   | 438         | 438   |
| Mean    | 304        | 306   | 847          | 813   | 584            | 589   | 549         | 555   |
| SD      | 59         | 62    | 137          | 127   | 87             | 60    | 67          | 71    |
| SE      | 24         | 25    | 56           | 52    | 35             | 25    | 27          | 29    |

**PRE Combined Training**


|      |     |     |     |     |     |     |     |     |
|------|-----|-----|-----|-----|-----|-----|-----|-----|
| RUI  | 312 | 308 | 904 | 881 | 588 | 671 | 651 | 679 |
| RAY  | 273 | 289 | 816 | 778 | 532 | 506 | 463 | 454 |
| FRE  | 267 | 265 | 843 | 788 | 562 | 597 | 562 | 561 |
| HUN  | 334 | 334 | 873 | 873 | 589 | 589 | 583 | 583 |
| JAG  | 359 | 372 | 748 | 683 | 687 | 657 | 534 | 511 |
| FLE  | 230 | 230 | 705 | 705 | 530 | 530 | 429 | 429 |
| Mean | 296 | 300 | 815 | 785 | 581 | 592 | 537 | 536 |
| SD   | 48  | 50  | 76  | 82  | 58  | 66  | 81  | 92  |
| SE   | 19  | 21  | 31  | 34  | 24  | 27  | 33  | 37  |

**POST Combined Training**


|      |     |     |     |     |     |     |     |     |
|------|-----|-----|-----|-----|-----|-----|-----|-----|
| RUI  | 320 | 325 | 954 | 959 | 625 | 701 | 647 | 714 |
| RAY  | 277 | 298 | 798 | 834 | 548 | 540 | 498 | 491 |
| FRE  | 293 | 290 | 887 | 853 | 603 | 602 | 563 | 590 |
| HUN  | 359 | 359 | 887 | 887 | 601 | 601 | 590 | 590 |
| JAG  | 382 | 396 | 807 | 704 | 752 | 684 | 520 | 505 |
| FLE  | 246 | 246 | 710 | 710 | 559 | 559 | 470 | 470 |
| Mean | 313 | 319 | 841 | 825 | 615 | 615 | 548 | 560 |
| SD   | 51  | 53  | 86  | 101 | 73  | 65  | 65  | 91  |
| SE   | 21  | 22  | 35  | 41  | 30  | 27  | 27  | 37  |

## APPENDIX G

Fig. 12. Mean ( $\pm$ SE) supine maximal exercise load for the three Phases.




a) Pre- versus post-training, kg-m. \*  $P < 0.05$  from pre-training.




b) After training, percent change. \*  $P < 0.05$  from zero.

Fig 13. Mean ( $\pm$ SE) supine maximal oxygen uptake for the three Phases.




a) Pre- versus post-training,  $\text{ml} \cdot \text{min}^{-1} \cdot \text{kg}^{-1}$ .




b) After training, percent change.

Fig. 14. Mean ( $\pm$ SE) supine maximal heart rate for the three Phases.




a) Pre- versus post-training, beats • min<sup>-1</sup>.




b) After training, percent change.

Fig. 15. Mean ( $\pm$ SE) supine maximal exercise tolerance for the three Phases.




a) Pre- versus post-training, min. \*  $P < 0.05$  from pre-training.




b) After training, percent change. \*  $P < 0.05$  from zero.



Fig. 16. Mean ( $\pm$ SE) resting pre-tilt heart rate for the three Phases.




a) Pre- versus post- training, beats  $\cdot$  min $^{-1}$ .  $P < 0.05$  from pre-training.




b) After training, percent change. \*  $P < 0.05$  from zero.

Fig. 17. Mean ( $\pm$ SE) resting pre-tilt systolic blood pressure for the three Phases.




a) Pre- versus post-training, mmHg. \*  $P < 0.05$  from pre-training.




b) After training, percent change. \*  $P < 0.05$  from zero.

Fig 18. Mean ( $\pm$ SE) resting pre-tilt diastolic blood pressure for the three Phases.




a) Pre- versus post-training, mmHg.




b) After training, percent change.

Fig. 19. Mean ( $\pm$ SE) resting pre-tilt mean arterial pressure for the three Phases.




a) Pre- versus post-training, mmHg.




b) After training, percent change.

Fig. 20. Mean ( $\pm$ SE) tilt-tolerance time for the three Phases.




a) Pre-versus post-training, min.




b) After training, percent change.

Fig. 21. Mean ( $\pm$ SE) heart rate at tilt-tolerance for the three Phases.




a) Pre- versus post-training, beats  $\cdot$  min $^{-1}$ .




b) After training, percent change.


Fig. 22. Mean ( $\pm$ SE) arterial pressure at tilt-tolerance for the three Phases.




a) Pre- versus post-training, mmHg.



b) After training, percent change.



**Fig 23.** Mean ( $\pm$ SE) cardiac R-R interval at rest and post-tilt, pre- and post-training for the three Phases. \*  $P < 0.05$  from corresponding rest value.



**Fig. 24.** Mean ( $\pm$ SE) stroke volume at rest and post-tilt, pre- and post-training for the three Phases. \*  $P < 0.05$  from corresponding rest value.






Fig. 25. Mean ( $\pm$ SE) end-diastolic volume at rest and post-tilt, pre- and post-training for the three Phases. \*  $P < 0.05$  from corresponding rest value.






Fig. 26. Mean ( $\pm$ SE) cardiac output at rest and post-tilt, pre- and post-training for the three Phases. \*  $P < 0.05$  from corresponding rest value.




**Fig 27.** Mean ( $\pm$ SE) Finapress arterial pressure at rest and post-tilt, pre- and post-training for the three Phases.




**Fig 28.** Mean ( $\pm$ SE) total peripheral resistance at rest and post-tilt, pre- and post-training for the three Phases. \*  $P < 0.05$  from corresponding rest value.

Fig 29. Mean ( $\pm$ SE) cardiovascular variables pre- and post-training for the three Phases.




a) Rest versus tilt-tolerance, percent change.




b) Pre-tilt, percent change.


**Fig 30. Mean ( $\pm$ SE) blood hemoglobin at rest and post-tilt for the three Phases.**



**a) Pre- versus post-training,  $g \cdot dl^{-1}$ . \*  $P < 0.05$  from corresponding rest value.**




**b) After training, percent change.**




**c) Pre- versus post-training, percent change. \*  $P < 0.05$  from zero.**


Fig. 31. Mean ( $\pm$ SE) raw hematocrit at rest and post-tilt for the three Phases.



a) Pre-versus post-training, percent units. \*  $P < 0.05$  from corresponding rest value.




b) After training, percent change.




c) Pre-versus post-training, percent change. \*  $P < 0.05$  from zero.


Fig 32. Mean ( $\pm$ SE) absolute plasma volume at rest and post-tilt for the three Phases.



a) Pre-versus post-training, ml. \*  $P < 0.05$  from corresponding rest value.




b) After training, percent change.




c) Pre-versus post-training, percent change. \*  $P < 0.05$  from zero.


Fig. 33. Mean ( $\pm$ SE) plasma sodium concentration at rest and post-tilt for the three Phases. Normal range is 136 – 145  $\text{mmol} \cdot \text{L}^{-1}$ .



a) Pre-versus post-training,  $\text{mmol} \cdot \text{L}^{-1}$ . #  $P < 0.05$  from comparable pre-passive; 1, 2, 3 are  $P < 0.05$  pairs of different values.




b) After training, percent change. \*  $P < 0.05$  from zero.




c) Pre-versus post-training, percent change.


**Fig. 34. Mean ( $\pm$ SE) plasma potassium concentration at rest and post-tilt for the three Phases. Normal range is  $3.6 - 5.6 \text{ mmol} \cdot \text{L}^{-1}$ .**



**a) Pre- versus post-training,  $\text{mmol} \cdot \text{L}^{-1}$ . 1, 2 are  $P < 0.05$  pairs of different values.**




**b) After training, percent change.**




**c) Pre- versus post-training, percent change.**




Fig. 35. Mean ( $\pm$ SE) plasma osmotic concentration at rest and post-tilt for the three Phases. Normal range is 285 – 295 mOsmol  $\cdot$  kg H<sub>2</sub>O<sup>-1</sup>.



a) Pre- versus post-training, mOsmol  $\cdot$  kg<sup>-1</sup>.




b) After training, percent change.




c) Pre- versus post-training, percent change. \*  $P < 0.05$  from zero.


**Fig. 36. Mean ( $\pm$ SE) plasma albumin concentration at rest and post-tilt for the three Phases. Normal range is  $4.6 - 6.7 \text{ g} \cdot \text{dl}^{-1}$ .**



**a) Pre-versus post-training,  $\text{g} \cdot \text{dl}^{-1}$ . \*  $P < 0.05$  from corresponding rest value. 1 – 8 are  $P < 0.05$  pairs of different values.**




**b) After training, percent change.**




**c) Pre-versus post-training, percent change. \*  $P < 0.05$  from zero.**


**Fig. 37. Mean ( $\pm$ SE) plasma total protein concentration at rest and post-tilt for the three Phases. Normal range is  $6 - 8 \text{ g} \cdot \text{dl}^{-1}$ .**



**a) Pre-versus post-training,  $\text{g} \cdot \text{dl}^{-1}$ . \*  $P < 0.05$  from corresponding rest value. 1, 2, 3 are  $P < 0.05$  pairs of different values.**




**b) After training, percent change.**




**c) Pre-versus post-training, percent change. \*  $P < 0.05$  from zero.**


**Fig. 38. Mean ( $\pm$ SE) plasma renin activity at rest and post-tilt for the three Phases. Normal range (supine) is  $3.2 \pm 1.0 \text{ ngAng1} \cdot \text{ml}^{-1} \cdot \text{hr}^{-1}$ .**



**a) Pre- versus post-training,  $\text{ngAng1} \cdot \text{ml}^{-1} \cdot \text{hr}^{-1}$ . \*  $P < 0.05$  from corresponding rest value. 1, 2 are  $P < 0.05$  pairs of different values.**




**b) After training, percent change.**




**c) Pre- versus post-training, percent change. \*  $P < 0.05$  from zero.**


Fig. 39. Mean ( $\pm$ SE) plasma aldosterone concentration at rest and post-tilt for the Passive and Exercise Phases. Normal range (upright) is 50 – 200  $\text{pg} \cdot \text{ml}^{-1}$ .



a) Pre-versus post-training,  $\text{pg} \cdot \text{ml}^{-1}$ . \*  $P < 0.05$  from corresponding rest value.




b) After training, percent change.




c) Pre-versus post-training, percent change. \*  $P < 0.05$  from zero.


Fig. 40. Mean ( $\pm$ SE) plasma vasopressin concentration at rest and post-tilt for the three Phases. Normal range is 1 – 3  $\text{pg} \cdot \text{mL}^{-1}$ .



a) Pre- versus post-training,  $\text{pg} \cdot \text{mL}^{-1}$ .




b) After training, percent change. \*  $P < 0.05$  from zero.




c) Pre- versus post-training, percent change.


Fig 41. Mean ( $\pm$ SE) plasma epinephrine concentration at rest and post-tilt for the three Phases.



a) Pre- versus post-training,  $\text{pg} \cdot \text{ml}^{-1}$ .




b) After training, percent change.




c) Pre- versus post-training, percent change. \*  $P < 0.05$  from zero.


Fig 42. Mean ( $\pm$ SE) plasma norepinephrine concentration at rest and post-tilt for the three Phases.



a) Pre- versus post-training,  $\text{pg} \cdot \text{ml}^{-1}$ . \*  $P < 0.05$  from corresponding rest value.




b) After training, percent change.




c) Pre- versus post-training, percent change. \*  $P < 0.05$  from zero.




Fig 43. Mean ( $\pm$ SE) plasma dopamine concentration at rest and post-tilt for the three Phases.



a. Pre- versus post-training,  $\text{pg} \cdot \text{ml}^{-1}$ .




b) After training, percent change. \*  $P < 0.05$  from zero.




c) Pre- versus post-training, percent change.

**Fig. 44. Mean ( $\pm$ SE) plasma growth hormone concentration at rest and post-tilt for the three Phases.**



**a) Pre- versus post-training by radioimmunoassay, ng • ml<sup>-1</sup>.**



**b) Pre- versus post-training by biological assay, ngHGH • ml<sup>-1</sup>. \* P < 0.001 from rest.**




Fig 45. Mean ( $\pm$ SE) urinary volume pre- and post-training for the three Phases,  $\text{ml} \cdot 24 \text{ hr}^{-1}$ . Top data are  $\text{ml} \cdot \text{min}^{-1}$ .







Fig 46. Mean ( $\pm$ SE) urinary creatinine excretion pre- and post-training for the three Phases,  $\text{mg} \cdot 24 \text{ hr}^{-1}$ .




**Fig 47.** Mean ( $\pm$ SE) urinary deoxyypyridinoline excretion pre- and post-training for the three Phases, nmol • 24 hr<sup>-1</sup>.




**Fig 48.** Mean ( $\pm$ SE) urinary deoxyypyridinoline/creatinine ratio pre- and post-training for the three Phases nmol/nmol.




**Fig 49.** Mean ( $\pm$ SE) urinary n-telopeptide pre- and post-training for the three Phases, nmol • 24 hr<sup>-1</sup>.




**Fig 50.** Mean ( $\pm$ SE) urinary n-telopeptide/creatinine ratio pre- and post-training for the three Phases, nmol/nmol.




**Fig 51.** Mean ( $\pm$ SE) urinary pyridinium crosslinks pre- and post-training for the three Phases,  $\text{nmol} \cdot 24 \text{ hr}^{-1}$ .




**Fig 52.** Mean ( $\pm$ SE) urinary pyridinium crosslinks/creatinine ratio pre- and post-training for the three Phases,  $\text{nmol/nmol}$ .



**Fig 53.** Mean ( $\pm$ SE) urinary hydroxyproline pre- and post-training for the three Phases,  $\mu\text{mol} \cdot \text{g}^{-1}$ .



**Fig. 54.** Mean ( $\pm$ SE) urinary hydroxyproline/creatinine ratio pre- and post-training for the three Phases,  $\mu\text{mol}/\mu\text{mol}$ .



**Fig 55.** Mean ( $\pm$ SE) urinary calcium excretion pre- and post-training for the three Phases,  $\text{mg} \cdot 24 \text{ hr}^{-1}$ .






Fig 56. Mean ( $\pm$ SE) left and right individual quadriceps muscle volumes (T1) for control (pre-training) and pre- and post-Combined training for the Combined Phase,  $\text{cm}^3$ . \*  $P < 0.05$  from pre-Combined training.




Fig 57. Regression of change in multiecho spin image (T2) on knee work capacity during pre-Combined testing, number of knee bends.

# REPORT DOCUMENTATION PAGE

*Form Approved*  
*OMB No. 0704-0188*

Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.

|                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                |                                                                     |                                                                                  |                                   |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------|----------------------------------------------------------------------------------|-----------------------------------|
| <b>1. AGENCY USE ONLY (Leave blank)</b>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                        | <b>2. REPORT DATE</b><br>Month 1998                                 | <b>3. REPORT TYPE AND DATES COVERED</b><br>Technical Memorandum                  |                                   |
| <b>4. TITLE AND SUBTITLE</b><br>Effect of Exercise Training and +Gz Acceleration Training on Men                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |                                                                     | <b>5. FUNDING NUMBERS</b><br><br>UPN 111-10-20                                   |                                   |
| <b>6. AUTHOR(S)</b><br>John E. Greenleaf, Shawn R. Siminson, Jodie M. Stocks, Joyce Evans, Charles F. Knapp, Stephenie A. Cowell, Kendra N. Bailey-Pemberton, Heather W. Wilson, Jamie M. Vener, Simon N. Evetts, Peter A. Hardy, Richard E. Grindland, Helmut Hinghofer-Szalkay, Scott M. Smith, Michael G. Ziegler, David R. Brown, David G. Evans, Fritz B. Moore, and David T. Quach                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                       |                                                                     |                                                                                  |                                   |
| <b>7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)</b><br><br>Ames Research Center<br>Moffett Field, CA 94035-1000                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                          |                                                                     | <b>8. PERFORMING ORGANIZATION REPORT NUMBER</b><br><br>A-00V0046                 |                                   |
| <b>9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)</b><br><br>National Aeronautics and Space Administration<br>Washington, DC 20546-0001                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |                                                                     | <b>10. SPONSORING/MONITORING AGENCY REPORT NUMBER</b><br><br>NASA/TM-2001-210926 |                                   |
| <b>11. SUPPLEMENTARY NOTES</b><br>Point of Contact: John Greenleaf, Ames Research Center, MS 221A-2, Moffett Field, CA 94035-1000<br>(650) 604-6604                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |                                                                     |                                                                                  |                                   |
| <b>12a. DISTRIBUTION/AVAILABILITY STATEMENT</b><br><br>Unclassified — Unlimited<br>Subject Category 52                      Distribution: Standard<br>Availability: NASA CASI (301) 621-0390                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |                                                                     | <b>12b. DISTRIBUTION CODE</b>                                                    |                                   |
| <b>13. ABSTRACT (Maximum 200 words)</b><br><br>Countermeasures for reduction in work capacity (maximal oxygen uptake and strength) during spaceflight and enhanced orthostatic intolerance during re-entry, landing and egress from the return vehicle are continuing problems. The purpose for this study was to test the hypothesis that passive-acceleration training; supine, interval, exercise plus acceleration training; and exercise combined with acceleration training would improve orthostatic tolerance in ambulatory men; and that addition of the aerobic exercise conditioning would not alter this improved tolerance from that of passive-acceleration training. Seven men (24-38 yr) underwent "Passive" training on the Ames human-powered centrifuge (HPC) for 30 min; "Exercise" training on the cycle ergometer with constant +Gz acceleration; and "Combined" exercise training at 40% to 90% of the HPC +Gz <sub>max</sub> exercise level. Maximal supine exercise loads increased significant (P<0.05) by 8.3% (Passive), 12.6% (Exercise), and by 15.4% (Combined) after training, but their post-training maximal oxygen uptakes and maximal heart rates were unchanged. Maximal time to fatigue (endurance) was unchanged with Passive but was increased (P<0.05) with Exercise and Combined training. Thus, the exercise in the Exercise and Combined training Phases resulted in greater maximal loads and endurance without effect on maximal oxygen uptake or heart rate. There was a 4% to 6% increase (P<0.05) in all four quadriceps muscle volumes (right and left) after post-Combined training. Resting pre-tilt heart rate was elevated by 12.9% (P<0.05) only after Passive training suggesting that the exercise training attenuated the HR response. Plasma volume (% Δ) was uniformly decreased by 8% to 14% (P<0.05) at tilt-tolerance pre- vs. post-training indicating essentially no effect of training on the level of hypovolemia. Post-training tilt-tolerance time and heart rate were increased (P<0.05) only with Passive training by 37.8% and by 29.1%, respectively. Thus, addition of exercise training appeared to attenuate the increased Passive tilt-tolerance. |                                                                     |                                                                                  |                                   |
| <b>14. SUBJECT TERMS</b><br><br>Acceleration, Exercise, Training, Deconditioning                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               |                                                                     | <b>15. NUMBER OF PAGES</b><br><br>138                                            | <b>16. PRICE CODE</b>             |
| <b>17. SECURITY CLASSIFICATION OF REPORT</b><br><br>Unclassified                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                               | <b>18. SECURITY CLASSIFICATION OF THIS PAGE</b><br><br>Unclassified | <b>19. SECURITY CLASSIFICATION OF ABSTRACT</b>                                   | <b>20. LIMITATION OF ABSTRACT</b> |