



# Space Station Views of African Sedimentary Basins — Analogs for Subsurface Patterns



**M. JUSTIN WILKINSON**  
Principal Geoscientist  
Jacobs Engineering  
NASA-Johnson Space Center  
Houston Texas



*4<sup>th</sup> Annual African Petroleum  
Forum, Mayfair, 2007*



- Inland deltas — *The Megafan Project* —
  - examples from astronaut/cosmonaut training
- Prediction
- Significance — Examples of hydrocarbon-charged megafans
- Exploration applications —
  - focus points – apexes
  - stream habit
  - shape
  - nesting patterns — on different continents
  - stratigraphic traps
- Coastal megafans —





**Earth Observations Station Message for 25-AUG-2006:GMT Day 237**  
 Message Generated 24-AUG-2006

Due to the planned launch of STS115 on August 27 (GMT 238), CEO will generate a target message tomorrow (GMT 237) for Saturday August 26 (GMT 238). CEO will resume generation of daily target messages on September 11 (GMT 254) assuming the current STS115 timeline does not change. CEO personnel will assist with image analysis operations during the STS115 mission.

While in X-POP altitude, flight rule constraints will be in effect for use of the Science Window. It is only available for use for ~1/4 of each orbit when it is in trail (facing into the RAM). This reduces the number of near-vertical targets that can be acquired with the Science Window. Targets are divided into (i) those that can be acquired with other windows, and (ii) those that cannot. These flight rules do not apply to

**25-AUG-2006:GMT Day 237**

**GMT Site**  
 06:25:41 Mud flow, Indonesia  
 Dynamic Event. Thousands of people on the eastern portion of Java (near Surabaya) have been forced from their homes by an ongoing flow of hot mud and gas that originates from a breach in an undersea volcanic pipeline project. The mud now flows south or approximately 20 kilometers. Look to the left or right of track (Fig. 1) for a widespread brown to dark brown coating on the ground surface and in stream channels.



Figure 1. Satellite data composite of eastern Java, Indonesia. Dashed circle indicates general target area. Red arrow indicates your approximate orbit track.

**GMT Site**  
 14:15:50 Oasis Impact Crater, Libya  
 Look to the right of track for this 18 km diameter impact crater. While the crater walls are not sharply defined, the circular impact structure is readily visible (Fig. 2).





Figure 2. Landsat image of the Oasis impact structure. Red arrow indicates your approximate orbit track.

**GMT Site**  
 17:02:27 Pliocene, Argentina  
 Look to either the left or right of track to map the channel courses of this river. The sunlight point will be to the left of track, and may be helpful in highlighting the river channel as it flows out from the Andes.

**GMT Site**  
 18:47:00 Tropical Storm Debby, Atlantic Ocean  
 Dynamic Event. This tropical storm is predicted to attain hurricane strength over the next two days. Look to the right of track for cloud structure and developing outflow bands.






*Modern Muglad basin — ~ 200 handheld images from several flights —*

- *revealed 11 nested megafans (excl. Sudd)*
- *> 57,000 km<sup>2</sup> of continuous megafan surface mapped (LANDSAT map)*





*Garonne-Arros compound  
megafan, southern France –  
Pyrenees STS51B-31-82*

*Khawr an Nil megafan  
central Sudan*







*Calanscio megafan, Western Desert, Egypt, Landsat ETM+ image -- with SRTM contours*

# Non-functional megafans —

Cunene megafan  
Etosha Pan



Okavango River megafan



## Global study of Megafans —

- river-made
- very smooth surfaces, of low slope
- fan-shaped, *cone of sediment* (convex contour (elevation lines))
- mean radius 100 – 300 km
- areas from 7000 – 200,000 km<sup>2</sup>
- *different from alluvial fans, floodplains, deltas*

*Kosi R. fan,  
India*




*Okavango R.  
inland delta,  
Botswana*



- “switching” behavior — Kosi River —


- cross entire surface of fan
- average rate ~19 yr between switching events




*Kosi R. avulsions*

## Distribution—

- >100 probable large fans identified worldwide, thus far —
- basin type —
  - foreland basins—56%
  - peri- and intracratonic basins—36%
  - rift basins—6%
  - interorogenic basins—2%
- occur in all climates



mapped from Space Shuttle photographs, other space-based imagery, maps (especially 1:1m ONC charts), various reports ©MJ Wilkinson



*Land's End*

*Edinburgh*

*Andes Mts*


*Megafans of Northern Argentina; Parana megafan, inset*

## Unexpected conclusions —

*Large fans are probably —*

- *A New class of landform feature on the planet (not the freak end point in the alluvial fan continuum)*
- *The Norm in all filling continental basins*

- *Prediction — successful prediction of location is now possible*



## Significance — hydrocarbons associated with megafans

### Paleogeography —


Continental fluvial fan,  
Upper Jurassic rocks,  
Colorado, Utah —

### *Large subsurface fan structure* —

- reconstructed from numerous wells
- hundreds of km in radius

### Oil and gas production in —

- Utah
- Colorado



after Jones et al. 2002



## Significance — hydrocarbons associated with megafans

Paleogeography —


Continental fluvial fan,  
Upper Jurassic rocks,  
Colorado, Utah —

*Large subsurface fan  
structure* —

- reconstructed from numerous wells
- hundreds of km in radius

Oil and gas production in —


- Utah
- Colorado



after Jones et al. 2002

# Significance — hydrocarbons associated with megafans

Paleogeography of a fluvial desert landscape —



Southern North Sea  
(Permian) —


- *one large fluvial feature*
- divergent drainage pattern
- on a scale of hundreds of km
- very likely a megafan

George & Berry 1997

- 160 TCF of natural gas

## Significance — hydrocarbons associated with megafans

Paleogeography of a fluvial desert landscape —



Southern North Sea  
(Permian) —

- *one large fluvial feature*
- divergent drainage pattern
- on a scale of hundreds of km
- very likely a megafan

... given the nature of terrigenous sedimentary systems at the basin margin, the potential for stratigraphic traps ... is substantially higher than for sequences such as aeolian sands ...

atural gas



— Moscariello, 2005

# Significance — hydrocarbons associated with megafans

## Paleogeography —

Fluvial continental sediments — mesoscale patterns in Rocky Mts foreland (Upper Cretaceous—Early Tertiary rocks)

- dominated by *several large fans*
- “gas saturated over a wide area”



Masters,  
1984








Camera OTH170

STS-116



01:47:36.149

**Debris Falls Aft (probably RCS paper) of the Right RCS Stinger**

JSC Image Science & Analysis Group



Camera OTH160


STS-116



01:47:36.291


**Debris Noted In Exhaust Plume After Liftoff**

JSC Image Science & Analysis Group










11

1















# Prediction —




*new megafan?*



*SRTM contour data*


# Prediction —

*new megafan?*




## Prediction —

- Reconstructed elevation contours —
- indicate a cone, subsequently twice incised
  - indicate slopes precisely within the range of Earth megafans



visual image




Elevation contours from MOLA data  
(Mars Orbiter Laser Altimeter)



## Prediction —

### Amazonis Planitia

- Flattest plains on Mars (Amazonis Planitia) —
- “outwash plains” made by rivers
- downstream of Marte Valles
- apparent fan radius 880 km





Amazonis Planitia, flattest terrain on Mars, Achp unit, “outwash” plains  
From MOLA data, apparent radius of zone of dispersive flow 885 km  
(after Fuller and Head 2002)



## Prediction —


Large fan on southern flank of  
Apollinaris Volcano —

- may be a lava fan
- or an outflow (fluvial) fan




## Exploration — Channel types —

- b.



## Exploration — Channel types —

- b.
- a.
- s
- s
- s



## Exploration — Channel types —

- *braided, straight, meandering and anastomosed*
- *stacked, braided* channels give best river sandstone body connectivity
- sequence of types differs downfan

### ISS imagery and a possible new river pattern

River patterns accepted in geology are : meandering, straight, braided, anastomosing (insets)


Detailed ISS image mosaic of the Bermejo River (Bermejo megafan, N Argentina), strongly suggests that a fourth pattern needs to be defined :

#### angular pattern

- > straight reaches alternating with tight (low radius) bends
- > floodplain almost without meander scars




straight



braided



meander and meander scars




Science Result uplinked to crew



## Exploration — Channel focus points —


- at
- p



*Exploration — Focus points and fan shape —*





*Okavango fan, Botswana*



*Cunene fan, Angola-Namibia*

# Exploration — shape relates to nesting patterns —



- fan shape
- “space sharing phenomenon” — crowding out alluvial fans
- basics of paleogeography




Exploration —

*nesting patterns* —

Contiguous megafans cover  
> 1.2m km<sup>2</sup> in S America



Exploration —  
*nesting patterns* —




*Logone & Chari megafans, Chad*

## Exploration — nesting patterns —

### Basin-and-swell topography — double- and single-margin basins



- double-margin basins —
    - most fans and fan clusters within rifted lowlands
    - triangle-shaped fans more frequent
  - single-margin basins —
    - fan distribution along basin circumferences
    - large fans at variable altitudes
      - on swell flanks mainly
      - sometimes on basin floors
      - sometimes on swell crests !
- larger sample required*
- diamond-shaped fans more frequent
  - clusters of fans in the T/T pattern



# Exploration —

## Paleogeography —

### Late Triassic Berkiné basin, Algeria —



Turner 2001

# Exploration —


## Paleogeography —

Largest known sandstone on Earth, with major plays —

North African and Arabian Cambro-Ordovician Quartz-rich Sandstones (NAACOQRS), Sahara and Saudi Arabia

- with significant hydrocarbon deposits

- *New theory: NAACOQRS probably composed of a series of multiple megafans*




from Burke et al., 2002



## Exploration —


### *Stratigraphic traps and megafans —*



*stacked megafans  
hundreds of km in radius  
thousands of metres in depth*


# Exploration —

## Coastal megafans —



Gulf of Mexico, looking east  
Astronaut image STS41C-51-241


Nile delta



STS57-73-75

# Coastal megafans —

Save and Limpopo River megafans,  
Mozambique —



*Pande & Temane  
gas fields*



Astronaut image  
STS55-151-20


# Coastal megafans —

Tana River megafan —

*Astronaut handheld images*




*Topography — SRTM mosaic  
(Shuttle Radar Topography Mission)*




# Lak Dera megafan —

*SRTM mosaic  
(Shuttle Radar  
Topography Mission)*

Juba River




*Astronaut  
imagery  
mosaic*



Astronaut images STS61C-46-66, 67, 68, STS41B-33-1360

## Paleogeographic implications —



- megafan rivers *commonly* hundreds of km long
- megafans *probably* extend offshore
- sweep angles need to be considered





Oil slick off southern Somalia —  
sunglint view from Space Shuttle

## Conclusions —



modern megafans —

- *analog*s for the subsurface
- new understanding for *reducing risk* —
  - large modern sample, worldwide
  - we can now predict location of buried, mesoscale continental sedimentary bodies









In memory of the crew of  
Space Shuttle *Columbia*, STS-107...

