

MARINE BOUNDARY LAYER CLOUD PROPERTIES FROM AMF POINT REYES SATELLITE OBSERVATIONS

Michael Jensen¹, Andrew M. Vogelmann¹, Edward Luke¹, Patrick Minnis², Mark A. Miller¹, Mandana Khaiyer³, Louis Nguyen² and Rabindra Palikonda^{2,3}

¹Brookhaven National Laboratory, Upton, New York

²NASA Langley Research Center, Hampton, VA

³Science Systems and Applications, Inc., Hampton, VA

Corresponding author: Mike Jensen, mjensen@bnl.gov, (631) 344-7021

1. OBJECTIVES

- Use satellite observations to place the AMF surface-based and aircraft observations into a larger-scale context relevant to GCM-sized grids (e.g. 300 x 300 km)
- Quantify the macro- and microphysical properties of California region marine boundary layer clouds.
- Quantify the diurnal cycle of MBL cloud properties from satellite observations

2. CLOUD SCREENING



- Identify GCM-sized boxes containing mainly MBL clouds (cloud fraction > 20%)
- Automated cloud identification algorithm screens to remove scenes containing overlying cirrus and other cloud types
- Compute scene-mean cloud macro- and microphysical properties
- For details see: Jensen et al, 2007: Investigation of regional and seasonal variations in MBL cloud properties from MODIS observations, *J. Climate*, (submitted).

3. MACROPHYSICAL PROPERTIES

- Mesoscale cloud structure is quantified using the effective cloud diameter:

$$C_D = \frac{4 \sum_i A_i}{\sum_i P_i}$$

- A_i = Area of a single cloud element, P_i = Perimeter of a single cloud element, N = number of cloud elements





SUMMARY

- C_D offers a simple measure of MBL cloud organization
- The diurnal cycle of cloud physical properties and C_D at Pt Reyes are consistent with previous work.
- The time series of C_D can be used to identify distinct mesoscale organization regimes within the Pt. Reyes observation period

4. CLOUD PROPERTIES

- Large values of C_D for solid cloud decks. Smaller values for scattered cloud scenes
- Large C_D generally accompanied by largest optical depth and liquid water path




5. DIURNAL CYCLE

- Max optical depth, LWP and R_e occur in the morning decreasing through the daytime
- Diurnal cycle of C_D is consistent with more solid clouds in the morning becoming more scattered in the afternoon.

6. TIME SERIES

- July shows 4-6 day cycle from larger C_D (more solid) to smaller C_D (more scattered)
- Large C_D tends to correlate with large optical depth, large liquid water path



Movie of C_D