



Operational and Research Musculoskeletal Summit

August 23 – 25, 2005

Space Center Houston
Club Conference Room

Summit Recommendations

Prepared By:

Richard A. Scheuring, D.O., M.S.
Advanced Projects/Flight Surgeon
University of Texas Medical Branch
NASA-Wyle Laboratories
and Life Science Systems

Marlei Walton, Ph.D., M.S.E.
Discipline Coordinating Scientist, Muscle and Exercise
NASA-Wyle Laboratories
and Life Science Systems

Janis Davis-Street, M.S.
Project Scientist - Exercise Countermeasures Project
Wyle Laboratories
Univ of Houston/NASA - Johnson Space Center

Todd J. Smaka, M.D.
Advanced Projects
NASA-Wyle Laboratories
and Life Science Systems

DeVon Griffin, Ph.D.
Human Health and Performance Office
Exploration Systems Division
NASA Glenn Research Center

Reviewed By:

Jeffrey Young, B.S.
Section Manager
Advanced Projects
Wyle Laboratories
and Life Science Systems

Approved By:

Kathy Johnson-Throop, Ph.D.
Project Manager
Exploration Medical Capability
NASA Johnson Space Center

January 17, 2006

To: NASA Space Medicine

Dear Dr. Duncan,

The Medical Informatics and Health Care Systems group in the Office of Space Medicine at NASA Johnson Space Center (JSC) has been tasked by NASA with improving overall medical care on the International Space Station (ISS) and providing insights for medical care for future exploration missions. To accomplish this task, a three day Operational and Research Musculoskeletal Summit was held on August 23-25th, 2005 at Space Center Houston. The purpose of the summit was to review NASA's a) current strategy for preflight health maintenance and injury screening, b) current treatment methods in-flight, and c) risk mitigation strategy for musculoskeletal injuries or syndromes that could occur or impact the mission. Additionally, summit participants provided a list of research topics NASA should consider to mitigate risks to astronaut health.

Prior to the summit, participants participated in a web-based pre-summit forum to review the NASA Space Medical Conditions List (SMCL) of musculoskeletal conditions that may occur on ISS as well as the resources currently available to treat them. Data from the participants were compiled and integrated with the summit proceedings. A summary of the meeting is attached. Summit participants included experts from the extramural physician and researcher communities, and representatives from NASA Headquarters, the astronaut corps, JSC Medical Operations and Human Adaptations and Countermeasures Offices, Glenn Research Center Human Research Office, and the Astronaut Strength, Conditioning, and Reconditioning (ASCR) group.

The following recommendations are based on a summary of summit discussions and the best possible evidence-based recommendations for musculoskeletal care for astronauts while on the ISS, and include recommendations for exploration class missions.

Musculoskeletal Summit Recommendations

1. General

1.1. Operations

- In the absence of other guidance, NASA should use the ground-based standard of care for both injuries and rehabilitation.
- Distribute ISS crew musculoskeletal data to panel members for assessment (*i.e.* data collection, collation, and de-attribution).

1.2. Research

- Apply results from human clinical data under unloaded conditions to bone healing in microgravity.

2. Back Pain

2.1. Operations

- Maintain current protocols with the addition of in-flight core stabilization program.
- Optimize the sleep position for comfort.
- Better evaluate pain, including timing, prior history, and objective assessment of site of pain.
- Based on history or symptoms, evaluate for prior conditions which may predispose to back pain (e.g. perimeningeal adhesions associated with vertebral fractures, pilots of high performance aircraft) and target for preflight conditioning.
- Interview ISS crew with respect to in-flight back pain (i.e. onset, character, alleviating factors, etc.).
- Consider axial loading (e.g. harness or artificial gravity) to reduce back pain.

2.2. Research

- Perform pre- and post-flight disk imaging (e.g. magnetic resonance imaging, MRI) to examine herniation of cervical and lumbar disks (at a minimum). Postflight image may be dependent on symptoms (should not be used for mission selection).
- For exploration class missions, a whole body MRI can serve as an objective baseline for establishing a body systems schematic. (This should not be used for mission selection.)
- Explore ultrasound to examine disk volume (validate protocols against the gold standard, currently MRI).
- Evaluate countermeasure efficacy in ground-based simulations that unload the back (gravity inversion boots and dry immersion).
- Determine whether there is an increased incidence of back pain or degenerative disk disease in individuals with known history of spine fracture or deformity.

3. Compartment Syndrome

3.1. Operations

- Use ultrasound to determine if fracture or hematoma is present; or increased compartment pressure; may use ultrasound to guide aspiration; splint and possibly wrap; may need to perform fasciotomy.

3.2. Research

- Since treating compartment syndrome during space flight could have significant risks during non-ISS missions, NASA should investigate the clinical use of:
 - hyperbaric treatment
 - ace wrap and splinting to prevent the onset of compartment syndrome
 - alternative minimally invasive treatment strategies.

4. Diagnostic Equipment

4.1. Operations

- Expand and examine ultrasound in the following areas:
 - diagnosis and distinction of specific injuries (e.g. compartment syndrome, fractures),
 - guidance of injections for pain management and/or fluid aspiration.

4.2. Research

- Bone mineral density and indices of bone “quality” of ISS astronauts should be rigorously examined using advanced ground based diagnostics (e.g. quantitative computerized tomography, finite element analysis, and magnetic resonance imaging) to better characterize the effects of space flight on fracture risk.
- Clinical utility of the above advanced ground based diagnostics should be established within 5 years. However, these techniques should be employed now both pre- and postflight since use of ISS is a time-limited opportunity.

5. Exercise/Conditioning

5.1. Operations

- Preflight
 - Better organize upper body, forearm, and hand regimens; avoid overuse injuries with exercise.
 - Protect crewmember exercise time.
 - Train regarding proper ergonomics with consideration of microgravity environment.
 - Add scapular stabilization to exercise program.
- In-flight
 - Optimize exercise prescriptions. (This depends on having functional hardware.)
 - Begin exercise as soon as possible even if full exercise protocols are not able to be implemented. Adding stretching protocols may benefit soft tissue and muscle.
 - Ensure that in-flight core stability exercises are included.
 - ISS exercise hardware reliability, capacity, and durability need to be improved.
 - Follow current terrestrial guidelines for in-flight injury rehabilitation:
 - begin with a limited range of motion without weight bearing,
 - introduce weight bearing exercise using splinting (currently available), aircasting (future inventory items), and Therabands.
- Proprioceptive training
 - Use appropriate exercise devices for improved proprioceptive training.

5.2. Research

- Evaluate preflight Physical Readiness Model data to determine program effectiveness.
- NASA should fund clinical research investigating
 - Intrinsic muscle fatigue associated with upper extremity and strength
 - Muscle recruitment for suit glove usage.
- Optimize ergonomic EVA suit-glove design.

- Build bone mass preflight and maintain in-flight using exercise and pharmacologic therapy.
- Confirm the hypothesis that a full 1 g daily load can be replaced by brief periods of microgravity exercise.
- Assess in-flight exercise equipment requirements for crew exploration vehicle (CEV), lunar surface asset module (LSAM), and lunar habitat.

6. Fracture Management

6.1. Operations

- For a simple fracture, treat the fracture with a splint (one joint above and below the fracture site) and provide analgesics as required. Pain control may require long acting local anesthetic (e.g. Marcaine) for a hematoma block.
- For some fractures, a traction splint may be necessary for acute management and the ground should be consulted.
- For open fractures, the same treatment should be employed as with some closed fractures, plus the appropriate wound care and antibiotics. Surgical procedures should rely on the appropriate (aseptic) wound care available.
- NASA should explore the possibility of training crewmembers in traction splinting to stabilize closed fractures.

6.2. Research

- Since previous data suggest that bones may not heal if they have had any significant microgravity exposure, the following areas should be investigated:
 - electrical, mechanical, and biological enhancement modalities for healing including hyperbaric treatment,
 - animal fracture healing in space and space analogs,
 - human fracture healing in space analog environments.

7. Musculoskeletal Trauma/Overuse

7.1. Operations

- In microgravity, initial compression of a joint should be sufficient to manage fluid accumulation without elevation because microgravity reduces peripheral fluid volume.
- If the injury is acute, short-term immobilization should be employed; splinting that is versatile, durable, and reusable (e.g. SAMS splint) is preferred for longer stabilization.
- NASA should explore the possibility of training crewmembers in steroid and/or long acting local anesthetic (e.g. Marcaine) injections for pain if such treatment is required to continue mission tasks.
- A targeted, specific exercise regimen should be used for upper extremity injuries.
- Crew members should learn modified grasp patterns during EVA to lessen fatigue.

7.2. *Research*

- Analyze muscle recruitment of load transfer in current glove design, and analyze future glove modifications.
- Thoroughly review Apollo missions for exploration data regarding how to effectively mitigate musculoskeletal issues.

8. **Astronaut Candidate Selection Standards**

8.1. *Operations*

- Vertebral fracture assessment (VFA) should be combined with DXA for a baseline measurement.
- Evaluation of Vitamin D status (screening for subclinical levels of 25-hydroxy vitamin D) should be performed as a non-exclusionary standard.
- The panel recommends the following changes to the current musculoskeletal selection standards (section 5.13):
 - No.11 Change sentence to read, “Symptomatic non-union of fractures.”
 - No.18 Under the waiver guideline regarding valgus of the heel, strike “...regardless of the presence or absence of symptoms.”
 - No.19 Change sentence to read, “Leg length discrepancy of more than 3.0 centimeters as determined by standardized radiographic measurement or CT scanogram.”
 - No.20 Criteria listed for osteoporosis and osteopenia should be re-evaluated against the new WHO guidelines, when they are released in 2006.
- The panel recommends the following changes to the current laboratory testing for musculoskeletal selection standards:
 - perform a 24-hour urine calcium measurement,
 - measure 25-hydroxy vitamin D.

9. **Mission Selection Standards**

9.1. *Operations*

- Orthopedic implants may increase the risk of localized osteoporosis or other problems. The effects of microgravity on these implants are unknown. Either the risk of implants should be quantified (see research recommendations below), or limiting crewmembers with implants to Low Earth Orbit (LEO) or lunar missions should be considered (*i.e.* exclude Mars).

9.2. *Research*

- Identify space flight analog populations with implants and determine the sequelae of such conditions.

10. **Musculoskeletal Loss**

10.1. *Operations*

- Measure real time bone changes.
- Use musculoskeletal force (isometric, concentric, eccentric, *etc.*) to drive muscle to anabolic state as soon as possible after entering microgravity.

10.2. Research

- NASA should continue and expand research in animal models to better understand musculoskeletal changes with unloading. The likely timescale for transitioning to human application is a minimum of ten years.
- Analyze the rate of individual astronaut bone change (e.g. bone mineral density (BMD), bone structure) pre- and post-flight from already existing pre- and post- mission evaluations.
- Characterize the density of cancellous and cortical bone components of the femoral neck pre- and postflight to quantify changes, with software designed to analyze geometry and size in QCT images.
- NASA should partner with industry for rapid and convenient methods to assess muscle loss markers in-flight, to evaluate response of markers to countermeasures.
- NASA should partner with NIH and other agencies to develop and evaluate:
 - chemically-based markers of musculoskeletal disease and bone loss,
 - other diagnostic indices of bone strength,
 - genetic, molecular and proteomic relationships between bone structure and fracture risk.
- A cooperative agreement should be developed between NASA and the NIH for the funding for bed rest studies.

11. Exercise/Nutrition

11.1. Operations

- Improve (better and more consistent) measurement of biomechanical parameters for on-orbit exercise equipment.
- Explore engineering options (e.g. using tether devices, therabands etc.) for flexibility and core stabilization exercise protocols.
- There is a strong recommendation for:
 - sodium limitation (not to exceed 5 grams daily),
 - increase supplementation of vitamin D3 to 1000 IU,
 - increase minimum fluid intake to 3 liters daily.

11.2. Research

- Establish ground based evaluations of exercise equipment both individually and combined. This is needed to validate effectiveness and optimize exercise prescriptions with respect to frequency, duration, intensity, and contraction mode.
- Evaluate efficacy of nutritional countermeasures as a muscle countermeasure in combination with resistance exercise. Ensure evaluation of other systems (e.g. bone).
- The effects of the following specific nutrients on bone health should be evaluated: calcium, animal protein, sodium, vitamin K, vitamin D, and eicosapentaenoic acid.
- The panel supports animal research to examine the effects of nutrition, exercise, and/or pharmacological countermeasures for musculoskeletal changes with unloading.

12. Pharmacology

12.1. Operations

- If bisphosphonates are administered:
 - fluid intake should never be restricted,
 - timing of intravenous administration before flight should permit monitoring of physiological changes,
 - ensure adequate intakes of calcium and vitamin D3.
- Fluid intake should be increased to a minimum of 3L to prevent dehydration and minimize renal stone risk.
- Consider potassium magnesium citrate as a renal stone countermeasure pending FDA approval.
- There is no concern regarding negative interactions between bisphosphonates and statins.

12.2. Research

- Further investigation of pharmacological countermeasures is needed:
 - anabolic drugs (including potential side effects),
 - bisphosphonates (effectiveness in microgravity and microgravity analogs, with and without exercise),
 - bisphosphonates (safety and interactions with current medications),
 - bisphosphonates (synergistic effects on gastrointestinal absorption),
 - Further evaluation of bisphosphonate impact on bone remodeling is needed in spaceflight analogs. Clinical data suggest that bisphosphonate suppression of bone turnover may lead to increased fracture risk in some individuals.
- Further investigation is needed to investigate the effects of statins on muscle metabolism.

13. Current Space Medicine Condition List (SMCL) Procedures (see Appendix A for an overview of recommendations)

14. ISS Post-flight Questionnaire: Musculoskeletal (see Appendix B for the reviewed and approved ISS musculoskeletal questionnaire)

Appendix A Current Space Medicine Condition List (SMCL) Procedures

Overview of recommendations

Space Medicine Condition List

Select Conditions- Recommendations to Checklists

Checklist	Change recommended	Rationale
4012 Back Pain	yes	-narcotics and Soma only at night for sleep as they may impair performance of duties -Toradol or morphine may be needed -consider heat/cold
4031 Fracture Dislocation	yes	-if open Fx then Abx dressing -pain relief -vital signs not needed -too broad
4039 Pain Relief	yes	-do not use demerol -several NSAIDs (personal choice)
7055 Sprain Strain	yes	-needs clarification
9300 Kidney Stones	yes	-no demerol -use morphine -toradol first choice for pain -IV fluids? 1L per hour x 4 hours, no limit on oral fluids

Recommendations to Inventory

Condition	Change recommended	Rationale
Dislocation (elbow)	yes	(see below)
Epicondylitis	yes	-long acting local anesthetic needed -use steroid injection with novocaine
Fracture (femur)	yes	traction mechanism needed
Fracture (ribs, multiple fractures)	yes	PPV, oxygen, intubation may be indicated
Injury (Acromioclavicular joint)	yes	(see below)
Injury (Back)	yes	cervical collar needed
Injury (Shoulder)	yes	(see below)
Intervertebral disc disorder (including herniation, resistant to treatment)	yes	consider soft back brace
Trauma (Major-limbs)	yes	-resuscitation needed prior to Fx Tx -consider preliminary life support issues
Compartment syndrome	yes	-Dx, Tx needs to be formalized better -Use ultrasound to determine if fracture or hematoma is present; or if there is increased compartment pressure -May use ultrasound to guide aspiration -Splint and possibly wrap -May need to perform fasciotomy
Amputation (finger(s))	no	
Amputation (foot or toes)	no	
Arthritis	no	
Carpal tunnel syndrome	no	
Dislocation (finger(s))	no	
Dislocation (hand, wrist)	no	
Dislocation (lunate)	no	
Dislocation (perilunate)	no	
Dislocation (toes)	no	
Felon	no	
Fracture (ankle/foot, non-displaced)	no	
Fracture (carpal bone(s))	no	
Fracture (clavicle)	no	
Fracture (elbow)	no	
Fracture (extremities)	no	
Fracture (hand and/or fingers, not requiring reduction)	no	
Fracture (hand and/or fingers, requiring reduction)	no	

Fracture (radius, ulna, displaced)	no	
Fracture (radius, ulna, non-displaced)	no	
Fracture (rib, single fracture)	no	
Fracture (tibia/fibula)	no	
Injury (Elbow)	no	
Injury (Extremity)	no	
Injury (Foot)	no	
Injury (Hip)	no	
Injury (Leg)	no	
Injury (Medial collateral ligament)	no	
Injury (Medial meniscus)	no	
Injury (Patellar)	no	
Injury (Spinal cord)	no	
Injury (Tarsal)	no	
Injury (Thumb)	no	
Injury (Wrist/hand)	no	
Internal derangement of knee (acute)	no	
Intervertebral disc disorder (including herniation, responsive to treatment)	no	
Laceration (Tendon rupture)	no	
Mallet finger	no	
Musculoskeletal pain	no	
Sprain (all other areas)	no	
Sprain (ankle, complete ligament rupture)	no	
Sprain (ankle, incomplete ligament rupture)	no	
Sprain (back)	no	
Sprain (finger(s))	no	
Sprain (thumb)	no	
Tendonitis	no	
Tenosynovitis	no	
Torn ligament (knee)	no	
Torn ligament (other than knee)	no	

Recommendations for condition additions and deletions

Condition	Action	Comment
Pelvic fracture	add to SMCL	-focus on fluid resuscitation -hemostasis -towel and duct tape -possible via crushing on EVA
Injury (various sites)	remove from SMCL	seems to be redundant to fracture, dislocation, sprain, and strain
Muscle rupture (non traumatic)	remove from SMCL	traumatic rupture is a higher risk on EVA

Additional musculoskeletal recommendations

Conditions	Comment
Fractures	open fractures will need wound care/debridement with hibiclens or equivalent
Shoulder and elbow injuries	need sling or immobilizer
Medications	need corticosteroid
Inventory	need more than one SAMS splint
Inventory	need ACE wraps
	manifest 6-inch wraps only, then cut to width
CMO assessment guide	add this
Evacuation timeframe	add this

Other recommendations

Conditions	Comment
Medications	would use as low a dose of ethinyl estradiol as possible to achieve desired effect, 0.030 mg may be fine for upper dose and 0.020 mg for lower dose range (if not degraded by radiation). Start a few months pre-flight to make sure cycles stop without breakthrough. Progestin change as appropriate.

Appendix B
ISS Postflight Questionnaire: Musculoskeletal

Section 1 of 3 (5 questions, 10 minutes): Back Discomfort

While back discomfort is common in the terrestrial environment, it is especially common in microgravity. In fact, many people who have never had terrestrial back discomfort will indeed have it in microgravity.

1.1A. During spaceflight prior to this flight, did you ever experience back discomfort?

Yes No If No, go to 1.2A.

1.1B. If yes, describe:

Which mission(s)	
Duration	
Frequency	
Location (lumbar, thoracic, cervical)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
What were you doing when it started?	
What made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	
What do you think caused it?	

1.2A. Prior to this flight, but excluding previous flights, did you ever experience back discomfort?

Yes No If No, go to 1.3A.

1.2B. If yes, describe:

Approximate dates	
Duration	
Frequency	
Location (lumbar, thoracic, cervical)	
Quality (sharp, dull, burning, etc.)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
What were you doing when it first started?	
What made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	
What do you think caused it?	

1.3A. During this flight, did you ever experience back discomfort?

Yes No If No, go to 1.4A.

1.3B. For each instance, describe (use extra sheets if necessary):

Flight Day Start – End	
Duration	
Location (lumbar, thoracic, cervical)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as	

numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
What were you doing when it started?	
What treatment(s) were given (also indicate duration)?	
Were the treatment(s) effective? (if more than one treatment, also indicate which helped most)	
What else made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	
What do you think caused it?	

1.4A. After your flight, did you ever experience back discomfort?

Yes No If No, go to 1.5.

1.4B. If yes, describe:

Approximate dates	
Duration	
Frequency	
Location (lumbar, thoracic, cervical)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
What were you doing when it first started?	
What made it better? (type of	

medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	
What do you think caused it?	

1.5. Regarding back discomfort, please tell us anything else you think may be important:

Section 2 of 3 (6 questions, 10 minutes): Bone and Joint Injuries/Discomfort

2.1A. Have you ever had a bone fracture?

Yes No If No, go to 2.2A.

2.1B. For each instance, describe (use extra sheets if necessary):

Location / name of bone	
Approximate dates (include flight day if during flight)	
Age at time of fracture	
What were the circumstances surrounding the fracture?	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	

2.2A. During spaceflight prior to this flight, did you ever experience joint injury or discomfort?

Yes No If No, go to 2.3A.

2.2B. If yes, describe:

Which mission(s)	
Duration	
Frequency	
Location (include side of body)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
How did you injure the joint? If unknown, what were you doing when it started hurting?	
What made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	

2.3A. Prior to your flight, but excluding previous flights, did you ever experience joint injury or discomfort?

Yes No If No, go to 2.4A.

2.3B. If yes, describe:

Approximate dates (be as specific as possible)	
Duration	
Frequency	
Location (include side of body)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as	

numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
How did you injure the joint? If unknown, what were you doing when it started hurting?	
What made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	

2.4A. During this flight, did you ever experience joint injury or discomfort?

Yes No If No, go to 2.5A.

2.4B. For each instance, describe (use extra sheets if necessary):

Flight Day Start – End	
Duration	
Location (include side of body)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
How did you injure the joint? If unknown, what were you doing when it started hurting?	
What treatment was given	
Was the treatment effective	
What else made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of	

day, etc.)	
What was the final diagnosis?	

2.5A. After your flight, did you ever experience joint injury or discomfort?

Yes No If No, go to 2.6.

2.5B. If yes, describe:

Approximate dates	
Duration	
Frequency	
Location (include side of body)	
Quality (sharp, dull, burning, etc.)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, etc.	
How did you injure the joint? If unknown, what were you doing when it started hurting?	
What made it better? (type of medication, body positions, stretching, massage, rest, etc.)	
What made it worse? (describe activities, body positions, times of day, etc.)	
What was the final diagnosis?	

2.6. Regarding joint injury or discomfort, please tell us anything else you think may be important:

Section 3 of 3 (7 questions, 10 minutes): Muscle Injury/Discomfort (including sprains/strains)

3.1A. During spaceflight prior to this flight, did you ever experience a muscle injury, sprain, or strain, or unusual muscle discomfort?

Yes No If No, go to 3.2A.

3.1B. If yes, describe:

Which mission(s)	
Duration	
Frequency	
Location (include side of body)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
How did you injure the muscle? If unknown, what were you doing when it started hurting?	
What made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	

3.2A. Prior to your flight, but excluding previous flights, did you ever experience a muscle injury, sprain, or strain, or unusual muscle discomfort?

Yes No If No, go to 3.3A.

3.2B. If yes, describe:

Approximate dates	
Duration	
Frequency	
Location (include side of body)	
Quality (sharp, dull, burning, etc.)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, etc.	
How did you injure the muscle? If unknown, what were you doing when it started hurting?	
What made it better? (type of medication, body positions, stretching, massage, rest, etc.)	
What made it worse? (describe activities, body positions, times of day, etc.)	
What was the final diagnosis?	

3.3A. During this flight, did you ever experience a muscle injury, sprain, or strain, or unusual muscle discomfort?

Yes No If No, go to 3.4A.

3.3B. For each instance, describe (use extra sheets if necessary):

Flight Day Start – End	
Duration	
Location (include side of body)	
Quality (sharp, dull, burning, etc.)	
Intensity (0-10 scale, where	

0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
How did you injure the muscle? If unknown, what were you doing when it started hurting?	
What treatment was given	
Was the treatment effective	
What else made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	

3.4A. After your flight, did you ever experience a muscle injury, sprain, or strain, or unusual muscle discomfort?

Yes No If No, go to 3.5A.

3.4B. If yes, describe:

Approximate dates	
Duration	
Frequency	
Location (include side of body)	
Quality (sharp, dull, burning, <i>etc.</i>)	
Intensity (0-10 scale, where 0= no discomfort whatsoever 10= worst discomfort imaginable)	
Related symptoms such as numbness, tingling, electrical sensations in other parts of the body, bruising, swelling, illness, <i>etc.</i>	
How did you injure the muscle? If unknown, what were you doing when it started hurting?	

What made it better? (type of medication, body positions, stretching, massage, rest, <i>etc.</i>)	
What made it worse? (describe activities, body positions, times of day, <i>etc.</i>)	
What was the final diagnosis?	

3.5A. At the time of any of your muscle injuries or discomfort, were you taking a medication for high cholesterol?

Yes No Never had injury or discomfort If No or Never, go to 3.6A.

3.5B. If yes, list the name of the medication, the dose you were taking, and approximately how long you had been taking the medication prior to your injury or discomfort:

3.6. Regarding muscle injury or discomfort, please tell us anything else you think may be important:
