

Retrieval with Infrared Atmospheric Sounding Interferometer and validation during JAIVE_x

Daniel K. Zhou¹, Xu Liu¹, Allen M. Larar¹, William L. Smith^{2,3} Jonathan P. Taylor⁴, Peter Schlüssel⁵, L. Larrabee Strow⁶, and Stephen A. Mango⁷

¹NASA Langley Research Center, Hampton, VA, USA

²Hampton University, Hampton, VA, USA

³University of Wisconsin-Madison, Madison, WI, USA

⁴Met Office, Exeter, Devon, UK

⁵EUMETSAT, Darmstadt, Germany

⁶University of Maryland Baltimore County, Baltimore, MD, USA

⁷NPOESS Integrated Program Office, Silver Spring, MD, USA

Talk Outline

- 1. LaRC IR-only Retrieval Algorithm Introduction**
- 2. Retrieval Demonstration: Global and Regional Cases**
- 3. Validation with Radiosondes and Dropsondes**
- 4. IASI, AIRS, and NAST-I Inter-comparison**
- 5. Summary and Future Work**

LaRC IR Retrieval Algorithm

PART A: REGRESSION RETRIEVAL (Zhou et al., GRL 2005)

Using **an all-seasonal-global training database** to diagnose 0-2 cloud layers from training relative humidity profile:

A single cloud layer is inserted into the input training profile. Approximate lower level cloud using opaque cloud representation.

Use parameterization of balloon and aircraft cloud microphysical data base to specify cloud effective particle diameter and cloud optical depth:

Different cloud microphysical properties are simulated for same training profile using random number generator to specify visible cloud optical depth within a reasonable range. Different habitats can be specified (Hexagonal columns assumed here).

Use LBLRTM/DISORT “lookup table” to specify cloud radiative properties:

Spectral transmittance and reflectance for ice and liquid clouds interpolated from multi-dimensional look-up table based on DISORT multiple scattering calculations.

Compute EOFs and Regressions from clear, cloudy, and mixed radiance data base:

Regress cloud, surface properties & atmospheric profile parameters against radiance EOFs.


PART B: 1-D VAR. PHYSICAL RETRIEVAL (Zhou et al., JAS 2007)

A one-dimensional (1-d) variational solution with the regularization algorithm (i.e., the minimum information method) is chosen for physical retrieval methodology which uses the regression solution as the initial guess.


Cloud optical/microphysical parameters, namely effective particle diameter and visible optical thickness, are further refined with the radiances observed within the 10.4 μm to 12.5 μm window region.

LaRC Algorithm Flowchart

HYBRID RETRIEVAL ALGORITHM FLOWCHART



Radiative Transfer Model (include clouds)



$$R = R_0 F_T \tau_{tc} + R_c \tau_{tc} + R_1 + R_1^\downarrow F_R \tau_{tc}$$

$$R_0 = \varepsilon B_s \tau_{cs} + \int_{\tau_{cs}}^1 B d\tau + (1 - \varepsilon) R_0^\downarrow \tau_{cs}$$

$$R_0^\downarrow = \tau_{cs} (R_1^\downarrow F_T + R_c) + \int_{\tau_{cs}}^1 B d\tau'$$

$$R_c = (1 - F_R - F_T) B(T_c)$$

$$R_1 = \int_{\tau_{tc}}^1 B d\tau$$

$$R_1^\downarrow = \int_{\tau_{tc}}^1 B d\tau'$$

R = upwelling spectral radiance at the top of atmosphere

F_T = cloud transmissive function

F_R = cloud reflective function

R_0 = upwelling emission below the cloud

R_0^\downarrow = downwelling emission below the cloud

R_c = emission from the cloud

R_1 = upwelling emission above the cloud

R_1^\downarrow = downwelling emission above the cloud

ε = surface emissivity

B = Planck function

τ = total transmittance from any given level to an upper boundary such as cloud level or the top of the atmosphere

τ' = the total transmittance from any given level to a lower boundary such as cloud level or the Earth's surface

τ_{cs} = transmittance between the cloud level and the Earth's surface

τ_{tc} = transmittance between the top of the atmosphere and cloud level

EOF Physical Regression Inversion

Statistics are formulated for one class of data which contains all cloud height conditions

and

2 other classes for which the cloud phase has been stratified to liquid and ice.

$$R = R_0 F_T \tau_{tc} + R_c \tau_{tc} + R_1 + R_1^\downarrow F_R \tau_{tc},$$

$$M_{ij} = \frac{1}{S} \sum_{k=1}^S \mathfrak{R}_{ki} \mathfrak{R}_{kj}$$

$$C_i = \sum_{j=1}^{nc} R_j E_{ji}$$

$$A_m = \sum_{i=1}^{n-1} K_{mi} C_i + K_{mn} P_s = \sum_{i=1}^{n-1} K_{mi} \left(\sum_{j=1}^{nc} R_j E_{ji} \right) + K_{mn} P_s$$

$$\Psi_i = \sum_{j=1}^5 \varepsilon_j e_{ji}$$

R = radiance

P_s = surface pressure

S = number of sample profiles

ℜ = radiance deviation from the mean

M = covariance matrix of ℜ

E = eigenvectors of M – EOFs

C = radiance EOF amplitudes

A = {T_s, ψ, T, q, ... Hcld, τcld, De, Pha} parameters

K = regression coefficients

ψ = emissivity EOF amplitudes

ε = emissivity

e = emissivity eigenvectors

F_T = cloud transmissive function {Hcld, τcld, De, Pha}

F_R = cloud reflective function {Hcld, τcld, De, Pha}

Hcld = cloud height

τcld = cloud optical depth

De = cloud particle diameter

Pha = cloud phase (ice or water cloud)

1-D Var. Physical Iterative Retrieval

$$Y = R_0 F_T \tau_{tc} + R_c \tau_{tc} + R_1 + R_1^\downarrow F_R \tau_{tc},$$

$$\delta Y = Y' \delta X$$

$$J(X) = [Y^m - Y(X)]^T E^{-1} [Y^m - Y(X)] + [X - X_0]^T (\gamma I) [X - X_0]$$

$$X_{n+1} = X_n + J''(X_n)^{-1} J'(X_n)$$

$$\delta X_{n+1} = (Y_n'^T E^{-1} Y_n' + \gamma I)^{-1} Y_n'^T E^{-1} (\delta Y_n + Y_n' \delta X_n)$$

$$\delta X_n = X_n - X_0$$

$$\delta Y_n = Y^m - Y(X_n)$$

$$\|Y[X(\gamma)] - Y^m\|^2 = \sigma^2$$

$$\gamma_{n+1} = q_n \gamma_n$$


each iteration by satisfying the following conditions:

$$q_1 = 1.0;$$

$$\text{if } \|Y(X_n) - Y^m\| < \sigma^2, \text{ then } q_n = 1.5;$$

$$\text{if } \|Y(X_n) - Y^m\| > \sigma^2, \text{ then } q_n = 0.5;$$

$$\text{if } \|Y(X_n) - Y^m\| = \sigma^2, \text{ then stop the iteration;}$$



Y = calculated Radiance

$X = \{T_s, T, q, o_3, co, \dots, Hcld, \tau_{cld}, De, Pha\}$

Y^m = observed Radiance

J = "Penalty function"







σ = total noise

E = error covariance matrix

γ = a smoothing factor


n = iteration number

Global Training for LaRC Algorithm




Channel Used in LaRC Retrieval Algorithm

IASI: 5008 channels for regression, 1697 channels for physical retrieval




AIRS: 1526 channels for regression, 943 channels for physical retrieval




Retrieval Parameters from this System

Brightness Temperature or Radiance Spectrum



Geophysical Parameters




Retrievals under clear conditions:

- Surface properties (skin temp and emissivity).
- Atmospheric temperature and moisture profiles.
- Atmospheric CO and O₃ abundances.


Retrievals under cloudy conditions:

- Atmospheric profile through optically thin cirrus clouds and above optically thick clouds.
- Effective cloud parameters (i.e., cloud top pressure, particle size, and optical depth).


IASI Retrieval Demo: Cloud Top Height




IASI vs. GOES-12: Cloud




IASI Retrieval Demo: Cloud Optical Depth



IASI Retrieval Demo: Cloud Particle Size




IASI Retrieval Demo: Surface Skin Temp






IASI Retrieval Demo: Moisture Distribution








Hurricane Dean (2007.8.20)




This image of Hurricane Dean is a combination of observations from NASA and NOAA satellites. The clouds were observed by a NOAA GOES-12 at 20:45 UTC on 20th August 2007. The land surface is a summertime image from the NASA Blue Marble image collection.


Hurricane Dean Observed with IASI



ΔTemp Horizontal Cross Section (K)






RH Horizontal Cross Section (%)




2007.8.20; ~15:14 UTC


Hurricane Dean Observed with AIRS



ΔTemp Horizontal Cross Section (K)



RH Horizontal Cross Section (%)



2007.8.20; ~07:23 UTC


IASI vs. AIRS: Cloud Top Height

Metop Satellite / IASI

Time: ~15:14 UTC

Eye Location: 17.8 N, 82.5 W

Cloud Top Height (km)




Aqua Satellite / AIRS




Time: ~07:23 UTC

Eye Location: 17.7 N, 80.0 W

Cloud Top Height (km)



Joint Airborne IASI Validation Exp.



Location/dates:

Ellington Field (EFD), Houston, TX, 14 Apr – 4 May, 2007.

Aircraft:

NASA WB-57 (NASt-I, NASt-M, S-HIS);

UK FAAM BAe146-301 (ARIES, MARSS, SWS; dropsondes; in-situ cloud phys. & trace species; etc.).

Satellites:

Metop (IASI, AMSU, MHS, AVHRR, HIRS).

A-train (Aqua AIRS, AMSU, HSB, MODIS; Aura TES; CloudSat; and Calipso).

Ground-sites:

DOE ARM CART ground site (radiosondes, lidar, etc.)

Participants:

include NASA, UW, MIT, IPO, NOAA, UKMO, EUMETSAT, ECMWF, ...






Case Study and Validation (2007.04.29)

IASI @ ~15:48 UTC

AIRS @ ~19:30 UTC


→ AIRS Interoperated to IASI FOV




IASI and AIRS inter-comparison at the same geophysical location and same horizontal resolution

IASI Retrieval: Cloud & Surface

Cloud Parameters:




Surface Properties:




IASI Cloud vs. GOES Image (4.29.2007)

IASI Cloud Top Height (km)




15:32 UTC GOES IR




IASI Retrieval: Δ Temp and RH Fields


Δ Temp Horizontal Cross Section (K)



RH Horizontal Cross Section (%)





IASI Retrievals vs. Radiosondes








Note:
 12:00 UTC = 07:00 Local
 15:48 UTC = 10:48 Local

Radiosonde and IASI retrieval comparison and statistical profiles over 20 radiosondes



High-Vertically-Resolved Retrievals





Drop

Drop







Raob

IASI Regression vs. Physical Retrieval

Statistical Regression Retrieval: RH(%)




Physical Retrieval: RH(%)




1. The retrieval improvement based on the EOF statistical regression through physical iterative retrieval is only contributed by IASI measurements as the minimum information methodology used.
2. A high-vertically-resolved atmospheric structure is captured very well by IASI measurements and/or retrievals; not only in the troposphere, but also in the boundary layer.

IASI (15:48 UTC) vs. AIRS (19:30 UTC)


IASI Retrieval




Temp Deviation from the Mean (K)




Relative Humidity (%)




AIRS Retrieval Interpolated to IASI FOV



Temp Deviation from the Mean (K)






Relative Humidity (%)





NAST-I: Connection between IASI and AIRS



Summary and Future Work

- 1. A state-of-the-art IR-only retrieval algorithm has been developed with an all-seasonal-global EOF Physical Regression and followed by 1-D Var. Physical Iterative Retrieval for IASI, AIRS, and NAST-I.**
- 2. The benefits of this retrieval are to produce atmospheric structure with a single FOV horizontal resolution (~15 km for IASI and AIRS), accurate profiles above the cloud (at least) or down to the surface, surface parameters, and/or cloud microphysical parameters.**
- 3. Initial case study and validation indicates that surface, cloud, and atmospheric structure (include TBL) are well captured by IASI and AIRS measurements. Coincident dropsondes during the IASI and AIRS overpasses are used to validate atmospheric conditions, and accurate retrievals are obtained with an expected vertical resolution.**
- 4. JAIVEx has provided the data needed to validated retrieval algorithm and its products which allows us to assess the instrument ability and/or performance.**
- 5. Retrievals with global coverage are under investigation for detailed retrieval assessment. It is greatly desired that these products be used for testing the impact on Atmospheric Data Assimilation and/or Numerical Weather Prediction.**

IASI vs. AIRS: Skin Temp.

Metop Satellite / IASI

Time: ~15:14 UTC


Eye Location: 17.8 N, 82.5 W

Aqua Satellite / AIRS

Time: ~07:23 UTC

Eye Location: 17.7 N, 80.0 W

Surf Temp (K)



Surf Temp (K)

