

Does the Hall Effect Solve the Flux Pileup Saturation Problem?

John C. Dorelli

NASA/GSFC, Code 673

Greenbelt, MD 20771

ABSTRACT: It is well known that magnetic flux pileup can significantly speed up the rate of magnetic reconnection in high Lundquist number resistive MHD, allowing reconnection to proceed at a rate which is insensitive to the plasma resistivity over a wide range of Lundquist number. Hence, pileup is a possible solution to the Sweet-Parker time scale problem. Unfortunately, pileup tends to saturate above a critical value of the Lundquist number, S_c , where the value of S_c depends on initial and boundary conditions, with Sweet-Parker scaling returning above S_c . It has been argued (see Dorelli and Birn [2003] and

Dorelli [2003]) that the Hall effect can allow flux pileup to saturate (when the scale of the current sheet approaches ion inertial scale, d_i) before the reconnection rate begins to stall. However, the resulting saturated reconnection rate, while insensitive to the plasma resistivity, was found to depend strongly on the d_i .

In this presentation, we revisit the problem of magnetic island coalescence (which is a well known example of flux pileup reconnection), addressing the dependence of the maximum coalescence rate on the ratio of d_i in the "large island" limit in which the following inequality is always satisfied: $l_\eta \ll d_i \ll \lambda$, where l_η is the resistive diffusion length and λ is the island wavelength.

Dorelli, J. C., Effects of Hall electric fields on the saturation of forced antiparallel magnetic field merging, *Phys. Plasmas*, 10, 3309-3314, 2003.

Dorelli, J. C., and J. Birn, Whistler-mediated magnetic reconnection in large systems:

Magnetic flux pileup and the formation of thin current sheets, *J. Geophys. Res.*, 108, 1133, doi:10.1029/2001JA009180, 2003.