

WPGM 2010
Taipei, Taiwan
22-25 June 2010

Hyperspectrally-resolved surface emissivity derived under optically thin clouds

Daniel K. Zhou¹, Allen M. Larar¹, Xu Liu¹, William L. Smith^{2,3}, L. Larrabee
Strow⁴, and Ping Yang⁵

¹NASA Langley Research Center, Hampton, VA, USA

²Hampton University, Hampton, VA, USA

³University of Wisconsin-Madison, Madison, WI, USA

⁴University of Maryland Baltimore County, Baltimore, MD, USA

⁵Texas A&M University, College Station, TX, USA

Surface spectral emissivity derived from current and future satellites can and will reveal critical information about the Earth's ecosystem and land surface type properties, which can be utilized as a means of long-term monitoring of global environment and climate change. Hyperspectrally-resolved surface emissivities are derived with an algorithm utilizes a combined fast radiative transfer model (RTM) with a molecular RTM and a cloud RTM accounting for both atmospheric absorption and cloud absorption/scattering. Clouds are automatically detected and cloud microphysical parameters are retrieved; and emissivity is retrieved under clear and optically thin cloud conditions. This technique separates surface emissivity from skin temperature by representing the emissivity spectrum with eigenvectors derived from a laboratory measured emissivity database; in other words, using the constraint as a means for the emissivity to vary smoothly across atmospheric absorption lines. Here we present the emissivity derived under optically thin clouds in comparison with that under clear conditions.