

## An Extended View of Ozone and Chemistry in the Atmosphere of Mars

Ramsey L. Smith<sup>1</sup>, Kelly E. Fast<sup>1</sup>, T. Kostiuk<sup>1</sup>, Franck Lefèvre<sup>2</sup>, Tilak Hewagama<sup>3</sup>, Timothy A. Livengood<sup>4</sup>

<sup>1</sup>*NASA Goddard Space Flight Center, Greenbelt, MD, USA*

<sup>2</sup>*LATMOS, CNRS, Paris, France*

<sup>3</sup>*University of Maryland, College Park, MD, USA,*

<sup>4</sup>*National Center for Earth and Space Science Education, Capitol Heights, MD, USA*

We present an ongoing effort to characterize chemistry in Mars' atmosphere in multiple seasons on timescales longer than spaceflight missions through coordinated efforts by GSFC's HIPWAC spectrometer and *Mars Express* SPICAM, archival measurements, and tests/application of photochemical models. The trace species ozone ( $O_3$ ) is an effective probe of Mars' atmospheric chemistry because it is destroyed by odd-hydrogen species ( $HO_x$ , from water vapor photolysis). Observed ozone is a critical test for specific predictions by 3-D photochemical models (spatial, diurnal, seasonal). Coordinated measurements by HIPWAC and SPICAM quantitatively linked mission data to the 23-year GSFC ozone data record and also revealed unanticipated inter-decadal variability of same-season ozone abundances, a possible indicator of changing cloud activity (heterogeneous sink for  $HO_x$ ). A detailed study of long-term conditions is critical to characterizing the predictability of Mars' seasonal chemical behavior, particularly in light of the implications of and the lack of explanation for reported methane behavior.