Global distribution of Mercury's neutrals from MESSENGER measurements combined with a tomographic method

Menelaos Sarantos^{1,2}, Bill McClintock³, Ron Vervack, Jr.⁴, Rosemary Killen⁵, Aimee Merkel³, James Slavin⁶, and Sean C. Solomon⁷

¹Heliophysics Science Division, NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA

²Goddard Planetary Heliophysics Institute, University of Maryland, Baltimore County, Baltimore, MD 21228, USA

³Laboratory for Atmospheric and Space Physics, University of Colorado, Boulder, CO 80303, USA

⁴Johns Hopkins University Applied Physics Laboratory, Laurel, MD 20723, USA

⁵Solar System Exploration Division, NASA Goddard Space Flight Center,

Greenbelt, MD 20771, USA

⁶Department of Atmospheric, Oceanic and Space Sciences, University of Michigan, Ann Arbor, Michigan, USA

⁷Department of Terrestrial Magnetism, Carnegie Institution of Washington, 5241 Broad Branch Road, N.W., Washington, DC 20015, USA

The MErcury Surface, Space ENvironment, GEochemistry, and Ranging (MESSENGER) spacecraft entered orbit about Mercury on March 18, 2011. Since then, the Ultraviolet and Visible Spectrometer (UVVS) onboard this spacecraft has been observing Mercury's collisionless exosphere. We present measurements by MESSENGER UVVS of the sodium, calcium, and magnesium distributions that were obtained during multiple passes through the tail over a period of one month. Global maps of the exosphere were constructed daily from such measurements using a recently developed tomographic technique. During this period, Mercury moved towards the Sun from being about 0.44 astronomical units (AU) to approximately 0.32 AU from the Sun. Hence, our reconstructions provide information about the three-dimensional structure of the exosphere, the source processes for these species, and their dependence with orbital distance during the entire in-leg of Mercury's orbit.