

Geophysical Research Abstracts  
Vol. 14, EGU2012-**PREVIEW**, 2012  
EGU General Assembly 2012  
© Author(s) 2012



## Using Satellite Remote Sensing and Modelling for Insights into NO<sub>2</sub> Air Pollution and NO<sub>x</sub> Emissions.

L.N. Lamsal (1), R.V. Martin (2), N.A. Krotkov (3), E.J. Bucsela (4), E.A. Celarier (1), A. van Donkelaar (2), and D. Parrish (5)

(1) USRA/NASA Goddard Space Flight Center, Atmospheric Chemistry and Dynamics, United States ([lok.lamsal@nasa.gov](mailto:lok.lamsal@nasa.gov)),

(2) Dalhousie University, Halifax, Canada, (3) NASA Goddard Space Flight Center, United States, (4) SRI International,

Menlo Park, CA, United States, (5) NOAA Earth System Research Laboratory, CO, United States

Nitrogen oxides (NO<sub>x</sub>) are key actors in air quality and climate change. Satellite remote sensing of tropospheric NO<sub>2</sub> has developed rapidly with enhanced spatial and temporal resolution since initial observations in 1995. We have developed an improved algorithm and retrieved tropospheric NO<sub>2</sub> columns from Ozone Monitoring Instrument. Column observations of tropospheric NO<sub>2</sub> from the nadir-viewing satellite sensors contain large contributions from the boundary layer due to strong enhancement of NO<sub>2</sub> in the boundary layer. We infer ground-level NO<sub>2</sub> concentrations from the OMI satellite instrument which demonstrate significant agreement with in-situ surface measurements. We examine how NO<sub>2</sub> columns measured by satellite, ground-level NO<sub>2</sub> derived from satellite, and NO<sub>x</sub> emissions obtained from bottom-up inventories relate to world's urban population. We perform inverse modeling analysis of NO<sub>2</sub> measurements from OMI to estimate "top-down" surface NO<sub>x</sub> emissions, which are used to evaluate and improve "bottom-up" emission inventories. We use NO<sub>2</sub> column observations from OMI and the relationship between NO<sub>2</sub> columns and NO<sub>x</sub> emissions from a GEOS-Chem model simulation to estimate the annual change in bottom-up NO<sub>x</sub> emissions. The emission updates offer an improved estimate of NO<sub>x</sub> that are critical to our understanding of air quality, acid deposition, and climate change.