

1 **DECADAL TRENDS IN ABUNDANCE, SIZE AND CONDITION OF ANTARCTIC TOOTHFISH IN MCMURDO**  
2 **SOUND, ANTARCTICA, 1972-2010**

3 **David G. Ainley<sup>1</sup>, Nadav Nur<sup>2</sup>, Joseph T. Eastman<sup>3</sup>, Grant Ballard<sup>2</sup>, Claire L. Parkinson<sup>4</sup>, Clive**  
4 **W. Evans<sup>5</sup> and Arthur L. DeVries<sup>6</sup>**

5 **<sup>1</sup>H.T. Harvey & Associates, Los Gatos CA 95032**

6 **<sup>2</sup>PRBO Conservation Science, Petaluma CA 94954**

7 **<sup>3</sup>Department of Biomedical Sciences, College of Osteopathic Medicine, Ohio University,**  
8 **Athens, OH 45701**

9 **<sup>4</sup>Cryospheric Sciences Branch/Code 614.1, NASA Goddard Space Flight Center, Greenbelt, MD**  
10 **20771**

11 **<sup>5</sup>School of Biological Sciences, University of Auckland, Auckland, NZ**

12 **<sup>6</sup>Department of Ecology, Ethology and Evolution, University of Illinois, Urbana, IL 61801**

13  
14 **Abstract**

15 We report analyses of a dataset spanning 38 years of near-annual fishing for Antarctic toothfish  
16 *Dissostichus mawsoni*, using a vertical setline through the fast ice of McMurdo Sound,  
17 Antarctica, 1972-2010. This constitutes one of the longest biological time series in the Southern  
18 Ocean, and certainly the longest for any fish. Fish total length, condition and catch per unit  
19 effort (CPUE) were derived from the >5500 fish caught. Contrary to expectation, length-  
20 frequency was dominated by fish in the upper half of the industrial catch. The discrepancy may  
21 be due to biases in the sampling capabilities of vertical (this study) versus benthic (horizontal)

22 fishing gear (industry long lines), related to the fact that only large Antarctic toothfish (>100 cm  
 23 TL) are neutrally buoyant and occur in the water column. Fish length and condition increased  
 24 from the early 1970's to the early 1990s and then decreased, related to sea ice cover, with lags  
 25 of 8 months to 5 years, and may ultimately be related to the fishery (which targets large fish)  
 26 and changes in the Southern Annular Mode through effects on toothfish' main prey, Antarctic  
 27 silverfish *Pleuragramma antarcticum*. CPUE was constant through 2001 and then decreased  
 28 dramatically, likely related to the industrial fishery, which began in 1996 and which  
 29 concentrates effort over the Ross Sea slope, where tagged McMurdo fish have been found. Due  
 30 to limited prey choices and, therefore, close coupling among mesopredators of the Ross Sea,  
 31 Antarctic toothfish included, the fishery may be altering the trophic structure of the Ross Sea.

32 **Keywords** Antarctic toothfish, Antarctic silverfish, climate change, change in fish condition,  
 33 change in fish abundance, Ross Sea, Southern Annular Mode, Southern Ocean

---

34	<b>Introduction</b>	<b>3</b>
35	<b>Methods</b>	<b>8</b>
36	Fish capture	8
37	Data analysis	10
38	<b>Results</b>	<b>19</b>
39	Characterization of the catch	19
40	Fish length	20
41	Fish condition	21
42	Catch per unit effort	23
43	Influence of environmental variables on fish metrics	25
44	<b>Discussion</b>	<b>31</b>
45	Toothfish length-frequency	31
46	Trends in toothfish length and condition	34
47	Change in toothfish prevalence	39
48	Recommendations	43
49	<b>References</b>	<b>45</b>

---

50

51

52 **Introduction**

53 Unlike any other marine ecosystem in the world, the high latitude waters of the Southern  
54 Ocean contain a single lineage of teleost fishes that dominates fish diversity, abundance and  
55 biomass (Eastman 2005). Tectonic, oceanographic and climactic changes over the past few tens  
56 of millions of years eliminated previous fish faunas and provided an opportunity for this lineage  
57 of perciform notothenioids to radiate opportunistically and to fill most benthic and pelagic  
58 niches, despite the lack of a swim bladder. For example, the 2- meter-long Antarctic toothfish  
59 *Dissostichus mawsoni*, and the 15–25 cm-long forage species, Antarctic silverfish *Pleuragramma*  
60 *antarcticum*, are key piscine predator and prey in the ecosystem in spite of the incongruity of  
61 being closely related sister species/genera (Balushkin 2000). As adults Antarctic toothfish are an  
62 important component of the community of mesopredators (also including penguins, seals, and  
63 cetaceans) that inhabit the Ross Sea, Southern Ocean; and most of them, too, depend on  
64 silverfish (Ballard et al. 2011). To date, the Ross Sea is the least anthropogenically altered  
65 stretch of ocean remaining on Earth (Halpern *et al.* 2008) and, thus, the interactions of this  
66 group of mesopredators and their prey are of particular interest as a model for how cold-water,  
67 continental shelf food webs once operated elsewhere.

68 In this system, productivity is highest and trophic interactions most intense during  
69 summer. Most of the mesopredators are present over the Ross Sea (shelf and slope) only  
70 during summer, moving north with daylight and the pack ice during winter; a few —Antarctic  
71 toothfish, emperor penguin *Aptenodytes forsteri* and Weddell seal *Leptonychotes weddellii* —  
72 remain year round. All feed during summer primarily on three prey species over the shelf:

73 Antarctic silverfish and two species of krill: Antarctic krill *Euphausia superba* and crystal krill *E.*  
74 *crystallorophias* (Ballard *et al.* 2011, Smith *et al.* 2011). Over the southern shelf, the silverfish by  
75 its numbers and diet is thought to be the main predator of crystal krill and the toothfish to be  
76 the main predator of silverfish— as measured in McMurdo Sound, 71.2% by frequency of  
77 occurrence and 89.2 % by dry weight (Eastman 1985b, La Mesa *et al.* 2004), consistent with the  
78 generalization that fish are the most important predators in most marine ecosystems (Sheffer  
79 *et al.* 2005). Off the shelf, the diet of these mesopredators is much more diverse (Ainley *et al.*  
80 1984, Fenaughty *et al.* 2003). Due to the reduced number of dietary items over the shelf, a tight  
81 trophic coupling has been observed among them, e.g. prey consumption of cetaceans  
82 negatively affects the foraging area and diet of penguins (Ainley *et al.* 2006), and the foraging of  
83 Weddell seals can negatively affect the prevalence of toothfish (Testa *et al.* 1985), although the  
84 consequences of this fish-seal interaction for local availability of silverfish remain unknown.

85         In this system, too, the apex predators are the Ross Sea killer whale *Orcinus [orca]*;  
86 thought to be a separate but yet-to-be-described species; Morin *et al.* 2010), which has been  
87 observed to feed substantially on toothfish (Ainley *et al.* 2009); and the larger killer whale  
88 “ecotype B” (*O. orca*), which feeds on seals and likely large penguins (Pitman and Ensor 2003,  
89 Ainley and Ballard 2011). Weddell seals also prey to a significant degree on toothfish (Ainley  
90 and Siniff 2009, and references therein; Kim *et al.* 2011), and along the Ross Sea shelf break and  
91 north, toothfish are fed upon by southern elephant seals *Mirounga leonina*, sperm whales  
92 *Physeter macrocephalus* and colossal squid *Mesonychoteuthis hamiltoni* (Yukhov 1970,  
93 Pinkerton *et al.* 2010, Ainley 2010, Smith *et al.* 2011).

94 To understand the trophodynamics of this system on the Ross Sea shelf requires  
95 detailed information on all of its predators and mesopredators. While much is known about the  
96 air-breathing members of the group, all of which have nowhere to hide from researchers, and  
97 indeed have been well researched in the Ross Sea since the mid-1900's, learning about  
98 toothfish life history and abundance has been a challenge. On the one hand, the morphology  
99 and physiology of Antarctic toothfish are among the best known of Southern Ocean species: it  
100 is one of just five neutrally buoyant nototheniids (Eastman and DeVries 1981), attaining this  
101 status at around 100 cm TL (Near *et al.* 2003), and ecologically dominates the Southern Ocean  
102 fish fauna in the sense that it is the major piscine predator (Eastman 1993). It has a suite of  
103 adaptations to allow presence at subzero temperatures including blood antifreeze (DeVries  
104 1988, DeVries and Cheng 2005) and other adaptations (Eastman 1993); it grows rapidly when a  
105 subadult, then its growth rate slows, and it can live to 50 yrs. On the other hand, the ecology  
106 and population dynamics of Antarctic toothfish have remained obscure. From a fishery, the  
107 "legal" portion of which began in 1996-97 (hereafter we will identify austral summers by the  
108 initial year, 1996 in this case), we have learned about the geographic aspects of size-frequency  
109 distribution (Hanchet *et al.* 2008, 2010), confirmed growth rates (cf. DeVries and Eastman 1998,  
110 Horn 2002, Brooks *et al.* 2010) and gained insights into variation in condition (Fenaughty *et al.*  
111 2008), age of recruitment (Parker and Grimes 2010) and diet of individuals in the deepest parts  
112 of the Ross Sea along its continental slope (Fenaughty *et al.* 2003). Otherwise, much remains to  
113 be learned about the Antarctic toothfish, unlike its only congener, the non-Antarctic Patagonian  
114 toothfish (*D. eleginoides*), about which much has been learned (Collins *et al.* 2010). Antarctic  
115 toothfish are thought to spawn during winter; no free eggs or larvae and rarely fish <50 cm

116 have been collected in the Ross Sea region and the spawning frequency, fecundity and aspect  
117 of larval and juvenile life history are unknown. This information gap is the result of severe  
118 restrictions on scientific investigation resulting from this Antarctic species' "preferred habitat":  
119 deep, cold, sea ice covered ocean. Any data from the fishery are limited to just the 3-4 months  
120 of ice-free summer (Dec-March).

121         Toward learning more about this important Southern Ocean mesopredator, we present  
122 results of analysis on data collected from >5500 fish captured, measured, marked and released,  
123 during an extended effort led by A. DeVries almost annually in McMurdo Sound, southern Ross  
124 Sea, 1972-2010. The data set thus constitutes one of the longest biological time series available  
125 for the Southern Ocean, the only long one for a fish species, and one that persisted through a  
126 period that has seen interesting changes in the Ross Sea, in large part driven by climatic forces.  
127 The Southern Annular Mode midway in the period (i.e, about 1986) shifted to mostly positive  
128 leading to a marked increase in winds, sea ice extent and persistence of the coastal polynyas  
129 that strongly affect Ross Sea processes (Parkinson 2002, Zwally *et al.* 2002, Jacobs *et al.* 2002,  
130 Stammerjohn *et al.* 2008, Jacobs 2006, Russell *et al.* 2006). Several of the air breathing  
131 mesopredators are known to have responded through changes mediated by sea ice (Ainley *et*  
132 *al.* 2005). In addition, minke whales *Balaenoptera bonaerensis* were heavily hunted in the late  
133 1970s-early 1980s, affecting penguin competitors, but since have recovered; Weddell seals in  
134 McMurdo Sound, too, were also severely hunted during that period but also appear to have  
135 mostly recovered (Ainley *et al.* 2007, Ainley 2010). In 1996, a toothfish fishery was inaugurated  
136 in the Ross Sea region, including to within ~60 km of McMurdo Sound (opposite side of Ross  
137 Island; FAO/CCAMLR Area 88), and this is now the largest fishery for toothfish south of the

138 Antarctic Polar Front (CCAMLR 2010). Therefore, in our analysis, given the close coupling  
139 observed among air breathing mesopredators, we hypothesized that we should see trends in  
140 toothfish condition, size and abundance that correlate with the above described environmental  
141 and anthropogenic changes. After examining trends over time, we explore the significance of  
142 environmental variables, especially those related to ice cover, to gain insight into the extent to  
143 which these factors could be explaining the observed temporal patterns in toothfish  
144 characteristics.

## 145 **Methods**

### 146 **Fish capture**

147 Scientific fishing occurred primarily from one site, about 4 km west of McMurdo Station, in the  
148 vicinity of 77° 51' S, 166° 40' E (Fig. 1); it was accessible from the Station by a ~10 min over-ice  
149 drive by tracked or wheeled vehicle. Note that most of the study period occurred before the  
150 age of GPS and the ability to precisely identify the location of the fish hut was limited. We know  
151 that the fishing site shifted slightly from year to year, e.g. to deal with pressure ridges, snow  
152 banks etc, and after 1999, to accommodate aircraft approach to the Sea Ice Runway. Both the  
153 runway and then the fish hut were repositioned anew each season, as noted. In 2000, the  
154 runway was shifted more than usual, this time northward, to be closer to and in view of  
155 McMurdo Station, forcing movement of the fishing site more west and south than usual.  
156 Nevertheless, all sites but one were within about a 2 km radius, with depth ranging 415 - 495  
157 m. Moreover, all sites were within the area in which the survey by Testa *et al.* (1985) logged  
158 high catch rates of Antarctic toothfish and the area within which Weddell seals have been

159 frequently seen with toothfish (Ainley and Siniff 2009), even to 2010. Beginning in 2008, the  
160 fishing was based out of Scott Base instead of McMurdo, requiring a longer drive by  
161 researchers.

162           Using a small hydro-winch with 3/32-inch wire and a 25 kg weight attached to the  
163 bottom end to keep the wire under tension (i.e., a vertical set line), a line was deployed through  
164 a hole drilled through the annual fast ice. A heated fish house was placed over the hole. Pulling  
165 the line, depending on number of fish caught, took 2-3 h. Initially 15-22 one-meter leaders with  
166 swiveled, stainless steel, long shank #10 hooks spaced 20-25 m apart were fished (thus  
167 sampling the lower ~300 m of water column, but not waters within 10-20 m of the bottom to  
168 avoid scavenging by benthic amphipods). Quickly it was found that most of the fish were caught  
169 in the lower half of the hook array. Thus leaders were shortened to 30 cm and hooks were  
170 spaced 3-5 m apart, starting 10 m from the bottom (thus the lower ~100m of water column was  
171 sampled but, again, not the bottom). For the first 3 years, live *Pagothenia borchgrevinki* were  
172 used for bait, but it was then found that dead bait worked just as well and so thereafter New  
173 Zealand yellow eyed mullet *Aldrichetta forsteri*, cut in half, was used. Sometimes the interval  
174 between deployments ran 12 h but the interval for the very large majority was 24 h sets. Some  
175 ran 48 h if poor weather prevented access to the fish house, but the fish caught were often  
176 exhausted after such prolonged time on hooks. Once this was realized, this long a soak time  
177 was avoided if possible. When the fishing became based out of Scott Base, this necessitated a  
178 much longer commute to the fish hut, and these sets often soaked for >45 h.



179 Captured fish not used for experiments were placed in a V-trough (with a seawater  
180 soaked cloth over their eyes), measured to the nearest cm; weighed to the nearest pound  
181 (converted to kg); tagged with a numbered "Floy" dart tag behind the 2<sup>nd</sup> ray of the 2nd dorsal  
182 fin and a tail locking tag; in many cases injected with tetracycline; and released. Tetracycline  
183 served as an otolith annuli marker in case the fish was recaptured. This procedure, lasting 3-5  
184 min, was done on the floor of the heated fish hut so that the fish neither warmed nor froze. The  
185 open sea surface was 0.5 m below the level of the floor so that the fish could be gently lifted by  
186 their gill covers and returned to the water without abrading their skin and causing scale loss.

187 Small fish on the line for any length of time appeared to have been prone to attack by  
188 large fish, as those retrieved often had many parallel teeth marks on both sides of their body as  
189 well as many missing scales. This is consistent with reports of small toothfish found in the  
190 stomachs of larger ones taken in the industrial fishery (Petrov and Tartarnikov 2010). Because  
191 standard, stainless steel, long-shank hooks were used (not short-shank with the gap of the tip  
192 being less than that at the lower part of the curve, as in the industrial fishery; CCAMLR 2008),  
193 most of the fish >160 cm were likely lost, given that some hooks were straightened out or  
194 broken. Although Weddell seals cannot entirely be ruled out as taking the bait or hooked fish, it  
195 certainly cannot be other fish species because there are none that are anywhere near the size  
196 of the toothfish in McMurdo Sound. When we retrieved toothfish in the presence of Weddell  
197 seals sharing the fishing hole, the seals paid little attention, apparently not recognizing the  
198 toothfish as prey in that context. [NOTE: We were always sufficiently far from the ice edge that  
199 killer whales would not have access to the fishing holes and we did not fish when the killer  
200 whales were present in the ship's channel.] On the few occasions when a large treble or

201 industrial-type hook was used, more of the larger specimens were caught. These hooks,  
202 however, left large wounds in the jaw likely leading to reduced survival. Therefore, their use  
203 was discontinued.

204

## 205 **Data analysis**

206 Fish metrics

207 *Overall strategy.* Our primary objectives with regard to analysis were two-fold: first to  
208 characterize change in fish metrics (length, condition, and capture rates) over the study period  
209 (1972-2010), and then to identify environmental variables influencing variation in these metrics  
210 among years and across decades. Analyses were conducted on trends in condition and length  
211 of toothfish (for condition,  $n = 5,403$  individuals, and for length,  $n = 5,437$ , among the 5,587  
212 caught among 27 yr of data), using both (1) mean condition index per year and (2) 50<sup>th</sup>, 75<sup>th</sup>,  
213 and 95<sup>th</sup> percentiles of total length (TL) in each year. (3) We confirmed the analysis of annual  
214 values of condition and length by analyzing trends with respect to the condition and length of  
215 individual fish. Finally (4), we assessed number of individuals caught per month and year as part  
216 of a “catch per unit effort” study (see “Catch per unit effort” below). Fish having partial data or  
217 measurements that clearly were wrongly recorded were excluded from the data.

218 We present results of toothfish length and condition in relation to variation among  
219 years, examining both linear and non-linear trends. To allow for non-linearity, we fit two types  
220 of models: (1) polynomials of second-order, and (2) results of “change-point” analyses using  
221 linear splines (Harrell 2001, pp. 18-19). In the latter case, two linear segments (each with its

222 own slope) are joined at a specified “knot”. This allowed us to estimate and test for changes in  
223 (linear) trend. Both the second-order polynomial and the “two-joined-linear-segments” model  
224 have two degrees of freedom and thus are easily compared with respect to fit (e.g., by  
225 minimization of deviance). We also explored other simple models, such as cubic polynomial and  
226 “step-function” models (where there is a disjunct step-wise increase or decrease). The favored  
227 function fit to the data, and presented here, was the one that optimized AIC (Akaike  
228 Information Criterion).

229           Because the toothfish appears to prey primarily on silverfish over the shelf (see  
230 Introduction), especially those of approximately 4-5 years of age (Eastman, pers. obs., see  
231 below), and because La Mesa *et al.* (2010) noted an effect of the amount of open water on the  
232 production and survival of silverfish larvae, we considered environmental variables with lagged  
233 effects of 1 year (specifically, 8 mo) to 5 years. Therefore, we looked at the effect of ice cover  
234 and its persistence previous to the current (ice-covered) fishing season.

235

236           *Condition.* We calculated a Fulton-type condition factor (Anderson and Gutreuter 1983),  
237 scaled to center around 1.0, as an index of weight per unit length, an indirect estimate of fish  
238 girth (Davidson and Marshall 2010) and a proxy for body shape:

$$239 \quad K = (W/TL^3) \times 10^2$$

240 where  $W$  = body weight in g and  $TL$  = total length in cm. Larger values for  $K$  indicate greater  
241 weight per unit length associated with a thicker body. We did not employ  $K$  as an indicator of  
242 body fat content. The resulting index varied from 0.577 to 2.99, with mean =  $1.265 \pm 0.149$  SD.

243 Fenaughty *et al.* (2008) identified an important condition threshold,  $K_{ah}$ , which designates “axe-  
244 handle” fish that are in particularly poor physiological condition,  $K_{ah} = 1.01248$ . Therefore, in  
245 addition to changes in mean condition, we analyzed the proportion of individuals that each year  
246 was below this threshold. The term is derived from the long-thin body with a large head of  
247 emaciated fish. Higher  $K$  is generally associated with better physiological condition in wild fish.

248 Individual condition data came almost entirely (99.9%) from captures in four months:  
249 September, October, November, and December; 28.1% of captures were in October, 50.0%  
250 were in November. Since captures were essentially confined to four months, we have analyzed  
251 monthly variation in condition and length by treating month as a factor with four levels (i.e., as  
252 a categorical variable). Thus, in controlling for the effect of month, we make no assumptions  
253 about how the dependent variables change from month to month. The sample size of captures  
254 per year differed greatly (Table 1).

255 Mean condition index,  $K$ , was calculated for each year, as was the proportion of fish  
256 below the axe-handle threshold. Total length was considered to be both an index of body size  
257 and a proxy of approximate age (Horn 2002, Brooks *et al.* 2010). To assess changes in the  
258 frequency distribution of fish length, with specific attention to fish that were of greatest length  
259 (and thus of greater age), we analyzed three metrics: the 50<sup>th</sup> percentile (i.e., median), 75<sup>th</sup>  
260 percentile (i.e., upper quartile), and 95<sup>th</sup> percentile. The latter was chosen as a more statistically  
261 robust measure of “especially large” individuals than the simple maximum; the 95<sup>th</sup> percentile  
262 reflects a larger sample (in fact, in 20 of 27 yr, at least 5 individuals were  $\geq 95^{\text{th}}$  percentile).

263 Variation in the year-specific percentile values was compared to the percentile values for the  
264 entire study.

265 For analysis of annual change in condition, we weighted the mean value by the inverse  
266 of the standard error of mean condition. Similarly, for annual change in length, we weighted  
267 the 50<sup>th</sup>, 75<sup>th</sup>, and 95<sup>th</sup> percentiles for a given year by the inverse of the standard error of mean  
268 length for that year. Thus the weighting reflected the sample size as well as the individual-level  
269 variability in the metric for that year (e.g., the less variable, the greater the weight).

270 *Catch per unit effort.* To analyze patterns in toothfish catch rate, i.e., catch per unit  
271 effort (CPUE), we analyzed the number of fish caught divided by the most appropriate measure  
272 of capture effort. Among effort metrics evaluated, the metric most closely reflecting the  
273 number of fish caught per year was the total number of hooks set in each year. Comparing  
274 results among years, the number of hook-hours did not correlate with the numbers of fish  
275 caught, once we controlled for the number of hooks set per year. We also analyzed effort as a  
276 separate covariate, where the dependent variable was the number of individuals caught.  
277 Results were similar using either method (dividing number caught by effort or statistically  
278 controlling for effort); we only report the former because it is more widely used.

279 We conducted the abundance analysis in two ways: (1) captures per 10 hooks, natural  
280 log-transformed, and (2) proportional change in abundance from one year to the next. Note  
281 that the proportional change in abundance from year  $t$ ,  $N_t$ , to abundance in year  $t+1$ ,  $N_{t+1}$ , is  
282 equal to the antilog of  $[\ln(N_{t+1})-\ln(N_t)]$ . Hence, we present graphically the results of capture per  
283 10 hooks in terms of ln-transformed capture rates.

284 Effort differed among months, and so did the capture rates with respect to month (see  
285 below; also Testa *et al.* 1985). We therefore analyzed between-year variation in captures per  
286 unit effort while controlling for month of capture, and present results that reflect the statistical  
287 adjustment with respect to month.

288 Furthermore, month of capture could confound observed patterns with respect to  
289 among-year variation in condition and length, given that monthly effort varied among years. To  
290 guard against such possible confounding, we repeated the analyses of condition and length,  
291 using individual observations instead, in relation to between-year patterns and with respect to  
292 the environmental variables. In all cases the pattern observed using annual measures of  
293 condition and length was confirmed as significant by the analyses of individuals, after  
294 controlling for month of capture.

295

296 Environmental variables

297 *Sea Ice.* Using satellite passive-microwave data for the Ross Sea sector from NASA  
298 Nimbus 7 satellite and the Defense Meteorological Satellite Program (e.g., Parkinson 2002,  
299 Zwally *et al.* 2002), ice area and ice extent were calculated daily or every other day for the  
300 period from January 1979 to December 2007, inclusive; no corresponding data were available  
301 prior to November 1978. Both ice variables to some degree measure the amount of open water  
302 over the shelf, one directly (area) and one indirectly (extent). Ice extent in the Ross Sea region  
303 varies with winter/spring wind strength and persistence, as does the extent of the coastal Ross  
304 Sea polynya (explained more fully in Ainley *et al.* 2005, 2010): more wind pushes the ice farther

305 north, but also frees the coastal area of ice. If there is little wind, there is little water among ice  
306 floes and ice area is maximized.

307 Statistical analyses were primarily conducted on average monthly values of ice area and  
308 ice extent. The maximum and minimum values for the 12 monthly averages were determined  
309 for each year. The minimum was always in February, but the maximum could occur in August,  
310 September, October, or November. In addition, we determined daily maximum and minimum  
311 values. However, results were not improved using daily rather than monthly minimum or  
312 maximum values, and hence we show only results using the month-specific values, whether by  
313 specified calendar month, e.g., "September", or with reference to maximum or minimum  
314 monthly value.

315 Analyses were also conducted on duration of the "ice season" or "open water season":  
316 (1) duration (number of days) that ice area exceeded an upper threshold value (3,037,000 km<sup>2</sup>)  
317 relative to the total number of days for which the metric was calculated for that year, and (2)  
318 duration (number of days) that ice area was below a lower threshold value (628,000 km<sup>2</sup>)  
319 relative to the total number of days calculated for that year. The upper threshold was chosen to  
320 be the 95<sup>th</sup> percentile value of ice area in the year with the lowest maximum value (1986). The  
321 lower threshold was chosen to be the 5<sup>th</sup> percentile value of ice area in the year with the  
322 highest minimum value (2003). However, these measures of duration above an upper threshold  
323 or below a lower threshold were no better predictor variables with respect to toothfish metrics  
324 than were the maximum or minimum monthly averages of ice area or ice extent explained  
325 above. Thus, the results with respect to relative ice season duration are not shown.

326 Additional ice metrics analyzed included: timing (day of year) of minimum and  
327 maximum, the difference in ice area or ice extent between minimum and the following  
328 maximum, difference between the minimum and the previous maximum in terms of ice area or  
329 ice extent, the difference in days between the minimum and the maximum, and the rate of  
330 change (difference in ice area or extent between maximum and minimum divided by number of  
331 days between minimum and maximum). Again, none of these variables proved superior in  
332 predicting fish metrics and so results with respect to these variables also are not shown.

333

334 *Other environmental variables.* We examined the importance of several environmental  
335 variables that by influencing ocean characteristics may in turn directly or indirectly affect  
336 toothfish, as they have other Ross Sea mesopredators (Testa *et al.* 1992, Wilson *et al.* 2001,  
337 Ainley *et al.* 2005, Rotella *et al.* 2009). These other environmental variables were the Southern  
338 Oscillation Index (SOI), the Southern Annular Mode (SAM; also called Antarctic Oscillation  
339 Index), and a measure of the size of the Ross Sea Polynya. It is well known that the state of the  
340 sea ice in this part of the Southern Ocean is affected by the atmospheric processes embedded  
341 in SOI and SAM (Jacobs 2006, Stammerjohn *et al.* 2008).

342 We used monthly SOI to calculate six metrics. The first three were: average annual SOI  
343 (January to December), the average for January to June, and the average for July to December.  
344 The other three were quadratic transformations of the corresponding SOI metric, designed to  
345 measure increasing effects of SOI associated with strong El Niño events, during which the SOI is  
346 strongly negative. The transformations were of the form  $(\text{SOI} - c)^2$  where  $c$  = maximum (positive



347 value) of the SOI metric (12-mo average or either of the 6-mo averages). (SOI – c) was thus  
348 always zero or negative and the stronger the El Niño the more negative was the value. Squaring  
349 (SOI – c) resulted in all non-negative values, with strong El Niño being associated with the  
350 largest positive values of the transformed metric and the strongest La Niña in the time series  
351 getting a score of zero (following the approach of Lee *et al.* 2007). In no case was a second-  
352 order polynomial (i.e., with 2 df) superior in terms of AIC to the respective quadratic  
353 transformation of SOI (i.e., with 1 df) when analyzing fish condition or length.

354 For all single-variable regression models, we report the regression coefficient ( $b$ , which  
355 is also commonly symbolized as  $\beta$ ). Where more than one environmental variable was analyzed  
356 in the same model, we report the multiple regression coefficient ( $b$ ) for the effect of a variable  
357 while controlling for all other variables in the model, symbolized. To aid in the comparison  
358 among variables (measured on very different scales, e.g., SOI and maximum ice area), we also  
359 present the standardized multiple regression coefficient (which we refer to as “*beta*”), which  
360 like a correlation coefficient, scales from -1 to +1 (Kutner *et al.* 2005).

361 SAM was represented by the value on 1 January each year (from  
362 <http://www.antarctica.ac.uk/met/gjma/sam.html>; Fig. 2) and is a proxy for strength of the  
363 Southern Ocean westerlies, which in turn affect the formation, extent and persistence of sea ice  
364 (Marshall *et al.* 2003, Russell *et al.* 2006, Stammerjohn *et al.* 2008). Using SAM values from 1  
365 January, about 9 to 11 months before toothfish were captured as part of this study, means that  
366 a lag of -1 was used; lag of 0 means SAM is calculated in the same calendar year as applies to

367 the sample of toothfish; lag of 1 means SAM is calculated between 1-plus years (21 to 23  
368 months) before the toothfish sample was obtained, etc.

369         Among these environmental variables, we examined condition, length, and numbers  
370 caught per unit effort in relation to same-year values, as well as with lags of 1, 2, 3, 4, and 5 yr.  
371 One rationale for examining lags was that condition and numbers of toothfish could well reflect  
372 availability of their prey, i.e., Antarctic silverfish. For example, in McMurdo Sound toothfish  
373 consume silverfish that are 81–195 mm SL with modal values of 100–150 mm (Eastman and  
374 DeVries, unpublished data) and, on the basis of Hubold and Tomo's (1989) aging studies for  
375 Weddell Sea silverfish, these specimens are 4+ years old. Thus, we reasoned ice (or other  
376 environmental) conditions in a given year may influence availability of prey 1 to 5 years later.

377

378

## 379 **Results**

### 380 **Characterization of the catch**

381 Between 1972 and 2010, 5587 toothfish were caught, including 835 kept for experiments  
382 and/or whose survival was in doubt. The largest fish caught was 193 cm (92 kg) and the  
383 smallest was 81 cm (4.5 kg); individuals between 120 and 170 cm predominated, with very few  
384 <90-100 cm (Fig. 3).

385         Of the 4752 fish tagged and released in the study, 17 have been recaptured (0.4%) with  
386 the annual growth rate being 2.0 cm in TL and 1 kg in mass per year (DeVries, unpubl. data).

387 These growth rates are slightly below those reported from analyses of the industrial catch (cf.  
388 Horn 2002, Brooks *et al.* 2010). Most of the recaptures (12) occurred at the McMurdo fishing  
389 site, but one was recaptured by the Russian vessel, *Yantar*, and four have been reported by the  
390 New Zealand fleet (DeVries, unpubl. data). One tagged individual was recaptured >1300 km  
391 north of McMurdo Sound, indicating migration north of the Ross Sea, possibly for spawning  
392 (Hanchet *et al.* 2008, 2010). Most recaptured McMurdo fish were encountered again 4-5 yr  
393 after release, with the longest interval being 18 yr. The recapture rate in the fishery during the  
394 most recent years, 2004-2009, for a select number of boats and not including fish recaptured  
395 the same year, is 4.96% (CCAMLR 2010). The discrepancy between our low recapture rate and  
396 this one could well be due to 1) McMurdo Sound fish being more transient than those on the  
397 main fishing grounds (continental slope) and/or 2) the fact that recaptures of our fish by the  
398 fishery were under-reported, with many vessels other than those from NZ (just 4 of 15-20  
399 permitted) reporting no recaptures at all (CCAMLR 2006). This would especially be true in early  
400 years of fishery, when no industry tagging program was in operation; and of course no data  
401 come from the Illegal, Unreported and Unregistered fleet. One of our tags was found on a  
402 vessel's processing room floor, indicating lack of vigilance.

403

#### 404 **Fish Length**

405 Annual variation in total length, considered an approximate proxy for age, was analyzed with  
406 respect to three metrics: 50<sup>th</sup>, 75<sup>th</sup> (i.e., upper quartile), and 95<sup>th</sup> percentiles. In general, the  
407 overall pattern during the study period was similar for each of these metrics.

408           The most parsimonious model, favored by AIC, describing the pattern of change in  
409 length across years was a two-part spline with change point (i.e., “knot”) at 1992, with  
410 increasing TL to 1992 and decreasing TL subsequently (Table 2, Fig. 4). These models were  
411 superior to a single linear trend, quadratic polynomial, or cubic polynomial model, or other  
412 simple transformation, as determined by AIC, and they accounted well for the pattern  
413 observed, except for 1972 being an outlier with respect to the predicted pattern. In regard to  
414 the 75<sup>th</sup> percentile, for the linear trend 1972-1992,  $b = 0.580 (\pm 0.123 \text{ SE}, t = 4.7, P < 0.0001)$  and  
415 for the linear trend 1992-2010,  $b = -0.526 (\pm 0.242, t = -2.2, P < 0.04)$ . The change in trend was  
416 significant (Table 2). In regard to the 95<sup>th</sup> percentile, for the line segment 1972-1992,  $b = 0.534$   
417  $(\pm 0.169, t = 3.6, P = 0.004)$  and for the line segment 1992-2010,  $b = -0.662 (\pm 0.331, t = -2.0, P =$   
418  $0.057)$ . The change in slope for the 95<sup>th</sup> percentile was also significant (Table 2). Finally, the 50<sup>th</sup>  
419 percentile (not shown) showed the same pattern, and was statistically significant with respect  
420 to analogous measures.

421           In order to assess whether or not the length statistics of the small sample of fish caught  
422 during 1991 and 1992 were mainly responsible for the observed 2-part trends, we also analyzed  
423 trends using individuals rather than years, analyzing with and without data from 1991 and  
424 1992. Models included effect of calendar month (analyzed as categorical variable; Table 3). The  
425 2-part spline model including 1991 and 1992, as expected, was highly significant (Table 3), as  
426 was the model excluding 1991 and 1992 data ( $P < 0.0001$ , Table 3). The coefficient,  $b$ , for the  
427 1972-1992 slope was 0.362 (SE = 0.041), and that for the 1972-1990 slope was 0.351 (SE =  
428 0.041; in both cases,  $P < 0.0001$ ). Similarly, the coefficient,  $b$ , for the 1992-2010 slope was -  
429 0.291 (SE = 0.104) and that for the 1997-2010 slope was -0.272 (SE = 0.104;  $P = 0.005$  and  $P =$

430 009, respectively). Finally, the change in slope from the earlier period (up to 1992) to the later  
431 period (from 1992) was significant whether or not 1991 and 1992 were included (Table 3).

432

### 433 **Fish Condition**

434 Condition ( $K$ ) of the Antarctic toothfish caught varied by month.  $K$  was relatively high in October  
435 and November, but lower in September and December (Fig. 5). Variation among months was  
436 significant ( $F_{3,5394} = 4.09$ ;  $P = 0.007$ ); the pattern and significance of monthly variation in  
437 condition were very similar after controlling for annual variation (year as factor).

438         Across all years, there was no significant linear trend in toothfish condition (Fig. 6).  
439 Instead there was a trend for mean  $K$  to increase from 1972 to 1991 and/or 1992 and decrease  
440 subsequently. Mean  $K$  was higher for 1991 and 1992 than in any other year; nevertheless, an  
441 increasing trend is evident from 1972 to 1990. The pattern evident indicated that change point  
442 analysis, with either 1991 or 1992 being the knot, would be most appropriate. Adjusted  $R^2$  was  
443 maximum and AIC was optimized when the change point was at 1992 rather than at 1991  
444 (Table 2). For the line segment 1972-1992,  $b = 0.00358$  ( $\pm 0.00094$ ,  $t = 3.8$ ,  $P = 0.001$ ), and for  
445 the line segment 1992-2010,  $b = -0.00927$  ( $\pm 0.00180$ ,  $t = -5.2$ ,  $P < 0.001$ ); the difference in  
446 slopes of the two lines was significant (Table 2).

447         AIC for a quadratic model for year was poorer than for the change-point analysis. More  
448 specifically, there is no evidence that the trend from 1972 to 1990 was anything but linear  
449 (linear trend through 1990:  $P < 0.0001$ , quadratic term for that time period,  $P > 0.2$ ) and,

450 therefore, the change-point analysis is preferred, with a change in trend either in 1991 or in  
451 1992 (Figure 6).

452 In order to assess whether or not mean condition of the small sample of fish caught  
453 during 1991 and 1991 were mainly responsible for the observed trends in condition, we also  
454 analyzed trends using individuals rather than mean values, analyzing with and without data  
455 from 1991 and 1992. As with length, models included effect of calendar month (analyzed as  
456 categorical variable; Table 3). The 2-part spline model including 1991 and 1992, as expected,  
457 was highly significant (Table 3), as was the model excluding 1991 and 1992 data (Table 3). The  
458 coefficient,  $b$ , for the 1972-1992 slope was 0.00338 (S.E. = 0.00039), and that for the 1972-1990  
459 slope was 0.00319 (S.E. = 0.00039 ;in both cases,  $P < 0.0001$ ). Similarly, the coefficient,  $b$ , for the  
460 1992-2010 slope was -0.00938 (S.E. = 0.00099 ) and that for the 1997-2010 slope was -0.00910  
461 (S.E. = 0.00099 ; $P < 0.001$  in both cases). Finally, the change in slope from the earlier period (up  
462 to 1992) to the later period (from 1992) was significant whether or not 1991 and 1992 were  
463 included (Table 3).

464

465 *Axe-handle fish.* The proportion of axe-handle fish decreased during the period 1972 to  
466 1992 from about 3 - 7% in the mid-1970's to 0 - 3% in the period 1987 to 1992, and then  
467 increased from 1992 on (Fig. 7). The one exception was 2010, in which there were no axe-  
468 handle fish, but only 7 total fish were caught that year. Among the set of models examined, the  
469 preferred model (as determined by AIC) was a two-part spline, with linear decrease up to 1992  
470 ( $b = -0.00167 \pm 0.00076$ ,  $t = -2.19$ ,  $P = 0.039$ ), and a linear increase since ( $b = 0.00420 \pm 0.00145$ ,

471  $t = 2.90, P = 0.008$ ). The linear trend in each time period was significant (see below) and the  
472 trends differ significantly from each other ( $F_{1,24} = 8.68, P = 0.007$ ).

473

#### 474 **Catch per Unit Effort**

475 Catch rate varied by month within each season (Fig. 5). The lowest capture rate occurred in  
476 September (even lower in August but only two days of fishing were available), but it then  
477 increased dramatically in October. Rates for November and December remained high, though  
478 with a slight downward trend from November to December. There was no significant variation  
479 in catch rates among October, November, December ( $P > 0.4$ ); but each of these differed from  
480 the September catch rate ( $P < 0.001$ ).

481 In regard to annual trends, unlike condition and length, there was no peak in CPUE  
482 midway during the study period. Instead, captures were fairly stable through 2001 (Fig. 8), with  
483 a slight, non-significant decline in capture rate evident late in this period, equivalent to a  
484 decline in CPUE of 1.24% per year (back-transformed geometric mean SE = 1.76%: analysis of In-  
485 transformed monthly capture rates, controlling for month of capture, slope not different from  
486 zero,  $P > 0.4$ ). The pattern then changed dramatically, but unfortunately only 3 years of effort  
487 data were available after 2001. For those 3 years, captures per 10 hooks deployed, summarized  
488 per year, ranged 0 - 0.4 fish, compared to 1975-2001 when the range was 1.0 - 3.4 fish. Though  
489 the information is not included in our analyses, fishing was undertaken by another researcher  
490 (G. Hofmann) in the same localities in 2006 for 4 weeks: no fish were caught but effort data are  
491 not available.

492 Realizing that data were available for only 3 of the 9 years after 2001, and that these 3  
493 years occurred at the end of this latter period, the best-fitting, most parsimonious model  
494 (optimizing AIC) indicated a step change to a different (and much lower) level of capture rate.  
495 This, however, must be viewed with caution with so few years since 2001. The pattern depicted  
496 is preferred by AIC when compared to the full set of competing models: (1) no change, (2) two  
497 linear segments joined at 1992, (3) two linear segments joined at 2001, (4) two linear segments  
498 joined at any year between 1992 and 2001, (5) two parallel lines with a step change occurring  
499 at or after 2001 (i.e., slope is non-zero but the same across all years; only the elevation changes  
500 at or after 2001), (6) inverse transformation of year, (7) year as a quadratic function, and (8)  
501 year as a cubic function. Nevertheless, with only three years of catch per unit effort available  
502 post-2001, it is not possible to adequately estimate the post-2001 trend. Essentially, all that can  
503 be said is that fishing success became dramatically reduced after 2001.

504

### 505 **Influence of environmental variables on fish metrics**

506 ***Fish Length.*** Exploratory analyses of the effects of the full set of candidate ice variables  
507 (see Methods) on our three TL metrics revealed four of particular interest: minimum ice area,  
508 minimum ice extent, maximum ice area, and maximum ice extent. Each of these was analyzed  
509 with respect to lags of 0 - 5 yr, yielding 24 ice variables (= 4 variables x 6 lags). For 95<sup>th</sup>  
510 percentile length, none of the 24 ice variables was significant. For 75<sup>th</sup> percentile length two  
511 variables were significant: maximum ice area with 4-yr lag and maximum ice area with 5-yr lag  
512 (Fig. 9); for 4-yr-lag, weighted regression,  $b = -0.0000084 (\pm 0.0000039, t = -2.15, P = 0.048;$



513 Table 4). We do not show results for 5-yr-lag because this relationship, though significant ( $b =$   
514  $0.0000089 \pm 0.0000041$ ,  $P = 0.049$ ), is driven entirely by results from 1991 and 1992.

515 Comparison of correlation coefficients between 75<sup>th</sup> percentile length and each of the  
516 two ice variables (maximum ice area with 4-yr lag vs 5-yr lag) is instructive, with and without  
517 1991 and 1992. For the 5-yr lag, the correlation coefficient with respect to 75<sup>th</sup> percentile  
518 length was  $r = -0.499$ , including the 2 years, but  $r = -0.050$  without those 2 years. For the 4-yr  
519 lag, the correlation coefficient with respect to 75<sup>th</sup> percentile length was  $r = -0.486$  including the  
520 2 years, but only dropped to  $r = -0.404$ , excluding the two years. Thus, the relationship  
521 between 75<sup>th</sup> percentile length and maximum ice area, with 4-yr-lag, does not hinge on these 2  
522 years having unusually large upper quartile values.

523 Median length (50% percentile) also demonstrated a significant relationship to  
524 maximum ice area 4 years previously (Table 4; Fig. 9), and one that was stronger than that  
525 shown for upper quartile length (for median length: correlation coefficient,  $r = -0.608$ ,  $P < 0.01$ ;  
526 compared to  $r = -0.486$  for upper quartile length and the same ice variable). This relationship is  
527 robust with respect to values for any individual year. Whereas 1991 is an influential  
528 observation, 1992 in this case is not (median length in 1992 = 136, compared to median length  
529 for entire sample = 134). Excluding 1991 still yields a relationship that is close to significant ( $P =$   
530  $0.075$ ).

531

532 ***Fish Condition.*** Two distinct ice variables provided the best statistical predictors of  
533 changes in  $K$ : minimum ice area (always in February) of the current year (8-10 mo prior to

534 fishing season) and maximum ice extent 3 yr previously (Fig. 10). While minimum ice area could  
535 also be a proxy for amount of open water, understanding maximum ice extent is far more  
536 complex, owing to the fact that large scale ice extent is a function largely of wind strength and  
537 persistence in the Ross Sea sector. Stronger winds (greater ice extent) also usually mean that  
538 the coastal polynyas are more prevalent (though not always, depending on temperature), a  
539 subject we shall return to in the Discussion. Regardless, these two variables (minimum ice area,  
540 maximum ice extent 3 yr previous) were relatively uncorrelated with each other ( $r = 0.324$ ).  
541 There were several other variables (such as length of ice season) that provided good predictive  
542 ability for explaining change in condition, though not quite as high, but also highly correlated  
543 with the two primary variables. Thus, as noted in Methods, we keep those results in mind but  
544 do not report them.

545         The relationship between toothfish condition and minimum ice area 8 - 10 mo earlier  
546 was non-linear: mean  $K$  declined with increasing minimum ice area (less open water than  
547 usual), but in an accelerating (down-turned) fashion. In a weighted regression, mean  $K$  in  
548 relation to minimum ice area, the relationship best reflected a quadratic transformation of  
549 minimum ice area (predictor variable = (minimum ice –  $c$ )<sup>2</sup> where  $c$  = smallest minimum ice area  
550 observed in any year = 131,448 in 1979) and was significant:  $b = -2.39 \times 10^{-13}$  ( $\pm 1.07 \times 10^{-13}$ ,  $t = -$   
551 2.23,  $P = 0.039$ ; Table 4).

552         When analyzing the data at the individual level, the effect of minimum ice area  
553 (quadratic transformed) on condition, while controlling for calendar month of capture, was also  
554 significant,  $b = -1.52 \times 10^{-13}$  ( $\pm 4.08 \times 10^{-14}$ ,  $t = -3.73$ ,  $P = 0.0002$ ). Furthermore, even excluding

555 the influential year 2003 (greatest minimum ice area and lowest mean condition index in the  
556 data set), the result of analyzing individual fish condition demonstrated a significant negative  
557 effect of minimum ice area (quadratic transformed),  $b = -1.14 \times 10^{-13}$  ( $\pm 4.24 \times 10^{-14}$ ,  $t = -2.69$ ,  $P =$   
558 0.007). Minimum ice area was a better predictor (as determined by AIC) of mean condition than  
559 was date of minimum or minimum ice extent.

560 The second primary ice variable to account for annual variation in mean condition was  
561 maximum ice extent 3 years earlier (Fig. 10); other lags showed no significant relationship, (0, 1,  
562 2, 4, or 5). Maximum ice extent was a better predictor than was ice extent in any specific  
563 month (e.g., October). In this case the relationship was linear. In a weighted regression, mean  $K$   
564 in relation to maximum ice extent with 3-yr-lag was:  $b = -7.18 \times 10^{-8}$  ( $\pm 3.19 \times 10^{-8}$ ,  $t = -2.23$ ,  $P =$   
565 0.04; Table 3).

566 There was no significant relationship ( $P > 0.15$ ) between proportion axe-handle fish and  
567 minimum ice area in the same year nor between proportion axe-handle fish and maximum ice  
568 extent 3 years previously, the two variables that correlated strongly with mean condition.  
569 However, there was a significant relationship between the proportion of axe-handle fish and  
570 maximum ice area 4 years previously, the same variable that predicted 50<sup>th</sup> percentile and 75<sup>th</sup>  
571 percentile length (Fig. 7). With increasing maximum ice area, the proportion of axe-handle fish  
572 increased 4 years later (adj  $r^2 = 0.2698$ ,  $F_{1,15} = 6.91$ ,  $P = 0.019$ ).

573

574 ***Change in Catch Rate (CPUE).*** There was no evidence that any environmental variable  
575 (ice variables, whether lagged or not, or SOI variables) affected toothfish CPUE (ln-transformed)

576 or annual changes in toothfish CPUE. Furthermore, there is no significant or AIC-justified  
577 association between CPUE (or changes in CPUE) and any of the ice variables, lagged or not,  
578 during the pre-crash period, 1972 to 2001. As noted, despite environmental variability, there  
579 was little change in capture rates from one year to the next up to 2001 (Figure 8).

580

581 ***Relationships to the Southern Oscillation.*** No significant relationship existed between  
582 SOI and the three types of fish metrics analyzed. This is not to say that SOI is unimportant, only  
583 that the effect of SOI is not direct. Instead, SOI during the months July-December is strongly  
584 associated with maximum ice area in the same year (this reaffirms previous analyses, i.e.,  
585 Wilson *et al.* 2001);  $b = 108,700 (\pm 2400, t = 4.52, P = 0.0001$ ; Fig. 11). A linear relationship  
586 between the two variables is best supported and confirmed by AIC (Table 3).

587 Because maximum ice area reflected SOI in July to December and maximum ice area  
588 predicted 75<sup>th</sup> percentile length of toothfish 4 years later (see above), there was consequently a  
589 modest, non-significant, negative correlation between SOI in July-December and 75<sup>th</sup> percentile  
590 length 4 years later ( $r = -0.255, P > 0.2$ ; Table 4). However, once we accounted for the effect of  
591 maximum ice area on length (with 4-yr-lag), the effect of SOI in July-December on 75<sup>th</sup>  
592 percentile length 4 years later was even less significant and reversed sign (partial regression  
593 coefficient,  $b = 0.585 \pm 0.721, t = 0.81, P > 0.4$ ; standardized partial regression coefficient  $\beta =$   
594  $0.225$ .), while the effect of maximum ice area on length 4 years later remained, even while  
595 controlling for SOI in July-December (partial regression coefficient,  $b = -0.0000107 \pm$   
596  $0.00000485, t = -2.21, P = 0.044$ ; standardized partial regression coefficient  $\beta = -0.613$ ).

597 Results were very similar to that found for 75<sup>th</sup> percentile length analyzing either 50<sup>th</sup> or  
598 95<sup>th</sup> percentile length in relation to the same two independent variables: SOI (July-Dec) with a  
599 4-year lag and maximum ice area with a 4-year lag. The effect of maximum ice area was  
600 significant, controlling for SOI, while the effect of SOI was not.

601 It was also the case that SOI influenced maximum ice extent and thus exerted an  
602 indirect effect on mean condition. However, just as with 75<sup>th</sup> percentile length, there was no  
603 direct effect of SOI on condition 3 years later, once we controlled for maximum ice extent  
604 (partial regression coefficient  $b = 0.0063 \pm 0.0082$ ,  $t = 0.73$ ,  $P > 0.4$ , standardized partial  
605 regression coefficient  $\beta = +0.217$ ). Note that SOI in the current year was not correlated with  
606 mean condition or any of the length metrics, nor was SOI associated with minimum ice area in  
607 the current year.

608

609 ***Relationships to Southern Annular Mode.*** There was little evidence of a direct effect of  
610 SAM (measured on a continuous scale) on toothfish length or condition, whether lagged or not.  
611 The regression of mean condition or the three percentiles of length on SAM was not significant  
612 in every case, with lags of 1 to 5 years.

613 However, the sign of SAM changed from predominantly negative prior to 1976, to  
614 increasingly positive from 1976 to the early 1990s, and then to mostly positive thereafter;  
615 during the period from 1972 (first year of this study) to 1992, 12 out of 21 years were  $\leq 0$ ,  
616 whereas from 1993 to 2011, 15 out of 19 years were positive (LRS = 5.60,  $df = 1$ ,  $P = 0.018$ ; Fig.  
617 2). In accord with the increasing windiness as a result of positive SAM, sea ice extent (and

618 related aspects of sea ice) changed as well (Fig. 11). During the more “negative” period of SAM  
619 (1972 to 1992), condition and length both increased. During the (mostly) “positive” period of  
620 SAM (1993 to 2011), condition and length both declined significantly.

621 While there was no relationship between condition or the length metrics and SAM as a  
622 quantitative variable, 75<sup>th</sup> percentile length was significantly greater in years in which SAM was  
623 positive 4 years earlier as compared with years in which SAM was negative ( $\leq 0$ ) 4 years earlier:  
624 145.85 ( $\pm 1.17$ ;  $n = 14$  years) vs. 142.14 ( $\pm 1.20$ ;  $n = 13$  years;  $P = 0.036$ ; results of weighted  
625 regression). Results were even stronger comparing 50<sup>th</sup> percentile lengths. The median fish  
626 length 4 years after a year with positive SAM was 136.39 (SE = 1.21) compared to median fish  
627 length of 131.84 (SE = 1.13) 4 years after a year with SAM of 0 or less ( $P = 0.011$ ).

628

## 629 **Discussion**

630 In this study we found that (1) the length-frequency of the McMurdo Sound catch was  
631 dominated by fish >100 cm and in the upper two-thirds of the overall distribution exhibited in  
632 the industrial toothfish catch; (2) McMurdo Sound fish length and condition increased from the  
633 early 1970’s to the early 1990s and then decreased, the proportion of “axe handle” fish (poor  
634 condition) in the catch changing inversely with mean fish condition; (3) fish length and  
635 condition decreased with increasing sea ice cover and/or inversely with the extent of open  
636 water, lagged by 8 mo to 5 yr; and (4) catch per unit effort, controlling for within-season  
637 variability, was constant through 2001 and then decreased dramatically. Below, we propose  
638 and discuss hypotheses explaining these patterns.

639

640 **Toothfish length-frequency**

641 Models to estimate stock size based entirely on what can be gleaned from the industrial catch,  
642 as is the case for the Ross Sea region toothfish fishery (CCAMLR 2005-2010), assume that after  
643 making certain statistical assumptions and corrections, e.g., by virtue of mark-recapture data,  
644 models are sufficient to inform the management strategy. One main assumption is that the  
645 size-age distribution in the catch, with adjustment, is representative of the full population. The  
646 disparity in the size distribution in the McMurdo Sound catch compared to the fishery catch is  
647 therefore of concern. The McMurdo Sound catch, ranging from 81-193 cm with a broad mode  
648 at 120-170 cm, reasonably mirrors the upper two-thirds of the commercial catch, i.e. "...50 to  
649 180 cm....In all seasons, there was a broad mode of adult fish at about 120–170 cm..." (CCAMLR  
650 2010, p. 6); ~15-20% of the industrial catch overall appears to be of fish <100 cm, and in years  
651 when there are more of these smaller fish the conclusion is that the fishery centered more over  
652 the Ross Sea shelf than in other years (CCAMLR 2010). In fact, with very few fish <100 cm (2.5%)  
653 in the McMurdo Sound catch, this disparity is even more apparent when compared to the size-  
654 frequency distributions in Hanchet *et al.* (2010). In the latter, only fish captured by the fishery  
655 over the shelf are displayed and >50% are <100 cm, with the fishery expending significant effort  
656 over the shelf (though less than on the slope) including waters close to the scientific fishing  
657 location (i.e., on the opposite side of Ross Island (Hanchet *et al.* 2008).

658           There are at least two factors that could explain the discrepancy. First, owing to lack of  
659 buoyancy in fish <100 cm (Eastman and Sidell 2002, Near *et al.* 2003), the vertical setline failed

660 to attract many of the smaller fish that remain on the bottom (Eastman and Barry 2002, Near *et*  
661 *al.* 2003), whereas the industrial longlines, deployed along the bottom, caught these fish in  
662 greater proportion. Second, the discrepancy could be due to the fact that large fish ate most of  
663 the smaller fish caught on the vertical set line. However, of the stomachs inspected in the  
664 McMurdo catch ( $n = 58$ ), evidence for cannibalism was absent (Eastman 1985a, b; La Mesa *et*  
665 *al.* 2004). That large toothfish, at least under ice cover, reside high in the water column, as  
666 indicated by the catch we report herein, is a fact confirmed by crittercams placed on Weddell  
667 seals as well as on a number of occasions by ROVs (to within 12 m of the surface: Fuiman *et al.*  
668 2002, Kim *et al.* 2011); that small toothfish do occur in McMurdo Sound is indicated by the  
669 catch of Weddell seals, who take the entire size range (cf. Ponganis and Stockard 2007, Ainley  
670 and Siniff 2009). In any case, it would appear that assumptions made in CCAMLR's fishery  
671 models about the representativeness of age-size structure as revealed by longlines need to be  
672 reconsidered.

673           In the context of the discrepancy between vertical and horizontal-bottom sampling it is  
674 interesting to consider why this species is one of the few neutrally buoyant Southern Ocean  
675 fish. The most parsimonious explanation is probably historical. Given the eradication of the  
676 previous taxonomically diverse Eocene fauna, the modern notothenioids diversified into a  
677 developing ecosystem with vacant water-column niches. As inferred from molecular sequence  
678 data, these niches were filled first by the neutrally buoyant clade about 24 million years ago  
679 (Near 2004). The water column contained few fishes and underexploited resources were  
680 available—the Antarctic toothfish diversified coincident with silverfish in response to this  
681 unusual situation. The questions remain: Is neutral buoyancy the means to exploit the high



682 prevalence of silverfish (another neutrally buoyant species), the means to escape cannibalism,  
683 or both (Pinkerton *et al.* 2007)? Is the lack of any water-column species in the diet of longline-  
684 caught fish (Fenaughty *et al.* 2003) a reflection of benthic longlines catching mainly fish that  
685 reside on or near the bottom? Why, supposedly, are toothfish only found mostly on the bottom  
686 in ice-free waters? Regardless, it appears that the clear ontogenetic shift in Antarctic toothfish  
687 from dwelling on the bottom as small fish to somewhere in the water column as large fish, at  
688 least in ice covered waters, is a major aspect of this species' natural history (DeVries and  
689 Eastman 1981, Near *et al.* 2003), with further implication of whether or not removing large fish  
690 results in the increase of small fish (through competitive release), i.e. the basis for modern  
691 fishery theory (cf. Constable *et al.* 2000, Longhurst 2010). In the hypothetical life history  
692 scenario proposed by Hanchet *et al.* (2008) no mention is made of the ontogenetic shift in  
693 buoyancy and its implications, other than to say that as fish mature they occupy waters having  
694 deeper bathymetry. The assumption is made that this species is entirely bottom dwelling  
695 (except perhaps larvae).

696           One further interesting question that arises from the size-frequency of McMurdo Sound  
697 fish compared to that of the industrial catch revolves around the age-related fecundity of this  
698 species. All large McMurdo Sound fish are sexually mature, but in gonadal resting stage; some  
699 females showed signs of previous spawning (Eastman and DeVries 2000). If the industrial catch  
700 underestimates the prevalence of large fish over the shelf, it would seem that estimates of  
701 age/size at maturity, which is based on the size-frequency seen in the industrial catch, needs to  
702 be revised once more (recently it was doubled from ~8 to ~16 y; Parker and Grimes 2010).

703

704 **Trends in toothfish length and condition**

705 Both total length and condition of toothfish in the McMurdo Sound catch increased from the  
706 mid 1970s to the early 1990s, and then switched to gradually decrease. Because of the low  
707 sample size for 1991 and 1992, and the lack of any data during 1993-1996, it is difficult to say  
708 when length and condition trends actually leveled off in the early to mid-1990's before  
709 decreasing. However, in the period up to and including 1990 there was no indication of leveling  
710 off. Instead, a linear trend was evident in condition and length from 1972 to 1990. Whether or  
711 not the decreasing trend is represented in the industrial catch, which only began (legally) in  
712 1996 (CCAMLR refers this as the 1997 fishing season), we do not know. Seemingly this would be  
713 difficult to ascertain in a sampling regime that is not only non-random but is geared to catch the  
714 largest fish as fast as possible before the season closes (Constable *et al.* 2000, Brooks 2008).

715         Regardless, it appears that growth rate in this species is not a constant but is subjected  
716 to environmental and potentially anthropogenic factors. According to fishery theory, as fish  
717 (especially large ones) are removed from the stock, fish size and condition should show a  
718 positive growth as smaller fish are released from competition. This is rarely demonstrated  
719 (Longhurst 2010) and is not evident in the McMurdo Sound data. What then are the factors  
720 that could account for the observed increases and then decreases in length and condition? If  
721 change in length is a function both of food availability and age, was there 1) a release from  
722 predation during the 1970-80s, thus allowing older fish to remain in the population longer,  
723 and/or 2) more favorable foraging related to competitive release or climatic factors that  
724 influenced silverfish?

725  
726           *Release from predation.* The southern Ross Sea was the scene for a number of “heroic”  
727 expeditions during the early half of the 1900s, and they all took large numbers of Weddell seals  
728 for human and dog food. The southern Ross Sea seal population was thought to have recovered  
729 by the 1960s (Stirling 1971), but then the NZ Antarctic Programme began to kill 50-100 seals  
730 annually to feed sled dogs and by the time this ended in the mid-1980s, the population, at least  
731 as measured in McMurdo Sound was halved, from 3000 to 1500 seals. Then over the next  
732 decade, it slowly recovered (partially) to ~2000+, where it has remained ever since  
733 (summarized in Ainley 2010). Given that Weddell seals eat a significant number of Antarctic  
734 toothfish (Ainley and Siniff 2009, and references therein; Kim *et al.* 2011), this marked decrease  
735 in Weddell seals in the southern Ross Sea is consistent with a lessening of predation pressure  
736 on Antarctic toothfish. Certainly, the lag that would occur between these two long-lived, late  
737 maturing species, predator versus prey, would be large. Added to this is the fact that another  
738 toothfish predator, sperm whales (admittedly at the periphery of their range), owing mainly to  
739 contraction of range as whalers depleted the stocks in the warmer parts of the Pacific during  
740 1700-mid 1900s, even now are a shadow of their former numbers in the Ross Sea sector  
741 (Whitehead 2000). Their numbers in the Ross Sea sector also may well have begun to recover in  
742 recent decades as they have elsewhere (see for instance, Whitehead *et al.* 1997, Branch and  
743 Butterworth 2001). The same is true for southern elephant seals, still another toothfish  
744 predator, which disappeared from the Ross Sea during the 1980s, their foraging range  
745 contracting as the Macquarie Island breeding population became severely reduced  
746 (summarized in Ainley 2010).

747  
748 *More favorable foraging related to competitive release or climatic factors.* The industrial  
749 take of minke whales during the 1970s-early 1980s (~20,000 taken from Ross Sea region), which  
750 appears to have resulted in the population increase of trophically competing Adélie Penguins at  
751 the time (Ainley *et al.* 2007), also could have benefited Antarctic toothfish, both the whales  
752 (and penguins) and the toothfish being significant predators of Antarctic silverfish (Ballard *et al.*  
753 2011). Furthermore, the time of the switch in toothfish length and condition trends from  
754 positive to negative corresponds to when the minke whale population was deemed to have  
755 reached recovery (roughly 1986-1990; Branch 2006, Ainley 2010).

756 There is also the possibility that the shifting climate of the Ross Sea region has affected  
757 (is affecting) the abundance of Antarctic silverfish, a subject about which we know even less.  
758 According to La Mesa *et al.* (2010), the prevalence of larval silverfish in the western Ross Sea  
759 was an order of magnitude higher in the 1999-2000 austral summer, when the Ross Sea  
760 polynyas opened early and large, compared to the austral summers of 1997-98 and 2003-04;  
761 with more open water, earlier, primary production is increased and, hypothetically, silverfish  
762 survival is enhanced owing to increased food. Silverfish spawn during the winter and their eggs  
763 can be found among the frazil ice associated with heavy pack or fast ice (Vacchi *et al.* 2004).  
764 Perhaps this is a protective strategy, and more ice at this time is beneficial, but once the eggs  
765 hatch the larvae need to forage on microbial organisms. In any case, though most of the  
766 silverfish contained in toothfish stomachs in McMurdo Sound during late spring are 100–150  
767 mm SL (Eastman and DeVries, unpublished data) and on the order of 4+ years old (Hubold and  
768 Tomo 1989), we did find that toothfish condition is positively affected by extensive open water,

769 and hence water open to insolation, during mid-summer. We also found the effect of a 3-yr-lag  
770 in ice extent on toothfish condition, and note that Ross Sea primary productivity cycles on a 2-4  
771 year period owing to a combination of nutrient and sea ice factors (Peloquin and Smith 2007).  
772 Consistent with the 4 yr and older age classes of silverfish that toothfish eat in mid-summer, we  
773 also found a negative relationship between toothfish length and maximum sea ice area 4-5  
774 years earlier. Greater ice area means less open water. That's not to say that we understand the  
775 interconnections, but these few factors and relationships would seem to implicate the  
776 availability (and possibly condition) of silverfish as being important to toothfish growth and  
777 condition. Obviously, we need to learn considerably more, not just about toothfish, but about  
778 silverfish as well, if we are to understand toothfish. Making what we can of the trends in axe  
779 handle fish, we note the consistency between the decreasing trend in prevalence while  
780 toothfish length and mean condition were increasing during the first half of the study period, as  
781 opposed to the increased prevalence of axe handles as length and condition began to decrease.

782         Another factor to consider is the Southern Annular Mode, a major driver of climate in  
783 the Southern Ocean (Thompson and Solomon 2002, Stammerjohn *et al.* 2008). During the  
784 period when toothfish length and condition were increasing, beginning in the mid 1970s SAM  
785 was increasingly becoming less negative and more positive until becoming almost entirely  
786 positive starting at the inflection point of both toothfish length and condition. Therefore, owing  
787 to increased windiness blowing south to north (as well as westerly winds in the north increasing  
788 Ekman transport), sea ice extent was expanding, early appearance of coastal polynyas was  
789 becoming more reliable, and the sea ice season was lengthening (Parkinson 2002, Stammerjohn  
790 *et al.* 2008). It would seem that more reliably present coastal polynyas in the Ross Sea would

791 favor silverfish production, but the decreasing annual period of open water would not. Other  
792 than the correspondence in the inflection points of length and condition and the onset of  
793 continually positive SAM, we again emphasize that we don't understand much about this  
794 predator-prey system, toothfish and silverfish. The correspondence in trends between those  
795 changes in trend and SAM, though, is noteworthy and deserving of closer study.

796         That the Southern Oscillation Index had no direct correlation to variation in any  
797 Antarctic toothfish parameters is perhaps not surprising. In the Patagonian toothfish, SOI  
798 correlates to egg and larval prevalence through its influence on ocean temperature but by the  
799 time fish recruit into the breeding population, the SOI signal has become obscured (Belchier  
800 and Collins 2008). Certainly many factors come into play between spawning and recruitment in  
801 a fish that matures 15 years later. Other, shorter-lived ground fish in the Scotia Sea have  
802 exhibited correlations to SOI, but as those authors note, these fish feed heavily on krill, a  
803 species known to track closely SOI changes to its ocean environment. Antarctic toothfish, as far  
804 as anyone knows, do not feed readily on krill; but then no one knows what toothfish eat during  
805 winter along the Ross Sea slope where krill and krill predators are abundant (Ballard *et al.*  
806 2011).

807

#### 808 **Change in toothfish prevalence**

809 *Monthly Pattern in CPUE.* The within-year pattern of apparent toothfish abundance (catch rate)  
810 was interesting but somewhat resists explanation given the lack of other information. The  
811 pattern shown in the 27 years of data was similar in form but with a lesser decrease in catch

812 rate after November than reported by Testa *et al.* (1985). In fact, the pattern apparent in our  
813 data was similar to one of the three years in the Testa *et al.* study; the other two showed a  
814 more extreme decrease for December. To explain the temporal pattern, one might be tempted  
815 to invoke some sort of migratory movement on the scale of shorebirds visiting the Arctic tundra  
816 to breed but then vacating it before food becomes unavailable, or for that matter the annual  
817 invasion and evacuation of the Ross Sea by Adélie penguins (Ballard *et al.* 2010). After all,  
818 McMurdo Sound tagged fish have been recaptured well to the north of McMurdo Sound, as  
819 noted (see Hanchet *et al.* 2008); southward movement into the southern Ross Sea by industry-  
820 tagged fish (along the slope) has been detected as well (Hanchet *et al.* 2010). On the other  
821 hand, toothfish could also seasonally change their position in the water column to be more  
822 vulnerable to our setline array. From personal experience with other fish species (ocean  
823 salmon), if you are not fishing at the depth where they occur you won't catch them, and their  
824 depth can change dramatically and day-to-day (D. Ainley, 100s of hours of pers. obs.).  
825 Therefore, we are inclined to explain the early spring increase in our catch rate on the basis of  
826 shifting position in the water column due to food availability. In early spring (August,  
827 September) with the sun first rising on 20<sup>th</sup> August, there would be no reason for silverfish, and  
828 therefore toothfish, to be very high above the bottom (where our hooks are located) given a  
829 phytoplankton concentration close to zero (indicated by visibility ~80 m; Barry 1988). Any prey  
830 for silverfish would be on or near the bottom and, indeed, crystal krill are known to feed on  
831 benthic detritus during winter (Nicol *et al.* 2004, Deibel and Daly 2007). Therefore, perhaps all  
832 toothfish, along with their prey, are found at or near the bottom at that time, and not inclined  
833 to find our baits. A couple of months later, when the plankton blooms begin and which are

834 sufficiently dense to alter the 1% light level from 54 m to just a few meters deep (visibility to <6  
835 m; Barry 1988, Arrigo *et al.* 1998), there is ample reason for zooplankton and fish to occupy the  
836 water column. Then, the depth of toothfish tracks that of silverfish in as noted in McMurdo  
837 Sound (Fuiman *et al.* 2002).

838         If there is a diminution of catch rate after November/December (perhaps in a few years;  
839 Testa *et al.* 1985), we hypothesize that this is related to a true decrease in toothfish abundance  
840 (especially large fish) owing to predation, which increases severely at that time as Weddell seals  
841 attempt to recover from breeding (beginning in late November), as a new cohort of seals begin  
842 to forage in the foodweb (late November), and as Ross Sea killer whales appear (early  
843 December; Ainley and Siniff 2009, Ainley *et al.* 2009). Dearborn (1965) and Calhaem and  
844 Christoffel (1969) reported toothfish eaten abundantly by seals as late as January in McMurdo  
845 Sound. Moreover, industrial vessels have been quite successful as well in January (Hanchet *et*  
846 *al.* 2010). Thus, the fish are present. Testa *et al.* (1985) showed that predation affects the  
847 spatial extent of toothfish abundance, so why not the temporal extent as well?

848         Indeed, according to Everson (1970), the movements and annual cycles of nototheniids  
849 in the Scotia Sea appear to have evolved to avoid predation by top predators. This brings us  
850 back to the question of why it seems that as encountered by the fishery, in ice-free waters,  
851 Antarctic toothfish are found on the bottom? Sperm whales and elephant seals (summer, ice-  
852 free visitors only) are known to dive to 2400 and 3000 m, respectively, Weddell seals to 700 m,  
853 and fish-eating killer whales to 350 m (Ballard *et al.* 2011); of course average foraging depths  
854 are proportionately much shallower. The fishery targets the fish in waters 1000-2000 m. Living



855 at that depth would be out of range for at least two of these predators. In shallower waters, the  
856 smaller fish, being also cryptically colored, appear to hide within the “forest” of benthic  
857 invertebrates (Eastman and Barry 2002). Of interest, in trophic studies that looked at fish  
858 caught in deep waters along the Ross Sea slope and over sea mounts to the north, 59-64% had  
859 empty or essentially empty (food “trace”) stomachs (Fenaughty *et al.* 2002). In contrast, surface  
860 waters over the slope are teeming with foraging predators, gorging on krill and fish (Ballard *et*  
861 *al.* 2011), and over the shelf in ice covered waters only 10% of toothfish stomachs have been  
862 found empty (Eastman 1985b). It would appear that the toothfish in the deep depths are biding  
863 their time, awaiting the return of sea ice cover and the seasonal departure of some of their  
864 predators. At this time they could be particularly vulnerable to baited hooks.

865

866 *Decadal Pattern in CPUE.* In regard to longer-term change in catch rate, no factor that  
867 we used in our analyses could explain the drop off in the CPUE of toothfish in McMurdo Sound  
868 beginning after 2001. One could hypothesize that the mega icebergs that blocked McMurdo  
869 Sound 2001-2005 were responsible (Arrigo *et al.* 2002), and one of the explanations favored by  
870 Hanchet *et al.* (2010, discussing a preliminary report of the McMurdo Sound data set), but  
871 other than a temporary effect, no other mesopredator was affected by the icebergs in a long-  
872 lasting manner (beyond 2005). In fact, Weddell seal numbers and pup production immediately  
873 returned to pre-iceberg levels (Siniff *et al.* 2008), Adélie penguin breeding population in the  
874 southwestern Ross Sea increased during this period and through to the present, and the  
875 prevalence of emperor penguins in McMurdo Sound increased by an order of magnitude  
876 (Landcare Research NZ, unpubl. data; Ainley pers. obs.). All of these species appear to have a

877 broadly similar diet during summer in waters of the southern shelf as do toothfish, principally  
878 through high consumption rates of silverfish (cf. La Mesa *et al.* 2004, Ballard *et al.* 2011). The  
879 fishing site did not change sufficiently to explain the trend either (see Hanchet *et al.* 2010).

880           Coincident with the decrease in the CPUE of toothfish, toothfish length (and condition)  
881 decreased as well. We can only conclude that the industrial fishery, which targets the largest  
882 fish, is reducing their prevalence in the southwestern Ross Sea. Could it also be reducing the  
883 prevalence of “high quality” individuals (ones that grow faster and mature quickly and are in  
884 better condition), as other fisheries do when targeting the largest, oldest fish (Longhurst 2010,  
885 Ainley *et al.* 2011)? Apparently this is happening in the heavily fished Patagonian toothfish of  
886 the Scotia Sea, although detected over a longer time period (Shust and Kozlov 2006). Given the  
887 inability of the industrial fishery to catch many large Antarctic toothfish over the Ross Sea shelf  
888 (Hanchet *et al.* 2008, 2010), fishery biologists dependent on CCAMLR data would not be aware  
889 of this decreasing prevalence of large fish. Indeed, Hanchet *et al.* (2010) state that trends are  
890 unlikely to be found in the industrial catch data owing to the high variability in catch rates and  
891 characteristics of individual vessels.

892           A change in the representation of large fish in the southern Ross Sea, indicative of the  
893 entire stock, would not be ecologically neutral (Longhurst 2010, Ainley *et al.* 2011 and  
894 references therein). How to detect this change is a challenge. Such a change was involved in the  
895 crash of Atlantic cod *Gadus morhua*, where the diminishment of the inshore portion of the  
896 stock occurred well before any signal was evident in the larger offshore stocks and prior to the  
897 eventual total collapse of the fishery (Longhurst 2010, and references therein). In accord with

898 the decreasing numbers of large fish, the Ross Sea killer whale prevalence in the southwestern  
899 Ross Sea continues to decrease (Ainley *et al.* 2009, unpubl. data), a pattern played out by the  
900 now endangered large-fish-eating, resident killer whales in waters of western Canada and Puget  
901 Sound (Ford *et al.* 2010). The killer whale social structure depends on the existence of large fish.  
902 The increase in penguin populations would be consistent as well with decreased competition  
903 with toothfish for silverfish prey.

904

### 905 **Recommendations**

906 We could suggest that this toothfish fishery in the Ross Sea be closed until the natural history of  
907 Antarctic toothfish, and its silverfish prey, is much better understood, given the observation of  
908 recent declines in condition, size, and CPUE for the Antarctic toothfish of McMurdo Sound, the  
909 only monitoring conducted independent of the fishery. An easier option to implement, given  
910 that New Zealand has taken 55% of the toothfish from the Ross Sea area (Ainley *et al.* 2011)  
911 and that the United States imports 40% of the total world catch of “Chilean sea bass” (including  
912 Ross Sea fish; US Dept of State website), is for NZ and US fishery agencies to immediately  
913 reconstitute the scientific fishing program described herein as it was up to 2001 (annual, entire  
914 season), as well as initiate other research, e.g. monitoring of dependent species, that would be  
915 independent of the unexplained biases in the industrial sampling. Besides McMurdo Sound,  
916 vertical longlines could be deployed in an analagous fashion through the seasonally long-lasting  
917 fast ice of Terra Nova Bay as well as Edisto-Moubray bays farther north along the Ross Sea  
918 coast. Doing so would be consistent for an agency, CCAMLR (Commisison for the Conservation

919 of Antarctic Marine Living Resources), that has made its name as setting the pace in ecosystem-  
920 based and precautionary management and which has a formal monitoring program ( CCAMLR  
921 Ecosystem Monitoring Program; CEMP) from which management of fisheries is supposed to  
922 become effectively informed (Constable *et al.* 2000, Croxall and Nicol 2004, Constable 2011).  
923 Trends in the “unofficial CEMP” based on changes in numbers of fish-eating killer whales, seals  
924 and penguins underway in the southern Ross Sea, by scientists engaged in pure rather than  
925 fishery-directed research, is pointing to the need for meaningful action.

## 926 **Acknowledgements**

927 The fishing effort described was funded for most of its tenure by the US National Science  
928 Foundation; logistics support was provided by the US Antarctic Program and Antarctica NZ.  
929 Data analysis and writing was funded by the Lenfest Ocean Program, and by the NSF Office of  
930 Polar Programs, Grant ANT-0440643. The views expressed herein do not necessarily reflect  
931 those of the National Science Foundation. We appreciate comments on the manuscript by C.  
932 Brooks, S. Hanchet, M. Pinkerton, J. Rotella, D. Siniff and xxxx; all improved it immensely. PRBO  
933 contribution # XXXX.

934

## 935 **REFERENCES**

- 936 Ainley, D.G. (2010) A history of the exploitation of the Ross Sea, Antarctica. *Polar Record* **46**,  
937 233-243.
- 938 Ainley, D.G. and Ballard, G. (2011) Non-consumptive factors affecting foraging patterns in  
939 Antarctic penguins: a review and synthesis. *Polar Biology*, DOI10.1007/s00300-011-1042-x.

- 940 Ainley, D.G., Ballard, G., Ackley, S., Blight, L.K., Eastman, J.T., Emslie, S.D., Lescroël, A.,  
941 Olmastroni, S., Townsend, S.E., Tynan, C.T., Wilson, P. and Woehler, E. (2007) Paradigm  
942 lost, or, is top-down forcing no longer significant in the Antarctic Marine Ecosystem?  
943 *Antarctic Science* **19**, 283-290.
- 944 Ainley, D.G., Ballard, G., Dugger, K.M. (2006) Competition among penguins and cetaceans  
945 reveals trophic cascades in the Ross Sea, Antarctica. *Ecology* **87**, 2080–2093.
- 946 Ainley, D.G., Ballard, G. and Olmastroni, S. (2009) An apparent decrease in the prevalence of  
947 “Ross Sea Killer Whales” in the southern Ross Sea. *Aquatic Mammals* **35**, 335-347.
- 948 Ainley, D.G., Brooks, C.M., Eastman, J.T. and Massaro, M. (2011) Unnatural selection of  
949 Antarctic toothfish in the Ross Sea, Antarctica. In: Huettmann F (ed) (2011) *Protection of*  
950 *the Three Poles*. Springer Tokyo, Japan, in press.
- 951 Ainley, D.G., Clarke, E.D., Arrigo, K., Fraser, W.R., Kato, A., Barton, K.J. and Wilson, P.R. (2005)  
952 Decadal-scale changes in the climate and biota of the Pacific sector of the Southern Ocean,  
953 1950s to the 1990s. *Antarctic Science* **17**, 171-182.
- 954 Ainley, D.G., O'Connor, E.F. and Boekelheide, R.J. (1984) *Ecology of seabirds in the Ross Sea,*  
955 *Antarctica*. AOU Monograph No 32, 1- 79.
- 956 Ainley, D.G., Russell, J., Jenouvrier, S., Woehler, E., Lyver, P. O'b., Fraser, W.R. and Kooyman  
957 G.L.. (2010) Antarctic penguin response to habitat change as earth's troposphere reaches  
958 2°c above pre-industrial levels. *Ecological Monographs* **80**, 49-66.
- 959 Ainley, D.G. and Siniff, D.B. (2009) The importance of Antarctic toothfish as prey of Weddell  
960 Seals in the Ross Sea: A Review. *Antarctic Science* **21**, 317-327.
- 961 Anderson, R.O. and Gutreuter, S.J. (1983) Length, weight, and associated structural indices. In:

- 962 Nielsen LA, Johnson DL (eds) *Fisheries techniques*. American Fisheries Society, Bethesda,  
963 MD, pp 283-333.
- 964 Arrigo, K.R., van Dijken, G. L., Ainley, D.G., Fahnestock, M.A. and Markus, T. (2002) The impact  
965 of the B-15 iceberg on productivity and penguin breeding success in the Ross Sea,  
966 Antarctica. *Geophysical Research Letters* **29**(7), 10.1029/2001GLO14160.
- 967 Arrigo, K.R., Robinson, D.H., Worthen, D.L., Schieber, B. and Lizotte, M.P. (1998) Bio-optical  
968 properties of the southwestern Ross Sea. *Journal of Geophysical Research* 103: 21,683-  
969 21,695.
- 970 Ballard, G., Jongsomjit, D. and Ainley, D.G. (2011) Coexistence of mesopredators in an intact  
971 polar ocean ecosystem: the basis for defining a Ross Sea Marine Protected Area. *Biological*  
972 *Conservation*, in press.
- 973 Ballard, G., Toniolo, V., Ainley, D.G., Parkinson, C.L., Arrigo, K.R. and Trathan, P.N. (2010)  
974 Responding to climate change: Adélie penguins confront astronomical and ocean  
975 boundaries. *Ecology* **91**, 2056–2069.
- 976 Balushkin, A.V. (2000) Morphology, classification, and evolution of notothenioid fishes of the  
977 Southern Ocean (Notothenioidei, Perciformes). *Journal of Ichthyology* **40** [Suppl 1]:S74–  
978 S109.
- 979 Barry, J.P. (1988). Hydrographic patterns in McMurdo Sound, Antarctica and their relationship  
980 to local benthic communities. *Polar Biology* **8**, 377-391.
- 981 Belchier, M. and Collins, M.A. (2008) Recruitment and body size in relation to temperature in  
982 juvenile Patagonian toothfish (*Dissostichus eleginoides*) at South Georgia. *Marine Biology*  
983 **155**, 493-503.

- 984 Branch, T.A. (2006) *Abundance estimates for Antarctic minke whales from three completed*  
985 *circumpolar sets of surveys, 1978/79 to 2003/04*. International Whaling Commission, *Paper*  
986 *SC/58/IA18*. Cambridge, UK.
- 987 Branch, T.A., and Butterworth, D.S. (2001) Estimates of abundance south of 60°S for cetacean  
988 species sighted frequently on the 1978/79 to 1997/98 IWC/IDCR-SOWER sighting surveys.  
989 *Journal of Cetacean Research and Management* **3**, 251-270.
- 990 Brooks, C.M. and Ashford, J.R. (2008) Spatial distribution and age structure of the Antarctic  
991 toothfish (*Dissostichus mawsoni*) in the Ross Sea, Antarctica. *CCAMLR WG-FSA-08-18*.  
992 Hobart; also available in: Brooks, C.M. (2008) *Radiometric age validation and spatial*  
993 *distribution of the Antarctic toothfish (Dissostichus mawsoni): Implications for a deep-sea*  
994 *Antarctic fishery*. M.Sc. thesis, Calif State Univ, Moss Landing Marine Labs, Moss Landing,  
995 CA. 109 pp
- 996 Brooks, C.M., Andrews, A.H., Ashford, J.R., Ramanna, N., Jones, C.D., Lundstrom, C.C. and  
997 Cailliet, G.M. (2010) Age estimation and lead–radium dating of Antarctic toothfish  
998 (*Dissostichus mawsoni*) in the Ross Sea. *Polar Biology*, DOI 10.1007/s00300-010-0883-z
- 999 CCAMLR (2005) Report of the Scientific Committee, Appendix F.  
1000 [http://www.ccamlr.org/pu/e/e\\_pubs/sr/05/toc.htm](http://www.ccamlr.org/pu/e/e_pubs/sr/05/toc.htm).
- 1001 CCAMLR (2006) Report of the Scientific Committee, Appendix F.  
1002 [http://www.ccamlr.org/pu/e/e\\_pubs/sr/06/toc.htm](http://www.ccamlr.org/pu/e/e_pubs/sr/06/toc.htm)
- 1003 CCAMLR (2008) [http://www.ccamlr.org/pu/e/e\\_pubs/fgl/fsa-08-60.pdf](http://www.ccamlr.org/pu/e/e_pubs/fgl/fsa-08-60.pdf)
- 1004 CCAMLR (2010) Report of the Scientific Committee, Appendix K.  
1005 [http://www.ccamlr.org/pu/e/e\\_pubs/sr/10/toc.htm](http://www.ccamlr.org/pu/e/e_pubs/sr/10/toc.htm).

- 1006 Calhaem, I. and Christoffel, D.A. (1969) Some observations of the feeding habits of a Weddell  
1007 seal, and measurements of its prey, *Dissostichus mawsoni*, at McMurdo Sound, Antarctica.  
1008 *New Zealand Journal of Marine and Freshwater Research* **3**, 181–190.
- 1009 Collins, M.A., Brickle, P., Brown, J. and Belchier, M. (2010) The patagonian toothfish: biology,  
1010 ecology and fishery. In: Lesser, M. (ed.) *Advances in marine biology* **58**, 227-300. Elsevier  
1011 Academic Press Inc, San Diego.
- 1012 Constable, A.J., de la Mare, W.K., Agnew, D.J., Everson, I. and Miller, D. (2000) Managing  
1013 fisheries to conserve the Antarctic marine ecosystem: practical implementation of the  
1014 Convention on the Conservation of Antarctic Marine Living Resources (CCAMLR). *ICES*  
1015 *Journal of Marine Science* **57**, 778–791.
- 1016 Constable, A.J. (2011) Lessons from CCAMLR on the implementation of the ecosystem approach  
1017 to managing fisheries. *Fish and Fisheries*, doi:10.1111/j.1467-2979.2011.00410.x
- 1018 Croxall, J.P. and Nicol, S. (2004) Management of Southern Ocean fisheries: global forces and  
1019 future sustainability. *Antarctic Science* **16**, 569–584.
- 1020 Davidson, D. and Marshall, T. (2010) Are morphometric indices accurate indicators of stored  
1021 energy in herring *Clupea harengus*? *Journal of Fish Biology* **76**, 913-929.
- 1022 Dearborn, J.H. (1965) Food of Weddell seals at McMurdo Sound, Antarctica. *Journal of*  
1023 *Mammalogy* **46**, 37–43.
- 1024 Deibel, D. and Daly, K.L. (2007) Zooplankton processes in Arctic and Antarctic polynyas. In:  
1025 Smith, W.O. and Barber, D.G. (eds.) *Polynyas: Windows to the World's Oceans*, pp 271-232.  
1026 Elsevier, San Diego.
- 1027 DeVries, A.L. (1988) The role of antifreeze glycopeptides and peptides in the freezing avoidance



- 1028 of Antarctic fishes. *Comparative Biochemistry and Physiology* **90B**, 611-621.
- 1029 DeVries, A.L. and Cheng, C-H.C. (2005) Antifreeze proteins and organismal freezing avoidance in  
1030 polar fishes. In: Farrell, A.P. and Steffensen, J.F. (eds) *The physiology of polar fishes*, vol 22.  
1031 Elsevier Academic Press, San Diego, pp. 155-201.
- 1032 DeVries, A.L. and Eastman, J.T. (1998) Brief review of the biology of *Dissostichus mawsoni*.  
1033 *CCAMLR Document WG-FSA-98/49*, Hobart, Australia.
- 1034 Eastman JT (1985a) *Pleuragramma antarcticum* (Pisces, Nototheniidae) as food for other fishes  
1035 in McMurdo Sound, Antarctica. *Polar Biology* **4**, 155–160.
- 1036 Eastman, J.T. (1985b) The evolution of neutrally buoyant notothenioid fishes: their  
1037 specializations and potential interactions in the Antarctic marine food web. In: Siegfried,  
1038 W.R., Condy, P.R. and Laws, R.M. (eds) *Antarctic Nutrient Cycles and Food Webs*. Springer-  
1039 Verlag, Berlin and Heidelberg, pp 430–436.
- 1040 Eastman, J. (1993) *Antarctic Fish Biology*. Academic Press, San Diego.
- 1041 Eastman, J.T. (2005) The nature of the diversity of Antarctic fishes. *Polar Biology* **28**, 93-107.
- 1042 Eastman, J.T. and Barry, J.P. (2002) Underwater video observation of the Antarctic toothfish  
1043 (*Dissostichus mawsoni* (Perciformes: Nototheniidae) in the Ross Sea, Antarctica. *Polar*  
1044 *Biology* **25**, 391-395.
- 1045 Eastman, J.T. and DeVries, A.L. (1981) Buoyancy adaptations in a swim-bladderless Antarctic  
1046 fish. *Journal of Morphology* **167**, 91-102.
- 1047 Eastman, J.T. and DeVries, A.L. (2000) Aspects of body size and gonadal histology in the  
1048 Antarctic toothfish, *Dissostichus mawsoni*, from McMurdo Sound, Antarctica. *Polar Biology*  
1049 **23**, 189–195.

- 1050 Eastman, J.T. and Sidell, B.D. (2002) Measurements of buoyancy for some Antarctic  
1051 notothenioid fishes from the South Shetland Islands. *Polar Biology* **25**, 753-760.
- 1052 Everson, I. (1970) The population dynamics and energy budget of *Notothenia neglecta* Nybelin  
1053 at Signy Island, South Orkney Islands. British Antarctic Survey Bulletin 23, 25–50.
- 1054 Fenaughty, J.M., Eastman, J.T. and Sidell, B.D. (2008) Biological implications of low condition  
1055 factor “axe handle” specimens of the Antarctic toothfish, *Dissostichus mawsoni*, from the  
1056 Ross Sea. *Antarctic Science* **20**, 537–551.
- 1057 Fenaughty, J.M., Stevens, D.W. and Hanchet, S.M. (2003) Diet of the Antarctic toothfish  
1058 (*Dissostichus mawsoni*) from the Ross Sea, Antarctica (CCAMLR Statistical Subarea 88.1).  
1059 *CCAMLR Science* **10**, 1-11.
- 1060 Ford, J.K.B., Ellis, G.M., Olesiuk, P.F., and Balcomb, K.C. (2010) Linking killer whale survival and  
1061 prey abundance: food limitation in the oceans’ apex predator? *Biology Letters* **6**, 139-142.
- 1062 Fuiman, L.A., Davis, R.W. and Williams, T.M. (2002) Behavior of midwater fishes under the  
1063 Antarctic ice: observations by a predator. *Marine Biology* **140**, 815–822.
- 1064 Halpern, B.S., Walbridge, S., Selkoe, K.A., Kappel, C.B., Micheli, F., D’Agrosa, C., Bruno, J.F.,  
1065 Casey, K.S., Ebert, C., Fox, H.E., Fujita, R., Heinemann, D., Lenihan, H.S., Madin, E.M.P.,  
1066 Perry, M.T., Selig, E.R., Spalding, M., Steneck, R. and Watson R. (2008) A global map of  
1067 human impact on marine ecosystems. *Science* **319**, 948–951.
- 1068 Hanchet, S.M., Mormede, S. and Dunn, A. (2010) Distribution and relative abundance of  
1069 Antarctic toothfish (*Dissostichus mawsoni*) on the Ross Sea shelf. *CCAMLR Science* **17**, 33-  
1070 51.

- 1071 Hanchet, S.M., Rickard, G.J., Fenaughty, J.M., Dunn, A. and Williams, M.J.H. (2008) Hypothetical  
1072 life cycle for Antarctic toothfish (*Dissostichus mawsoni*) in the Ross Sea region *CCAMLR*  
1073 *Science* **15**, 35–53.
- 1074 Harrell, F. E., Jr. (2001) *Regression Modeling Strategies: With Applications to Linear Models,*  
1075 *Logistic Regression, and Survival Analysis.* Springer, New York.
- 1076 Horn, P.L. (2002) Age and growth of Patagonian toothfish (*Dissostichus eleginoides*) and  
1077 Antarctic toothfish (*D. mawsoni*) in waters from the New Zealand subantarctic to the Ross  
1078 Sea, Antarctica. *Fishery Research* **56**, 275–287.
- 1079 Hubold, G. and Tomo, A.P. (1989) Age and growth of Antarctic silverfish *Pleuragramma*  
1080 *antarcticum* Boulenger, 1902, from the southern Weddell Sea and Antarctic Peninsula.  
1081 *Polar Biology* **9**, 205-212.
- 1082 Jacobs, S. (2006) Observations of change in the Southern Ocean. *Philosophical Transactions of*  
1083 *the Royal Society, Series A* **364**, 1657–1681.
- 1084 Jacobs, S.S., Giulivi, C.F. and Mele, P.A. (2002) Freshening of the Ross Sea during the late 20th  
1085 century. *Science* **297**, 386–389.
- 1086 Kim, S.Z., Ainley, D.G., Pennycook, J. and Eastman, J.T. (2011) Antarctic toothfish heads found  
1087 along tide cracks of the McMurdo Ice Shelf. *Antarctic Science,*  
1088 doi:10.1017/S095410201100040X.
- 1089 Kutner, M., Nachtsheim, C., Neter, J. and Li, W. (2005) *Applied Linear Statistical Models, 5<sup>th</sup>*  
1090 *edition.* McGraw Hill, New York
- 1091 La Mesa, M., Eastman, J.T. and Vacchi, M. (2004) The role of notothenioid fish in the food web  
1092 of the Ross Sea shelf waters: a review. *Polar Biology* **27**, 321–338.

- 1093 La Mesa, M., Catalano, B., Russo, A., Greco, S., Vacchi, M and Assali, M. (2010) Influence of  
1094 environmental conditions on spatial distribution and abundance of early life stages of  
1095 Antarctic silverfish, *Pleuragramma antarcticum* (Nototheniidae), in the Ross Sea. *Antarctic  
1096 Science* **22**, 243-255.
- 1097 Lee, D. E., Nur, N., and Sydeman, W.J. (2007) Climate and demography of the planktivorous  
1098 Cassin's auklet *Ptychoramphus aleuticus* off northern California: implications for population  
1099 change. *Journal of Animal Ecology* **76**, 337-347.
- 1100 Longhurst, A. (2010) *Mismanagement of Marine Fisheries*. Cambridge University Press,  
1101 Cambridge, UK.
- 1102 Marshall, G.J. (2003) Trends in the Southern Annular Mode from observations and reanalysis.  
1103 *Journal of Climate* **16**, 4134-4143.
- 1104 Morin, P.A., Archer, F.I., Foote, A.D., Vilstrup, J., Allen, E.E., Wade, P., Durban, J., Parsons, K.,  
1105 Pitman, R., Li, L., Bouffard, P., Abel Nielsen, S.C., Rasmussen, M., Willerslev, E., Gilbert,  
1106 M.T.P. and Harkins, T. (2010) Complete mitochondrial genome phylogenetic analysis of  
1107 killer whales (*Orcinus orca*) indicates multiple species. *Genome Research*, doi:  
1108 10.1101/gr.102954.109.
- 1109 Near, T.J. (2004) Estimating divergence times of notothenioid fishes using a fossil-calibrated  
1110 molecular clock. *Antarctic Science* **16**, 37-44.
- 1111 Near, T.J., Russo, S.E., Jones, C.D. and DeVries, A.L. (2003) Ontogenetic shift in buoyancy and  
1112 habitat in the Antarctic toothfish, *Dissostichus mawsoni* (Perciformes: Nototheniidae).  
1113 *Polar Biology* **26**, 124–128.

- 1114 Nicol, S., Virtue, P., King, R., Davenport, S.R., McGaffin, A.F. and Nichols, P. (2004) Condition of  
1115 *Euphausia crystallorophias* off East Antarctica in winter in comparison to other seasons.  
1116 *Deep Sea Research II* **51**, 2215-2224.
- 1117 Parker, S.J. and Grimes, P. J. (2010). Length and age at spawning of Antarctic toothfish  
1118 *Dissostichus mawsoni* in the Ross Sea. *CCAMLR Science* **17**, 53-73.
- 1119 Parkinson, C.L. (2002) Trends in the length of the Southern Ocean sea ice season, 1979–99.  
1120 *Annals of Glaciology* **34**, 435–440.
- 1121 Peloquin, J. A. and Smith, W.O., Jr. (2007) Phytoplankton blooms in the Ross Sea, Antarctica:  
1122 Interannual variability in magnitude, temporal patterns, and composition. *Journal of*  
1123 *Geophysical Research* **112**, C08013, doi: 10.1029/2006JC003816.
- 1124 Petrov, A.F. and Tatarnikov, V.A. (2010) New data on migrations of Antarctic toothfish  
1125 *Dissostichus mawsoni* in the Dumont d’Urville Sea in the 2008/2009 Season. *Journal of*  
1126 *Ichthyology* **50**, 140–141.
- 1127 Pinkerton, M., Hanchet, S. and Bradford-Grieve, J. (2007) Finding the role of Antarctic toothfish  
1128 in the Ross Sea ecosystem. *Water and Atmosphere* **15**, 20-21.
- 1129 Pinkerton, M., Bradford-Grieve, J. and Hanchet, S. (2010) A balanced model of the food web of  
1130 the Ross Sea, Antarctica. *CCAMLR Science* **17**, 1-31.
- 1131 Pitman, R.L., and Ensor, P. (2003) Three forms of killer whales (*Orcinus orca*) in Antarctic waters.  
1132 *Journal of Cetacean Research and Management* **5**, 1–9.
- 1133 Ponganis, P.J. and Stockard, P.K. (2007) The Antarctic toothfish: how common a prey for  
1134 Weddell seals. *Antarctic Science* **19**, 41-42.

- 1135 Rotella, J.A., Link, W., Nichols, J.D., Hadley, G., R.A. Garrott and Proffitt, K.M. (2009) An  
1136 evaluation of density-dependent and density-independent influence on population growth  
1137 rates in Weddell seals. *Ecology* **90**, 975-984.
- 1138 Russell, J.L., Dixon, K.W., Gnanadesikan, A., Stouffer, R.J. and Toggweiler, J.R. (2006) The  
1139 Southern Hemisphere westerlies in a warming world: propping open the door to the deep  
1140 ocean. *Journal of Climate* **19**, 6382–6390.
- 1141 Scheffer, M., Carpenter, S. and Young, B. (2005) Cascading effects of overfishing marine  
1142 systems. *Trends in Ecology and Evolution* **20**, 579-581.
- 1143 Shust, K.V and Kozlov, A.N. (2006) Changes in size composition of the catches of toothfish  
1144 *Dissostichus eleginoides* as a result of longterm long-line fishing in the region of South  
1145 Georgia and Shag Rocks. *Journal of Ichthyology* **46**, 752-758.
- 1146 Siniff, D.B., Garrott, R.A., Rotella, J.J., Fraser, W.R. and Ainley, D.G. (2008) Projecting the effects  
1147 of environmental change on Antarctic seals. *Antarctic Science* **20**, 425-435.
- 1148 Smith, W.O., Jr., Ainley, D.G., Cattaneo-Vietti, R. and Hofmann, E.E. (2011) The Ross Sea  
1149 continental shelf: regional biogeochemical cycles, trophic interactions, and potential future  
1150 changes. In: *Antarctica: An Extreme Environment in a Changing World*. J. Wiley and Sons,  
1151 London.
- 1152 Stammerjohn, S.E., Martinson, D.G., Smith, R.C., Yuan, X. and Rind, D. (2008) Trends in Antarctic  
1153 annual sea ice retreat and advance and their relation to El Nino Southern Oscillation and  
1154 Southern Annular Mode variability. *Journal of Geophysical Research* **113**, C03S90,  
1155 doi:10.1029/2007JC004269.

- 1156 Stirling, I. (1971) Population aspects of Weddell seal harvesting at McMurdo Sound, Antarctica.  
1157 *Polar Record* **15**, 652–667.
- 1158 Testa, J.W., Oehlert, G., Ainley, D.G., Bengtson, J.L., Siniff, D.B., Laws, R.M. and Rounsevell, D.  
1159 (1991) Temporal variability in Antarctic marine ecosystems: periodic fluctuations in the phocid  
1160 seals. *Canadian Journal of Fisheries and Aquatic Science* **48**, 631-639.
- 1161 Testa, J.W., Siniff, D.B., Ross, M.J. and Winter, J.D. (1985) Weddell seal-Antarctic cod  
1162 interactions in McMurdo Sound, Antarctica. In: Siegfried, W.R., Condy, P.R. and Laws, R.M.  
1163 (eds) *Antarctic Nutrient Cycles and Food Webs*. Springer-Verlag, Berlin and Heidelberg, pp  
1164 561–565.
- 1165 Thompson, D.W.J., and Solomon, S. (2002) Interpretation of recent Southern Hemisphere  
1166 climate change. *Science* **296**, 895–899.
- 1167 Vacchi, M., La Mesa, M., Dalu, M. and Macdonald, J. (2004) Early life stages in the life cycle of  
1168 Antarctic silverfish, *Pleuragramma antarcticum* in Terra Nova Bay, Ross Sea. *Antarctic  
1169 Science* **16**, 299-305.
- 1170 Whitehead, H. (2000) Density dependent habitat selection and the modeling of sperm whale  
1171 (*Physeter macrocephalus*) exploitation. *Canadian Journal of Fisheries and Aquatic Science*  
1172 **57**, 223-230.
- 1173 Whitehead, H., Christal, J. and Dufault, S. (1997) Past and present whaling and the rapid decline  
1174 of sperm whales off the Galapagos Islands. *Conservation Biology* **11**, 1387-1396.
- 1175 Wilson, P.R., Ainley, D.G., Nur, N. Jacobs, S.S. Barton, K.J. Ballard G. and Comiso, J.C. (2001)  
1176 Adélie Penguin population change in the Pacific Sector of Antarctica: Relation to sea-ice

- 1177 extent and the Antarctic Circumpolar Current. *Marine Ecology Progress Series* **213**, 301-
- 1178 309.
- 1179 Yukhov, V.L. (1970) New data on the distribution and biology of *Dissostichus mawsoni* Norm. in
- 1180 Antarctic high latitudes. *Journal of Ichthyology* **10**, 422-424.
- 1181 Zwally, H.J., Comiso, J.C., Parkinson, C.L., Cavalieri, D.J. and Gloersen, P. (2002) Variability of
- 1182 Antarctic sea ice 1979–1998. *Journal of Geophysical Research* **107**, doi:
- 1183 10.1029/2000JC000733.
- 1184
- 1185



1186  
 1187 **Table 1** Summary of data used to calculate changes in fish length, condition and catch per unit effort  
 1188 (based on hooks deployed). If the field “total hooks fished” is blank, then no effort data were available  
 1189 for that year. Other aspects of “standardizing” the effort, include: vessel = same each year (fish hut);  
 1190 vessel operator = same (DeVries); fish measurer = same (DeVries); gear = same (vertical set line, number  
 1191 of hooks varied, as indicated); location = same, i.e. within 2 km radius off Cape Armitage, except in a  
 1192 couple of years as noted in Fig 1; depth = 415-495 m in all years.

Year	Dates	24 h fishing periods	N fish caught (measured)	Total hooks fished	Fish per hook
1972	9/15-12/13	90	96 (96)		
1973	9/11-12/6	86	184 (184)		
1974	9/8-12/19	105	470 (470)		
1975	9/16-12/29	105	346 (331)	2525	0.137
1976	12/1-1/2	33	167 (160)	485	0.344
1977			-		
1978	10/15-12/21	67	349 (341)	1113	0.314
1979	10/18-12/11	55	261 (255)	938	0.278
1980	10/20-12/15	57	202 (184)	1373	0.147
1981	10/13-12/20	69	268 (255)	1986	0.135
1982	10/12-12/22	71	499 (218)	2939	0.170
1983	10/13-12/20	69	553 (546)	3155	0.175
1984	10/1-12/8	69	205 (200)	1338	0.153
1985	10/15-12/20	67	318 (315)	1690	0.188
1986	11/2-12/30	58	151 (141)	783	0.193
1987	10/11-12/10	60	412 (407)	1388	0.297
1988	9/4-12/11	98	294 (294)		
1989	9/7-12/6	90	182 (178)	1185	0.154
1990	8/31-12/8	100	283 (265)	1190	0.238
1991	11/19-11/30	11	7 (7)		
1992	9/9-11/18	70	32 (31)	231	0.139
1993			-		
1994			-		
1995			-		
1996			-		
1997	10/11-12/14	64	72 (72)	442	0.163
1998	10/9-12/9	61	243 (243)		
1999			-		
2000			-		
2001	11/14-12/11	27	76 (76)	499	0.144
2002	10/18-12/15	59	91 (91)		
2003	10/3-12/22	80	17 (17)		
2004	10/11-12/22	74	19 (19)		

2005	1/10-1/14	5	0		
2006	*	*	*		
2007	10/18-11/30	73	2 (0)	660	0.003
2008			-		
2009	12/1-12/20	20	0	117	0.000
2010	10/26-12/3	38	10 (7)	162	0.043

1193 \* Another researcher (G. Hofmann, pers. comm.) at site off Cape Armitage (same location as most others  
 1194 shown in Fig 1) in late Nov 2001, 10 24-h sets (15 hooks/set), caught 71 fish (included in the 2001 total); in  
 1195 late Nov 2006, approximately same site, 7 24-h sets (12 hooks/set), caught 0 fish.

1196

1197

1198

1199

1200

1201

**Table 2** Weighted regression analysis of percentile total length (A, B) and mean condition index, *K* (C); shown are statistics for the two-part spline model and for the difference in slope 1972 to 1992 and from 1992 to 2010 (N = number of years analyzed).

	N	Df	F	P	Adj $r^2$
A) TL 75 <sup>th</sup> percentile					
2-part spline model	27	2, 24	11.07	0.004	0.436
Test of difference in slope		1, 24	11.75	0.002	
B) TL 95 <sup>th</sup> percentile					
2-part spline model	27	2, 24	5.09	0.014	0.24
Test of difference in slope		1, 24	7.32	0.012	
C) Mean condition, <i>K</i>					
2-part spline model	27	2, 24	13.77	0.001	0.496
Test of difference in slope		1, 24	27.07	0.0001	

1202

1203

1204 **Table 3** Regression analysis of total length (A, B) and condition index,  $K$  (C, D) using individuals  
 1205 (N) and comparing data sets with (A, C) and without (B, D) 1991 and 1992. Shown are results of  
 1206 models for 2-part spline (estimated slope up to 1992 plus estimated slope since 1992, with knot  
 1207 at 1992) and for the difference in the two slopes. All models included month of capture (df = 3,  
 1208 for September, October, November, and December).

	N	Df	F	P	Adj $R^2$
A) Total Length, all years					
full model (includes month)	5437	5, 5431	30.15	0.0001	0.0261
2-part spline component		2, 5431	41.42	0.0001	
Test of difference in slope		1, 5431	25.89	0.0001	
B) Total Length, without 1991 & 1992					
full model (includes month)	5399	5, 5393	29.34	0.0001	0.0256
2-part spline component		2, 5393	38.48	0.0001	
Test of difference in slope		1, 5393	23.25	0.0001	
C) Condition, all years					
full model (includes month)	5403	5, 5397	24.66	0.0001	0.0214
2-part spline component		2, 5397	55.94	0.0001	
Test of difference in slope		1, 5397	108.61	0.0001	
D) Condition, without 1991 & 1992					
full model (includes month)	5365	5, 5359	22.98	0.0001	0.0201
2-part spline component		2, 5359	50.66	0.0001	
Test of difference in slope		1, 5359	99.46	0.0001	

1209  
 1210  
 1211  
 1212


1213 **Table 4** Summary of statistics for regression analyses with respect to fish length (75<sup>th</sup> and 50<sup>th</sup>  
 1214 percentiles) and mean condition in relation to environmental variables, and relationship  
 1215 between SOI and Max ice. N = number of years analyzed.

	N	df	F	P	Adj $r^2$
TL 75 <sup>th</sup> percentile-Max ice area 4 yr lag	17	1, 16	4.63	0.048	0.1849
TL 50 <sup>th</sup> percentile-Max ice area 4 yr lag	17	1, 16	8.78	0.009	0.3273
Condition – Min ice area 8 mo lag	20	1, 19	4.99	0.039	0.1734
Condition – Max ice extent 3 yr lag	18	1, 17	4.98	0.040	0.1895
TL 75 <sup>th</sup> percentile – SOI (July-Dec) 4 yr lag	25	1, 24	1.59	0.219	0.0242
SOI – Max ice area same year	29	1, 28	20.5	0.0001	0.4101


1216

1217

1218



1219 **Figure 1** Southern McMurdo Sound showing locations where the set line was positioned  
 1220 seasonally, 1972-2010. Numeral 1 indicates position of fishing site in 2009 and 2010, as well as  
 1221 site of “Penguin Ranch”, where Ponganis and Stockard (2007) recorded numerous seals  
 1222 capturing toothfish in 2003 and 2004; numeral 2 is alongside positions where fishing was  
 1223 otherwise conducted; movement of sites in part forced by shifting location of the sea ice  
 1224 runways (only the two most common runway positions shown). The dashed lines separate  
 1225 zones of fishing success attained by Testa *et al.* (1985; fish per day) as a function of distance  
 1226 from the major Weddell seal breeding location, which is in top right corner of map.  
 1227  
 1228




1229

1230 **Figure 2** Variation in Southern Annular Mode, 1965 – 2011, values on 1 Jan each year (data  
 1231 from <http://www.antarctica.ac.uk/met/gjma/sam.html>).

1232

1233




1234

1235 **Figure 3** Length-frequency of Antarctic toothfish caught in southern McMurdo Sound, 1972-  
1236 2010. Dashed line is the locally weighted regression (lowess, using running-line least squares) of  
1237 length on year.

1238




1239



1240 **Figure 4** Left, change in fish length in the McMurdo Sound catch, expressed as the 75<sup>th</sup>  
1241 percentile of length; right, change in fish length in the McMurdo Sound catch, expressed as the  
1242 95<sup>th</sup> percentile of length, 1972-2010.  
1243

1244


1245



1246 **Figure 5** Left, mean toothfish condition and standard error among fish caught in McMurdo  
1247 Sound, monthly, 1972-2010 (n >5500 fish); right, mean catch per unit effort and standard error  
1248 by month over the period 1975-2010.

1249

1250




1251

1252 **Figure 6** Trends in mean fish condition, K, among fish caught in McMurdo Sound, 1972-2010

1253


1254



1255 **Figure 7** Left, change in prevalence of axe handle fish in the McMurdo Sound toothfish catch,  
1256 1972-2010; right, relationship between proportion of axe handle fish in the catch and maximum  
1257 ice area 4 years earlier.

1258

1259




1260

1261 **Figure 8** Annual variation in fish captures (ln-transformed) per 10 hooks set in McMurdo  
1262 Sound, 1972-2010. Monthly variation in capture rates, after adjusting for month of capture, as  
1263 well as best-supported model for annual variation (stair-step), are shown.

1264


1265



1266 **Figure 9** Fish length as affected by maximum ice area, 4 years previous, for both 75th and 50th  
1267 percentile.

1268

1269




1270 **Figure 10** Left, mean condition index in relation to minimum ice area 8-10 months previously,  
1271 1979-2007; right, fish condition as affected by ice extent three years previously, 1979-2007.

1272

1273

1274



1275 **Figure 11** Left, relationship between maximum ice area and SOI during July-December of same  
1276 year; right, trend in maximum ice extent 3 years previously during the study period (consistent  
1277 with Zwally *et al.* 2002, though a longer time series).