

# 1 Assessment of Gravity Wave Momentum Flux Measurement Capabilities by 2 Meteor Radars Having Different Transmitter Power and Antenna 3 Configurations 4

D. C. Fritts<sup>1</sup>, D. Janches<sup>2</sup>, W. K. Hocking<sup>3</sup>, N. J. Mitchell<sup>4</sup>, and M. J. Taylor<sup>5</sup> <sup>1</sup>NorthWest Research Associates, Colorado Research Associates division, Boulder, CO <sup>2</sup>NASA/Goddard Space Flight Center, Space Weather Laboratory, Greenbelt, MD <sup>3</sup>Department of Physics, University of Western Ontario, London, ONT, Canada <sup>4</sup>Centre for Space, Atmos. & Oceanic Science, Dept. of Electronic & Electrical Engineering, University of Bath, UK <sup>5</sup>Department of Physics, Utah State University, Logan, UT <sup>12</sup> <sup>13</sup> <sup>14</sup> Corresponding author: D. Fritts <sup>16</sup> <sup>17</sup> Email: [dave@cora.nwra.com](mailto:dave@cora.nwra.com) <sup>18</sup> Phone: 303-415-9701, ext. 205 <sup>19</sup>

<sup>21</sup> <sup>22</sup> <sup>23</sup> Submitted to Journal of Geophysical Research – Atmospheres <sup>24</sup> 15 November 2011

## 27 Abstract

28 Measurement capabilities of five meteor radars are assessed and compared to determine 29 how well radars having different transmitted power and antenna configurations perform in 30 defining mean winds, tidal amplitudes, and gravity wave (GW) momentum fluxes. The five 31 radars include two new-generation meteor radars on Tierra del Fuego, Argentina ( $53.8^{\circ}\text{S}$ ) and on 32 King George Island in the Antarctic ( $62.1^{\circ}\text{S}$ ) and conventional meteor radars at Socorro, New 33 Mexico ( $34.1^{\circ}\text{N}$ ,  $106.9^{\circ}\text{W}$ ), Bear Lake Observatory, Utah ( $\sim 41.9^{\circ}\text{N}$ ,  $111.4^{\circ}\text{W}$ ), and Yellowknife, 34 Canada ( $62.5^{\circ}\text{N}$ ,  $114.3^{\circ}\text{W}$ ). Our assessment employs observed meteor distributions for June of 35 2009, 2010, or 2011 for each radar and a set of seven test motion fields including various 36 superpositions of mean winds, constant diurnal tides, constant and variable semidiurnal tides, 37 and superposed GWs having various amplitudes, scales, periods, directions of propagation, 38 momentum fluxes, and intermittencies. 39 Radars having higher power and/or antenna patterns yielding higher meteor counts at 40 small zenith angles perform well in defining monthly and daily mean winds, tidal amplitudes, 41 and GW momentum fluxes, though with expected larger uncertainties in the daily estimates. 42 Conventional radars having lower power and a single transmitting antenna are able to describe 43 monthly mean winds and tidal amplitudes reasonably well, especially at altitudes having the 44 highest meteor counts. They also provide qualitative estimates of GW momentum fluxes at the 45 altitudes having the highest meteor counts; however, these estimates are subject to uncertainties 46 of  $\sim 20$  to  $50\%$  and uncertainties rapidly become excessive at higher and lower altitudes. 47 Estimates of all quantities degrade somewhat for more complex motion fields. 48 49 50 **1**

## Introduction

51 Gravity waves (GWs) play significant roles in atmospheric dynamics, chemistry, 52 microphysics, and plasma processes extending from Earth's surface into the thermosphere and 53 ionosphere. At lower altitudes, GWs contribute to boundary layer and mesoscale structures, 54 organize convection, induce turbulence and mixing of relevance to weather prediction and 55 aircraft at flight altitudes. In the middle stratosphere, GWs induce polar stratospheric clouds that 56 contribute to ozone destruction. Throughout the atmosphere, but especially at higher altitudes, 57 GWs systematically influence the large-scale circulation and thermal structure via energy and 58 momentum transport and deposition. These influences are typically slow and systematic at lower 59 altitudes, but may be local and strong accompanying severe events. In the mesosphere and lower 60 thermosphere (MLT), GWs lead to closure of the mesospheric jets, large departures from 61 radiative equilibrium, strong turbulence and mixing of heat, momentum, and constituents, and 62 significant interactions with, and influences on, tides and planetary waves. GWs are expected to 63 have similar, but still largely unknown, effects higher in the thermosphere and ionosphere. 64 Indeed, the vertical transport of horizontal momentum by GWs, and its deposition accompanying 65 GW dissipation, account for the major influences of these motions on the mean and large-scale 66 atmospheric circulation, structure, and variability. Reviews of these various dynamics 67 emphasizing the higher altitudes are provided by *Hines* [1960], *Lighthill* [1978], *McIntyre* 68 [1989], *Hocke and Schlegel* [1996], *Nappo* [2002], *Fritts and Alexander* [2003], *Kim et al.* 69 [2003], *Fritts et al.* [2006a], and *Fritts and Lund* [2011]. 70 The various responses to momentum transport by GWs noted above make this a key 71 quantity in understanding and modeling atmospheric structure at all altitudes. Consequently, 72 significant efforts have addressed these dynamics employing theoretical, modeling, and

73 observational methods, typically focusing on the higher altitudes where the effects of these  
74 dynamics are most pronounced [Fritts and Alexander, 2003]. Further quantification of these 75 dynamics  
throughout the atmosphere also remain a major research need, given the important of 76 their proper  
parameterization for weather and climate forecasting [Kim et al., 2003]. 77 Prior to new meteor radar capabilities,  
direct measurements of GW momentum fluxes 78 have only been possible with Doppler radars or lidars having  
symmetric, relatively narrow, off79 zenith coplanar beam pairs [Vincent and Reid, 1983; Reid and Vincent 1987;  
Fritts and Vincent, 80 1987; Fukao et al., 1988; Reid et al., 1988; Fritts and Yuan, 1989; Fritts et al., 1990, 1992, 81  
2006b; Fritts and Janches, 2008; Sato, 1990, 1993, 1994; Tsuda et al., 1990; Wang and Fritts, 82 1990, 1991;  
Hitchman et al., 1992; Nakamura et al., 1993; Murayama et al., 1994; Murphy and 83 Vincent, 1993, 1998; Acott et  
al., 2009] or via *in situ* measurements by aircraft at lower altitudes 84 or chaff measurements in the MLT [e.g., Lilly  
and Kennedy, 1973; Lilly et al., 1982; Brown, 85 1983; Meyer et al., 1989; Nastrom and Fritts, 1992; Smith et al.,  
2008]. Indirect methods have 86 nevertheless contributed to quantification of, or constraints on, momentum fluxes  
employing 87 airglow [e.g., Swenson et al., 1999; Fritts et al., 2002; Espy et al., 2004, 2006], balloon [e.g., 88  
Hertzog and Vial, 2001; Hertzog et al., 2008], and satellite measurements [e.g., Ern et al., 2004]. 89 These various  
measurements have revealed a variety of responses, including systematic seasonal90 mean momentum fluxes  
accounting for large-scale forcing at lower and higher altitudes, 91 responses to specific sources and source regions,  
episodic and strong forcing accompanying 92 GWs that are transient and localized, but achieve large amplitudes, and  
significant filtering and 93 interactions with larger-scale mean, tidal, and planetary wave (PW) motions. 94 Ideally,  
routine momentum flux measurements would provide continuous sensitivity to 95 the magnitudes that are  
dynamically important and also capture the temporal variations

96 accompanying strong, but transient, events that may nevertheless contribute significantly to  
97 hourly, daily, monthly, and seasonal means. Improved understanding and parameterization of 98 GW  
influences also impose a need for such measurements at a range of locations spanning 99 representative source  
regions for GWs and the environments in which they propagate. 100 Unfortunately, those systems that do (or  
did) have such measurement capabilities are very 101 limited, their costs are very high (typically ~\$1 to 10M,  
with several much more costly), very 102 few measure continuously, and only two (to our knowledge), sodium  
resonance lidars at the 103 Arctic Lidar Observatory for Middle Atmosphere Research (ALOMAR) at  $69.3^{\circ}\text{N}$  in  
Norway 104 and the Andes Lidar Observatory (ALO) at  $30^{\circ}\text{S}$  in Chile, are even at the edges of regions of 105  
identified high seasonal GW activity extending into the stratosphere and MLT. So there are 106 considerable  
motivations for exploring other measurement techniques that may offer the 107 potential to significantly expand  
such measurements with much cheaper systems. Such systems 108 should also have the potential for continuous  
measurements; optical systems, whether active 109 (lidar) or passive (airglow, FPI, etc.) cannot provide full  
diurnal and annual measurements, 110 independent of weather, suggesting that only radar measurements are  
likely to address this need. 111 The Southern Argentina Agile MEteor Radar (SAAMER) [*Fritts et al.*, 2010a,  
hereafter 112 *F10a*] was specifically designed to add a potential for GW momentum flux measurements to the  
113 conventional meteor radar capabilities for mean, tidal, and PW measurements for the reasons 114 noted  
above. This was accomplished by devising a radar beam pattern that yielded a majority of 115 meteors at zenith  
angles  $<50^{\circ}$  and employing high power to achieve as high a meteor rate as 116 possible at these angles.  
SAAMER was installed at Rio Grande on Tierra del Fuego, Argentina 117 ( $53.8^{\circ}\text{S}$ ,  $67.7^{\circ}\text{W}$ ) and became  
operational in May 2008. SAAMER momentum flux measurement 118 capabilities were evaluated using  
observed meteor distributions for September 2008, a series of

119 test motion fields including prescribed mean winds, diurnal and semidiurnal tide winds, and  
120 GWs having various spatial and temporal scales [Fritts *et al.*, 2010b, hereafter *F10b*], and the 121 analysis  
procedure suggested by Hocking [2005] employing a “full-field” fit to the various 122 winds, variances, and  
covariances characterizing the superposed mean winds, tides, and GWs. 123 These suggested that for real wind fields  
having similar composition and character to the test 124 fields, SAAMER should be expected to provide very good  
definition of the mean and tidal winds 125 and reasonably good definition of GW momentum fluxes where meteor  
count rates are sufficient. 126 The apparent success of SAAMER in measuring GW momentum fluxes in addition  
to 127 mean, tidal, and PW winds was the motivation for a SAAMER clone, the Drake Antarctic Agile 128 MEteor  
Radar (DrAAMER) that was installed at the Brazilian Antarctic Comandante Ferraz 129 Base ( $62.1^{\circ}\text{S}$ ,  $58.7^{\circ}\text{W}$ ) in  
March 2010. Comparisons of the mean and tidal winds and the GW 130 momentum fluxes measured by SAAMER  
and DrAAMER by Fritts *et al.* [2011, hereafter *F11*] 131 demonstrated close correspondence between mean and  
tidal winds at the two sites, as well as 132 significant similarities between the inferred GW momentum fluxes  
between years and sites from 133 April to June of 2010 and 2011. This agreement suggested significant confidence  
in the GW 134 momentum flux measurement potential for both radars, given the tests performed with SAAMER 135  
data. Nevertheless, we believe it is important to subject DrAAMER to the same tests applied to 136 SAAMER. We  
also believe that applications of these tests to additional meteor radars having 137 different characteristics  
(particularly power and beam configuration) may help us identify what 138 radar capabilities are required to achieve  
GW momentum flux measurements that are sufficiently 139 accurate to be valuable to the community. For example,  
the Hocking [2005] analysis method was 140 tested with meteor radar data from Socorro, NM and Resolute Bay,  
Canada and judged to 141 provide reasonable two-month estimates (though without validation). Other meteor radars  
have

142 also been employed for GW variance, momentum flux, and tidal modulation studies and suggest  
143 that more radars may provide enhanced measurement capabilities with testing and suitable 144 analysis  
techniques [Kumar et al., 2007; Antonita et al., 2008; Clemesha et al., 2009; Mitchell 145 and Beldon, 2009; Placke  
et al., 2011]. A related study by *Vincent et al.* [2010] employed a 146 Monte Carlo analysis of measurement  
capabilities assuming a conventional meteor radar beam 147 pattern and expected radial velocity and angle-of-arrival  
uncertainties and concluded that 148 accurate momentum flux estimates with such systems would likely require long  
averages. 149 Our goals in this paper are to evaluate SAAMER and DrAAMER GW momentum flux 150  
measurement capabilities relative to several other conventional meteor radars in order to 151 determine 1) what radar  
parameters are required to enable such measurements, 2) whether other 152 radars can also provide credible GW  
momentum flux measurements in their current 153 configurations, or 3) whether system upgrades are needed to  
enable these capabilities. The other 154 radars employed for this assessment are at Socorro (SRO), New Mexico  
(34.1°N, 106.9°W), Bear 155 Lake Observatory (BL), Utah (~41.9°N, 111.4°W), and Yellowknife (YKF), Canada  
(62.5°N, 156 114.3°W) (see *Hocking*, 2005, and *Hocking and Kishore Kumar*, 2011, respectively, for more 157  
detailed descriptions of the SRO and YKF radars). To address these goals, our paper is structured 158 as follows.  
Section 2 provides a summary of the characteristics of the radars employed for this 159 assessment and the analysis  
methods used to evaluate the performance of each. Test motion 160 fields, which include mean winds, diurnal and  
semidiurnal tides, and/or GWs having various 161 characteristics are described in Section 3. The performance of the  
five radars for the various tests 162 employing real spatial and temporal sampling for each is described in Section 4.  
Section 5 163 summarizes our findings and discusses measurement accuracies relative to other assessments. 164 Our  
conclusions are presented in Section 6.

### Characteristics of Radars Employed for this Study

166 We employ five different meteor radars for our assessment of GW momentum flux 167 measurement capabilities in this study. Two of these were specifically designed intending to 168 provide this capability (*i.e.*, SAAMER and DrAAMER); the other three are conventional meteor 169 radars having lower peak power and a single transmitting antenna that had not anticipated such 170 measurements (*i.e.*, the SKiYMET meteor radars previously installed at Socorro, NM, Bear Lake 171 Observatory, UT, and Yellowknife, Canada; see Table 1). The relative locations of the five 172 radars and the meteor distributions for each are shown for one day during the month in Figure 1. 173 Meteor distributions in altitude, by day throughout the month considered, and by hour 174 throughout a composite day for each radar are shown for comparison in Figure 2. The meteor 175 distributions in Figure 1 all exhibit nulls in the polar diagrams that arise from excluding meteors 176 at range increments corresponding to the pulse repetition frequency (PRF) in order to exclude 177 contamination due to ground clutter. SAAMER and DrAAMER use a smaller PRF, and thus 178 have these nulls at larger spacing. The statistics in Figure 2 include meteors at zenith angles from 179 15 to 60° for all radars. As expected, meteor counts are significantly larger for the radars having 180 higher power, and higher at smaller zenith angles for SAAMER and DrAAMER than for the 181 radars with single-antenna TX systems. They also vary significantly throughout the month, 182 especially at SAAMER and DrAAMER, and exhibit large diurnal variations at all radars. 183 The GW momentum flux measurement potential of SAAMER was previously evaluated 184 for a number of test fields by *F10b*. Comparisons of SAAMER and DrAAMER measurements 185 for April, May, and June of 2010 and 2011 were presented by *F11*, and suggested that 186 DrAAMER momentum flux estimates provided a consistent picture of GW momentum fluxes 187 accompanying strong GW sources over the Drake Passage, but did not evaluate the DrAAMER



188 momentum flux measurement capability directly. An initial application of the method employed  
189 here was also used by *Hocking* [2005] to estimate two-monthly GW momentum fluxes by 190 SkiYMET meteor  
radars at Socorro, NM and Resolute Bay, Canada. These estimates were seen 191 to be roughly consistent with  
expected values, but were not evaluated in detail. However, a 192 systematic evaluation of the relative momentum  
flux measurement potential of these various 193 radars with specific test fields has not been performed. This is the  
goal here, and the 194 characteristics of the five radars are described for comparison in Table 1. We employ meteor  
195 distributions obtained during June 2011 for KGI and TdF, June 2010 for SRO and BL, and June 196 2009 for  
YKF, for which meteor counts tend to be larger at northern than at southern latitudes. 197 **3 Specification of Test**  
**Motion Fields** 198 We showed in *F10b* that the SAAMER beam pattern and meteor counts enable relatively 199  
high-precision measurements of mean winds and tides over fairly short intervals, and that GW 200 momentum fluxes  
can be estimated with reasonable accuracies where meteor counts are high, 201 even when the large- and small-scale  
motion field is variable on multiple time scales. The first 202 measurements over DrAAMER and comparisons with  
those over SAAMER by *F11* suggest that 203 the same can likely be said for DrAAMER. Here we repeat the tests  
previously applied to 204 SAAMER for DrAAMER and the three conventional meteor radars at Socorro, MN, Bear  
Lake 205 Observatory, UT, and Yellowknife, Canada in order to evaluate DrAAMER capabilities more 206  
completely and determine whether, and under what conditions, useful GW momentum flux 207 measurements may  
also be possible with the other conventional meteor radars. 208 Following *F10b*, we employ real meteor spatial and  
temporal distributions observed by 209 SAAMER, DrAAMER, and the three other meteor radars during various  
months to evaluate the 210 measurement capabilities of each. Radial velocities at each meteor location and time for  
each

211 radar and each month assessed are specified by each of seven test velocity fields. Each test field 212 includes superposed mean, tidal, and/or GW velocity fields with constant and/or spatially- and 213 temporally-variable tidal and GW amplitudes that are intended to be representative of the scales, 214 amplitudes, and momentum fluxes in the MLT over SAAMER and DrAAMER. Mean, tidal, and 215 GW parameters defining these motion fields for each case are listed in Table 2. The test fields 216 are those employed by *F10b*, range from highly idealized and stationary to spatially and 217 temporally modulated at large and small scales, and have the following components: 218  $U(x,y,z,t) = U_M + U_D(z,t) \sin(2\pi t/T_D) + U_{SD}(z,t) \sin(2\pi t/T_{SD})$  219  $+ U_{GW1}(x,y,z,t) \sin(k_1x + l_1y + m_1z - 2\pi t/T_{GW1})$  220  $+ U_{GW2}(x,y,z,t) \sin(k_2x + l_2y + m_2z - 2\pi t/T_{GW2})$  221  $+ U_{GW3}(x,y,z,t) \sin(k_3x + m_3z)$  (1) 222

$V(x,y,z,t) = V_M - V_D(z,t) \cos(2\pi t/T_D) - V_{SD}(z,t) \cos(2\pi t/T_{SD})$  223  $+ V_{GW1}(x,y,z,t) \sin(k_1x + l_1y + m_1z - 2\pi t/T_{GW1})$  224  $+ V_{GW2}(x,y,z,t) \sin(k_2x + l_2y + m_2z - 2\pi t/T_{GW2})$  225  $+ V_{GW4}(x,y,z,t) \sin(l_4y + m_4z)$  (2) 226  $W(x,y,z,t) =$

$W_{GW1}(x,y,z,t) \sin(k_1x + m_1z - 2\pi t/T_{GW1})$  227  $+ W_{GW2}(x,y,z,t) \sin(k_2x + l_2y + m_2z - 2\pi t/T_{GW2})$  228  $+ W_{GW3}(x,y,z,t)$  229  $\sin(k_3x + m_3z) + W_{GW4}(x,y,z,t) \sin(l_4y + m_4z)$  (3)

230 Note that we employ the same test cases defined by F10b, but these Eqs. (1-3) correct several  
231 typos appearing in the equations presented in *F10b*. 232 Each test field includes some or all of the following  
components: 233 a. zonal and meridional mean winds,  $U_M$  and  $V_M$ , 234 b. diurnal and semidiurnal tides having zonal  
and meridional amplitudes of  $(U_D, V_D)$  235 and  $(U_{SD}, V_{SD})$  assumed to rotate counterclockwise with time (assuming  
a southern 236 hemisphere location), and which may have either constant or varying amplitudes with 237 increasing  
altitude, 238 c. traveling GWs having amplitudes  $(U_{GW}, V_{GW}, W_{GW})$ , zonal, meridional, or oblique 239  
propagation, spatial and temporal variability, correlated horizontal and vertical 240 motions, and constant or variable  
momentum fluxes, and 241 d. stationary mountain waves (MWs) having zonal and meridional propagation, only 242  
spatial variability, correlated horizontal and vertical motions, and constant 243 momentum fluxes. 244 In each case,  
the test field amplitudes were chosen to correspond roughly to measured 245 values over SAAMER and DrAAMER  
on TdF and KGI, which include large semidiurnal tides 246 and GW momentum fluxes [*F10a*, *F10b*, *F11*]. Large  
semidiurnal tide amplitudes at these sites 247 are consistent with expectations of the most recent version of the  
Global-Scale Wave Model 248 (GSWM-09) [*Zhang et al.*, 2010a, b, *F11*]. Large GW momentum fluxes are  
suggested by the 249 major global hotspot of GW activity in the stratosphere and lower mesosphere centered over  
the 250 Southern Andes, Drake Passage, and Antarctic Peninsula [*F10a*, and references therein]. 251 **4 Evaluation  
of Radar Measurement Capabilities**

252 We describe here the performance of each of the five radars for the seven test cases 253 described above. Mean winds, diurnal and semidiurnal tidal amplitudes, and GWs having 254 prescribed spatial and/or temporal variability and momentum fluxes propagating zonally, 255 meridionally, or at other azimuths are defined by Eqs. (1-3) for the seven cases listed in Table 2. 256 Specified and recovered (*i.e.*, the velocity fields inferred from our S-transform tidal fits and the 257 Hocking statistical analysis) profiles of the mean winds, diurnal and semidiurnal tides, and GWs 258 momentum fluxes, and their daily and composite-day hourly variability, where appropriate, are 259 discussed separately for each case below. 260 ***b.1. Case 1*** 261 Specified fields, and those recovered using observed meteor distributions in space and 262 time for each radar for June 2009, 2010, or 2011 as described above, are shown as monthly 263 means for Case 1 in Figure 3. In this case, tides have only temporal variations and GWs have 264 only horizontal variations, with 50 and 100 km wavelengths in the zonal and meridional motions, 265 respectively (see Table 2). 266 As seen in Figure 3, agreement between specified and recovered fields is highly variable 267 among the five radars. As might be expected based on meteor counts, mean winds and diurnal 268 and semidiurnal tide amplitudes are recovered quite well at KGI, TdF, and BL. Mean winds and 269 diurnal tide amplitudes are recovered within a few percent at each of these sites, with slightly 270 greater uncertainties at BL than at the other two. Semidiurnal tide amplitudes are systematically 271 less than specified at all three sites, but only by ~2-3% at TdF and BL, and by ~5% at KGI. 272 Mean winds and tidal amplitudes exhibit similar tendencies and accuracies at SRO and YKF at 273 the central altitudes, but increasing errors at the lowest and highest altitudes where meteor counts 274 are small.

275 GW momentum flux estimates at KGI, TdF, and SRO are likewise accurate within ~10%, 276 except at KGI and TdF below 79.5 km. In contrast, SRO and YKF exhibit errors of ~20% or 277 larger at intermediate altitudes, again with larger errors where meteor counts are small. 278 **b.2. Case 2** 279 Case 2 includes no mean winds or tides and only a zonal stationary GW with prescribed 280 horizontal and vertical phase variations with wavelengths of 50 and 15 km (not physical, but 281 simple, see Table 2). Referring to Figure 4, we see that, as in Case 1, the recovered mean and 282 tidal motions are quite accurate at KGI, TdF, and BL, typically within  $\sim 2 \text{ ms}^{-1}$  or better, except 283 for the tidal amplitudes at BL above  $\sim 92 \text{ km}$ . Estimates at SRO and YKF are slightly less 284 accurate, within  $\sim 5 \text{ ms}^{-1}$ , but nevertheless quite reasonable, except at the highest and lowest 285 altitudes, especially at YKF. 286 Unlike Case 1, all the radars except YKF provided accurate estimates of GW momentum 287 fluxes in this case, with the largest departures of only  $\sim 5 \text{ m}^2 \text{ s}^{-2}$  at the highest and lowest altitudes 288 shown. However, meteor counts at YKF were apparently too small to yield adequate statistics, 289 except at the central altitudes having the highest meteor counts, from 82.5 to 88.5 km. The 290 significant differences in the accuracies of the GW momentum fluxes at all sites and in the mean 291 wind and tidal amplitudes between Cases 1 and 2 suggest a strong influence of large and variable 292 winds on the quantification of GW momentum fluxes, even for persistent and well-defined GWs. 293 These influences will be quantified further below and discussed in some detail in Section 5. 294 **b.3. Case 3** 295 Case 3 considers a motion field varying from that of Case 2 only in having two 296 orthogonal GWs, each of which has no phase variation in altitude or time and with the GW 297 propagating in the zonal plane having a large vertical velocity of  $10 \text{ ms}^{-1}$  (again not physical, but

298 simple). Results displayed in Figure 5 show that mean motions, tidal amplitudes, and GW  
299 momentum fluxes are again described well with the meteor distributions for KGI, TdF, and BL, 300 and the  
mean motions and GW momentum fluxes are described reasonably at SRO and at YKF, 301 but with increasing  
errors in momentum fluxes at the higher altitudes. However, tidal amplitudes 302 now exhibit larger errors at SRO  
and YKF than seen in Case 2, with these motions reasonably 303 defined at ~85 to 92 km at YKF and in the zonal  
component at SRO, but poorly defined at lower 304 and higher altitudes at both these sites and even at central  
altitudes in the meridional components 305 at SRO. This cannot be a result of insufficient sampling of the mean and  
tidal motions, as they 306 are all zero. Instead, the cause appears to be the larger GW vertical velocities that are not  
307 sufficiently averaged in defining large-scale (horizontal) winds, despite the validity of the 308 momentum flux  
estimates for stationary GWs at these sites (see further discussion in Section 5). 309 **b.4. Case 4** 310 We now  
consider a more complex superposition of mean, tidal, and GW fields given by 311 the sum of the mean and tidal  
motions in Case 1 and both stationary and propagating GWs in 312 both the zonal and meridional planes (see Table  
2). This case differs from the previous cases in 313 that both zonal and meridional GW momentum fluxes now have  
contributions from stationary 314 and propagating GWs having different spatial structures (with horizontal  
wavelengths of 30, 40, 315 50, and 100 km). Case 4 also includes both large semidiurnal tide amplitudes and GW  
vertical 316 velocities that were suggested above to contribute to errors in the estimates of mean and tidal 317  
amplitudes and GW momentum fluxes. Depending on the phases of the four GWs, vertical 318 motions may be ~15  
ms<sup>-1</sup> or larger in this case. 319 Results displayed in Figure 6 show that errors in estimates of the mean winds and tidal  
320 amplitudes are comparable (and small) for Cases 1 and 4 at KGI, TdF, and BL. Semidiurnal tide

321 amplitudes are very slightly underestimated, with the largest errors of ~5% at KGI (note the  
322 different scales in the third columns of Figures 3 and 6). As seen in Case 1, mean wind and tidal 323 amplitude  
estimates at SRO and YKF for Case 4 are also reasonable at central altitudes, but 324 exhibit greater errors at lower  
and/or higher altitudes and at central altitudes in the meridional 325 diurnal tide amplitudes. 326 Turning now to  
GW momentum flux estimates for Case 4, we see that these are very 327 good at KGI, TdF, and BL, with  
comparable errors ( $\sim$ a few  $\text{m}^2 \text{s}^{-2}$ ) except at the highest and 328 lowest altitudes where they are somewhat larger.  
Similar estimates are also obtained at SRO and 329 YKF at central altitudes (with errors for Case 4 of  $\sim 10 \text{ m}^2 \text{ s}^{-2}$  or  
less), with much larger errors 330 occurring at lower and higher altitudes. Apparently, mean winds and large tidal  
amplitudes do 331 not prevent accurate estimates of GW momentum fluxes when meteor counts are sufficiently 332  
high and GWs are sufficiently sustained (whether stationary or propagating) to enable sampling 333 of all phases  
throughout the month. Large GW amplitudes (especially vertical velocities), 334 however, do cause errors in  
estimates of means winds and tidal amplitudes where meteor counts 335 are not sufficiently high. 336 **b.5. Case 5**  
337 Case 5 explores a superposition of larger mean winds, a larger, but constant, diurnal tide, 338 a semidiurnal tide  
that exhibits both a 10-day amplitude modulation and amplitude growth with 339 altitude, and four superposed GWs.  
The latter include 1) stationary GWs having zonal and 340 meridional orientations, constant amplitudes and  
momentum fluxes, and horizontal wavelengths 341 of 30 and 40 km and 2) propagating GWs having zonal and  
meridional orientations, horizontal 342 wavelengths of 50 and 100 km, respectively, and amplitude (and momentum  
flux) modulation 343 by, and anti-correlations with, the semidiurnal tide amplitudes.

Results for Case 5 are displayed in four formats. Figure 7 shows monthly mean profiles of mean winds, tidal amplitudes, and GW momentum fluxes as discussed for Cases 1 to 4 above. Figure 8 displays daily estimates of mean winds and diurnal tide amplitudes throughout the month in the presence of variable semidiurnal tide and GW amplitudes. Figure 9 compares daily estimates of semidiurnal tide amplitudes with the specified variations. Hourly estimates of GW momentum fluxes throughout the composite day are compared with the specified variations in Figure 10. Results shown in Figure 7 reveal mean wind estimates (first column) that are very accurate at all radars (within  $\sim 1 \text{ ms}^{-1}$  except at the lowest and highest altitudes at SRO and YKF). Diurnal tide amplitude estimates (second column) are likewise very good at TdF, nearly as good at KGI and BL (within  $\sim 1$  to  $2 \text{ ms}^{-1}$ ), and reasonable at central altitudes at YKF (within  $\sim 2 \text{ ms}^{-1}$ ). Somewhat larger errors ( $\sim 5 \text{ ms}^{-1}$ ) are again seen in the meridional diurnal amplitude at SRO. Mean semidiurnal tide amplitude estimates (third column) are seen to be very precise at TdF and BL, to exhibit slight underestimates at KGI, and to have comparable errors about the specified mean amplitudes at SRO. Larger errors are seen at YKF (as large as  $\sim 5$ - $10 \text{ ms}^{-1}$ ), but the amplitude growth with altitude is still captured at the central altitudes.

Daily mean wind and diurnal tide amplitude estimates shown for Case 5 in Figure 8 are most accurate at TdF and BL (with RMS uncertainties of  $\sim 1 \text{ ms}^{-1}$  or less) and somewhat less accurate at KGI (with RMS uncertainties of  $\sim 2 \text{ ms}^{-1}$ ), except at the lowest altitude at KGI and the lowest and highest altitudes at BL where meteor counts are small. Daily mean estimates are considerably less accurate at SRO and YKF due to the significantly smaller meteor counts at all altitudes and the asymmetric meteor distributions seen in Figure 1. RMS errors of these estimates range from a few  $\text{ms}^{-1}$  for the zonal components at central altitudes at SRO to  $\sim 100\%$  or greater



367 uncertainties at YKF that render the latter useless in defining day-to-day variability in these  
368 quantities. 369 Estimates of semidiurnal tide zonal and meridional amplitudes for each day in Case 5 at 370  
3-km altitude intervals are displayed in the left and right panels of Figure 9. These estimates are 371 seen to be in  
close agreement with the specified values (dashed lines) at all but the lowest 372 altitude at TdF and the lowest and  
highest altitudes for KGI and BL. Estimates at SRO and YKF, 373 in contrast, are reasonably accurate at the central  
four altitudes, apart from sporadic departures of 374  $\sim 20$  to  $60 \text{ ms}^{-1}$  at these altitudes, primarily at YKF and in the  
meridional component at SRO. 375 Hourly estimates of zonal and meridional GW momentum fluxes for the Case 5  
376 composite day at 3-km altitude intervals are shown in the left and right panels of Figure 10. 377 These are seen  
to be close approximations to the specified values at TdF for all but the lowest 378 and highest altitudes, and even to  
follow the semidiurnal momentum flux modulations at these 379 altitudes, but with larger uncertainties. The results  
for KGI and BL are comparable and also very 380 good, but exhibit somewhat larger fluctuations about the specified  
values than seen at TdF at the 381 central four altitudes altitudes and increasing uncertainties at KGI at 76.5, 79.5,  
and 97.5 km and 382 at BL at 76.5, 94.5, and 97.5 km. Composite day hourly momentum flux estimates for SRO and  
383 YKF are seen to occasionally follow the specified semidiurnal modulation at 85.5, 88.5, and 91.5 384 km, but  
with very large uncertainties. At higher and lower altitudes, these estimates exhibit very 385 large errors. 386 **b.6.**

**Case 6** 387 We now compare the ability of the five radars to define mean and tidal motions and GW 388 momentum  
fluxes for the motion field defined by Case 6 in Table 2. This case includes mean and 389 tidal motions  
representative of higher latitudes and two transient GW packets having propagation 390 to the east and north and that  
occur randomly for 3 and 4 hr each day. The two GWs have zonal 391 and meridional propagation, daily mean  
momentum fluxes of 50 and  $25 \text{ m}^2 \text{ s}^{-2}$ , and periods of 20 392 and 30 min, respectively. Both also have horizontal and  
vertical phase variations with 393 wavelengths given in Table 2. Inferred monthly mean winds, tidal amplitudes, and  
GW 394 momentum fluxes obtained with each radar are shown in Figure 11. Hourly estimates of zonal 395 and  
meridional GW momentum fluxes for a Case 6 composite day at 3-km altitude intervals are 396 shown in the left and  
right panels of Figure 12. Note here that the composite day momentum flux 397 variations are different for each  
radar in each component. Case 6 differs from Cases 1 and 4 in 398 having 1) different, but comparable, mean

motions, 2) stochastic rather than uniform (and 399 stationary) GWs, and 3) vertical phase variations of the GWs.

400 Monthly estimates of mean winds and tidal amplitudes shown in the three left columns of 401 Figure 11 have accuracies and biases that are very similar to those seen previously in Cases 1 402 and 4. Mean wind and diurnal tide amplitude estimates are again very accurate at TdF and nearly 403 as accurate at KGI and BL. Somewhat greater uncertainties are observed at SRO and YKF, but 404 even these are not larger than  $\sim 2 \text{ ms}^{-1}$  except at the lowest and highest altitudes. Semidiurnal tide 405 amplitude estimates are also nearly identical to those obtained in Cases 1 and 4. Amplitudes are 406 again under-estimated by  $\sim 5\%$  at KGI, SRO, and YKF, and by  $\sim 2\%$  at TdF and BL. The similar 407 accuracies achieved in these three cases suggest that the greater intermittency of the larger 408 amplitude GWs in Case 6 does not impact the ability of these radars to provide reasonably 409 accurate definitions of the monthly mean and tidal fields.

410 GW momentum flux estimates shown in the right column of Figure 11 are seen to be 411 relatively more accurate at KGI, TdF, and BL, where uncertainties are  $\sim 10\%$  or less at central 412 altitudes, but approach 20% at KGI and BL at the lowest and/or highest altitudes. Momentum

413 flux estimates are also relatively accurate at SRO and YKF at the central 2 to 4 altitudes having  
414 the highest meteor counts, but exhibit large errors at lower and higher altitudes where meteor 415 counts are low.  
Accuracies at KGI, TdF, and BL are comparable to those seen in Case 1 for 416 which GW amplitudes and  
momentum fluxes are smaller. However, accuracies are not as good 417 as seen in Case 4 where the zonal  
momentum flux is largely determined by the stationary, large 418 amplitude GW contributing the majority of the  
momentum flux. 419 Diurnal variability of the composite day zonal and meridional momentum fluxes for each  
420 radar shown in Figure 12 reinforce the statements above about the monthly mean profiles. The 421 TdF and BL  
radars capture the specified momentum flux distributions extremely well, but with 422 slightly greater uncertainties  
at TdF at the highest altitude, at TdF and BL at the lowest altitude, 423 and a greater loss of sensitivity at the highest  
altitudes at BL. Momentum flux estimates at KGI 424 are slightly less precise than either TdF or BL at central and  
lower altitudes, but are somewhat 425 better than at BL at the higher altitudes. In contrast, momentum flux estimates  
at SRO and YKF 426 are only accurate at the central ~4 and 2 altitudes, respectively, and are susceptible to very  
large 427 uncertainties below and above. These results suggest that high meteor counts (and more uniform 428  
sampling of the motion field) are essential in defining momentum fluxes when GWs are variable 429 and tides  
achieve large amplitudes. 430 In order to assess whether momentum flux estimates are negatively influenced by large  
431 tidal amplitudes, we also performed the same mean, tidal, and GW momentum flux assessments 432 for Case 6,  
but with zero mean winds and tidal amplitudes. This yielded 1) very accurate mean 433 wind estimates (except at the  
lower altitudes at YKF), 2) diurnal and semidiurnal tide amplitude 434 estimates of  $\sim 1 \text{ ms}^{-1}$  at KGI, TdF, and BL and  
 $\sim 2 \text{ ms}^{-1}$  at the central altitudes at SRO and YKF, 435 and 3) momentum flux estimates comparable to or slightly more  
accurate than with the tides

436 present, with the greatest improvements at SRO. Hourly estimates for the composite day without  
437 mean winds and tides are likewise of comparable accuracy to those for the full field specified for 438 in Case 6  
in Table 2. 439 **b.7. Case 7** 440 Case 7 is very similar to Case 6, with the same mean and tidal winds, but including  
two 441 intermittent large-amplitude GWs having propagation to the NE and SE. The GW propagating to 442 the NE  
occurs for 2 hr twice daily at random times; that propagating to the SE occurs for 1 hr 3 443 times daily at random  
times. The GWs have periods of 20 and 15 min and both horizontal and 444 vertical phase variations. Amplitudes  
and wavelengths of each are listed in Table 2. Together, 445 they result in mean zonal and meridional momentum  
fluxes of 75 and  $-25 \text{ m}^2 \text{ s}^{-2}$ , respectively. 446 Inferred monthly mean winds, tidal amplitudes, and GW momentum  
fluxes obtained with each 447 radar for Case 7 are shown in Figure 13. Figure 14 shows hourly estimates of zonal  
and 448 meridional GW momentum fluxes for a Case 7 composite day at 3-km altitude intervals. 449 Monthly mean  
wind, tidal amplitude, and GW momentum flux estimates for Case 450 7 are very similar to those seen in Case 6  
above. Mean wind and diurnal tide amplitude estimates 451 are very accurate at KGI, TdF, and BL but exhibit larger  
errors at SRO and YKF at the highest 452 and lowest altitudes, respectively. Semidiurnal tides are under-estimated  
by  $\sim 5\%$  at KGI, SRO, 453 and YKF, and by  $\sim 2\%$  at TdF and BL, as in Case 6. However, larger errors are seen at  
SRO and 454 YKF at the highest and lowest altitudes, respectively, as seen for the mean winds and diurnal tide 455  
amplitudes. GW momentum flux estimates also exhibit comparable accuracies to those in Case 456 6, suggesting that  
GW packet duration does not have a strong influence on the ability to define 457 these quantities. Finally, as for Case  
6, we also performed these assessments with mean winds 458 and tidal amplitudes specified to be zero. Similar to the  
results for Case 6, this yielded 1) 459 extremely accurate mean wind estimates (except at the lower altitudes at YKF),  
2) diurnal and 460 semidiurnal tide amplitude estimates of  $\sim 1\text{-}2 \text{ ms}^{-1}$  at KGI, TdF, and BL and  $\sim 2\text{-}5 \text{ ms}^{-1}$  at the 461  
central altitudes at SRO and YKF, and 3) momentum flux estimates comparable to or slightly 462 more accurate than  
with the tides present. As in Case 6, hourly estimates for the composite day 463 without mean winds and tides are of  
comparable accuracy to those for the full field specified for 464 in Case 7 in Table 2. 465 **5 Summary and**  
**Discussion** 466 We have assessed the relative measurement accuracies of meteor radars having similar 467  
frequencies (32.55 to 36.9 MHz) but different antenna configurations, peak power, and sampling 468 modes. Our

evaluation was performed with meteor distributions observed by five radars 469 extending from  $62.1^{\circ}$  S to  $62.5^{\circ}$  N obtained during June of 2009, 2010, or 2011 for which meteor 470 statistics tend to be better at northern than southern latitudes. Two of these radars, SAAMER on 471 Tierra del Fuego and DrAAMER on King George Island ( $53.8$  and  $62.1^{\circ}$  S, respectively) were 472 specifically designed for enhanced measurement capabilities for large-scale motions and GW 473 momentum fluxes. This was accomplished in each case with an 8-Yagi transmitting array 474 directing power into 8 lobes centered at  $35^{\circ}$  zenith angles separated by  $45^{\circ}$  in azimuth. The other 475 three radars are conventional meteor radars at northern latitudes (see Figure 1). Our purpose was 476 to determine the relative abilities of different meteor radar configurations to recover specified 477 mean winds, diurnal and semidiurnal tide amplitudes, and GW momentum fluxes. 478 Seven test motion fields included various superpositions of constant mean winds, 479 constant and/or variable tidal amplitudes, and GWs having varying amplitudes, momentum 480 fluxes, scales, periods, propagation directions, and intermittencies. The test fields were sampled 481 according to the observed meteor distributions throughout the month for each radar assuming no 482 radial velocity, range, or angular uncertainties in the measurements. Only meteors at zenith 483 angles between  $15$  and  $60^{\circ}$  were used, due to the large altitude uncertainties accompanying  $\sim 1^{\circ}$  484 zenith angle uncertainties in angle-of-arrival estimates at larger zenith angles. The various fields 485 were estimated in 3-km altitude bins (centered from 76.5 to 94.5 km) using the method described 486 by *F10b*, in which mean winds and tidal amplitudes are determined by S-transform fits and 487 removed from each radial velocity distribution before application of the *Hocking* [2005] method. 488 For each case mean wind, tidal amplitude, and GW momentum flux profiles were determined for 489 monthly and/or daily intervals (as appropriate). Also estimated in cases having diurnal variations 490 in GW momentum fluxes were hourly profiles throughout a composite day. 491 Our evaluation of relative radar performance revealed the following: 492 1) measurement accuracies depend strongly on meteor counts and antenna beam patterns; 493 accuracies improve with higher meteor counts, smaller zenith angles, and more 494 symmetric beam patterns enabled by crossed antennas, 495 2) SAAMER (on TdF), having  $\sim 12,000$  meteors/day between  $15$  and  $60^{\circ}$  zenith angles 496 ( $\sim 120$  meteors/hr in a 3-km altitude bin at  $\sim 90$  km) performed best overall, yielding quite 497 accurate mean wind,

tidal amplitude, and GW momentum flux estimates for all test cases, 498 3) the BL meteor radar and DrAAMER (on KGI) performed comparably and very well, 499 despite DrAAMER having ~30% smaller meteor counts (~8,000 meteors/day compared 500 to ~12,000 total at BL, or ~80 and 120 meteors/hr in a 3-km altitude bin near the peak of 501 the meteor distribution); however, DrAAMER yielded a consistently larger under502 estimate of semidiurnal tide amplitudes (~5% rather than the ~2% seen at BL), and 503 4) the SRO and YKF meteor radars having 6 kW peak power and antenna with only one 504 polarization (having ~ 5,000 and 3,000 meteors/day, respectively, or 50 and 30 505 meteors/hr in 3 km) provided less accurate measurements and over more limited 506 altitudes; they nevertheless described monthly mean winds and tidal amplitudes 507 adequately, with the best results at central altitudes; monthly mean and hourly composite508 day GW momentum flux estimates were generally poor at lower and higher altitudes, but 509 reasonable at ~85 to 90 km. 510 More general conclusions obtained from inter-comparisons of our various cases include 511 the following: 512 5) momentum flux estimates are more accurate when tidal amplitudes are small (or zero) 513 and when GW amplitudes and momentum fluxes are large, 514 6) momentum flux estimates are not significantly impacted by complexity of the GW field, 515 including superposition, multiple scales and frequencies, and intermittency, 516 7) well configured radars (those having symmetric beam patterns (*e.g.*, crossed Yagis) 517 and/or more meteors at smaller zenith angles) capture daily and composite-day hourly 518 variability of total winds and GW MFs well at altitudes where meteor count are high, and 519 8) radars that have asymmetric beam patterns (crossed Yagis) and/or low meteor counts 520 (less than ~25 meteors/hr in 3 km) will likely exhibit significant biases in estimates of 521 daily mean winds and tidal amplitudes; they will likely also exhibit significant biases in 522 estimates of hourly composite-day GW momentum fluxes and variability of these fields 523 on longer time scales. 524 Our assessment of meteor radar performance above assumed no radial velocity, range, or 525 angle-of-arrival measurement uncertainties. Introducing such uncertainties would of course 526 degrade measurement accuracies to some degree in all cases. Here we assess the likely impacts

527 of such measurement uncertainties for the magnitudes of the mean, tidal, and GW motions and  
528 momentum fluxes considered above. 529 If we assume that RMS radial velocity uncertainties are  $\sim 1$  to  $2 \text{ ms}^{-1}$   
(which is a 530 reasonable estimate for SAAMER), then these contributions to measured radial velocities are 531  
significantly less than typical radial velocity magnitudes, given the mean winds and tidal and 532 GW amplitudes  
employed for our assessment. For meteors at the largest zenith angles,  $\sim 60^\circ$ , 533 these uncertainties would contribute  
little to GW momentum flux assessments given their small 534 implied errors in vertical velocity contributions to  
radial velocity variance relative to the large 535 contributions by large horizontal velocities. These uncertainties at  
large zenith angles would also 536 contribute very small errors in estimates of mean and tidal motions, which are  
obtained by 537 averaging large horizontal motions rather than effectively differencing radial velocity variances 538  
for GW momentum flux assessments. 539 At smaller zenith angles, there is a greater potential for influences of radial  
velocity 540 uncertainties, given that these angles provide greater sensitivity to the differential radial 541 velocities  
with azimuth that contribute to the momentum flux determination. The most sensitive 542 radars to such  
uncertainties will be those focusing transmitted power at small zenith angles, 543 specifically SAAMER and  
DrAAMER, both of which achieve maximum meteor counts at  $\sim 35^\circ$  544 zenith angles. 545 To assess qualitatively  
the impacts of radial velocity uncertainties on measurements of 546 GW momentum fluxes, we assume that all  
meteors occur at the  $35^\circ$  zenith angle of maximum 547 meteor counts for SAAMER and DrAAMER. We then  
evaluate the relative uncertainty of the 548 momentum flux assessment using the “dual-beam” analysis of *Vincent  
and Reid* [1983] for a GW

549 propagating zonally with superposed tidal winds and nominal radial velocity measurement

550 uncertainties. The east and west radial velocities are given by 551  $V_E = (u' + U_D + U_S)\sin\theta + w'\cos\theta \pm \Delta v_E'$   
(4) 552 and 553  $V_W = -(u' + U_D + U_S)\sin\theta + w'\cos\theta \pm \Delta v_W'$  (5) 554 where  $u'$  and  $w'$  are the horizontal and vertical  
GW velocities,  $U_D$  and  $U_S$  are the zonal diurnal 555 and semidiurnal tidal motions (assuming negligible vertical tidal  
motions and no horizontal 556 phase variations),  $\theta = 35^\circ$ ,  $\Delta v_E'$  and  $\Delta v_W'$  are the radial velocity uncertainties, and  
angle 557 brackets denote temporal averaging. Assuming all of the velocities in Eqs. (4) and (5) are 558 uncorrelated  
except for  $u'$  and  $w'$ , which are in phase or antiphase, squaring, subtracting Eq. (5) 559 from Eq. (4), and  
rearranging, the measured momentum flux may be written as 560  $\langle u'w' \rangle_M = (V_E^2 - V_W^2 + \Delta v_E'^2 - \Delta v_W'^2)/2\sin(2\theta)$   
(6) 561  $= \langle u'w' \rangle_T + (\Delta v_E'^2 - \Delta v_W'^2)/2\sin(2\theta)$  562 where  $\langle u'w' \rangle_T$  is the true GW zonal momentum flux for the  
specified motion field. The 563 fractional uncertainty in the measured momentum flux is then 564  $\Delta \langle u'w' \rangle / \langle u'w' \rangle_T$   
 $= (\Delta v_E'^2 - \Delta v_W'^2) / [\langle u'w' \rangle_T 2\sin(2\theta)]$  (7) 565 Thus, fractional uncertainties are minimized if radial velocity  
uncertainty variances are small 566 relative to the true GW momentum flux, as  $2\sin(2\theta) = 1.88$  for the zenith angle  
specified. The 567 largest magnitude of the numerator occurs if the radial velocity uncertainty variance is negligible  
568 in one beam. But for the uncertainties assumed above, this is  $\Delta v'^2 \sim 3 \text{ m}^2 \text{ s}^{-2}$ , with likely values 569 significantly  
smaller. By comparison, mean GW momentum fluxes in the MLT are typically  $\sim 10 \text{ m}^2 \text{ s}^{-2}$ , with peak values  $\sim 3$  to  
10 times larger, at most locations, with values over the Drake 571 Passage “hotspot” expected (and measured, see  
*F10b*) to be  $\sim 2$  to 5 times larger. These values



572 suggest a maximum fractional uncertainty of the momentum flux due to radial velocity  
573 uncertainties of ~15% at typical sites and ~5% in GW source hotspots. 574 We anticipate range uncertainties  
comparable to the range resolution of 2 km and angle575 of-arrival uncertainties of  $\sim 1^\circ$  in zenith angle and  $\sim 1.5^\circ$  in  
azimuth. These imply that altitude 576 uncertainties are defined by zenith angle uncertainties near zenith angles of  
 $\sim 60^\circ$  and by range 577 uncertainties at smaller zenith angles. At all zenith angles, however, these values suggest  
altitude 578 uncertainties comparable to or smaller than our chosen 3-km altitude bin. Thus range and angle579  
of-arrival uncertainties appear unlikely to significantly impact measurements of mean and tidal 580 winds having  
large vertical scales or GW momentum fluxes that rely most on radial velocities 581 occurring at smaller zenith  
angles. These results suggest that measurement uncertainties are 582 likely to contribute much less to momentum  
flux uncertainties than inadequate definition of the 583 mean, tidal, and GW fields due to low meteor counts arising  
due to low radar sensitivity or small 584 averaging intervals and/or range bins. 585 Several previous studies have  
assessed the potential for momentum flux measurement 586 uncertainties due to inadequate sampling of the motion  
field. *Kudeki and Franke* [1998] and 587 *Thorsen et al.* [2000] evaluated momentum flux measurements employing  
the “dual-beam” 588 technique and concluded that very long averaging intervals were required to achieve statistical  
589 significance. *Kudeki and Franke* [1998] assumed the GW velocity fields to be defined by 590 Gaussian  
distributions with a net momentum flux equal to 1% of that assuming a single GW was 591 present and inferred a  
required integration time of 16 days. This assumption, however, is in 592 contradiction to the strong correlations  
among component velocities that often occur when one or 593 several large-amplitude GWs dominate the  
high-frequency motion field and the total momentum 594 flux, as is often observed. In such cases, the averaging time  
implied by the *Kudeki and Franke*

595 [1998] would decrease by 2 decades or more. Examples of dual-beam or multiple-beam  
596 measurements employing MF, VHF, and UHF radars that provide clear evidence for significant 597 and variable  
GW momentum fluxes (magnitudes as large as  $\sim 70 \text{ m}^2 \text{ s}^{-2}$ ) occurring on time scales 598 as short as a few hours or less  
include the following: 599 1) MLT measurements with the MF radar at Buckland Park, Australia showing clear  
diurnal 600 tide modulation of GW momentum fluxes with a modulation amplitude of  $\sim 30 \text{ m}^2 \text{ s}^{-2}$  601 [*Fritts and*  
*Vincent*, 1987], 602 2) MLT measurements with the SOUSY VHF radar at Andoya, Norway revealing 603  
momentum fluxes (per unit density) of  $\sim 66 \text{ m}^2 \text{ s}^{-2}$  over  $\sim 3$  hours [*Reid et al.*, 1988], 604 3) troposphere and  
stratosphere measurements using the MU VHF radar at Shigaraki, Japan 605 for multi-beam measurements of GW  
momentum fluxes over 6 days that exhibit clear 606 episodic enhancements as short as an hour simultaneously in  
multiple beams and 607 spanning multiple altitudes [*Fritts et al.*, 1990], 608 4) MLT measurements employing the  
former VHF radar at Poker Flat, Alaska that revealed 609 clear anti-correlations between diurnal and semidiurnal  
tide winds and GW momentum 610 fluxes over 4 and 16 day intervals exhibiting stronger momentum flux  
modulations ( $\sim 5$  to 611  $10 \text{ m}^2 \text{ s}^{-2}$ ) accompanying larger tidal amplitudes [*Wang and Fritts*, 1991], 612 5) MLT  
measurements spanning two 10-day intervals with the Jicamarca VHF radar in Peru 613 that exhibit daily mean  
momentum fluxes of  $\sim 10 \text{ m}^2 \text{ s}^{-2}$  and maxima of  $\sim 30 \text{ m}^2 \text{ s}^{-2}$ , with 614 hourly profiles exhibiting significant coherence in  
altitude and time and maximum 615 magnitudes of  $\sim 60 \text{ m}^2 \text{ s}^{-2}$  [*Fritts et al.*, 1992],

616 6) MLT measurements with the MF radar at Buckland Park, Australia exhibiting clear  
617 diurnal modulation of GW momentum fluxes by a large-amplitude 2-day wave [Murphy 618 and Vincent, 1998],  
619 7) MLT measurements employing the dual-beam Arecibo UHF radar that revealed variable 620 GW momentum  
fluxes often near zero, but exhibiting occasional maxima as large as  $\sim 50$  621  $\text{m}^2 \text{s}^{-2}$ , that were largely anti-correlated  
with the large-scale wind field [Fritts *et al.*, 622 2006b], and 623 8) MLT measurements spanning  $\sim 8$  hr employing  
the Arecibo UHF radar that revealed very 624 significant GW activity, but with momentum fluxes that were very  
small, suggesting 625 ducting [Fritts and Janches, 2008]. 626 A more recent study by Vincent *et al.* [2010]  
employed numerical simulations of test GW 627 fields with a statistical model of radial velocity and angle-of-arrival  
uncertainties to evaluate the 628 accuracy of meteor radar measurements of mean winds and GW velocity variances  
and 629 momentum fluxes as functions of the meteor rate within a 2-km range bin. These authors 630 concluded that  
mean winds could be determined with relatively few meteors, but that estimation 631 of GW variances, and  
especially momentum fluxes, with small uncertainties required 632 considerable averaging enabling large meteor  
counts. The meteor counts employed to assess 633 measurement capabilities in the study ranged from 10 to 200. 634  
By comparison, our assessment above assumed a need for significant meteor counts, with 635 the typical averaging  
interval for definition of mean and tidal winds and GW momentum fluxes 636 being a month, but with mean wind  
and semidiurnal tide assessments also evaluated for one day 637 or for one hour of a composite day for the month  
(hence 24 or 30 hr of data). Employing real 638 meteor distributions observed during a test month for each of five  
radars (June of 2009, 2010, or

639 2011), these yielded meteor counts in a 3-km altitude bin at the peak of the meteor distribution  
640 for the month of ~80,000 at TdF, ~70,000 at BL, ~40,000 at KGI, and ~25,000 at SRO and YKF. 641 Meteor  
counts at the upper and lower edges of these distributions varied by radar, but were 642 typically smaller by ~2 to 4  
times. Daily and composite-day hourly assessments employed ~3 643 and 4% of these meteor counts, respectively.  
Thus, in all cases, our meteor counts for each radar 644 were significantly larger than those assessed by *Vincent et al.*  
[2010] near the peak of the meteor 645 distribution. 646 Assuming that accuracies vary as the signal-to-noise ratio  
(S/N) and thus as  $N^{1/2}$ , where N 647 is the meteor count, we can compare our inferred accuracies with those of  
*Vincent et al.* [2010]. 648 As noted above, the radars at BL, TdF, and KGI have peak monthly meteor counts of N  
~40,000 649 to 80,000 in a 3-km altitude bin, with N ~10,000 to 25,000 at the edges of the distributions. 650 These  
imply reductions by factors of ~15 to 25 in the mean wind measurement errors at the 651 meteor distribution peak  
relative to the  $5 \text{ ms}^{-1}$  uncertainties displayed by *Vincent et al.* [2010] in 652 Table 1, with reductions by factors of ~10  
at the highest and lowest altitudes. Inspection of the 653 monthly mean winds for the seven cases considered above  
suggests that our results are largely 654 consistent with that expectation. Similarly, we expect error reductions of ~4  
and 2 in the center 655 and edge regions for daily mean winds, implying errors of ~1 to  $3 \text{ ms}^{-1}$ . These are largely 656  
consistent with the daily mean winds inferred for our Case 5 test fields displayed in Figure 8. 657 Finally, meteor  
counts at SRO and YKF suggest monthly mean wind uncertainties smaller than 658 the  $\sim 5 \text{ ms}^{-1}$  uncertainties of  
*Vincent et al.* [2010] by ~12 and 5 in the center and edge regions. In 659 these cases, our uncertainties seem to be  
comparable to, or somewhat larger than, those expected. 660 We note, however, that the antenna patterns implied by  
the meteor locations shown in Figure 1 661 suggest significant asymmetries that likely also contribute to  
measurement errors.

Applying the same factors to our momentum flux estimates, we expect to see errors of 663 ~10 and 25% for the monthly means at the center and the edges of these distributions. Again, our 664 results are consistent with these expectations, though with variations that depend on the 665 complexity of our motions fields and the meteor distribution of each radar. Note, in particular, 666 that *Vincent et al.* [2010] did not include tidal winds in their assessment, but that variable tidal 667 winds can induce apparent mean winds and GW momentum fluxes if the meteor distribution is 668 not approximately uniform in space and time. Our results also are relatively more accurate for 669 cases having constant GW character throughout the month; they are less accurate for Cases 6 and 670 7 for which GWs are randomly distributed and likely more similar to the GW spectra specified 671 by *Vincent et al.* [2010]. We cannot assess the contributions of the assumed radial velocity and 672 angle-of-arrival uncertainties employed by *Vincent et al.* [2010] to the overall uncertainties 673 displayed in their Table 1. However, our own assessment of radial velocity uncertainties above 674 suggests that the majority of the measurement errors seen in our results can be attributed 675 primarily to the expected dependence on meteor counts. 676 677 **6**

**Conclusions** 678 Our assessment of meteor radar measurement capabilities presented in previous sections 679 has demonstrated both 680 1) clear capabilities for quantitative measurements of mean winds, tides, and GW 681 momentum fluxes by radars that achieve sufficiently high meteor counts and provide 682 sensitivity to radial velocities at sufficiently small zenith angles, and 683 2) strong dependence of these measurement capabilities on radar power, beam geometry and 684 sensitivity, and spatial and temporal meteor distributions.

685 Meteor radars that combine high power, high meteor counts at small zenith angles, and  
686 symmetric beam patterns exhibit the smallest measurement errors. Those having low power, 687 acquiring a  
majority of meteor detections at large zenith angles, and/or experiencing beam or 688 meteor detection asymmetries  
exhibit much larger measurement uncertainties, thus either 689 requiring longer averaging intervals or precluding the  
more challenging measurements, 690 especially estimations of GW momentum fluxes. 691 Monthly mean winds  
can typically be measured to within  $1 \text{ ms}^{-1}$  over  $\sim 15$  to  $20 \text{ km}$ , with 692 maximum accuracies between  $\sim 85$  and  $90$   
 $\text{km}$ , for radar frequencies near  $35 \text{ MHz}$  and having 693 higher power than conventional meteor radars. These radars  
can also provide daily mean winds 694 having accuracies of  $\sim 1$  to  $3 \text{ ms}^{-1}$  where meteor counts are sufficient, even in  
the presence of 695 large tidal motions. Monthly mean diurnal and semidiurnal tide amplitudes are also recovered  
696 very well by higher-power meteor radars, but typically with amplitude under-estimates of a few 697 %. Mean  
winds and tidal amplitudes are recovered much more accurately, however, by meteor 698 radars that achieve high  
meteor counts at small zenith angles compared to meteor radars 699 achieving the same meteor counts, but at larger  
zenith angles. 700 GW momentum flux estimates are challenging for meteor radars because of the large 701  
number of meteors required to adequately resolve differences in radial velocity variances at 702 opposite azimuths.  
Nevertheless, accurate estimates over sufficiently long time scales (typically 703 averages over  $\sim 1$  to  $30$  days) appear  
to be possible with sufficient meteor counts ( $\sim 2,000$  to 704  $60,000$  in an altitude bin). Accuracies are also enhanced  
with high meteor counts at small zenith 705 angles, given the greater sensitivity of these radial velocities to the  
correlations between 706 horizontal and vertical velocities within the GW field. Importantly, meteor radars appear to  
707 measure momentum fluxes for stationary and propagating GWs, for complex GW 708 superpositions, and for  
GW fields exhibiting significant intermittency equally well on monthly 709 time scales. 710 **Acknowledgments**

The research described here was performed under NSF grants ATM711 0634650, ATM-0824742, and  
OPP-0839084. We are grateful for the valuable assistance of 712 personnel at Estacion Astronomica Rio Grande  
(EARG) with the operations and maintenance of 713 SAAMER. We are also grateful to the Secretaria for the  
Interministerial Commission of Sea 714 Resources (SECIRM), the Brazilian Antarctic Program (PROANTAR), the  
National Institute for 715 Science and Technology, Antarctic Environmental Research (INCT-APA), and the 716

ATMANTAR/MCT/CNPq project, FAPERJ, for their support of this research and visits to 717 Ferraz Station to install and service DrAAMER. 718 719 **References** 720 Acott, P., 2009: Mesospheric momentum flux studies over Fort Collins, CO, Ph.D. dissertation, 721 Colorado State University, 2009. 722 Antonita, T. M., G. Ramkumar, K. K. Kumar, and V. Deepa, 2008: Meteor wind radar 723 observations of gravity wave momentum fluxes and their forcing toward the Mesospheric 724 Semiannual Oscillation, *J. Geophys. Res.*, 113, D10115, doi:10.1029/2007JD009089. 725 Clemesha, B. R., P. Batista, R. A. Buriti da Costa, and N. Schuch, 2009: Seasonal variations in 726 gravity wave activity at three locations in Brazil, *Ann. Geophys.*, 27, 1059–1065. 727 Brown, P. R. A., 1983: Aircraft measurements of mountain waves and their associated 728 momentum flux over the British Isles, *Q. J. Roy. Met. Soc.*, 109, 849-865. 729 Ern, M., P. Preusse, M. J. Alexander, and C. D. Warner, 2004: Absolute values of gravity wave 730 momentum flux derived from satellite data, *J. Geophys. Res.*, 109, D20103,

731 doi:10.1029/2004JD004752. 732 Espy, P. J., R. E. Hibbins, G. R. Swenson, J. Tang, M. J. Taylor, D. M. Riggin,  
and D. C. Fritts, 733 2006: Regional variations of mesospheric gravity-wave momentum flux over Antarctica, 734  
Ann. Geophys., 24, SRef-ID: 1432-0576/ag/2006-24-81, 81–88. 735 Espy, P. J., G. O. L. Jones, G. R. Swenson,  
and M. J. Taylor, 2004: Tidal modulation of the 736 gravity-wave momentum flux in the Antarctic mesosphere,  
Geophys. Res. Lett., 31, L11111, 737 doi:10.1029/2004GL019624. 738 Fritts, D. C., and M. J. Alexander, 2003: A  
review of gravity wave dynamics and effects in the middle 739 atmosphere, *Rev. Geophys.*, **41**,  
doi:10.1029/2001RG000106. 740 Fritts, D. C., and D. Janches, 2008: Dual-beam measurements of gravity wave  
momentum fluxes 741 over Arecibo: Re-evaluation of wave structure, dynamics, and momentum fluxes, J. 742  
Geophys. Res., 113, D05112, doi:10.1029/2007JD008896. 743 Fritts, D. C., D. Janches, and W. K. Hocking, 2010b:  
Southern Argentina agile meteor radar 744 (SAAMER): Initial assessment of gravity wave momentum fluxes, J.  
Geophys. Res., 115, 745 D19123, doi:10.1029/2010JD013891. 746 Fritts, D. C., D. Janches, H. Iimura, W. K.  
Hocking, and J. V. Bageston, and N. M. Pene, 2011: 747 Drake Antarctic Agile Meteor Radar (DrAAMER) First  
Results: Configuration and 748 Comparison of Mean and Tidal Wind and Gravity Wave Momentum Flux  
Measurements 749 with SAAMER, J. Geophys. Res., submitted. 750 Fritts, D. C., D. Janches, H. Iimura, W. K.  
Hocking, N. J. Mitchell, B. Fuller, B. Vandeppeer, J. 751 Hormaechea, C. Brunini, and H. Levato, 2010a: Southern  
Argentina agile meteor radar 752 (SAAMER): System design and initial measurements of large-scale winds and  
tides, J. 753 Geophys. Res., 115, D18112, doi:10.1029/2010JD013850.



754 Fritts, D. C., D. Janches, D. M. Riggan, R. G. Stockwell, M. P. Sulzer, and S. Gonzalez, 2006b:  
755 Gravity waves and momentum fluxes in the MLT using 430 MHz dual-beam measurements 756 at Arecibo: 2.  
Frequency spectra, momentum fluxes, and variability, *J. Geophys. Res.*, 111, 757 D18108,  
doi:10.1029/2005JD006883. 758 Fritts, D. C., and T. Lund, 2011: Gravity wave influences in the thermosphere and  
ionosphere: 759 Observations and recent modeling, *Aeronomy of the Earth's Atmosphere and Ionosphere*, M. 760  
Abdu and D. Pancheva, Eds., Springer, 109-130. 761 Fritts, D. C., T. Tsuda, T. E. VanZandt, S. A. Smith, T. Sato, S.  
Fukao, and S. Kato, 1990: 762 Studies of velocity fluctuations in the lower atmosphere using the MU radar: II.  
Momentum 763 fluxes and energy densities. *J. Atmos. Sci.*, 47, 51-66. 764 Fritts, D. C., S. L. Vadas, K. Wan, and J.  
A. Werne, 2006a: Mean and variable forcing of the 765 middle atmosphere by gravity waves, *J. Atmos. Solar-Terres.*  
*Phys.*, 68, 247-265. 766 Fritts, D. C., S. A. Vadas, and Y. Yamada, 2002: An estimate of strong local gravity wave  
body 767 forcing based on OH airglow and meteor radar observations, *Geophys. Res. Lett.*, 29 (10), 768  
10.1029/2001GL013753. 769 Fritts, D. C., and R. A. Vincent, 1987: Mesospheric momentum flux studies at  
Adelaide, 770 Australia: Observations and a gravity wave/tidal interaction model, *J. Atmos. Sci.*, 44, 605771 619.  
772 Fritts, D. C., and L. Yuan, 1989: Measurement of momentum fluxes near the summer 773 mesopause at Poker  
Flat, Alaska, *J. Atmos. Sci.*, 46, 2569-2579. 774 Fritts, D. C., L. Yuan, M. H. Hitchman, L. Coy, E. Kudeki, and R.  
F. Woodman, 1992: 775 Dynamics of the equatorial mesosphere observed using the Jicamarca MST radar during  
June 776 and August 1987, *J. Atmos. Sci.*, 49, 2353-2371.

777 Fukao, S., T. Sato, T. Tsuda, and S. Kato, 1988: VHF Doppler Radar Determination of the  
778 Momentum Flux in the Upper Troposphere and Lower Stratosphere: Comparison between 779 the Three- and  
Four-Beam Methods, *J. Atmos. Ocean. Tech.*, 5(1), 57-69. 780 Hertzog, A., G. Boccara, R. A. Vincent, F. Vial, P.  
Cocquerez, 2008: Estimation of Gravity Wave 781 Momentum Flux and Phase Speeds from Quasi-Lagrangian  
Stratospheric Balloon Flights. Part II: 782 Results from the Vorcore Campaign in Antarctica, *J. Atmos., Sci.*, 65,  
3056-3070, 783 doi:10.1175/JAS2710.1. 784 Hertzog, A., and F. Vial, 2001: A study of the dynamics of the  
equatorial lower stratosphere by use 785 of ultra-long-duration balloons, *J. Geophys. Res.*, 106, 22,745-22,761. 786  
Hitchman, M. H., K. W. Bywaters, D. C. Fritts, L. Coy, and E. Kudeki, 1992: Mean winds and 787  
momentum fluxes over Jicamarca, Peru during June and August 1987, *J. Atmos. Sci.*, 49, 788 2372-2383. 789 Hines,  
C. O., 1960: Internal gravity waves at ionospheric heights, *Can. J. Phys.* 38(11): 1441– 790 1481, 1960. 791 Hocke,  
K. and K. Schlegel, 1996: A review of atmospheric gravity waves and traveling 792 ionospheric disturbances:  
1982–1995, *Ann. Geophys.*, 14, 917–940. 793 Hocking, W. K., 2005: A new approach to momentum flux  
determinations using SKiYMET 794 meteor radars, *An. Geophys.*, 23, 1–7. 795 Hocking, W.K., 2005: A new  
approach to momentum flux determinations using SKiYMET 796 meteor radars, *Ann. Geophysicae*, 23, 2433-2439.  
797 Hocking, W.K. and G. Kishore Kumar, 2011: Long Term Behaviour of the MLT Quasi 7 day 798 Wave at two  
Radar-sites at Northern Polar Latitudes, *J. Atmos. Solar Terr. Phys.*, 73, 1616– 799 1628. 800 Kim, Y.-J., S. D.  
Eckermann, and H.-Y. Chun, 2003: A overview of the past, present and future

801 of gravity-wave drag parameterization for numerical climate and weather prediction models,  
802 Atmos. Ocean, 41, 65-98. 803 Kudeki, E., and S. J. Franke, 1998: Statistics of momentum flux estimation, J.  
Atmos. Solar804 Terres. Phys., 60, 1549-1553. 805 Kumar, K. K., T. M. Antonita, and S. T. Shelbi, 2007: Initial  
results from SKiYMET meteor 806 radar at Thumba (8.5°N, 77°E): 2. Gravity wave observations in the MLT region,  
Radio Sci., 807 42, RS6009, doi:10.1029/ 2006RS003553. 808 Lighthill, M. J., 1978: Waves in Fluids, Cambridge  
University Press, Cambridge, England. 809 Lilly, D. K., and P. J. Kennedy, 1973: Observations of a stationary  
mountain wave and its 810 associated momentum flux and energy dissipation, J. Atmos. Sci., 30(6), 1135-1152.  
811 Lilly, D. K., J. M. Nicolls, P. J. Kennedy, J. B. Klemp, and R. M. Chervin, 1982: Aircraft 812 measurements of  
wave momentum fluxes over the Colorado Rocky Mountains, Q. J. Roy. 813 Met. Soc., 108, 625-642. 814 McIntyre,  
M. E., 1989: On dynamics and transport near the polar mesopause in summer. J. 815 Geophys. Res., 94,  
14,617-14,628. 816 Meyer, W., R. Siebenmorgen, and H.-U. Widdel, 1989: Estimates of gravity wave momentum  
817 fluxes in the winter and summer high mesosphere over Scandinavia, J. Atmos. Terres., Phys., 818 51(4),  
311-319. 819 Mitchell, N. J., and C. L. Beldon, 2009: Gravity waves in the mesopause region observed by 820  
meteor radar: 1. A simple measurement technique, J. Atmos. Sol. Terr. Phys., 71, 866–874. 821 Murayama, Y., T.  
Tsuda, and S. Fukao, 1994: Seasonal variation of gravity wave activity in the 822 lower atmosphere observed with  
the MU radar, J. Geophys. Res., 99(D11), 23,057–23,069.

823 Murphy, D. J., and R. A. Vincent, 1993: Estimates of momentum flux in the mesosphere and  
824 lower thermosphere over Adelaide, Australia, from March 1985 to February 1986, *J. Geophys. Res.*,  
98(D10), 18,617-18,638. 826 Murphy, D. J., and R. A. Vincent, 1998: Mesospheric momentum fluxes over Adelaide  
during 827 the 2-day wave: Results and interpretation, *J. Geophys. Res.*, 103(D22), 28,627–28,636, 828  
doi:10.1029/1998JD200001. 829 Nakamura, T., T. Tsuda, M. Yamamoto, S. Fukao, and S. Kato, 1993:  
Characteristics of Gravity 830 Waves in the Mesosphere Observed With the Middle and Upper Atmosphere Radar 1.  
831 Momentum Flux, *J. Geophys. Res.*, 98(D5), 8899–8910. 832 Nappo, C. J., 2002: An Introduction to  
Atmospheric Gravity Waves, Academic Press. 833 Nastrom, G. D., and D. C. Fritts, 1992: Sources of mesoscale  
variability of gravity waves, I: 834 Topographic excitation, *J. Atmos. Sci.*, 49, 101-110. 835 Placke, M., P.  
Hoffmann, E. Becker, C. Jacobi, W. Singer, and M. Rapp, 2011: Gravity wave 836 momentum fluxes in the  
MLT—Part II: Meteor radar investigations at high and midlatitudes 837 in comparison with modeling studies, *J.*  
*Atmos. Solar-Terres. Phys.*, 73, 911-920. 838 Reid, I. M., and R. A. Vincent, 1987: Measurements of mesospheric  
gravity wave momentum 839 fluxes and mean flow accelerations at Adelaide, Australia. *J. Atmos. Terres. Phys.*, 49,  
443840 460. 841 Reid, I. M., R. Rüster, P. Czechowsky, and G. Schmidt, 1988: VHF radar measurements of 842  
momentum flux in the summer polar mesosphere over Andenes (69° N, 16° E), Norway, 843 *Geophys. Res. Lett.*,  
15, 1263-1266. 844 Sato, K., 1990: Vertical Wind Disturbances in the Troposphere and Lower Stratosphere 845  
Observed by the MU Radar, *J. Atmos. Sci.*, 47(23), 2803-2817.

846 Sato, K., 1993: Small-Scale Wind Disturbances Observed by the MU Radar during the Passage  
847 of Typhoon Kelly, *J. Atmos. Sci.*, 50(4), 518-537. 848 Sato, K., 1994: A statistical study of the structure,  
saturation and sources of inertio-gravity waves 849 in the lower stratosphere observed with the MU radar, *J. Atmos.*  
*Terres. Phys.*, 56 (6), 755-850. 851 Smith, R. B., B. K. Woods, J. Jensen, W. A. Cooper, J. D. Doyle, Q. Jiang,  
and V. Grubisic, 852 2008: Mountain waves entering the stratosphere, *J. Atmos. Sci.*, 65, 2543-2562, 853  
doi:10.1175/2007JAS2598.1. 854 Swenson, G. R., R. Haque, W. Yang, and C. S. Gardner, 1999: Momentum and  
energy fluxes of 855 monochromatic gravity waves observed by an OH imager at Starfire Optical Range, New 856  
Mexico, *J. Geophys. Res.*, 104(D6), 6067–6080, doi:10.1029/1998JD200080. 857 Thorsen, D., S. J. Franke, and E.  
Kudeki, 2000: Statistics of momentum flux estimation using the 858 dual coplanar beam technique, *Geophys. Res.*  
*Lett.*, 27, 3193-3196. 859 Tsuda, T., Y. Murayama, M. Yamamoto, S. Kato, and S. Fukao, 1990: Seasonal variation  
of 860 momentum flux in the mesosphere observed with the MU radar. *Geophys. Res. Lett.*, 17, 861 725-728. 862  
Vincent, R. A., S. Kovalam, I. M. Reid, and J. P. Younger, 2010: Gravity wave flux retrievals 863 using meteor  
radars, *Geophys. Res. Lett.*, 37, L14802, doi:10.1029/2010GL044086. 864 Vincent, R.A., I. M. Reid, 1983: HF  
Doppler measurements of mesospheric momentum fluxes. 865 *Journal of Atmospheric Science* 40, 1321–1333. 866  
Wang, D.-Y., and D. C. Fritts, 1990: Mesospheric momentum fluxes observed by the MST radar 867 at  
Poker Flat, Alaska, *J. Atmos. Sci.*, 47, 1511-1521.

86 Wang, D.-Y., and D. C. Fritts, 1991: Evidence of gravity wave-tidal interaction observed near  
8 the summer mesopause at Poker Flat, Alaska, J. Atmos. Sci.,48, 572-583.  
86  
9 Zhang, X.,J. M. Forbes, and M. E. Hagan, 2010a: Longitudinal variation of tides in the MLT  
87  
0 region: 1. Tides driven by tropospheric net radiative heating, J. Geophys. Res. (Space  
87  
1 Physics), 115, 6316, doi:10.1029/2009JA014897.  
87  
2 Zhang, X.,J. M. Forbes, and M. E. Hagan, 2010b: Longitudinal variation of tides in the MLT  
87  
3 region: 2. Relative effectis of solar radiative and latent heating, J. Geophys. Res. (Space  
87  
4 Physics), 115, 6317, doi:10.1029/2009JA014898.  
87  
5  
87  
6  
87  
7  
87  
8

radar	location (lat/long)	frequency, bandwidth	peak power	resol.	TX/RX ants.	PRF (Hz)	coding, integr.	meteors /day
DrAAMER (KGI)	(62.1oS, 58.7oW)	36.9 MHz, 35-125 kHz	30 kW	2 km	8/5, 3-ele. cross. Yagis	1730	2 bit, 4 samp.	~9,000
SAAMER	(53.8oS,	32.55 MHz,	60 kW	2 km	8/8, 3-ele.	2144,	2 bit, 4	~14,000

86 the summer mesopause at Poker Flat, Alaska, J. Atmos. Sci.,48, 572-583.  
9  
87  
0 Zhang, X.,J. M. Forbes, and M. E. Hagan, 2010a: Longitudinal variation of tides in the MLT  
879 87  
1 region: 1. Tides driven by tropospheric net radiative heating, J. Geophys. Res. (Space  
87  
2 Physics), 115, 6316, doi:10.1029/2009JA014897.  
87  
3 Zhang, X.,J. M. Forbes, and M. E. Hagan, 2010b: Longitudinal variation of tides in the MLT  
87  
4 region: 2. Relative effectis of solar radiative and latent heating, J. Geophys. Res. (Space  
87  
5 Physics), 115, 6317, doi:10.1029/2009JA014898.  
87  
6

**Table 1.** Radar characteristics for the five radars employed in this evaluation of measurement 880 capabilities. Meteor counts for zenith angles between 15 and 50° are ~50% of the total counts 881 (right column) for KGI and TdF and ~30% of the total counts for the other radars.

882 883 **Table 2.** Mean, tidal, and GW parameters used for test cases evaluating radar measurement 884 capabilities

employing real meteor distributions and test motion fields. Mean GW momentum

86 Wang, D.-Y., and D. C. Fritts, 1991: Evidence of gravity wave-tidal interaction observed near  
8  
86 the summer mesopause at Poker Flat, Alaska, J. Atmos. Sci.,48, 572-583.  
9  
87 Zhang, X.,J. M. Forbes, and M. E. Hagan, 2010a: Longitudinal variation of tides in the MLT  
0  
87 region: 1. Tides driven by tropospheric net radiative heating, J. Geophys. Res. (Space  
1  
87 Physics), 115, 6316, doi:10.1029/2009JA014897.  
2  
87 Zhang, X.,J. M. Forbes, and M. E. Hagan, 2010b: Longitudinal variation of tides in the MLT  
3  
87 region: 2. Relative effectis of solar radiative and latent heating, J. Geophys. Res. (Space  
4  
87 Physics), 115, 6317, doi:10.1029/2009JA014898.  
5  
87  
6  
87  
7  
87  
8

radar	location (lat/long)	frequency, bandwidth	peak power	resol.	TX/RX ants.	PRF (Hz)	coding, integr.	meteors /day
DrAAMER (KGI)	(62.1oS, 58.7oW)	36.9 MHz, 35-125 kHz	30 kW	2 km	8/5, 3-ele. cross. Yagis	1730	2 bit, 4 samp.	~9,000
SAAMER (TdF)	(53.8oS, 67.7oW)	32.55 MHz, 50-250 kHz	60 kW	2 km	8/8, 3-ele. cross. Yagis	2144, 1730	2 bit, 4 samp.	~14,000
Socorro, NM (SOC)	(34.1oN, 106.9oW)	35.24 MHz, 35-125 kHz	6 kW	2 km	1/5, 2-ele. Yagis	2140	1 bit, 8 samp.	~5,000
Bear Lake, UT (BLO)	(41.9oN, 111.4oW)	35.24 MHz, 50-250 kHz	12 kW	2 km	1/5, 2-ele. cross. Yagis	2140	1 bit, 4 samp.	~11,000
Yellowknife, Canada (YK)	(62.5oN, 114.3oW)	35.65 MHz, 50-125 kHz	6 kW	2 km	1/5, 2-ele. Yagis	2140	1 bit, 8 samp.	~2,500
Parameter	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6	Case 7	
UM, VM	20, 10	0, 0	0, 0	20, 10	40, -20	-20, -10	-20, -10	
UD, VD	10, 10	0, 0	0, 0	10, 10	20, 20	10, 10	10, 10	
USD, VSD	50, 50	0, 0	0, 0	50, 50	20+2(z - 80)sin2( $\pi t/T_M$ )	50, 50	50, 50	
UGW1	10 0 5	20 0 5	20 0	10 0 5	20 abs[sin(2 $\pi t/T_M$ )]	40F6(t) 0	30F7(t)	
VGW1	2 $\pi$ /50	2 $\pi$ /50	-10	2 $\pi$ /50	*sin(2 $\pi t/T_{SD}$ ) 0 -10	20F6 (t)	30F7(t)	
WGW1 k1 l1	0 0 20	0	2 $\pi$ /30	0 0 20	abs[sin(2 $\pi t/T_M$ )]	2 $\pi$ /50 0	10F7 (t)	
m1 TGW1		2 $\pi$ /15 $\infty$	0 0 $\infty$		*abs[sin(2 $\pi t/T_{SD}$ )] 2 $\pi$ /50 0 0 20	2 $\pi$ /15 20	2 $\pi$ /40 2 $\pi$ /40 2 $\pi$ /15 20	
UGW2	0 20 2	--	0 10 2	0 20 2	0 20 abs[sin(2 $\pi t/T_M$ )]	0 30G6	30G7(t)	
VGW2	0		0	0	*cos(2 $\pi t/T_{SD}$ ) 5	(t) 10G6	-30G7(t)	
WGW2 k2 l2	2 $\pi$ /100		2 $\pi$ /40	2 $\pi$ /100	abs[sin(2 $\pi t/T_M$ )]	(t) 0	20G7(t)	
m2 TGW2	0 30		0 $\infty$	0 30	*abs[cos(2 $\pi t/T_{SD}$ )] 0 2 $\pi$ /100 0 30	2 $\pi$ /100 2 $\pi$ /20 30	2 $\pi$ /50 2 $\pi$ /50 2 $\pi$ /20 15	
UGW3	--	--	--	20 -10	20 -10 2 $\pi$ /30 0 $\infty$	--	--	
WGW3 k3				2 $\pi$ /30				



40 885

fluxes for  
each case  
are shown  
at the  
bottom.

Units for  
velocities,  
wavenum  
bers, and  
periods

are 886  
 $\text{ms}^{-1}$ ,  $\text{km}^{-1}$ ,  
and min,  
and  $T_M =$

10 days  
and  $T_{SD} =$   
12 hrs.

GWs in  
Case 6 are  
modulated  
by 887  
amplitude  
functions

$F_6(t) = 1(t$

= 0-3 h +

21R<sub>1</sub> hr)

and F<sub>6</sub>(t)

= 0

otherwise,

and G<sub>6</sub>(t)

= 1 (t =

0-4 888 h

+ 20R<sub>2</sub> hr)

and G<sub>6</sub>(t)

= 0

otherwise,

with R<sub>1</sub>

and R<sub>2</sub>

random

variables

between 0

and 1

chosen

889

separately

for each

of the 30

days of

the test

month.

GWs in

Case 7 are

modulated

by

amplitude

890

functions

$F_7(t) = 1$  (t

= 0-2 and

8-10 h +

14R<sub>3</sub> hr)

and  $F_7(t)$

= 0

otherwise,

and  $G_7(t)$

= 1 (t =

0-1 h 891

and 6-7 h

and 10-11

h and

19-20 h +

4R<sub>3</sub> hr)

and  $G_7(t)$

= 0

otherwise,

with  $R_3$

and  $R_4$

random

892

variables

between 0

and 1 as

above.

893 894

895 896

897 898

899 900

901 902

903 904

905 906

**Figure**

**Captions**

907

**Figure 1.**

Daily

meteor

distributio

ns

illustratin

g the  
beam  
patterns  
for the  
five radars  
employed  
908 in this  
assessment  
t of  
measurem  
ent  
capabilitie  
s. The  
radar  
locations  
and  
frequencie  
s are listed  
in 909  
each  
panel. The  
radars at  
BL, TdF,  
and KGI  
use

crossed  
Yagi  
antennas  
and have  
nearly 910  
symmetric  
beam  
patterns;  
The radars  
at YKF  
and SRO,  
however,  
use  
uncrossed  
(linearly-  
911  
polarized)  
Yagis  
oriented  
north-south  
and  
east-west,  
respective  
ly,  
resulting

in the  
asymmetri  
c 912  
meteor  
distributio  
ns  
favoring  
east-west  
and  
north-sout  
h  
measurem  
ents,  
respective  
ly, at  
those 913  
sites.  
Nulls at  
various  
radii in  
the meteor  
detections  
are due to  
removal  
of meteor

detections

that 914

may have

ground

clutter

contamina

tion. 915

**Figure 2.**

Accepted

meteors

employed

for mean

and tidal

wind and

momentu

m flux

estimates

916 for

each radar

(labeled at

left).

Meteor

distributio

ns are

shown as



(left)  
monthly  
counts in  
each 917  
500 m  
range bin,  
(center)  
counts per  
day, and  
(right)  
counts per  
hour for a  
composite  
day.  
These 918  
statistics  
include  
meteors at  
zenith  
angles  
between  
15 and 60°  
for all  
radars  
examined

here. 919

**Figure 3.**

Monthly  
mean

(left)

winds,

(second

column)

diurnal

tide

amplitude

s, (third

column)

920

semidiurn

al tide

amplitude

s, and

(right)

GW

momentu

m fluxes

for Case

1. Solid

and

dashed  
921 lines  
denote  
zonal and  
meridiona  
l profiles  
and the  
radars are  
designated  
at the left  
of each  
row. 922  
Vertical  
dashed  
lines show  
specified  
values.

923

**Figure 4.**

As in  
Figure 3,  
but for  
Case 2.

924

**Figure 5.**

As in  
Figure 3,  
but for  
Case 3.

925

**Figure 6.**

As in  
Figure 3,  
but for  
Case 4.

926

**Figure 7.**

As in  
Figure 3,  
but for  
Case 5.

927 **Figure 8.** Daily (left) zonal mean winds, (second column) meridional mean winds, (third 928 column) zonal diurnal tide amplitudes, and (right) meridional diurnal tide amplitudes for Case 5. 929 Vertical dashed lines show specified values. 930 **Figure 9.** Daily (left) zonal and (right) meridional semidiurnal tide amplitudes for Case 5 for the 931 five radars (labeled at the top of the left panels). Colored lines show measured amplitudes from 932 (black) 76.5 to (red) 97.5 km, dashed lines show specified amplitudes. 933 **Figure 10.** Hourly (left) zonal and (right) meridional GW momentum fluxes for Case 5 for a 934 composite day for the five radars (labeled as in Figure 9). Colored lines show measured 935 amplitudes from (black) 76.5 to (red) 97.5 km. Dashed lines show specified amplitudes. 936 **Figure 11.** As in Figure 3 for Case 6. 937 **Figure 12.** As in Figure 10, but for Case 6. 938 **Figure 13.** As in Figure 3 for Case 7. 939 **Figure 14.** As in Figure 10, but for Case 7.

Bear Lake (BL)  $f = 35.24$  MHz Yellowknife (YKF)  $f = 36.65$  MHz Socorro (SRO)  $f = 35.24$  MHz

300

300

20  
0

200

200