

The Kinematic and Microphysical Control of Storm Integrated Lightning Flash Extent

Lawrence D. Carey¹, William J. Koshak², Harold S. Peterson³, Elise V. Schultz¹, Retha Mathee¹, Christopher J. Schultz^{1,2}, Walter A. Petersen⁴, Lamont Bain¹



¹ University of Alabama in Huntsville, Department of Atmospheric Science, Huntsville, AL

² Earth Science Office, NASA, Marshall Space Flight Center (MSFC), Huntsville, AL

³ Universities Space Research Association (USRA), Huntsville, AL

⁴ NASA GSFC/Wallops Flight Facility, Wallops Island, VA




Objective


To investigate the kinematic and microphysical control of lightning properties, particularly those that may govern the production of nitrogen oxides (NO_x) in thunderstorms, such as flash rate, type (intracloud [IC] vs. cloud-to-ground [CG]) and extent.

Data and Methodology

- NASA MSFC Lightning Nitrogen Oxides Model (LNOM) is applied to North Alabama Lightning Mapping Array (NALMA) and Vaisala National Lightning Detection Network™ (NLDN) observations following ordinary convective cells through their lifecycle.
- LNOM provides estimates of flash type, channel length distributions, lightning segment altitude distributions (SADs) and lightning NO_x production profiles (Koshak et al. 2012).



- LNOM lightning characteristics are compared to the evolution of updraft and precipitation properties inferred from dual-Doppler (DD) and polarimetric radar analyses of UAHuntsville Advanced Radar for Meteorological and Operational Research (ARMOR, C-band, polarimetric) and KHTX (S-band, Doppler).



Radars: Multi-Doppler (30°), ARMOR: 100 km, KHTX/Hytop: 100 km


NALMA: NALMA sensors, 150, 250 km range rings, NA-LMA Altitude errors (m)

Center location of LNOM analysis cylinders shown at 1602, 1633, and 1657 UTC. Note storm on edge of DD lobes at 1602 UTC.


3 April 2007: Ordinary Convective Cell

- LNOM is applied in a Lagrangian sense (i.e., storm following) to well isolated thunderstorm cell on 3 April 2007 over Northern Alabama. Pulse severe (1" hail) at 1637 UTC.


LNOM Analysis Cylinders (LAC's), ARMOR Reflectivity (4 km), NALMA Flash Origins




Time Evolution of Flash Properties




Select Individual Flash and Radar Structure



Time-Height Cross Sections



Storm Integrated SAD and NO_x over 1-Hour Lifecycle



Summary and Discussion

- LNOM successfully run in Lagrangian mode for isolated, ordinary (pulse severe) thunderstorm
- Vertical SAD and NO_x production similar to long term
- Like flash rate, flash extent is generally correlated with production of precipitation ice and updraft < -5°C (when measured well by Doppler network)
- Descent of precipitation ice mass (graupel and small ice) associated with peak in CG rate and extent (1610-1616 UTC)
- Similar descent of hail core associated with lower CG rates and extent, especially at low levels (1634-1642 UTC)
- Updraft volume, precipitation type and process (coalescence-freezing) at T < -5°C modulated flash (and charging) vertical extent.
- Lofting of supercooled drops to -10°C and colder common even when > 5 m s⁻¹ updrafts less widespread (e.g., 1616 UTC). Z₀ columns were typically lightning minimum.
- Large reflectivity gradient at heights above -10°C (limit vertical extent of precip. ice) resulted in narrow (yet active) charging and lightning zones at later times (e.g., 1627 UTC)
- 1" hail reported at 1637 UTC with relatively suppressed activity. Efficient wet growth of frozen drops?