

This is the author's final, peer-reviewed manuscript as accepted for publication. The publisher-formatted version may be available through the publisher's web site or your institution's library.

Obesity classification in military personnel: a comparison of body fat, waist circumference, and body mass index measurements

Katie M. Heinrich, Nattinee Jitnarin, Richard R. Suminski, LaVerne Berkel, Christine M. Hunter, Lisa Alvarez, Antionette R. Brundige, Alan L. Peterson, John P. Foreyt, C. Keith Haddock, Walker S. C. Poston

How to cite this manuscript

If you make reference to this version of the manuscript, use the following information:

Heinrich, K. M., Jitnarin, N., Suminski, R. R., Berkel, L., Hunter, C. M., Alvarez, L., . . . Poston, W. S. C. (2008). Obesity classification in military personnel: A comparison of body fat, waist circumference, and body mass index measurements. Retrieved from <http://krex.ksu.edu>

Published Version Information

Citation: Heinrich, K. M., Jitnarin, N., Suminski, R. R., Berkel, L., Hunter, C. M., Alvarez, L., . . . Poston, W. S. C. (2008). Obesity classification in military personnel: A comparison of body fat, waist circumference, and body mass index measurements. *Military Medicine: International Journal of AMSUS*, 173(1), 67-73.

Copyright: Copyright © 2008 AMSUS – Association of Military Surgeons of the U.S.

Publisher's Link: <http://www.ingentaconnect.com/content/amsus>

This item was retrieved from the K-State Research Exchange (K-REx), the institutional repository of Kansas State University. K-REx is available at <http://krex.ksu.edu>

1 **Pages:** 26
 2 **Words:** 2778
 3 **Tables:** 3
 4 **Figures:** 3
 5 **References:** 28
 6 **Contact:** Katie M. Heinrich, Ph.D.
 7 **Email:** katiemh@hawaii.edu
 8 **Guarantor:** Katie M. Heinrich, Ph.D.

9
 10 **Obesity classification in military personnel: a comparison of body fat, waist**
 11 **circumference, and body mass index measurements**

12 **Katie M. Heinrich, Ph.D.***
 13 **Nattinee Jitnarin, M.S.†**
 14 **Richard R. Suminski, Ph.D.‡**
 15 **LaVerne Berkel, Ph.D. †**
 16 **Christine M. Hunter, LCDR USPHS§**
 17 **Lisa Alvarez, Ph.D.¶**
 18 **Antionette R. Brundige, M.A. ||**
 19 **Alan L. Peterson, LTCOL, USAF-Retired || ****
 20 **John P. Foreyt, Ph.D. ¶**
 21 **C. Keith Haddock, Ph.D. †**
 22 **Walker S.C. Poston, Ph.D. †**

23
 24 *Department of Public Health Sciences, University of Hawaii at Manoa, 1960 East-West
 25 Road, Biomed D104B, Honolulu, HI 96822

26 †Department of Informatic Medicine and Personalized Health, University of Missouri-
 27 Kansas City School of Medicine, Western Missouri Mental Health Center, 1000 East
 28 24th Street, 5th Floor, Kansas City, MO 64108

29 ‡Department of Physiology, Kansas City University of Medicine and Biosciences, 1750
 30 Independence Ave., 468 Strickland Education Pavilion, Kansas City, MO 64106

31 §Division of Diabetes, Endocrinology, and Metabolic Diseases, NIDDK, Two Democracy
 32 Place, 605, 6707 Democracy BLVD, Bethesda, MD 20892

33 ¶Behavioral Medicine Research Center, Department of Medicine, Baylor College of
 34 Medicine, One Baylor Plaza, Houston, TX 77030

35 || Wilford Hall Medical Center, United States Air Force, Lackland AFB, TX 78236

36 **Department of Psychiatry and Behavioral Science, University of Texas Health Science
 37 Center in San Antonio, 7000 Fannin Street, Houston, TX 77030

38
 39
 40
 41
 42

43 **Keywords:** military personnel, obesity, body composition, body fat percentage, body
 44 mass index, waist circumference, ROC curves

ABSTRACT

45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61

Objective: To evaluate obesity classifications from body fat percentage (BF%), body mass index (BMI), and waist circumference (WC).

Methods: 451 overweight/obese active duty military personnel completed all three assessments.

Results: Most were obese (men=81%; women=98%) using National Institutes of Health (NIH) BF% standards (men>25%; women>30%). Using the higher World Health Organization (WHO) BF>35% standard, 86% of women were obese. BMI (55.5% and 51.4%) and WC (21.4% and 31.9%) obesity rates were substantially lower for men and women, respectively; $p<0.05$. BMI/WC were accurate discriminators for BF%-obesity (Θ for all comparisons >0.75 , $p<0.001$). Optimal cut-points were lower than NIH/WHO standards; WC=100cm and BMI=29 maximized sensitivity and specificity for men, and WC=79cm and BMI=25.5 (NIH) or WC=83cm and BMI=26 (WHO) maximized sensitivity and specificity for women.

Conclusion: Both WC and BMI measures had high rates of false negatives compared to BF%. However, at a population-level, WC/BMI are useful obesity measures, demonstrating fair-to-high discriminatory power.

62

INTRODUCTION

63 For military personnel with fitness requirements, it is important to accurately
64 determine body composition. High body fat percentage (BF%), or excess adipose tissue
65 is of particular interest because it is related to increased morbidity and mortality risk (1).
66 Dual-energy X-ray absorptiometry (DEXA) often is used as a reference method for body
67 composition analysis and is considered the “gold standard” (2). DEXA is a multi-
68 compartment model technique that examines both segmental and whole lean body
69 mass and BF (2-3). However, the use of DEXA is costly and impractical for field use in
70 large studies (2,4).

71 A comparable body composition measurement device is the Tanita foot-to-foot
72 bioelectrical impedance analyzer. Besides body weight, the Tanita measures resistance
73 to a small electrical current to estimate BF% (5). Tanita analyzers have compared
74 favorably with DEXA ($r=0.94$, $p<0.001$) (5,6-7), and are inexpensive and simple to use
75 in the field (8).

76 Other prediction techniques estimate body composition and BF distribution (5).
77 For example anthropometric waist circumference (WC) measurements assess the
78 regional distribution of BF. In one study (9), WC measurements correlated with BF mass
79 for men and women and with BF% for women, but no associations were found between
80 WC and BF% for men. In addition, the correlations between WC and trunk fat were
81 higher than those between WC and total BF (9).

82 BMI, a ratio of weight to height, is commonly used in population studies. Although
83 BMI is a simple and widely used estimate of weight status in population studies,
84 numerous investigations have questioned its validity because it cannot distinguish

85 between fat and fat free mass (e.g., 10
86 http://www.consumerfreedom.com/news_detail.cfm?headline=2764,11). Individuals
87 with greater muscle mass, such as athletes and military personnel, may be classified as
88 overweight or obese, while individuals who have excess fat, but not excess weight may
89 be misclassified as having lower health risks based on their misleading 'healthy' BMIs.
90 The rate of false negatives also increases with age, as older individuals tend to have
91 higher body fat percentages than younger individuals with the same BMI (12-13).

92 Often WC or BMI measurements are used to screen for overweight and obesity
93 in military personnel (14). Individuals exceeding measurement thresholds may have to
94 undergo retraining programs or even dismissal (8). Because military populations are
95 more active and younger than the general population, the validity of BMI for categorizing
96 obesity and estimating BF% has been questioned (15). The purpose of this study was
97 to evaluate the relationships between bioelectrical impedance determined BF%, WC,
98 and BMI determined obesity in a military sample.

99 **METHOD**

100 **Participants**

101 A total of 451 participants were recruited and randomized in the parent weight gain
102 prevention clinical trial (16). Inclusion criteria for this study included the following: 1) within
103 five pounds below or equal or exceeding their Maximum Allowable Weight according to
104 USAF Weight and Height tables (AFI 40-502, which was deactivated when the new USAF
105 Fitness Program [AFI 10-248] went into effect in January 2004); 2) access to personal
106 computer with Internet access; 3) plan to remain in the local area for one year; and 4) male
107 or female between ages 18-55 years. Exclusion criteria included: 1) pregnant, breast-

108 feeding, planning to become pregnant or became pregnant; 2) weight loss of >10 pounds
109 in last 3 months; 3) use of a prescription or nonprescription weight-loss medication during
110 the 6 months prior to screening; 4) on any military medical profile; or 5) meeting a specific
111 exclusion such as history of myocardial infarction, stroke, or cancer in the last 5 years,
112 diabetes, angina, and orthopedic or joint problems that would prohibit exercise. Study
113 procedures were approved by the Institutional Review Boards of Wilford Hall Medical
114 Center, Baylor College of Medicine, and U.S. Army Medical Research and Materiel
115 Command Fort Detrick, Maryland. All participants gave informed consent.

116 **Measures**

117 Demographics: Each participant was asked to complete basic demographic
118 information to include rank, age, race/ethnicity, marital status, years of education, years
119 of military service, and whether they planned to retire from the military after at least 20
120 years of service.

121 Body Fat Percentage Estimation –BF% was estimated using a field-based Tanita
122 Body Composition Analyzer foot-to-foot with scale (Tanita corp., Tokyo, Japan). Each
123 military member wore an approved uniform of the day or standard physical training
124 uniform or gym clothes without shoes or socks. The National Institutes of Health (NIH)
125 cut-points to indicate obesity were used (i.e., >25% for men and >30% for women) (17).

126 Waist Circumference – WC was measured with a Gulick tape according to
127 procedures outlined in AFI 10-248 and DoD Instruction 1308.3 (Figure 1.) using
128 standardized anatomical landmarks, i.e., the iliac creast and the umbilicus, for women
129 and men (18). Cut-points of WC >88cm (35in) for women and WC >102cm (40in) for
130 men were used to indicate obesity (18).

131 <<<<Figure 1>>>>

132 Weight and Height – Weight and height were assessed at military Health and
133 Wellness Centers and followed the procedures outlined in AFI 40-502 (The Weight and
134 Body Fat Management Program and DoD Instruction 1308.3). Height was measured
135 with the military member standing on a flat surface without shoes. Weight also was
136 measured in an approved uniform or gym clothes without shoes. Measurements were
137 made on calibrated scales and recorded to the nearest pound.

138 Body Mass Index (BMI) – BMI was determined by dividing weight in kilograms by
139 height in meters². Both the National Institutes of Health (NIH) and World Health
140 Organization (WHO) have BMI guidelines estimating body fatness and corresponding
141 with morbidity and mortality risks. Current NIH guidelines were used to categorize
142 individuals with a BMI of 25 to 29.9 as overweight and those with BMIs ≥ 30 as obese
143 (17). For women, the more liberal WHO cut-point of 35% was also used to indicate
144 obesity (19).

145 Blood Pressure – Blood pressure was measured following the standard
146 epidemiological protocol, i.e., five minutes of rest in a seated position and then three
147 separate blood pressure measurements using alternating arms with a mercury
148 sphygmomanometer separated by 2 minutes between each reading (20). The first
149 reading was omitted and the last two averaged to obtain each subject's blood pressure.
150 Participants had not smoked for at least one hour prior to the measurement session.

151 **Statistical Analysis**

152 Statistical analyses were performed using SPSS[®] (version 14.0; SPSS Inc.,
153 Chicago, IL, USA). Means \pm standard deviation scores or percentages were calculated

154 for all baseline demographic variables stratified by each obesity criterion (i.e., BF%,
155 WC, and BMI) and gender. The NIH (17) obesity standards based on BF%, WC and
156 BMI were used for the primary analyses. Additionally, we recomputed all analyses for
157 women using the more liberal WHO standards presented by DeLorenzo and colleagues
158 (21), i.e., BF% >35% for women, because these values more closely match the WHO
159 standards for obesity (19). Sensitivity, specificity, rates of false positives, false
160 negatives, and accuracy also were computed.

161 Receiver Operating Characteristic (ROC) curves were computed for WC and BMI
162 using BF% as the criterion. ROC curves assess the ability of diagnostic or screening
163 tests, such as WC and BMI, to correctly classify disease status or health outcomes. The
164 area under the curve (AUC) is a quantitative method of evaluating test accuracy in
165 discriminating between diseased or not (or healthy or not). Conventionally, the AUC
166 (often expressed as Θ , or theta) is expressed as a single number and ranges from 0.5
167 (no accuracy) to 1.0 (perfect accuracy) (22-23). AUC Guidelines for any test are 0.5-
168 0.7=none to low discriminatory power, 0.7-0.8=fair discriminatory power, 0.8-0.9=good
169 discriminatory power, and > 0.9=high discriminatory power. Tests can be compared to
170 one another on one criterion by examining their 95% confidence intervals for overlap or
171 non-overlap with overlapping intervals indicating that the methods are statistically
172 similar.

173 RESULTS

174 **Table 1** provides data on the demographic characteristics of the men and women
175 who participated in the study.

176 <<<<Table 1>>>>

177 On average, the sample consisted of an equal number of men and women in their early
178 to mid thirties who were married, educated and planning to retire from the military. Over
179 half of male participants were White, 15.3% were African American, 19.8% were
180 Hispanic, 3.6% were Asian or Pacific Islander, and 3.6% were of other ethnicities, while
181 53.3% of female participants were White, 30.6% were African American, 11.8% were
182 Hispanic, 1.3% were Asian or Pacific Islander, and 3.1% were of other ethnicities.
183 Obesity classification by BF% (>25% for men and >30% for women using the NIH [17]
184 standard) indicated that 81.1% of the men and 98.3% of the women were obese.
185 Obesity prevalence estimates based on WC and BMI also are provided in Table 1.

186 Next, we examined obesity prevalence for women using the alternate and higher
187 BF% cutoff (>35%) suggested by DeLorenzo and colleagues (21). Using this criterion,
188 only 85.6% were classified as obese, rather than the 98.3% found using the more
189 stringent NIH standard (17). Correlations between BF%, WC, and BMI were high and
190 statistically significant for both men ($r_{BF\%-WC}=0.629$; $r_{BF\%-BMI}=0.759$; $r_{WC-BMI}=0.741$; all
191 $p<0.001$) and for women ($r_{BF\%-WC}=0.626$; $r_{BF\%-BMI}=0.691$; $r_{WC-BMI}=0.665$; all $p<0.001$),
192 respectively.

193 Obesity rates, as estimated by WC and BMI were much lower than those derived
194 from BF% (see Table 1; $p<0.05$ for both women and men) In addition, a higher
195 percentage of the men were classified as obese according to WC and BMI standards as
196 compared to the women. For women, and to a lesser extent men, BF% tended to be
197 somewhat high for the groups classified as obese using WC and BMI. For example,
198 women designated as obese using BMI standards had an average BF% of 43.9% which
199 exceeds the 40% BF% used to define morbid obesity (24).

200 As shown in **Table 2**, WC and BMI methods were accurate and statistically
201 significant for discriminating between BF%-defined obesity.

202 <<<<Table 2>>>>

203 While WC only showed fair discriminatory power in men, it demonstrated high
204 discriminatory power in women (i.e., $\Theta > 0.900$). For both men and women, BMI
205 displayed good discriminatory power for accurately predicting obesity, with both AUC
206 (Θ) values exceeding 0.800. However, both methods were statistically similar as
207 evidenced by the overlap in the 95% confidence intervals for each gender. The AUC
208 also was computed using the alternate BF% criterion for women (i.e., $>35\%$). As can be
209 seen in Table 2, using this criterion, both WC and BMI demonstrated good
210 discriminatory power and were statistically equivalent. However, in both men and
211 women (using both BF% criteria), the cutpoints for WC and BMI for optimizing
212 detection of obesity were somewhat lower than the NIH (17) standards. For example,
213 for men, a WC=100cm and BMI=29 maximized both sensitivity and specificity (i.e., both
214 $>65\%$). For women, a WC=79cm and BMI=25.5 maximized detection of obesity using
215 the NIH (17) BF% criterion while a WC=83cm and BMI=26 maximized detection using
216 the alternate criterion of BF% >35 .

217 The sensitivity, specificity, and accuracy of WC and BMI for predicting BF%-
218 based obesity using the NIH (17) criterion and alternate standard for women
219 (BF% $>35\%$) are provided in **Table 3**. **Figure 2** presents the false positive and false
220 negative rates.

221 <<<<Table 3>>>>

222 <<<<Figure 2>>>>

223 More non-obese men (BF% < 25%) were misclassified as obese using the BMI method
224 than the WC method (35% vs. 21%). All non-obese women (BF% < 30%) were
225 correctly identified as such by both methods (specificity = 100%). WC and BMI were
226 not very accurate for correctly identifying obese subjects, especially obese women
227 using the NIH standard for BF% (17). Specifically, 78% of BF%-defined obese women
228 were misclassified as non-obese (i.e., false negatives) using BMI standards and 68%
229 were misclassified as non-obese using WC standards. In BF%-defined obese men,
230 35% and 42% were misclassified with the BMI and WC methods, respectively. The
231 proportion of men correctly identified as being obese or not obese using the WC and
232 BMI methods was two- to three-fold greater than the proportion of women correctly
233 identified; i.e., accuracy was 62.0% for the WC method in men versus 34.0% in women
234 and 69.0% for the BMI method in men versus 23.0% in women. Using the alternate
235 BF% standard of >35% for women (19) only minimally improved accuracy and
236 sensitivity (see Table 3).

237 WC was an acceptable predictor of BMI-based obesity in men and women. As
238 shown in **Figure 3**, only 26% of the men and 22% of the women were misclassified
239 using WC to predict BMI-based obesity.

240 <<<<Figure 3>>>>

241 The percentages of individuals misclassified by WC compared to the BMI criterion were
242 evenly distributed between false negatives (22.5%) and false positives (25.5%).

243 **DISCUSSION**

244 The purpose of this study was to examine relationships among BF%, WC and
245 BMI in a sample of military personnel. Like similar investigations (e.g., 21), we found

246 that both WC and BMI underestimated obesity compared with BF%. Whereas
247 approximately 50% of men and 21% to 32% of women were classified as obese using
248 WC or BMI methods, 80% of men and up to 98% of women were classified as obese
249 utilizing the BF% cutoffs for men and women, respectively. This finding is more
250 noteworthy when examined in light of the actual BF% values for the individuals
251 classified as obese in this military sample, which tended to be high. In addition, this
252 also is notable because the field method we used to determine BF% tends to
253 underestimate BF% when compared with DEXA (7). This pattern of results indicates
254 that regardless of method used to assess or estimate BF%, a significant number of the
255 military personnel in this study had rates of excess body fat that put them at higher risk
256 for cardiovascular disease and other obesity-related comorbidities.

257 Although both WC and BMI tended to underestimate obesity compared to BF%,
258 both exhibited statistically acceptable discriminatory power for accurately predicting
259 obesity in women and WC displayed fair discriminatory power for predicting obesity in
260 men based on AUC statistics. Even so, 68% to 78% of BF%-defined obese women
261 were misclassified as non-obese according to WC and BMI standards, respectively.
262 Accuracy improved somewhat for men, with only 35% (WC) and 42% (BMI)
263 misclassified. These results should inform policy or practice in how WC and BMI
264 derived data are utilized. For example, concerns about the potential for false positives,
265 or a high number of athletic individuals with low percentages of body fat being
266 misclassified as obese in the military population, were not supported by our data.
267 Rather, what emerged from these data, consistent with similar investigations, has been
268 a higher rate of false negatives, or individuals whose weight to height ratio would

269 suggest minimal risk for morbidity or mortality, but whose BF% suggests otherwise.
270 This study also found that the cutpoints for WC and BMI for optimizing detection of
271 obesity were lower than national standards, echoing previously stated concerns about
272 the accuracy of these cutpoints specifically related to age, gender, and ethnicity (e.g.,
273 4,12,14).

274 Although not ideal, both WC and BMI were statistically equivalent and reasonably
275 accurate in predicting BF% obesity in the present study. Of the two, BMI is preferred
276 from a field perspective because it is less intrusive (does not require subjects to remove
277 or raise their clothing), is more comparable across studies (compared with WC, which
278 can be measured in a number of different ways, yielding varied results), and is simpler
279 to obtain and report (i.e., it does not require special training). The ideal field situation,
280 however, may be to combine WC and BMI data to improve prediction (25).

281 The findings of this study should be considered in light of its limitations. First, the
282 instrument used to measure BF% was the Tanita bioelectrical impedance measure, a
283 good field measure (4), but not the “gold standard.” Second, the participants were not a
284 random sample from the general military population, but rather a group active duty U.S.
285 Air Force personnel who volunteered to participate in a weight gain prevention study.
286 The results, therefore, may not generalize to the larger military population. Research
287 also has shown differences to exist between BF%, BMI and WC for different genders,
288 ages, and ethnicities (26-28). Future studies with larger samples would be able to
289 conduct ROC analyses among racial subgroups in order to determine if differences exist
290 in areas under the curve for different racial groups in the military.

291 In conclusion, BF%, WC, and BMI, were significantly correlated for both men and
292 women, although obesity rates varied substantially depending on the method of
293 determination. When using BF% as the criterion, obesity identified using WC and BMI
294 was more accurate for men than for women. For this sample, optimal cutpoints for
295 identifying obese men were lower than the national standards (WC=100cm vs 102cm,
296 and BMI=29 vs 30), as were optimal cutpoints for identifying obese women (WC=79cm
297 vs 88cm and BMI=26 vs 30). While both WC and BMI tended to underestimate obesity
298 rates as compared to BF%, both are adequate for use in large clinical and population
299 studies, with BMI being the preferred, less-intrusive method. Future research should
300 examine a larger, more representative military sample using all three obesity
301 measurements.

ACKNOWLEDGEMENTS

Trial Registry: ClinicalTrials.gov: Preventing overweight in USAF personnel: Minimal contact program, NCT00417599

This work was supported by a grant from the U.S. Army Medical Research and Material Command's Peer Review Medical Research Program (DAMD17-02-2-0018).

REFERENCES

1. Mokdad AH, Ford ES, Bowman BA, Dietz WH, Vinicor F, Bales VS, Marks JS. Prevalence of obesity, diabetes, and obesity-related health risk factors, 2001. *JAMA* 2003;289(1):76-79.
2. van der Ploeg GE, Withers RT, Laforgia J. Percent body fat via DEXA: comparison with a four-compartment model. *J Appl Physiol* 2003;94:499-506.
3. Watts K, Naylor LH, Davis EA, Jones TW, Beeson B, Bettenay F, Siafarikas A, Bell L, Ackland T, Green DJ. Do skinfolds accurately assess changes in body fat in obese children and adolescents? *Med Sci Sports Exerc* 2006;38(3):439-444.
4. Deurenberg-Yap M, Schmidt G, van Staveren WA, Deurenberg P. The paradox of low body mass index and high body fat percentage among Chinese, Malays and Indians in Singapore. *Int J Obes* 2000;24:1011-1017.
5. Jebb SA, Cole TJ, Doman D, Murgatroyd PR, Prentice AM. Evaluation of the novel Tanita body-fat analyser to measure body composition by comparison with a four-compartment model. *Br J Nutr* 2000;83:115-122.
6. Nunez C, Gallagher D, Visser M, Pi-Sunyer FX, Wang Z, Heymsfield SB. Bioimpedance analysis: evaluation of leg-to-leg system based on pressure contact foot-pad electrodes. *Med Sci Sports Exerc.* 1997;29(4):524-531.
7. Rubiano F, Nunez C, Heymsfield SB. A comparison of body composition techniques. *Ann N Y Acad Sci* 2000;204:335-338.
8. Prentice AM, Jebb SA. Beyond body mass index. *Obes Rev* 2001;2(3):141-147.

9. Wang J, Thornton JC, Bari S, Williamson B, Gallagher D, Heymsfield SB, Horlick M, Kotler D, Laferrere B, Mayer I, Pi-Sunyer FX, Pierson RN Jr. Comparisons of waist circumference measured at 4 sites. *Am J Clin Nutr* 2003;77(2):379-384.
10. Consumerfreedom.com. Unsportsmanlike BMI deserves huge technical foul. http://www.consumerfreedom.com/news_detail.cfm?headline=2764; 2005
Accessed August 30, 2006.
11. Poston WSC, Foreyt JP. Body mass index: uses and limitations. *Strength Cond J* 2002;24(4):15-17.
12. Gallagher D, Visser M, Sepulveda D, Pierson RN, Harris T, Heymsfield SB. How useful is body mass index for comparison of body fatness across age, sex, and ethnic groups? *Am J Epidemiol* 1996;143(3):228-239.
13. Movsesyan L, Tanko LB, Larsen PH, Christiansen C, Svendsen OL. Variations in percentage of body fat within different BMI groups in young, middle-aged and old women. *Clin Physiol Funct Imaging* 2003;23:130-133.
14. Deurenberg P, Deurenberg-Yap M. Validity of body composition methods across ethnic population groups. *Acta Diabetol* 2003;40:S246-S249.
15. Haddock CK, Poston WS, Klesges RC, Talcott GW, Lando H, Dill PL. An examination of body weight standards and the association between weight and health behaviors in the United States Air Force. *Mil Med* 1999;164:51-4.
16. Hunter CM, Alvarez L, Brundige AR, Poston WS, Peterson A, Van Brunt D, Haddock C, Foreyt J. Controlled trial of an interactive internet weight loss program: six-month results. *Obes Res* 2005;13:A2-A3.

17. National Institutes of Health. Clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults: the evidence report. U.S. Department of Health and Human Services. NIH Publication No. 98-4083; 1998.
18. Lohman TG, Roche AF, Martorell R. Anthropometric Standardization Reference Manual. Champaign, IL: Human Kinetics; 1988
19. World Health Organization. Obesity: preventing and managing the global epidemic. Report of a WHO consultation. Geneva: World Health Organization; 2004.
20. National Heart, Lung, and Blood Institute. JNC 7 Express: the seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. U.S. Department of Health and Human Services. NIH Publication No. 03-5233, 2003.
21. De Lorenzo A, Deurenberg P, Pietrantuono M, Di Daniele N, Cervelli V, Andreoli A. How fat is obese? *Acta Diabetol* 2003;40:S254-S257.
22. Hanley JA, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiol* 1982;143(1):29-36.
23. Zweig MH, Campbell G. Receiver operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. *Clin Chem* 1993;39(4):561-577.
24. National Institutes of Health. The practical guide: identification, evaluation, and treatment of overweight and obesity in adults. U.S. Department of Health and Human Services, NIH Publication No. 00-4084; 2000.

25. Booth M, Hunter C, Gore C, Bauman A, Owen N. The relationship between body mass index and waist circumference: implications for estimates of the population prevalence of overweight. *Int J Obes* 2000;24:1058-1061.
26. Wagner DR, Heyward VH. Measures of body composition in blacks and whites: a comparative review. *Am J Clin Nutr* 2000;71:1392-1402.
27. Ko GTC, Tang J, Chan JCN, Sung R, Wu MMR, Wai HPS, Chen R. Lower BMI cut-off value to define obesity in Hong Kong Chinese: an analysis based on body fat assessment by bioelectrical impedance. *Br J Nutr* 2001;85:239-242.
28. Okusun IS, Tedders SH, Choi S, Dever GE. Abdominal adiposity values associated with established body mass indexes in white, black and Hispanic Americans. A study from the third national health and nutrition examination survey. *Int J Obes Relat Metab Disord* 2000;24(10):1279-85.

Table 1. Demographic variables classified by gender.¹

Men (N = 222)								
Variable	Obesity classified by BF% ²		Obesity classified by WC ³		Obesity classified by BMI ⁴			
	No	Yes	No	Yes	No	Yes		
Number (%)	42 (18.9)	180 (81.1)	108 (48.6)	114 (51.4)	100 (45.0)	122 (55.0)		
Age (years)	36.1±8.0	35.1±6.4	34.0±6.9	36.5±6.4	35.2±7.0	35.4±6.5		
Percent Caucasian	66.7	55.6	53.7	61.4	67.0	50.0		
Percent Married	81.0	82.8	83.3	81.6	84.0	81.1		
Percent with ≥ Some College	97.6	96.7	95.4	98.2	97.0	96.7		
Percent Enlisted	69.0	77.8	79.6	72.8	76.0	76.2		
Years of Service	13.8±7.9	13.9±6.3	12.8±6.6	14.8±6.5	13.6±6.8	14.0±6.2		
Percent Planning to Retire from AF	76.2	86.0	81.3	86.8	85.0	83.5		
BMI (kg/m ²)	28.6±1.6	31.4±3.0	29.3±1.5	32.4±2.8	28.6±0.9	32.7±2.3		
Waist Circumference (cm)	97.1±6.9	104.0±7.3	96.6±4.3	108.5±5.4	98.2±5.9	106.4±7.0		
Body Fat %	23.3±1.7	30.6±4.3	26.8±3.4	31.6±4.9	26.1±2.7	31.8±4.7		
Women (N = 229)								
Variable	Obesity classified by BF% ²		Obesity classified by BF% ⁵		Obesity classified by WC ³		Obesity classified by BMI ⁴	
	No	Yes	No	Yes	No	Yes	No	Yes
Number (%)	4 (1.7)	225 (98.3)	33 (14.4)	196 (85.6)	156 (68.1)	73 (31.9)	180 (78.6)	49 (21.4)
Age (years)	30.5±6.0	32.5±7.7	32.3±9.1	32.5±7.4	32.0±7.8	33.5±7.2	32.1±7.7	34.0±7.3
Percent Caucasian	25.0	53.8	63.6	51.5	51.9	56.2	55.0	46.9
Percent Married	75.0	60.4	57.6	61.2	59.6	63.0	61.1	59.2

Percent with \geq Some College	50.0	89.3	93.9	88.8	92.9	82.2	90.6	85.7
Percent Enlisted	75.0	81.3	69.7	83.2	77.6	89.0	78.3	91.8
Years of Service	9.6 \pm 6.4	11.5 \pm 6.4	10.0 \pm 7.1	11.7 \pm 6.2	10.8 \pm 6.4	12.8 \pm 6.2	10.9\pm6.4	13.4\pm5.8
Percent Planning to Retire from AF	33.3	75.9	68.8	76.4	73.4	79.5	71.9	87.8
BMI (kg/m ²)	25.5\pm1.0	28.0\pm2.4	25.8\pm1.0	28.3\pm2.4	27.1\pm1.9	29.9\pm2.4	27.0\pm1.5	31.6\pm1.5
Waist Circumference (cm)	77.2\pm2.4	86.3\pm6.5	80.2\pm4.6	87.1\pm6.3	82.6\pm3.9	93.7\pm4.1	84.2\pm5.2	93.4\pm5.7
Body Fat %	28.8\pm1.0	39.7\pm4.0	32.6\pm2.0	40.6\pm3.3	38.1\pm3.9	42.5\pm3.4	38.3\pm3.6	43.9\pm3.5

¹Bolded values indicate statistically significant differences, $p < 0.05$.

²NIH standards for obesity based on body fat (BF) percentage are $>25\%$ for men and $>30\%$ for women (17).

³NIH standards for obesity based on waist circumference (WC) are $>88\text{cm}$ (35in) for women and $>102\text{cm}$ (40in) for men (17).

⁴NIH standards for obesity based on body mass index (BMI) are ≥ 30 (17).

⁵Standards for obesity based on body fat (BF) percentage $>35\%$ for women suggested by DeLorenzo et al. (25).

Table 2. Area under the curve (AUC) using waist circumference (WC) and BMI as the predictors of body fat percentage-based obesity stratified by gender.

Men Using NIH¹ BF% Criterion				
Variables	AUC² (SE)³	P-Value	95% Confidence Interval	
			Lower Bound	Upper Bound
WC	0.761 (0.040)	< 0.001	0.683	0.839
BMI	0.841 (0.034)	< 0.001	0.774	0.907
Women Using NIH¹ BF% Criterion				
Variables	AUC² (SE)³	P-Value	95% Confidence Interval	
			Lower Bound	Upper Bound
WC	0.925 (0.033)	0.004	0.861	0.990
BMI	0.826 (0.069)	0.026	0.690	0.961
Women Using WHO⁴ BF% Criterion				
Variables	AUC² (SE)³	P-Value	95% Confidence Interval	
			Lower Bound	Upper Bound
WC	0.807 (0.037)	< 0.001	0.735	0.880
BMI	0.833 (0.029)	< 0.001	0.777	0.889

¹NIH (17) standards for obesity based on body fat (BF) percentage are >25% for men and >30% for women.

²AUC=Area Under the Curve

³Standard Error

⁴Standards for obesity based on BF% are >25% for men and >35% for women (25).

Table 3. Specificity, Sensitivity and Accuracy using waist circumference (WC) and BMI as predictors of BF%-based obesity and using WC as a predictor of BMI-based obesity.

Men Using NIH¹ BF% Criterion			
Variable	Specificity² (%)	Sensitivity³ (%)	Accuracy⁴ (%)
WC as a predictor of BF%-based obesity	79.0	58.0	62.0
BMI as a predictor of BF%-based obesity	65.0	65.0	69.0
WC as a predictor of BMI-based obesity	75.0	73.0	74.0
Women Using NIH¹ BF% Criterion			
Variable	Specificity² (%)	Sensitivity³ (%)	Accuracy⁴ (%)
WC as a predictor of BF%-based obesity	100.0	32.0	34.0
BMI as a predictor of BF%-based obesity	100.0	22.0	23.0
WC as a predictor of BMI-based obesity	76.0	80.0	79.0
Women Using WHO⁵ BF% Criterion			
Variable	Specificity² (%)	Sensitivity³ (%)	Accuracy⁴ (%)
WC as a predictor of BF%-based obesity	100.0	37.0	46.0%
BMI as a predictor of BF%-based obesity	100.0	25.0	36.0

¹NIH standards for obesity based on body fat (BF) percentage are >25% for men and >30% for women.

²Specificity is defined as is the proportion of true negatives identified by the screening test divided by all those free of the disease or disorder.

³Sensitivity is defined as is the proportion of true positives identified by the screening test divided by all those with the disease or disorder.

⁴Accuracy is defined as the proportions of individuals correctly screened as having or not having the disease divided by the total sample population.

⁵Standards for obesity based on body fat (BF) percentage are >25% for men and >35% for women suggested by DeLorenzo et al. (2003).

Figure Legends

Figure 1.

Not Applicable

Figure 2.

■ False Negative □ False Positive

Figure 3.

■ False Negative □ False Positive

Figure 1. Waist Circumference Measurement Procedure

A7.3.3. A seamstress tape measure will be used for the abdominal circumference.

A7.3.4. Member stands looking straight ahead, arms down to sides.

A7.3.5. Examiner is positioned at right side of the member.

A7.3.6. Measurement is taken on bare skin; examiner feels to locate the upper hipbone and top of the right iliac crest.

A7.3.7. A horizontal landmark is located just above the uppermost border of the right iliac crest.

A7.3.8. The tape is placed in a horizontal plane around the abdomen at the level of this landmark. Examiner ensures that the plane of the tape is parallel to the floor and that the tape is snug, but does not compress the skin. Measurement is taken at the end of a normal respiration.

A7.3.9. Take the circumference measure three times and record each measurement to the nearest $\frac{1}{2}$ inch. If any of the measures differ by more than one inch from the other two, take an additional measurement. Add the three closest measurements, divide by 3, and round down to the nearest $\frac{1}{2}$ inch. Record this value as the abdominal circumference measure.

Measuring Tape Position for Abdominal Circumference.






Figure 2. Percentage of false negatives* and false positives for the classification of obesity using waist circumference and body mass index with NIH body fat percentage as the criterion in military men and women and DeLorenzo et al. (2003) body fat percentage criterion for women only.**




The study was conducted at a military medical and research center and three military bases in the Southwest US between the dates of June 2003 and October 2005.

*False negatives are defined as the proportion of individuals who have the disease but are screened as not having the disease.

**False positives are defined as the proportion of individuals who do not have the disease but are screened as having the disease.

Figure 3. Percentage of false negatives* and false positives for the classification of obesity using NIH waist circumference with NIH body mass index as the criterion in military men and women.**



The study was conducted at a military medical and research center and three military bases in the Southwest US between the dates of June 2003 and October 2005.

*False negatives are defined as the proportion of individuals who have the disease but are screened as not having the disease.

**False positives are defined as the proportion of individuals who do not have the disease but are screened as having the disease.