

Acta Crystallographica Section E

**Structure Reports****Online**


ISSN 1600-5368

Editors: **W.T.A. Harrison, J. Simpson and M. Weil**

## 4-Carbamoylpiperidinium 5-nitrosalicylate

**Graham Smith and Urs D. Wermuth***Acta Cryst.* (2011). **E67**, o122

This open-access article is distributed under the terms of the Creative Commons Attribution Licence <http://creativecommons.org/licenses/by/2.0/uk/legalcode>, which permits unrestricted use, distribution, and reproduction in any medium, provided the original authors and source are cited.



*Acta Crystallographica Section E: Structure Reports Online* is the IUCr's highly popular open-access structural journal. It provides a simple and easily accessible publication mechanism for the growing number of inorganic, metal-organic and organic crystal structure determinations. The electronic submission, validation, refereeing and publication facilities of the journal ensure very rapid and high-quality publication, whilst key indicators and validation reports provide measures of structural reliability. In 2007, the journal published over 5000 structures. The average publication time is less than one month.

Crystallography Journals **Online** is available from [journals.iucr.org](http://journals.iucr.org)

## 4-Carbamoylpiperidinium 5-nitrosalicylate

Graham Smith\* and Urs D. Wermuth

Faculty of Science and Technology, Queensland University of Technology, GPO Box 2434, Brisbane, Queensland 4001, Australia  
Correspondence e-mail: g.smith@qut.edu.au


Received 29 November 2010; accepted 30 November 2010

Key indicators: single-crystal X-ray study;  $T = 200$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.040;  $wR$  factor = 0.092; data-to-parameter ratio = 12.9.

In the crystal structure of the title compound,  $\text{C}_6\text{H}_{13}\text{N}_2\text{O}^+ \cdot \text{C}_7\text{H}_4\text{NO}_5^-$ , the isonipecotamide cations and the 5-nitrosalicylate anions form hydrogen-bonded chain substructures through head-to-tail piperidinium–carboxylate  $\text{N}-\text{H} \cdots \text{O}$  hydrogen bonds and through centrosymmetric cyclic head-to-head amide–amide hydrogen-bonding associations [graph set  $R_2^2(8)$ ]. These chains are cross-linked by amide–carboxylate  $\text{N}-\text{H} \cdots \text{O}$  and piperidinium–nitro  $\text{N}-\text{H} \cdots \text{O}$  associations, giving a sheet structure.

### Related literature

For structural data on isonipecotamide salts, see: Smith *et al.* (2010); Smith & Wermuth (2010*a,b,c,d*). For structures of 5-nitrosalicylates, see: Smith *et al.* (2005). For hydrogen-bonding graph-set and motif classification, see: Etter *et al.* (1990); Allen *et al.* (1998).



### Experimental

#### Crystal data

$\text{C}_6\text{H}_{13}\text{N}_2\text{O}^+ \cdot \text{C}_7\text{H}_4\text{NO}_5^-$   
 $M_r = 311.30$   
Monoclinic,  $P2_1/n$   
 $a = 15.0442$  (10) Å  
 $b = 5.5851$  (3) Å  
 $c = 17.1939$  (10) Å  
 $\beta = 91.466$  (6)°

$V = 1444.22$  (15) Å<sup>3</sup>  
 $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.12$  mm<sup>-1</sup>  
 $T = 200$  K  
 $0.40 \times 0.25 \times 0.16$  mm

#### Data collection

Oxford Diffraction Gemini-S CCD-detector diffractometer  
Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)  
 $T_{\min} = 0.912$ ,  $T_{\max} = 0.980$   
9191 measured reflections  
2833 independent reflections  
1850 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.031$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.040$   
 $wR(F^2) = 0.092$   
 $S = 0.95$   
2833 reflections  
219 parameters  
H atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.11$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.17$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$\text{N1A}-\text{H11A} \cdots \text{O12}^i$	1.00 (2)	1.71 (2)	2.688 (2)	164.2 (18)
$\text{N1A}-\text{H12A} \cdots \text{O11}$	0.95 (2)	1.80 (2)	2.747 (2)	173.9 (17)
$\text{N41A}-\text{H41A} \cdots \text{O52}^{ii}$	0.83 (2)	2.39 (2)	3.216 (2)	170.8 (19)
$\text{N41A}-\text{H42A} \cdots \text{O41A}^{iii}$	0.99 (2)	1.91 (2)	2.873 (2)	164.8 (18)
$\text{O2}-\text{H2} \cdots \text{O12}$	0.96 (2)	1.58 (2)	2.4897 (18)	156 (2)

Symmetry codes: (i)  $x, y + 1, z$ ; (ii)  $x + \frac{1}{2}, -y + \frac{3}{2}, z + \frac{1}{2}$ ; (iii)  $-x + 1, -y, -z + 2$ .

Data collection: *CrysAlis PRO* (Oxford Diffraction, 2010); cell refinement: *CrysAlis PRO*; data reduction: *CrysAlis PRO*; program(s) used to solve structure: *SIR92* (Altomare *et al.*, 1994); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008) within *WinGX* (Farrugia, 1999); molecular graphics: *PLATON* (Spek, 2009); software used to prepare material for publication: *PLATON*.

The authors acknowledge financial support from the Australian Research Council, the Faculty of Science and Technology and the University Library, Queensland University of Technology, and the School of Biomolecular and Physical Sciences, Griffith University.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5426).

### References

- Allen, F. H., Raithby, P. R., Shields, G. P. & Taylor, R. (1998). *Chem. Commun.*, pp. 1043–1044.
- Altomare, A., Cascarano, G., Giacovazzo, C., Guagliardi, A., Burla, M. C., Polidori, G. & Camalli, M. (1994). *J. Appl. Cryst.* **27**, 435.
- Etter, M. C., MacDonald, J. C. & Bernstein, J. (1990). *Acta Cryst.* **B46**, 256–262.
- Farrugia, L. J. (1999). *J. Appl. Cryst.* **32**, 837–838.
- Oxford Diffraction (2010). *CrysAlis PRO* Oxford Diffraction Ltd, Yarnton, Oxfordshire, England.
- Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
- Smith, G., Hartono, A. W., Wermuth, U. D., Healy, P. C., White, J. M. & Rae, A. D. (2005). *Aust. J. Chem.* **58**, 47–52.
- Smith, G. & Wermuth, U. D. (2010*a*). *Acta Cryst.* **C66**, o609–o613.
- Smith, G. & Wermuth, U. D. (2010*b*). *Acta Cryst.* **C66**, o614–o618.
- Smith, G. & Wermuth, U. D. (2010*c*). *Acta Cryst.* **E66**, o3162.
- Smith, G. & Wermuth, U. D. (2010*d*). *Acta Cryst.* **E66**, o3260.
- Smith, G., Wermuth, U. D. & Young, D. J. (2010). *Acta Cryst.* **E66**, o3160–o3161.
- Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## **supplementary materials**

*Acta Cryst.* (2011). E67, o122 [ doi:10.1107/S1600536810050129 ]

## 4-Carbamoylpiperidinium 5-nitrosalicylate

G. Smith and U. D. Wermuth

### Comment

The structures of a number of salts of the amide piperidine-4-carboxamide (isonipecotamide, INIPA) with a range of carboxylic acids, mainly aromatic, are now known (Smith & Wermuth, 2010*a*, 2010*b*, 2010*c*, Smith & Wermuth, 2010*d*; Smith *et al.*, 2010). The title compound  $C_6H_{13}N_2O^+ C_7H_4NO_5^-$  (I) was obtained from the 1:1 stoichiometric reaction of 5-nitrosalicylic acid with INIPA in methanol and the structure is reported here.

In (I) (Fig. 1) the cations and anions form hydrogen-bonded chain substructures through head-to-tail piperidinium  $N-H\cdots O_{\text{carboxyl}}$  hydrogen bonds and through centrosymmetric cyclic head-to-head amide–amide hydrogen-bonding associations [graph set  $R^2_2(8)$  (Etter *et al.*, 1990)]. These chains are cross linked by amide  $N-H\cdots O_{\text{carboxyl}}$  and piperidinium  $N-H\cdots O_{\text{nitro}}$  associations to giving a two-dimensional sheet structure (Fig. 2). The amide–amide dimer association [the 'amide motif' (Allen *et al.*, 1998)] is relatively common among the INIPA salts (Smith & Wermuth, 2010*b*; Smith *et al.*, 2010).

The 5-nitrosalicylate anions are essentially planar [torsion angles for the carboxyl group (C2–C1–C11–O11), 178.30 (16)° and the nitro group (C4–C5–N5–O52), -175.57 (16)°], which is the usual conformation for this anion in its proton-transfer compounds (Smith *et al.*, 2005).

### Experimental

The title compound was synthesized by heating together under reflux for 10 minutes, 1 mmol quantities of piperidine-4-carboxamide (isonipecotamide) and 5-nitrosalicylic acid in 50 ml of methanol. After concentration to *ca* 30 ml, partial room temperature evaporation of the hot-filtered solution gave pale yellow prisms of the title compound from which a specimen was cleaved for the X-ray analysis.

### Refinement

Hydrogen atoms involved in hydrogen-bonding interactions were located by difference methods and their positional and isotropic displacement parameters were refined. Other H-atoms were included in the refinement at calculated positions using a riding-model approximation [C–H = 0.93–0.98 Å] and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ .

### Figures




Fig. 1. Molecular configuration and atom naming scheme for the INIPA cation and the 5-nitrosalicylate anion in (I). The inter-species hydrogen bond is shown as a dashed line and displacement ellipsoids are drawn at the 40% probability level.




Fig. 2. The hydrogen-bonded chain substructures in (I) showing the cyclic  $R^2_2(8)$  amide–amide and cation–anion associations. Non-associative H atoms are omitted and hydrogen bonds are shown as dashed lines. For symmetry codes, see Table 1.

## 4-carbamoylpiperidine 2-hydroxy-5-nitrobenzoate

### Crystal data

$C_6H_{13}N_2O^+ \cdot C_7H_4NO_5^-$

$M_r = 311.30$

Monoclinic,  $P2_1/n$

Hall symbol:  $-P\ 2yn$

$a = 15.0442\ (10)\ \text{\AA}$

$b = 5.5851\ (3)\ \text{\AA}$

$c = 17.1939\ (10)\ \text{\AA}$

$\beta = 91.466\ (6)^\circ$

$V = 1444.22\ (15)\ \text{\AA}^3$

$Z = 4$

$F(000) = 656$

$D_x = 1.432\ \text{Mg m}^{-3}$

Melting point: 463 K

Mo  $K\alpha$  radiation,  $\lambda = 0.71073\ \text{\AA}$

Cell parameters from 3270 reflections

$\theta = 3.6\text{--}28.7^\circ$

$\mu = 0.12\ \text{mm}^{-1}$

$T = 200\ \text{K}$

Prism, pale yellow

$0.40 \times 0.25 \times 0.16\ \text{mm}$

### Data collection

Oxford Diffraction Gemini-S CCD-detector diffractometer

Radiation source: Enhance (Mo)X-ray source graphite

Detector resolution:  $16.077\ \text{pixels mm}^{-1}$

$\omega$  scans

Absorption correction: multi-scan (*CrysAlis PRO*; Oxford Diffraction, 2010)

$T_{\min} = 0.912$ ,  $T_{\max} = 0.980$

9191 measured reflections

2833 independent reflections

1850 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.031$

$\theta_{\max} = 26.0^\circ$ ,  $\theta_{\min} = 3.6^\circ$

$h = -18 \rightarrow 18$

$k = -6 \rightarrow 6$

$l = -12 \rightarrow 21$

### Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.040$

$wR(F^2) = 0.092$

$S = 0.95$

2833 reflections

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H atoms treated by a mixture of independent and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0486P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

219 parameters

$$\Delta\rho_{\max} = 0.11 \text{ e } \text{\AA}^{-3}$$

0 restraints

$$\Delta\rho_{\min} = -0.17 \text{ e } \text{\AA}^{-3}$$

### Special details

**Geometry.** Bond distances, angles *etc.* have been calculated using the rounded fractional coordinates. All su's are estimated from the variances of the (full) variance-covariance matrix. The cell e.s.d.'s are taken into account in the estimation of distances, angles and torsion angles

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
O41A	0.44900 (9)	0.0250 (2)	0.90133 (7)	0.0575 (5)
N1A	0.39663 (10)	0.5656 (3)	0.69549 (9)	0.0452 (6)
N41A	0.54984 (11)	0.2658 (4)	0.96084 (10)	0.0480 (6)
C2A	0.49383 (12)	0.5253 (3)	0.70538 (12)	0.0545 (7)
C3A	0.51390 (12)	0.3391 (3)	0.76686 (10)	0.0516 (7)
C4A	0.47324 (11)	0.4101 (3)	0.84420 (9)	0.0423 (6)
C5A	0.37359 (11)	0.4529 (3)	0.83207 (10)	0.0451 (6)
C6A	0.35478 (11)	0.6369 (3)	0.76956 (10)	0.0463 (6)
C41A	0.48930 (12)	0.2177 (3)	0.90488 (10)	0.0435 (6)
O2	0.40541 (10)	-0.1530 (2)	0.42604 (8)	0.0605 (5)
O11	0.32739 (8)	0.1529 (2)	0.63160 (7)	0.0507 (4)
O12	0.39253 (9)	-0.1469 (3)	0.57010 (7)	0.0600 (5)
O51	0.16400 (10)	0.7176 (3)	0.34253 (8)	0.0729 (6)
O52	0.17024 (9)	0.7584 (2)	0.46713 (8)	0.0562 (5)
N5	0.19077 (10)	0.6528 (3)	0.40756 (9)	0.0510 (6)
C1	0.32654 (11)	0.1565 (3)	0.49283 (9)	0.0388 (6)
C2	0.35489 (12)	0.0445 (3)	0.42422 (10)	0.0460 (6)
C3	0.32936 (13)	0.1368 (4)	0.35198 (10)	0.0550 (7)
C4	0.27649 (13)	0.3354 (4)	0.34625 (10)	0.0529 (7)
C5	0.24816 (11)	0.4452 (3)	0.41392 (9)	0.0423 (6)
C6	0.27300 (11)	0.3580 (3)	0.48665 (9)	0.0396 (6)
C11	0.35061 (12)	0.0510 (3)	0.57113 (10)	0.0443 (6)
H4A	0.50110	0.55900	0.86250	0.0510*
H11A	0.3874 (12)	0.689 (4)	0.6540 (12)	0.065 (6)*
H12A	0.3699 (12)	0.422 (4)	0.6763 (11)	0.063 (6)*
H21A	0.52260	0.67460	0.72000	0.0650*
H22A	0.51770	0.47330	0.65630	0.0650*
H31A	0.57780	0.32230	0.77390	0.0620*
H32A	0.49000	0.18580	0.75010	0.0620*
H41A	0.5753 (13)	0.398 (4)	0.9620 (11)	0.054 (7)*
H42A	0.5568 (13)	0.147 (4)	1.0029 (12)	0.075 (7)*

## supplementary materials

---

H51A	0.34480	0.30340	0.81770	0.0540*
H52A	0.34870	0.50710	0.88050	0.0540*
H61A	0.29100	0.65250	0.76110	0.0560*
H62A	0.37790	0.79110	0.78630	0.0560*
H2	0.4119 (15)	-0.184 (4)	0.4809 (14)	0.109 (9)*
H3	0.34850	0.06260	0.30700	0.0660*
H4	0.25970	0.39640	0.29770	0.0630*
H6	0.25380	0.43450	0.53130	0.0470*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
O41A	0.0736 (9)	0.0473 (8)	0.0500 (8)	0.0056 (7)	-0.0280 (7)	-0.0127 (6)
N1A	0.0518 (10)	0.0467 (10)	0.0368 (9)	-0.0015 (8)	-0.0070 (7)	-0.0078 (8)
N41A	0.0472 (10)	0.0463 (10)	0.0495 (10)	0.0124 (9)	-0.0208 (8)	-0.0123 (9)
C2A	0.0482 (12)	0.0624 (13)	0.0531 (12)	0.0009 (10)	0.0056 (9)	-0.0120 (10)
C3A	0.0439 (11)	0.0576 (12)	0.0530 (12)	0.0122 (9)	-0.0014 (9)	-0.0134 (10)
C4A	0.0421 (10)	0.0420 (10)	0.0420 (10)	0.0090 (8)	-0.0125 (8)	-0.0149 (8)
C5A	0.0422 (10)	0.0580 (11)	0.0348 (10)	0.0155 (9)	-0.0066 (8)	-0.0066 (9)
C6A	0.0429 (10)	0.0561 (11)	0.0396 (10)	0.0127 (9)	-0.0066 (8)	-0.0080 (9)
C41A	0.0418 (10)	0.0454 (11)	0.0426 (10)	0.0189 (9)	-0.0139 (8)	-0.0195 (9)
O2	0.0834 (10)	0.0489 (8)	0.0501 (8)	-0.0077 (8)	0.0216 (8)	-0.0104 (7)
O11	0.0597 (8)	0.0621 (8)	0.0303 (7)	-0.0163 (7)	-0.0002 (6)	-0.0097 (6)
O12	0.0762 (10)	0.0563 (8)	0.0474 (8)	0.0017 (8)	0.0024 (7)	-0.0007 (7)
O51	0.0929 (11)	0.0800 (11)	0.0455 (8)	-0.0009 (8)	-0.0025 (8)	0.0174 (7)
O52	0.0630 (9)	0.0544 (8)	0.0516 (9)	-0.0081 (7)	0.0075 (7)	-0.0025 (7)
N5	0.0580 (10)	0.0546 (10)	0.0406 (10)	-0.0209 (9)	0.0067 (8)	0.0033 (8)
C1	0.0423 (10)	0.0414 (10)	0.0331 (10)	-0.0211 (9)	0.0064 (7)	-0.0067 (8)
C2	0.0590 (12)	0.0407 (10)	0.0390 (11)	-0.0221 (9)	0.0148 (9)	-0.0066 (9)
C3	0.0810 (15)	0.0528 (12)	0.0322 (11)	-0.0223 (11)	0.0224 (10)	-0.0069 (9)
C4	0.0715 (14)	0.0585 (12)	0.0291 (10)	-0.0264 (11)	0.0099 (9)	0.0020 (9)
C5	0.0483 (11)	0.0438 (11)	0.0352 (10)	-0.0184 (9)	0.0073 (8)	0.0017 (8)
C6	0.0430 (10)	0.0457 (10)	0.0302 (9)	-0.0230 (9)	0.0056 (7)	-0.0079 (8)
C11	0.0453 (11)	0.0505 (11)	0.0369 (11)	-0.0208 (9)	0.0001 (8)	-0.0070 (9)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

O41A—C41A	1.236 (2)	C2A—H22A	0.9700
O2—C2	1.340 (2)	C2A—H21A	0.9700
O11—C11	1.243 (2)	C3A—H31A	0.9700
O12—C11	1.273 (2)	C3A—H32A	0.9700
O51—N5	1.233 (2)	C4A—H4A	0.9800
O52—N5	1.228 (2)	C5A—H51A	0.9700
O2—H2	0.96 (2)	C5A—H52A	0.9700
N1A—C6A	1.489 (2)	C6A—H62A	0.9700
N1A—C2A	1.485 (2)	C6A—H61A	0.9700
N41A—C41A	1.335 (2)	C1—C6	1.386 (2)
N1A—H12A	0.95 (2)	C1—C11	1.505 (2)
N1A—H11A	1.00 (2)	C1—C2	1.411 (2)

N41A—H41A	0.83 (2)	C2—C3	1.390 (3)
N41A—H42A	0.99 (2)	C3—C4	1.367 (3)
N5—C5	1.448 (2)	C4—C5	1.392 (2)
C2A—C3A	1.508 (3)	C5—C6	1.385 (2)
C3A—C4A	1.530 (2)	C3—H3	0.9300
C4A—C5A	1.527 (2)	C4—H4	0.9300
C4A—C41A	1.513 (2)	C6—H6	0.9300
C5A—C6A	1.509 (2)		
C2—O2—H2	102.4 (13)	C5A—C4A—H4A	109.00
C2A—N1A—C6A	112.29 (14)	C3A—C4A—H4A	109.00
C2A—N1A—H12A	108.5 (12)	C4A—C5A—H52A	109.00
C6A—N1A—H11A	111.8 (12)	C6A—C5A—H51A	109.00
H11A—N1A—H12A	106.5 (17)	C4A—C5A—H51A	109.00
C6A—N1A—H12A	109.7 (11)	C6A—C5A—H52A	109.00
C2A—N1A—H11A	107.8 (11)	H51A—C5A—H52A	108.00
H41A—N41A—H42A	122.7 (18)	H61A—C6A—H62A	108.00
C41A—N41A—H42A	116.8 (12)	C5A—C6A—H61A	110.00
C41A—N41A—H41A	120.1 (13)	C5A—C6A—H62A	110.00
O51—N5—O52	122.14 (16)	N1A—C6A—H61A	109.00
O51—N5—C5	118.93 (15)	N1A—C6A—H62A	110.00
O52—N5—C5	118.93 (14)	C2—C1—C11	120.22 (15)
N1A—C2A—C3A	111.33 (15)	C6—C1—C11	120.82 (14)
C2A—C3A—C4A	110.68 (14)	C2—C1—C6	118.90 (15)
C3A—C4A—C41A	110.79 (14)	O2—C2—C1	121.94 (15)
C5A—C4A—C41A	110.14 (14)	O2—C2—C3	118.03 (16)
C3A—C4A—C5A	109.56 (13)	C1—C2—C3	120.02 (16)
C4A—C5A—C6A	111.69 (14)	C2—C3—C4	120.82 (17)
N1A—C6A—C5A	110.57 (14)	C3—C4—C5	119.17 (16)
O41A—C41A—N41A	122.38 (17)	N5—C5—C6	119.77 (14)
N41A—C41A—C4A	116.62 (16)	C4—C5—C6	121.25 (16)
O41A—C41A—C4A	120.99 (15)	N5—C5—C4	118.98 (15)
N1A—C2A—H21A	109.00	C1—C6—C5	119.84 (15)
N1A—C2A—H22A	109.00	O11—C11—C1	120.16 (15)
C3A—C2A—H21A	109.00	O12—C11—C1	115.81 (15)
C3A—C2A—H22A	109.00	O11—C11—O12	123.99 (16)
H21A—C2A—H22A	108.00	C2—C3—H3	120.00
C2A—C3A—H32A	110.00	C4—C3—H3	120.00
C4A—C3A—H31A	109.00	C3—C4—H4	120.00
C2A—C3A—H31A	110.00	C5—C4—H4	120.00
C4A—C3A—H32A	109.00	C1—C6—H6	120.00
H31A—C3A—H32A	108.00	C5—C6—H6	120.00
C41A—C4A—H4A	109.00		
C6A—N1A—C2A—C3A	56.83 (19)	C6—C1—C2—C3	0.2 (3)
C2A—N1A—C6A—C5A	-56.01 (18)	C11—C1—C2—O2	-1.8 (3)
O51—N5—C5—C4	4.5 (2)	C11—C1—C2—C3	177.28 (17)
O51—N5—C5—C6	-174.74 (16)	C2—C1—C6—C5	0.2 (2)
O52—N5—C5—C4	-175.57 (16)	C11—C1—C6—C5	-176.88 (16)
O52—N5—C5—C6	5.2 (2)	C2—C1—C11—O11	178.30 (16)



## supplementary materials

---

N1A—C2A—C3A—C4A	-56.37 (19)	C2—C1—C11—O12	-3.8 (2)
C2A—C3A—C4A—C5A	55.43 (18)	C6—C1—C11—O11	-4.7 (3)
C2A—C3A—C4A—C41A	177.15 (14)	C6—C1—C11—O12	173.26 (16)
C5A—C4A—C41A—O41A	49.2 (2)	O2—C2—C3—C4	178.81 (18)
C5A—C4A—C41A—N41A	-132.02 (17)	C1—C2—C3—C4	-0.3 (3)
C3A—C4A—C41A—O41A	-72.2 (2)	C2—C3—C4—C5	0.0 (3)
C3A—C4A—C5A—C6A	-55.62 (18)	C3—C4—C5—N5	-178.83 (17)
C41A—C4A—C5A—C6A	-177.72 (14)	C3—C4—C5—C6	0.4 (3)
C3A—C4A—C41A—N41A	106.61 (18)	N5—C5—C6—C1	178.76 (15)
C4A—C5A—C6A—N1A	55.72 (18)	C4—C5—C6—C1	-0.5 (3)
C6—C1—C2—O2	-178.84 (16)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

<i>D</i> —H $\cdots$ <i>A</i>	<i>D</i> —H	H $\cdots$ <i>A</i>	<i>D</i> $\cdots$ <i>A</i>	<i>D</i> —H $\cdots$ <i>A</i>
N1A—H11A $\cdots$ O12 <sup>i</sup>	1.00 (2)	1.71 (2)	2.688 (2)	164.2 (18)
N1A—H12A $\cdots$ O11	0.95 (2)	1.80 (2)	2.747 (2)	173.9 (17)
N41A—H41A $\cdots$ O52 <sup>ii</sup>	0.83 (2)	2.39 (2)	3.216 (2)	170.8 (19)
N41A—H42A $\cdots$ O41A <sup>iii</sup>	0.99 (2)	1.91 (2)	2.873 (2)	164.8 (18)
O2—H2 $\cdots$ O12	0.96 (2)	1.58 (2)	2.4897 (18)	156 (2)
C2A—H22A $\cdots$ O2 <sup>iv</sup>	0.97	2.57	3.450 (2)	150
C6A—H61A $\cdots$ O11 <sup>v</sup>	0.97	2.60	3.263 (2)	126
C6A—H62A $\cdots$ O41A <sup>i</sup>	0.97	2.58	3.417 (2)	145

Symmetry codes: (i)  $x, y+1, z$ ; (ii)  $x+1/2, -y+3/2, z+1/2$ ; (iii)  $-x+1, -y, -z+2$ ; (iv)  $-x+1, -y, -z+1$ ; (v)  $-x+1/2, y+1/2, -z+3/2$ .

Fig. 1




Fig. 2

