

DRAFT JANUARY 2002

1 A review of farm-scale nutrient budgets for organic farms as a tool for management of  
2 soil fertility.

3

4 C.A. Watson<sup>1\*</sup>, H. Bengtsson<sup>2</sup>, M. Ebbesvik<sup>3</sup>, A-K. Løes<sup>3</sup>, A. Myrbeck<sup>2</sup>, E. Salomon<sup>4</sup>,  
5 J. Schroder<sup>5</sup> & E.A. Stockdale<sup>6</sup>

6

7 1 Environment Division, SAC, Craibstone Estate, Aberdeen AB21 9YA, UK

8 2 Swedish University of Agricultural Sciences, Department of Soil Sciences,  
9 P.O. Box 7014, S-750 07 Uppsala, Sweden

10 3 Norwegian Centre for Ecological Agriculture (NORSØK) N-6630 Tingvoll,  
11 Norway

12 4 Swedish Institute of Agricultural and Environmental Engineering Box 7033,  
13 SE-750 07 Uppsala, Sweden

14 5 Plant Research International, Wageningen, The Netherlands

15 6 IACR-Rothamsted, Harpenden, Herts, AL5 2JQ, UK

16

17 \* Corresponding author. Fax: +44 1224 714591, E-mail [c.watson@ab.sac.ac.uk](mailto:c.watson@ab.sac.ac.uk)

18 (C.Watson)

19

20

1 **Abstract.** On organic farms, where the importation of materials to build/maintain soil  
2 fertility is restricted, it is important that a balance between inputs and outputs of  
3 nutrients is achieved to ensure both short-term productivity and long-term  
4 sustainability. This paper considers different approaches to nutrient budgeting on  
5 organic farms and evaluates the sources of bias in the measurements and/or estimates  
6 of the nutrient inputs and outputs. The paper collates 88 nutrient budgets compiled at  
7 the farm scale in 9 temperate countries. All the nitrogen (N) budgets showed an N  
8 surplus (average 83.2 kg N ha<sup>-1</sup> year<sup>-1</sup>). The efficiency of N use, defined as  
9 outputs/inputs, was highest (0.9) and lowest (0.2) in arable and beef systems  
10 respectively. The phosphorus (P) and potassium (K) budgets showed both surpluses  
11 and deficits (average 3.6 kg P ha<sup>-1</sup> year<sup>-1</sup>, 14.2 kg K ha<sup>-1</sup> year<sup>-1</sup>) with horticultural  
12 systems showing large surpluses resulting from purchased manure. The estimation of  
13 N fixation and quantities of nutrients in purchased manures may introduce significant  
14 errors in nutrient budgets. Overall, the data illustrate the diversity of management  
15 systems in place on organic farms, and suggest that used together with soil analysis,  
16 nutrient budgets are a useful tool for improving the long-term sustainability of organic  
17 systems.

18

19 **Keywords:** nutrient budgets, organic farms, nutrient use efficiency, nutrient surplus

20

## INTRODUCTION

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25

In organic farming, the farm is considered as an integrated whole, recognising that complex relationships exist between resource flows on the farm and the many environmental factors that influence them. Organic farming systems emphasise reliance on ecological interactions and biological processes over direct intervention. As a result, the use of imported materials to build/maintain soil fertility is restricted. Achieving a balance between inputs and outputs of nutrients within the farm system is critical to ensure both short-term productivity and long-term sustainability. Nutrient management must be understood, planned and managed over periods of longer than a single crop or growing season (Watson *et al.* 2002).

Nutrient budgets are becoming increasingly accepted as a tool to describe nutrient flows within farming systems and to assist in the planning of the complex and coincident spatial and temporal nutrient management within rotational cropping and mixed farming systems (Watson & Stockdale 1997). In this paper, therefore, we consider different approaches to nutrient budgeting and evaluate the sources of bias in the measurements and/or estimates of the inputs and outputs used to compile budgets that are particularly pertinent to organic farming systems. Depending on the farm management and the balance of inputs and outputs of nutrients, N, P and K budgets have been shown to range from deficit to surplus in organic farming systems (e.g. Fagerberg *et al.* 1996; Nolte & Werner 1994; Wieser *et al.* 1996). We have brought together 88 nutrient budgets compiled at the farm scale from research and commercial organic farms of different types in nine countries with temperate climates. Our aim is to examine relationships between nutrient budgets and estimates of nutrient use efficiency derived from them, and management practices and/or farm type.



1 1997). This approach, therefore, is inappropriate for the compilation of nutrient  
2 budgets relevant to organic farms (Watson & Atkinson 1999). However, this type  
3 of nutrient budget has been used widely in policy analysis.

4 2) *Surface budgets* consider the difference between total inputs and removal in crop  
5 and/or animal offtake. These budgets include uncontrollable inputs but do not  
6 usually provide information on the fate or origin of any budget surplus *i.e.*  
7 whether it is lost from the system or 'stored' in the soil. Soil surface budgets are  
8 used to determine crop nutrient requirements (particularly P and K) from  
9 fertilisers and manures at a field scale (MAFF 2000).

10 3) *System budgets* give detailed information on inputs, outputs, losses and internal  
11 flows, usually for a number of compartments *e.g.* soil, crop, livestock, manures.  
12 Aarts *et al.* (1992) presented changes in storage, transfers and nutrient surpluses of  
13 dairy systems in the Netherlands using this approach. Such budgets need larger  
14 data inputs than 1) and 2) above but the increasing availability of relevant  
15 computer models can reduce the need for additional measurements.

16 There is no one correct approach to the compilation of nutrient budgets, instead  
17 appropriate methodology should be chosen depending on the purpose/question which  
18 is driving the compilation of the budget and the nutrient or nutrients being considered  
19 (Oenema & Heinen 1999).

20

#### 21 *System definition*

22 The delineation of system boundaries in both space and time is a critical step in the  
23 compilation of nutrient budgets. In order to allow useful interpretation of the data, the  
24 definition of the system boundary also needs to be made explicit when the budget is  
25 presented.

1 Working within the horizontal dimension, including all the land within the farm  
2 boundary i.e. including woodland, tracks etc., provides a complete picture of the  
3 whole farm environment. More commonly only the managed land is included, so that  
4 for example, field margins are not included in estimates of field size. Another  
5 fundamental issue is the definition of boundaries in the vertical dimension; rooting  
6 depth is commonly used as the lower boundary.

7 Temporally, the question arises as to whether the budget should consider a single  
8 growing season, a calendar year (in which case where does it begin and end in relation  
9 to cropping pattern) or a complete crop rotation over several years? The decision will  
10 depend on the type of system and the purpose of the budget. For example, where P is  
11 applied once in 5 years on a rotational basis, budgets for a single year will not be  
12 either typical or useful. The use of data that describe complete rotations is critical for  
13 the compilation of nutrient budgets in organic farming systems, particularly where  
14 data is used to examine their likely environmental impact. For example, leaching  
15 losses have been shown to be large immediately following ploughing of leys but when  
16 averaged over whole farms and rotations losses are likely to be much lower (Philipps  
17 *et al.* 1998; Stopes *et al.* 2002). Occasionally longer-term records have been kept  
18 (Nolte & Werner 1994; Fagerberg *et al.* 1996), which allow the variation between  
19 years to be elucidated. This allows for climatic variation and its influence on crop  
20 establishment and yield, as illustrated for the stockless organic system at ADAS  
21 Terrington (Table 1). Budgets calculated across rotations can also reveal variation  
22 caused by farm management practices, such as batch rearing of animals, which do not  
23 match to an annual time step. For example, Kaffka & Koepf (1989) present farm-gate  
24 balances over the period 1952-81 for the biodynamic farm at Talhof as well as

1 considering rotational means. In such cases, interpretation of the nutrient budgets is  
2 assisted by the availability of long-term data sets for soil chemical properties.

### 3 *Presentation*

4 Nutrient budgets are generally presented in tabular form for an annual time step on  
5 the basis of  $\text{kg ha}^{-1} \text{ year}^{-1}$  or  $\text{kg farm}^{-1} \text{ year}^{-1}$ , and many published studies present only  
6 annual means. There is of course no inherent reason why nutrient budgets should be  
7 calculated over any particular time step or presented in any particular way. The  
8 methodology (as described earlier) can apply to any system whose temporal  
9 boundaries can be much longer or shorter than a year. The methodology may also  
10 relate to a unit of livestock or the production of a given number of calories for human  
11 consumption rather than a farmed area (e.g. Watson & Atkinson 1999; Jarvis 1999).  
12 Nutrient budgets may also be presented as flow diagrams *e.g.* putting numbers on the  
13 arrows of Figure 1 for a specific farm or rotation. The presentation of nutrient budgets  
14 is often closely related to the purpose of the study, and in some cases, e.g. in  
15 education, diagrams which simply show the major nutrient flows can be as useful as  
16 actually putting numbers on all the arrows (Watson & Stockdale 1999).

17

## 18 **QUANTIFICATION OF INPUTS AND OUTPUTS IN ORGANIC FARMING** 19 **SYSTEMS**

20

21 The major input and output flows for N, P and K in organic farming systems,  
22 where the spatial system boundary is defined as managed land on the farm considered  
23 to rooting depth are illustrated in Figure 2. This is the system for which we have  
24 compiled budgets from the literature and it can be described as a farm-scale surface  
25 budget. Oenema & Heinen (1999) have recently reviewed sources of bias in nutrient

1 budgets. We will not repeat their analysis but highlight additional concerns  
2 particularly relevant for the measurement/estimation of each of these flows in organic  
3 farming systems.

4

#### 5 *Purchased inputs and sold outputs*

6 Purchased inputs in feeds and supplementary fertilisers, *e.g.* rock phosphate, are  
7 permitted under organic standards (EC 2092/91). The nutrient imports in these  
8 materials are relatively easy to quantify from farm records of amounts purchased and  
9 manufacturers' information on product composition. Seed inputs are also relatively  
10 easy to quantify from quantity purchased and average percentage composition of  
11 seeds. At commonly used seed rates (Lampkin & Measures 1999), field beans (*Vicia*  
12 *faba*) could be expected to contribute 10 kg N ha<sup>-1</sup>, cereals 3 kg N ha<sup>-1</sup> and grass  
13 clover mix about 1 kg N ha<sup>-1</sup>. In general seed contributes relatively little to the  
14 nutrient input on a whole farm basis, except for seed potatoes, which can import  
15 substantial quantities of K.

16 Many studies have relied on published standard/average values for the N, P and K  
17 contents of inputs, crop and animal products. Analytical data of this type is readily  
18 available for conventional systems *e.g.* Agricultural Research Council 1976;  
19 Fagerberg *et al.* 1993; Holland *et al.* 1991 *etc.* However, it may not be appropriate to  
20 use these values in organic agriculture. Indeed even within conventional systems the  
21 range in nutrient contents measured for any material due to season and site differences  
22 may be large (Jarvis *et al.* 1996) and may invalidate the use of simple average values  
23 for detailed nutrient management planning. Where measured values for an individual  
24 site are used in place of literature derived standard values, nutrient budgets can  
25 change substantially. For two different sites in NE Scotland, Table 2 illustrates the



1 difference in the nutrient budget for a six-course rotation using literature and then  
2 measured values for the K content of straw, silage and grain. At one site the balance  
3 changes from a negative annual value to a positive one, potentially changing any  
4 management recommendations from this budgeting exercise.

5 Annual applications of manures or composts in organic systems are limited to the  
6 equivalent of 170 kg ha<sup>-1</sup> year<sup>-1</sup> of N over the entire holding (Directive 91/676/EC).  
7 Application rates on individual areas of land as high as 250 kg ha<sup>-1</sup> year<sup>-1</sup> are however  
8 permitted (as per the DEFRA Code of Good Agricultural Practice). Inputs from  
9 manure are difficult to quantify since accurate measurements of both quality and  
10 quantity are rarely available on commercial farms. The nutrient content varies greatly  
11 depending on the type of animal, its diet, the nature and amount of bedding material,  
12 the degree of separation of solids and liquids, dilution by rain water and storage  
13 conditions (Shepherd *et al.* 1999). Mean N contents of cattle FYM, 5 kg N t<sup>-1</sup> on a  
14 fresh weight basis, (range of 2 to 10 kg N t<sup>-1</sup>) and cattle slurry, 2.5 kg N m<sup>-3</sup>, (range of  
15 1.1 to 4.1 m<sup>-3</sup>) collected in organic farming systems have been shown to be about 15%  
16 lower than manure from conventional systems (Dewes & Hunsche, 1998; Shepherd *et al.*  
17 *et al.* 1999; Steineck *et al.* 1999). The differences are less clear for P and K, Dewes and  
18 Hunsche (1998) found that the K content of cattle manure was higher from organic  
19 farms but Steineck *et al.* (1999) found no significant differences between the P and K  
20 content of manures from the two systems. Composted municipal and green household  
21 waste is occasionally, but increasingly, used in organic farming systems. These  
22 typically contain 9 to 17 kg N t<sup>-1</sup> dry weight and can also supply significant quantities  
23 of P and K (Berner *et al.* 1995; Rodrigues *et al.* 1995). However, like FYM, composts  
24 are variable in composition, depending not only on source but also on batch.

1 In general, nutrient data for organic crops and manures produced on organic farms  
2 is becoming available for use in simple nutrient budgeting exercises for organic  
3 systems. However, where the budgets are to be used to make detailed management  
4 recommendations, data should be collected on that farm and ideally over a number of  
5 seasons so that site and seasonal variability can be taken into account especially in  
6 relation to changes in soil nutrient status.

7

### 8 *Deposition*

9 Deposition of nutrients is rarely measured, even as part of detailed nutrient  
10 budgeting studies, instead data is usually taken from national figures. Figures for N  
11 deposition are increasingly available as maps of deposition, *e.g.* Stanners & Bourdeau  
12 (1995) and National Expert Group on Transboundary Air Pollution (2001). However,  
13 substantial local variation can occur due to the impact of ammonia volatilisation from  
14 housed livestock. Inputs of P and K from deposition are generally very low, except in  
15 systems receiving the influence of seas spray, where K inputs are increased (Review  
16 Group on Acid Rain 1997). While deposition is likely to represent a larger  
17 proportional nutrient input to organic than to conventional systems, there is little that  
18 can be done within organic systems to manage or adjust this input. In contrast,  
19 conventional farming systems might adjust fertilisation strategy – quantities and/or  
20 timing.

21

### 22 *Nitrogen fixation*

23 Nitrogen fixation represents a major input of N into organic farming systems. The  
24 amount of N fixed by leguminous crops is notoriously variable, being dependent on  
25 the climate, soil pH, available N, P and K, age of legume, species, cultivar and strain

1 of symbiotic rhizobium (Cowling 1982; Ledgard & Steele 1992). White clover  
2 (*Trifolium repens*) is the most common legume in mixed organic systems in temperate  
3 regions, where it is usually grown with grass and utilised for grazing. Estimates of the  
4 amount of nitrogen fixed average 150 kg N ha<sup>-1</sup> year<sup>-1</sup> (range 80 to 250 kg N ha<sup>-1</sup>  
5 year<sup>-1</sup>) and 85 kg N ha<sup>-1</sup> year<sup>-1</sup> (range 50 to 130 kg N ha<sup>-1</sup> year<sup>-1</sup>) for 1-2 year old and  
6 older leys respectively (Kristensen *et al.* 1995). This decline in fixation is believed to  
7 be due to the build up of soil available N causing a decline in the proportion of clover  
8 (Crush 1987; Evans *et al.* 1995; Fisher 1996). Grazing has been shown to reduce  
9 fixation by 14-21% through the effect of higher soil N and greater grass competition  
10 (Eriksen *et al.* 1996). A sole crop of red clover (*Trifolium pratense*) is estimated to fix  
11 240 kg N ha<sup>-1</sup> year<sup>-1</sup> (Lampkin 1990; Schmidt *et al.* 1999). Other legumes grown in  
12 organic rotations, either as fodder or as green manures, include lucerne, vetches,  
13 lupins and trefoils. Estimates of N fixation range from 200 to 500kg ha<sup>-1</sup> year<sup>-1</sup> for  
14 lucerne (Lampkin 1990) and 150 to 200kg ha<sup>-1</sup> year<sup>-1</sup> for vetch (Nutman 1976; Sprent  
15 & Bradford 1977). However these species are often more difficult to manage and are  
16 less widely used, being confined to particular soil types or rotations.

17 Grain legumes obtain only 50% of their N from the atmosphere, compared with  
18 90% by forage legumes. Nevertheless, the annual fixation by field beans has been  
19 estimated at between 150 and 280 kg N ha<sup>-1</sup> (Nutman 1976; Kopke 1987), with peas  
20 fixing between 100 and 250 kg N ha<sup>-1</sup> (Jensen 1989; Fisher 1996). However, much of  
21 the fixed N is removed when the grain is harvested. Sylvester-Bradley & Cross (1991)  
22 have estimated that the effect of the nitrogen residue from combined peas or beans is  
23 equivalent to only 20 to 25 kg N ha<sup>-1</sup> year<sup>-1</sup> applied as fertiliser. There may even be a  
24 net removal of nitrogen by grain legumes under some conditions (Fisher 1996; Jensen  
25 1989).

1 It is unlikely that farmers and/or their advisors will make direct measurements of N  
2 fixation to check the assumptions made within budgets (unlike measurements of  
3 nutrient contents of inputs etc.). However, a number of empirical relationships have  
4 been proposed for estimating N fixation by legumes (Barry *et al.* 1993; Kirchmann *et*  
5 *al.* 1988; Kristensen *et al.* 1995; Watson & Goss 1997; Haraldsen *et al.* 2000;  
6 Korsaaeth & Eltun 2000). It is clear from the range of factors that these different  
7 authors included in their relationship (Table 3) that not all of them are suited to  
8 practical application using the type of information that is routinely available on farms.  
9 The use of grass-only reference crops or non-nodulating legumes for comparison will  
10 never be practical. Better quantification and record-keeping with regard to cutting and  
11 grazing management, *e.g.* yields of swards (both cut and grazed), and legume contents  
12 of swards, should however allow farmers and systems researchers to improve  
13 estimates of N fixation in combination with continued improvement and validation of  
14 practical models of N fixation.

15

## 16 **NUTRIENT BUDGETS FOR ORGANIC FARMING SYSTEMS**

17

### 18 *Data sources*

19 Following a literature search in refereed journals and English language conference  
20 proceedings, papers that detailed the compilation of nutrient budgets on biodynamic  
21 and organic farms were collated. Nitrogen, P and K budgets were included in this  
22 study where inputs and outputs of N, P or K were detailed separately on an annual  
23 basis. Some additional budgets published in theses or unpublished reports have also  
24 been included. The literature sources are summarised in Table 4. Most farm types are  
25 represented but dairy farms dominate those studied, particularly due to the Swedish

1 survey of 37 organic dairy farms. In total 88 farms were included and N budgets were  
2 the most commonly reported (88 farms) followed by P (71 farms) and K (70 farms).  
3 There are few published data on the use of other nutrients on organic farms although  
4 Mg, S and Zn budgets are reported by Nolte & Werner (1994), Nguyen *et al.* (1995)  
5 and Öborn *et al.* (2001) respectively.

6

### 7 *Data manipulation and analysis*

8 Nutrient budgets were compiled by considering all the inputs and outputs of  
9 nutrients as described in the papers (Table 4) to compile a surface budget at farm-  
10 scale (Figure 2). Inputs have been separated into purchased inputs excluding manure,  
11 purchased manure, fixation (N only) and deposition (N only) to allow the dependence  
12 of the farms on different input sources to be derived. Where no values were given for  
13 N in deposition, these have been obtained from national information (*e.g.* Stanners &  
14 Bourdeau 1995). In the published nutrient budgets surveyed here, only two papers  
15 made direct measurements of nitrogen fixation; Patriquin *et al.* (1981) used the  
16 acetylene reduction technique and Granstedt (1992) the difference method. Four of  
17 the studies did not include any estimate of N fixation (Kaffka & Koepf 1989; Fowler  
18 *et al.* 1993; Watson *et al.* 1994; Wieser *et al.* 1996), one was based on an estimated  
19 annual value (Nguyen *et al.* 1995) and the remainder were based on empirical  
20 relationships. Where no values were given for symbiotic N fixation, but information  
21 was provided on the areas growing leguminous and non-leguminous crops, an annual  
22 fixation value was derived from literature estimates (Whitehead 1995).

23 The resulting nutrient surplus or deficit ( $\Delta$  nutrient) for each farm is the difference  
24 between nutrients sold in plant and animal produce and nutrient inputs in feed, seed,  
25 supplementary nutrients, fixation and deposition (N only). The value of  $\Delta$  nutrient

1 represents the amalgamation of any nutrient losses from the system and any change in  
2 the storage of nutrients within the system. Some of the budgets included had also  
3 made measurements/estimates to allocate the nutrient surplus between losses  
4 (volatilisation, leaching, denitrification) and 'storage'. However, to give the maximum  
5 data set for comparisons between farms, only surface budgets at the farm-scale are  
6 considered in this paper. Nutrient use efficiency was also calculated; it was defined as  
7 nutrients exported in sales divided by net nutrient imports.

8 Statistical analysis of the data was based on examination of correlations, scatter  
9 plots and multiple linear regression.

10

### 11 *Results*

12 All of the N budgets showed an N surplus. Averaging over all farm types the  
13 surplus was 86.2 kg N ha<sup>-1</sup> year<sup>-1</sup> (Table 5). However, the efficiency of N use was  
14 relatively low (Table 5; average 0.3), except in the arable systems studied (where it  
15 was 0.8 and 1.0). The high efficiency of N use by the arable farms is interesting  
16 although it is difficult to draw conclusions from such a small data set. The data  
17 presented in Table 6 is from a subset of those dairy farms in Table 5 where a more  
18 detailed dataset was readily available. Across all the dairy farms studied in detail, N  
19 inputs averaged 118 kg N ha<sup>-1</sup> year<sup>-1</sup> (SE 7.5; Table 6). On average, 62% of the N  
20 inputs were derived from N fixation (range 19-87%, SE 2.8) and 25% on average in  
21 purchased feed and bedding (range 0-65%, SE 2.7). Only 4 of the 47 farms studied  
22 imported any manure, these also had some crop production on the farm (dairy farms  
23 14, 20, 33 and 47; Table 6). N outputs in products were also variable between farms  
24 (average 26 kg N ha<sup>-1</sup> year<sup>-1</sup>, SE 1.8; Table 6). However, across all dairy farms there  
25 was no significant increase in N in the products sold with increasing total N input

1 (Table 6) *i.e.* there was neither an increase in milk yield, crop outputs nor the N  
2 concentration in these products with increasing N input. Consequently there was a  
3 highly significant linear correlation between total N input and the N surplus  
4 ( $r^2=0.9455$ ,  $n=47$ ). Jarvis (1999) also found a highly significant correlation between N  
5 applied in fertiliser (the dominant N input in conventional farms) and the N surplus in  
6 conventional dairy farms. From farm-scale surface budgets the calculated N surplus is  
7 an indicator of the potential losses from the system. On dairy farms in the UK, it has  
8 been estimated that 75% of the surplus is lost, split roughly evenly between the  
9 processes of leaching, denitrification and volatilisation (Jarvis *et al.* 1996). In other  
10 studies larger proportions of the surplus are assumed to be lost; Aarts *et al.* (1992) in  
11 the Netherlands suggested that most if not all ( $> 94\%$ ) of the N surplus would be lost  
12 from the system.

13 The P and K budgets calculated show both surpluses and deficits (Table 5, average  
14  $3.6 \text{ kg P ha}^{-1} \text{ year}^{-1}$ ,  $14.2 \text{ kg K ha}^{-1} \text{ year}^{-1}$ ). The horticultural systems studied all  
15 imported significant quantities of manure to the system and showed the highest  
16 average P and K surplus. However these systems also showed the greatest range in the  
17 nutrient budgets due to differences in crop rotation, management and yields achieved  
18 (Table 5). Very high efficiency values were obtained for P and K in systems operating  
19 with very low to no inputs and showing nutrient budgets in deficit (Table 5). Inputs of  
20 P and K from accumulated reserves prior to conversion to organic farming and  
21 weathering of soil parent materials are excluded from the budgets as compiled, which  
22 may represent significant inputs to the system (*e.g.* Goulding & Loveland 1987).  
23 However, in many soils such high efficiencies coupled to negative nutrient budgets  
24 indicate that the system is not sustainable in the long-term. Greater attention should be

1 paid to the long-term capacity of the soil to supply nutrients in the design of  
2 appropriate site-specific rotations in organic farming systems.

3 Across all the dairy farms studied, P and K inputs averaged 8 kg P ha<sup>-1</sup> year<sup>-1</sup> and  
4 26 kg K ha<sup>-1</sup> year<sup>-1</sup> (SE 1.1 and 1.8 respectively; Table 6). Only 4 farms purchased  
5 manure, which made up 52 and 55 % of the P and K inputs respectively on those  
6 farms. Other purchased inputs *e.g.* in animal feed, bedding material and  
7 supplementary fertilisers made up 87 and 94% on average of the P and K inputs  
8 respectively. The level of P and K inputs were highly significantly correlated ( $r^2 =$   
9 0.6807, n=47). P and K outputs averaged 5 kg P ha<sup>-1</sup> year<sup>-1</sup> and 7 kg K ha<sup>-1</sup> year<sup>-1</sup> (SE  
10 0.4 and 0.7 respectively; Table 6). There was a significant linear relationship between  
11 P output in products and P inputs for the dairy farms ( $r^2 = 0.4858$ ; n=47) where the  
12 average efficiency calculated from the gradient of the relationship was 0.23. However,  
13 there was no significant relationship for K. Three farms (17,19,47) whose output had  
14 a significant component of crop products could be identified from the K:P ratio of the  
15 outputs (3.5 on average) whereas for all other farms the ratio was c 1.5 and within the  
16 ratio seen for milk (Holland *et al.* 1991). For both P and K there was a highly  
17 significant linear correlation between total input and the surplus ( $r^2=0.9127$  and  
18 0.9205 for P and K respectively, n=47).

19

20

## CONCLUSIONS

21 The nutrient budgets shown (Tables 5 and 6) demonstrate the considerable range in  
22 nutrient budgets not only between farm types but also within any farm type. Future  
23 research needs to address the scope for increasing nutrient use efficiency through  
24 management practices. The data highlight the importance of balancing P and K  
25 offtake in organic produce with P and K inputs from organically acceptable sources.



1 This is particularly important for the long-term maintenance of soil fertility and  
2 yields. Farmgate budgets are unable to reveal whether surplus nutrients are  
3 accumulated in soil organic matter or lost to the environment. However, the large  
4 surpluses of N on some farms suggest that the effects of management practice on the  
5 environmental impact of organic farming warrants further investigation. The data  
6 presented here also suggest some cause for concern in relation to the sustainability of  
7 organic dairy systems because of their dependence on imported feedstuffs and  
8 bedding for P and K, and for N on the very variable fixation by legumes or imports of  
9 manure or compost. This is in agreement with the findings of Goulding *et al.* (2000).  
10 Longer-term studies, particularly those including the monitoring of soil nutrient pools,  
11 are critical if we are to increase our understanding of the sustainability of nutrient  
12 management in organic farming systems.

13

14

#### **ACKNOWLEDGEMENTS**

15 The authors wish to thank Graham Horgan (BioSS) for statistical advice and Sue  
16 Fowler (University of Wales, Aberystwyth) for advice on typing of farms. SAC  
17 receives financial support from SEERAD and IACR-Rothamsted receives support  
18 from the BBSRC. DEFRA also funded some of the projects contributing to this paper.  
19 We also wish to acknowledge support from the Swedish Research Programme Food  
20 21- Sustainable Food Production.

21

22

#### **REFERENCES**

23 Aarts HFM Biewinga EE & Van Keulen H 1992. Dairy farm management based on  
24 efficient nutrient management. Netherlands Journal of Agricultural Science 40,  
25 285-299.

- 1 Agricultural Research Council 1976. The nutrient requirements of farm livestock.  
2 No4. Composition of British Feedingstuffs. Agricultural Research Council  
3 London.
- 4 Bachinger J & Stein-Bachinger K 2000. Organic farming on large farms with special  
5 reference to Eastern Germany. In: Soil quality, sustainable agriculture and  
6 environmental security in Central and Eastern Europe, eds MJ Wilson & B  
7 Maliszewska-Kordybach, Kluwer Academic Publishers Dordrecht pp 125-137.
- 8 Barry DAJ Goorahoo D & Goss MJ 1993. Estimation of nitrate concentrations in  
9 groundwater using a whole farm nitrogen budget. *Journal of Environmental*  
10 *Quality* 22, 767-775.
- 11 Berner A Scherrer D & Niggli U 1995. Effect of different organic manures and garden  
12 waste compost on the nitrate dynamics in soil N, N uptake and yield of winter  
13 wheat. *Biological Agriculture and Horticulture*, 11, 289-300.
- 14 Berry PM Stockdale EA Sylvester-Bradley R Philipps L Smith KA Lord EI Watson  
15 CA & Fortune S 2002. N, P and K budgets for crop rotations on nine organic  
16 farms in the UK. *Soil Use and Management* (this volume)
- 17 Björklund J & Salomon E 1995. Plant nutrient flows in organic agriculture - results  
18 from the Tingvall farm (In Swedish). Technical and practical information sheet  
19 soil/plants no 6, Swedish University of Agricultural Sciences, Uppsala.
- 20 Committee on Long Range Soil and Water Conservation (CLSWC) 1993. Nitrogen  
21 and phosphorus mass balances: methods and interpretation. In: *Soil and water*  
22 *quality: An Agenda for Agriculture*, National Research Council National  
23 Academy Press Washington DC pp 431-477.

- 1 Cowling DW 1982. Biological nitrogen fixation and grassland production in the  
2 United Kingdom. Philosophical Transactions of the Royal Society B. 296,397-  
3 404.
- 4 Crush JR 1987. Nitrogen Fixation. In: White clover, eds MJ Baker & WM Williams,  
5 CAB International Wallingford pp 185-201.
- 6 Cuttle S & Bowling P 1997. The nutrient status on farms converting to organic milk  
7 production. New Farmer & Grower Autumn 1997, 26-27.
- 8 Dewes T & Hunsche E 1998. Composition and microbial degradability in the soil of  
9 farmyard manure from ecologically-managed farms. Biological Agriculture and  
10 Horticulture 16, 251-258.
- 11 Ebbesvik M 1998. Økologisk eng. Viktige faktorer for avlingsnivå. Norwegian Centre  
12 for Ecological Agriculture Report, 44 pages.
- 13 Eriksen J Askegaard M & Vinther FP 1996. Nitrogen cycling in an organic dairy crop  
14 rotation. Effects of organic manure type and livestock density. In: Symbiotic  
15 nitrogen fixation in crop rotations with manure fertilisation, ed J Raupp,  
16 Proceedings 3<sup>rd</sup> meeting AIR3-CT94-1940, pp 74-83.
- 17 Evans J Chalk PM & O'Connor GE 1995. Potential for increasing N<sub>2</sub> fixation on field  
18 pea through soil management and genotype. Biological Agriculture and  
19 Horticulture 12, 97-112.
- 20 Fagerberg B Salomon E & Steineck S 1993. The computer program NPK-FLO. Users  
21 manual for calculation of plant nutrient balances in the farm and in the soil.  
22 Swedish University of Agricultural Sciences, Department of Crop Production  
23 Sciences, Internal publications 9, Uppsala.

- 1 Fagerberg B Salomon E & Jonsson S 1996. Comparisons between conventional and  
2 ecological farming systems at Öjebyn. Swedish Journal of Agricultural  
3 Research 26, 169-180.
- 4 Fisher NM 1996. The potential of grain and forage legumes in mixed farming  
5 systems. In: Legumes in sustainable farming systems, ed D Younie, BGS  
6 Occasional Symposium No. 30, British Grassland Society Reading pp 290-299.
- 7 Fowler SM Watson CA & Wilman D 1993. N, P and K on organic farms – herbage  
8 and cereal production, purchases and sales. Journal of Agricultural Science,  
9 Cambridge 120, 353-360.
- 10 Goodlass G Halberg N & Vershuur G 2002. Input output accounting systems in the  
11 European Community- an appraisal of their usefulness in raising awareness of  
12 environmental problems. European Journal of Agronomy, in press.
- 13 Goss MJ & Goorahoo D 1995. Nitrate contamination of groundwater: Measurement  
14 and prediction. Fertiliser Research 42, 331-338.
- 15 Goulding KWT & Loveland P 1987. The classification and mapping of potassium  
16 reserves in soils of England and Wales. Journal of Soil Science 37, 555-565.
- 17 Goulding KWT Stockdale EA Fortune S & Watson CA 2000. Nutrient cycling on  
18 organic farms. Journal of the Royal Agricultural Society of England 161, 65-75.
- 19 Granstedt A 1992. Case studies on the flow and supply of nitrogen in alternative  
20 farming in Sweden.1. Skilleby farm 1981-1987. Biological Agriculture and  
21 Horticulture 9, 15-63.
- 22 Hansen S 1995. Effects of soil compaction and manure level on utilization of nitrogen  
23 in cattle slurry. In: Crop development for the cool and wet regions of Europe -  
24 Nitrogen supply and nitrogen fixation of crops for cool and wet climates. eds R

- 1           Samuelson B Solsheim K Pithan E Watten-Melwaer, Proceedings of the COST  
2           814 Workshop, Tromso, Norway, pp 135-142.
- 3   Haraldsen TK Asdal A Grasdalen C Nesheim L & Ugland TN 2000. Nutrient  
4           balances and yields during conversion from conventional to organic cropping  
5           systems on silt loam and clay soils in Norway. *Biological Agriculture &*  
6           *Horticulture* 17, 229-246.
- 7   Holland B Welch AA Unwin ID Buss DH Paul AA & Southgate DAT 1991.  
8           McCance & Widdowson's *The composition of food*. 5<sup>th</sup> Edition. The Royal  
9           Society of Chemistry Cambridge.
- 10   Jarvis SC 1999. Accounting for nutrients in grassland: challenges and needs. In:  
11           Accounting for nutrients: A challenge for grassland farmers in the 21st Century.  
12           ed A.J. Corrall, BGS Occasional Symposium 33, British Grassland Society  
13           Reading pp 3-12.
- 14   Jarvis SC Wilkins RJ & Pain BF 1996. Opportunities for reducing the environmental  
15           impact of dairy farming management: a systems approach. *Grass and Forage*  
16           *Science* 52, 21-31.
- 17   Jensen ES 1989. The role of pea cultivation in the nitrogen economy of soils and  
18           succeeding crops. In: *Legumes in farming systems*, eds P Plancquaert & R  
19           Haggar, Kluwer Academic Publishers Dordrecht pp 3-15.
- 20   Kaffka S & Koepf H 1989. A case study of the nutrient regime in sustainable farming.  
21           *Biological Agriculture and Horticulture* 6, 89-106.
- 22   Kirchmann H Torssell B & Roslon E 1988. A simple model for nitrogen balance  
23           calculations of temporary grassland-ruminant systems. *Swedish Journal of*  
24           *Agricultural Research* 18, 3-8.

- 1 Kopke U 1987. Symbiotische Stickstoff Fixierung und Vorfuchtkirkung von  
2 Ackerbohnen (*Vicia faba*). Habilitation Thesis University of Gottingen.
- 3 Korsæth A & Eltun R 2000. Nitrogen mass balances in conventional, integrated and  
4 ecological cropping systems and the relationship between balance calculations  
5 and nitrogen runoff in an 8-year field experiment in Norway. *Agriculture,*  
6 *Ecosystems and Environment* 79, 199-214.
- 7 Kristensen ES Høgh-Jensen H & Kristensen IS 1995. A simple-model for estimation  
8 of atmospherically-derived nitrogen in grass-clover systems. *Biological*  
9 *Agriculture and Horticulture* 12, 263-276.
- 10 Lampkin N 1990. *Organic Farming*. Farming Press Ipswich, UK.
- 11 Lampkin N & Measures M (eds) 1999. *Organic farm management handbook* 3<sup>rd</sup>  
12 Edition, University of Wales Aberystwyth.
- 13 Ledgard SF & Steele KW 1992. Biological nitrogen fixation in mixed legume/grass  
14 pastures. *Plant and Soil* 141, 137-153.
- 15 Løes AK & Øgaard AF 1997. Changes in the nutrient content of agricultural soil on  
16 conversion to organic farming, in relation to farm level nutrient balances and  
17 soil contents of clay and organic matter. *Acta Agriculturae. Scandinavica,*  
18 Section B, Soil and Plant Science. 47, 201-214.
- 19 MAFF 2000. *Fertiliser recommendations for agricultural and horticultural crops*  
20 (RB209). The Stationery Office Norwich.
- 21 Meisinger JJ & Randall GW 1991. Estimating Nitrogen budgets for soil-crop systems.  
22 In: *Managing nitrogen for groundwater quality and farm profitability*, eds RF  
23 Follett DR Keeney & RM Cruse. SSSA Madison, Wisconsin pp 85-124.

- 1 Munters PJAL 1997. The Dutch Manure Policy: MINAS (Nutrient accounting  
2 system) Report from Dutch Dept. of Agriculture of the Ministry of Agriculture,  
3 Nature Management and Fisheries.
- 4 Myrbeck Å 1999. Växtnäringsflöden och -balanser på gårdar med olika  
5 driftsinriktning - En studie av 1300 svenska gårdar. Meddelande från  
6 Jordbearbetningsavdelningen nr 30. Department of Soil Sciences, Swedish  
7 University of Agricultural Sciences Uppsala.
- 8 National Expert Group on Transboundary Air Pollution (NEGTAP) 2001.  
9 Transboundary Air Pollution. Acidification, Eutrophication and Ground Level  
10 Ozone in the UK. 1st Report. March 2001.
- 11 Nauta WJ van der Burgt GJ & Baars T 1999. Partner farms: A participatory approach  
12 to collaboration between specialised organic farms. In: Designing and testing  
13 crop rotations for organic farming, eds Olesen JE Eltun R Gooding MJ Jensen  
14 ES & Kopke U, DARCOF Tjele pp 149-158.
- 15 Nguyen ML Haynes RJ & Goh KM 1995. Nutrient budgets and status in three pairs of  
16 conventional and alternative mixed cropping farms in Canterbury, New  
17 Zealand. *Agriculture Ecosystems and Environment* 52, 149-162.
- 18 Nolte C & Werner W 1994. Investigations on the nutrient cycle and its components of  
19 a biodynamically-managed farm. *Biological Agriculture and Horticulture* 10,  
20 235-254.
- 21 Nutman PS 1976. I.B.P. field experiments on nitrogen fixation by nodulated legumes.  
22 In: *Symbiotic nitrogen fixation in plants*, ed PS Nutman, Cambridge University  
23 Press Cambridge pp 211-237.
- 24 Öborn I Gustafson G Salomon E Lundgren A Bengtsson H Jonsson S Holmqvist J  
25 Nilsson I & Andersson A 2001. Farm gate and farm balances of P, K and Zn in

- 1 organic and conventional dairy farming at the Öjebyn Farm in Northern  
2 Sweden. In: Element balances as a sustainability tool. JTI-Rapport Lantbruk &  
3 Industri 281, pp 25-26.
- 4 Oenema O & Heinen M 1999. Uncertainties in nutrient budgets due to biases and  
5 errors. In: Nutrient disequilibria in agro-ecosystems, concepts and case studies,  
6 eds EMA Smaling O Oenema L Fresco, CAB International Wallingford pp 72-  
7 95.
- 8 Patriquin DG Burton D & Hill N 1981. Strategies for achieving self sufficiency in  
9 nitrogen on a mixed farm in Eastern Canada based on use of the faba bean. In:  
10 Genetic engineering of symbiotic nitrogen fixation and conservation of fixed  
11 nitrogen, eds JM Lyons *et al.*, Plenum Publishing Corporation, New York pp  
12 651- 671.
- 13 Philipps L Stockdale EA & Watson CA 1998. Nitrogen leaching losses from mixed  
14 organic farming systems in the UK. In: Mixed farming systems in Europe, eds  
15 H Van Keulen *et al.*, APMinderhoudhoeve-reeks nr.2, Lanbouwniversiteit  
16 Wageningen pp 165-170.
- 17 Review Group on Acid Rain 1997. Acid Deposition in the United Kingdom 1992-  
18 1994. Fourth Report June 1997. AEA Technology PLC Abingdon, UK.
- 19 Rodrigues MS Lopez-Real JM & Lee HC 1995. Composted organic waste as a soil  
20 amendment in a pot experiment. In: Soil Management in Sustainable  
21 Agriculture, eds HF Cook & HC Lee, Wye College Press Ashford pp 461-465.
- 22 Schmidt H Philipps L Welsh JP & von Fragstein P 1999. Legume breaks in stockless  
23 organic farming rotations: nitrogen accumulation and influence on the following  
24 crops. Biological Agriculture and Horticulture 17, 159-170.



- 1 Shepherd MA Bhogal A Lennartsson M Rayns F Jackson L Philipps L & Pain B  
2 1999. The environmental impact of manure use in organic farming. MAFF  
3 commissioned review.
- 4 Sprent JI & Bradford AM 1977. Nitrogen fixation in field beans (*Vicia faba*) as  
5 affected by population density, shading and its relationship with soil moisture.  
6 *Journal of Agricultural Science, Cambridge* 88, 303-310.
- 7 Stanners D & Bourdeau P (eds) 1995. Europe's environment: The Dobriš assessment.  
8 European Environment Agency. Copenhagen, Denmark.
- 9 Steineck S Gustafson G Andersson A Tersmeden M & Bergström J 1999. Plant  
10 nutrients and trace elements in livestock wastes in Sweden. Rapport 5111,  
11 Swedish Environmental Protection Agency, Stockholm.
- 12 Stopes C Lord EI Philipps L & Woodward L 2002. Nitrate leaching from organic  
13 farms and conventional farms following best practice. *Soil Use & Management*  
14 18,
- 15 Sylvester-Bradley R & Cross RB 1991. Nitrogen residues from peas and beans and  
16 the response of the following cereal to applied nitrogen. *Aspects of Applied*  
17 *Biology* 27, 293-298.
- 18 Van Noordwijk M 1999. Nutrient cycling in ecosystems versus nutrient budgets of  
19 agricultural systems In: *Nutrient disequilibria in agro-ecosystems, concepts and*  
20 *case studies*, eds EMA Smaling O Oenema L Fresco, CAB International  
21 Wallingford pp 1-26.
- 22 Vereijken P. 1986. Maintenance of soil fertility on the biodynamic farm at Nagele. In:  
23 *The importance of biological agriculture in a world of diminishing resources*,  
24 eds H Vogtmann E Boehncke & I Fricke, Vergleichsgruppe Witzehausen pp 23-  
25 30.

- 1 Watson CA & Atkinson D 1999. Using nitrogen budgets to indicate nitrogen use  
2 efficiency and losses from whole farm systems: a comparison of three  
3 methodological approaches. *Nutrient Cycling in Agroecosystems* 53, 259-267.
- 4 Watson CA Atkinson D Gosling P Jackson L Rayns FW 2002. Managing soil fertility  
5 in organic farming systems. *Soil Use & Management* 18,.
- 6 Watson CA Fowler SM & Wilman D 1994. Purchases and sales of N, P and K, soil  
7 inorganic N and nitrate leaching on an organic horticultural holding. *Biological  
8 Agriculture and Horticulture* 10, 189-195.
- 9 Watson CA & Goss MJ 1997. The estimation of nitrogen fixation by grass/white  
10 clover mixtures in cut and grazed swards. *Soil Use & Management* 13, 165-167.
- 11 Watson CA & Stockdale EA 1997. Using nutrient budgets to evaluate the  
12 sustainability of farming systems. *Newsletter of the European Network on  
13 Organic Farming*, Number 5, October 1997, 16-19.
- 14 Watson CA & Stockdale EA 1999. Whole farm nutrient management: What do  
15 budgets tell us? In: *Accounting for nutrients: A challenge for grassland farmers  
16 in the 21st Century*. ed A.J. Corral, BGS Occasional Symposium 33, British  
17 Grassland Society Reading pp 35-40.
- 18 Watson CA Younie D Stockdale EA & Cormack WF 2000. Yield and nutrient  
19 balances of stocked and stockless organic rotations. *Aspects of Applied Biology*  
20 62, 261-268.
- 21 Whitehead DC 1995. *Grassland nitrogen*. CAB International Wallingford UK.
- 22 Wieser I Hess J & Lindenthal T 1996. Nutrient balances on organically managed  
23 farms in Upper Austria. *Bodenkultur* 47, 81-88.

- 1 Wood M 1996. Nitrogen fixation: How much and at what cost? In: Legumes in
- 2 sustainable farming systems, ed D Younie, BGS Occasional Symposium No.
- 3 30, British Grassland Society Reading pp 26-35

1 TABLE HEADINGS

2 Table 1 Annual surface N budget for the period 1995-1999 for the stockless organic  
3 system at ADAS Terrington, UK. (Rotation comprises red clover, potatoes,  
4 winter wheat, winter beans, spring wheat).

5 Table 2 Surface K budgets calculated for ley/arable rotations at SAC Farms Tulloch  
6 and Woodside. The estimated budget uses standard literature values for the K  
7 content of crop products. The corrected budget uses analytical values for  
8 crop products.

9 Table 3 Parameters used in a number of empirical relationships to predict N fixation

10 Table 4 Data sources for the compilation of nutrient budgets at farm-scale for organic  
11 farms.

12 Table 5 Summary of farm-scale nutrient budgets by farm type

13 Table 6 Simplified nutrient budgets for 47 farms where dairy production is  
14 considered to be the major enterprise, but which also may have some  
15 cropping on farm (mixed) listed in order of increasing total N input to the  
16 farm system. n/a = information not available.

1 FIGURE HEADINGS

2

3 Figure 1. Simple diagrammatic representation of nutrient flows that may occur on a  
4 farm. Where the boundary of the system is drawn will determine which  
5 flows represent inputs, outputs and internal flows for that system.

6 ----- might represent the farm boundary, including cropped and uncropped land

7 \_\_\_\_\_ might represent the crop rotation boundary, including soil to rooting depth

8

9 Figure 2. Surface budget at the farm-scale used for the N, P and K budgets  
10 presented. The farm system boundaries are the cropped land to rooting depth.  
11  $\Delta$  nutrient (*i.e.* the budget surplus or deficit) calculated in this way represents  
12 the amalgamation of any nutrient losses from the system and any change in  
13 the storage of nutrients within the system.

14



Table 1

	1995	1996	1997	1998	1999
Deposition	30	30	30	30	30
Seed	4	4	4	4	4
Fixation (winter beans, red clover)	24	20	45	37	35
<b>Inputs - total</b>	<b>58</b>	<b>54</b>	<b>79</b>	<b>71</b>	<b>69</b>
<b>Crop output</b>	<b>92</b>	<b>110</b>	<b>71</b>	<b>81</b>	<b>89</b>
Balance	-34	-56	8	-10	-20

Table 2

	Woodside		Tulloch	
	Estimated <sup>a</sup>	Corrected	Estimated <sup>a</sup>	Corrected
<b>INPUTS</b>				
Deposition	2.1	2.1	2.1	2.1
Seed	3.2	3.2	0.7	0.7
Manure	50.5	50.5	53.1	53.1
Grazing	40.5	45.8	36.0	36.0
<b>OUTPUTS</b>				
Products	36.0	32.0	25.5	23.2
Straw	16.7	10.9	8.8	3.7
Silage	38.7	42.6	70.2	64.3
Liveweight gain	12.0	12.0	10.6	10.6
<b>BALANCE</b>	<b>-7.0</b>	<b>4.1</b>	<b>-23.2</b>	<b>-10.0</b>

<sup>a</sup>From Watson *et al.* (2000)



Table 3

Reference	Legumes studied	Variables									
		Yield of legume + grass	Yield of grass-only reference crop	% legume in mixture	Years after establishment	N content of legumes	N content of legume + grass	N content of grass-only reference crop	% legume N derived from fixation	Correction for stubble/root N	Sward management
Barry <i>et al.</i> (1993)	Alfalfa	√	√								√
Haraldsen <i>et al.</i> (2000)	Soybean			√		√			√		
Kirchmann <i>et al.</i> (1988)	Grass-clover	√	√			√	√	√	√	√	
Korsaeth & Eltun (2000) <sup>a</sup>	Grass-white clover	√		√						√	
	Grass-red clover										
	Grey peas										
	Common vetch										
Kristensen <i>et al.</i> (1995)	Grass-clover (red and white mix)			√	√					√	
Watson & Goss (1997)	Grass-white clover	√	√	√						√	√

<sup>a</sup> Modified version of Hansen (1995)

Table 4

Country	Farm types	Years of data compiled	N	P	K	Reference
Austria	9 dairy	1	√ <sup>b</sup>	√	√	Wieser <i>et al.</i> (1996)
Canada	1 arable, 1 dairy, 1 pig	1	√			Goss & Goorahoo (1995)
Canada	1 poultry	1	√ <sup>b</sup>			Patriquin <i>et al.</i> (1981)
Germany	2 mixed	3	√			Bachinger & Stein-Bachinger (2000)
Germany	1 mixed <sup>a</sup>	1	√ <sup>b,c</sup>	√	√	Kaffka & Koepf (1989)
Germany	1 mixed <sup>a</sup>	3	√	√	√	Nolte & Werner (1994)
Netherlands	1 dairy <sup>a</sup>	4	√	√	√	Vereijken (1986)
Netherlands	1 dairy, 1 arable	?	√	√	√	Nauta <i>et al.</i> (1999)
New Zealand	3 mixed	1	√ <sup>b</sup>	√	√	Nguyen <i>et al.</i> (1995)
Norway	9 dairy 2 dairy <sup>a</sup> 1 sheep 1 mixed <sup>a</sup>	1-6	√	√	√	Ebbesvik (1998) Løes & Øgaard (1997)
Sweden	1 dairy	5	√ <sup>b</sup>	√	√	Fagerberg <i>et al.</i> (1996)
Sweden	1 dairy	2	√	√	√	Björklund & Salomon (1995)
Sweden	37 dairy	1	√	√	√	Myrbeck (1999)
Sweden	1 dairy <sup>a</sup>	7	√			Granstedt (1992)
UK	2 dairy 3 beef	2 on two farms; 1 on 3 farms	√ <sup>b,c</sup>	√	√	Fowler <i>et al.</i> (1993)
UK	1 dairy	3	√	√	√	Cuttle & Bowling (1997)
UK	2 horticultural	2 on one holding; 1 on one holding	√ <sup>b,c</sup>	√	√	Watson <i>et al.</i> (1994)
UK	1 beef 1 dairy 1 horticulture	1	√	√	√	Goulding <i>et al.</i> (2000)
UK	1 beef	3	√	√	√	Watson & Atkinson (1999)

<sup>a</sup> Biodynamic<sup>b</sup> No deposition data <sup>c</sup> No fixation data

Table 5

Farm type	n	Surplus (Input-Output) kg ha <sup>-1</sup> year <sup>-1</sup>			Efficiency (Output/Input)		
		Mean	SE	Range	Mean	SE	Range
N							
Arable	2	25.6	24.4	1.2- 50.0	0.9	0.1	0.8-1.0
Beef	5	112.0	25.6	18.4-164.0	0.2	0.03	0.1-0.2
Dairy	67	82.1	6.7	2.1-217.0	0.3	0.02	0.0-0.9
Horticulture	3	194.2	100.7	91.0-395.6	0.3	0.1	0.1-0.4
Mixed	8	54.6	8.6	21.0- 91.6	0.4	0.05	0.2-0.5
Mean		83.2					
P							
Arable	1	-6.0			1.3		
Beef	4	-1.8	1.4	-6 - 0	2.8	1.4	1.0- 7.0
Dairy	56	3.1	0.9	-6.5 - +36.0	2.1	1.2	0.3-66
Horticulture	3	38.9	26.0	1.7 - +89.0	0.4	0.2	0.1- 0.9
Mixed	6	-2.4	1.3	-6.9 - + 4.0	13.6	9.8	0.6-70
Mean		3.6					
K							
Arable	1	57.0			0.8		
Beef	4	3.0	3.4	-4.5 - + 12.0	2.8	2.4	0.2- 10
Dairy	58	9.6	2.0	-26.5 - + 58.0	5.3	4.6	0.1-266
Horticulture	3	122.0	88.0	-23.0 - +281.0	0.7	0.4	0.1- 1.6
Mixed	3	-2.2	1.2	-4.4 - - 0.3	1.6	0.3	1.1- 2.0
Mean		14.2					



Table 6

Country	Robust type	Farm size (ha)	Stocking rate (lu ha <sup>-1</sup> ) <sup>a</sup>	Nutrient flows kg ha <sup>-1</sup>									
				N fixation	Total N input	N in product sold	N surplus	Total P input	P in product sold	P surplus	Total K input	K in product sold	K surplus
1 Sweden	Mixed	55	0.49	26	36	34	2	1	1	-5	2	2	-7
2 Sweden	Mixed	33	n/a	21	37	22	15	2	5	-2	2	5	-3
3 Sweden	Mixed	101	0.57	18	39	13	26	2	3	-1	2	4	-2
4 Sweden	Mixed	180	n/a	22	42	15	27	2	3	-1	11	4	7
5 Sweden	Mixed	72	0.72	37	49	37	12	4	7	-4	7	10	-6
6 Sweden	Mixed	160	0.55	30	52	30	22	4	6	-2	5	8	-3
7 Sweden	Mixed	106	0.47	42	56	16	40	3	3	0	2	4	-2
8 Sweden	Mixed	132	0.66	44	64	22	42	4	5	-1	5	5	0
9 Sweden	Mixed	125	0.56	17	67	26	41	18	5	13	25	7	18
10 Sweden	Mixed	130	n/a	51	69	13	56	2	3	-1	7	4	3
11 Sweden	Mixed	64	0.48	40	73	20	53	7	4	3	32	5	27
12 Sweden	Mixed	67	1.03	45	73	31	42	10	6	4	11	6	5
13 Sweden	Mixed	185	n/a	35	75	24	51	5	5	0	11	6	5
14 Sweden	Mixed	71	n/a	47	78	15	64	3	3	0	5	5	0
15 Sweden	Mixed	44	1.07	64	85	20	64	3	4	-1	6	5	1
16 Sweden	Mixed	175	0.69	52	88	17	71	8	3	5	4	5	-1
17 Sweden	Mixed	83	0.87	33	89	38	51	8	6	2	12	18	-6
18 Norway	Dairy	14	n/a	52	97	35	62	9	7	2	53	10	43
19 Netherlands	Mixed	22	n/a	75	99	42	57	0	7	-7	0	27	-27
20 Sweden	Mixed	97	0.80	45	102	12	90	1	2	-1	2	3	-1
21 Sweden	Mixed	85	0.74	78	106	17	89	5	4	1	10	4	6
22 Sweden	Mixed	112	0.61	38	107	27	80	11	9	2	10	8	2
23 Norway	Dairy	9	n/a	46	107	34	73	19	7	12	62	23	39
24 Sweden	Mixed	129	n/a	76	108	18	90	5	4	1	5	6	-1
25 UK	Mixed	63	1.55	71	122	31	91	10	6	4	22	9	13
26 Sweden	Mixed	107	0.92	86	130	37	93	6	7	-1	10	10	0
27 Sweden	Mixed	30	0.80	80	131	18	113	6	3	3	15	5	10
28 Sweden	Mixed	60	0.68	111	135	17	117	3	3	0	4	4	0
29 Sweden	Mixed	25	1.38	69	137	33	104	20	7	13	50	9	41
30 Sweden	Mixed	59	0.47	65	142	41	101	15	9	6	24	11	13
31 Sweden	Mixed	83	0.34	33.7	144	28	116	21	6	15	66	8	58
32 Sweden	Mixed	60	1.25	113	147	18	128	3	4	-1	4	5	-1
33 Sweden	Mixed	137	0.77	27	147	31	117	18	6	12	31	8	23
34 UK	Mixed	99	1.90	123	157	24	133	7	5	2	10	5	5
35 UK	Dairy	56	n/a	117	159	37	122	3	7	-4	12	9	3
36 Netherlands	Dairy	52	n/a	80	160	49	111	11	10	1	43	16	27
37 Austria	Dairy	16	n/a	150	172	14	157	6	3	3	3	3	0
38 Austria	Dairy	30	n/a	150	173	15	158	2	4	-2	3	3	0
39 Austria	Dairy	20	n/a	150	177	21	156	4	5	-1	5	4	1
40 Austria	Dairy	13	n/a	150	179	23	156	3	5	-2	5	5	0
41 Austria	Dairy	32	n/a	150	180	18	162	2	3	-1	24	4	20
42 Austria	Dairy	52	n/a	150	181	14	167	4	3	1	19	5	14
43 Austria	Dairy	32	n/a	150	181	22	159	10	5	5	7	5	2
44 Austria	Dairy	37	n/a	150	183	8	175	3	2	1	5	2	3
45 Sweden	Mixed	37	1.19	57	186	69	117	29	13	16	30	11	19
46 Austria	Dairy	15	n/a	150	186	16	170	6	4	2	16	4	12
47 UK	Mixed	233	2.20	86	247	51	196	31	10	21	66	13	53

<sup>a</sup> lu=livestock units

Figure 1.




Figure 2

