

Short report

## First report of *Metarhizium anisopliae* IP 46 pathogenicity in adult *Anopheles gambiae* s.s. and *An. arabiensis* (Diptera; Culicidae)

Ladslaus L Mnyone<sup>\*1,2,3</sup>, Tanya L Russell<sup>1,7</sup>, Issa N Lyimo<sup>1,4</sup>,  
Dickson W Lwetoijera<sup>1,6</sup>, Matthew J Kirby<sup>1,2</sup> and Christian Luz<sup>1,5</sup>

Address: <sup>1</sup>Biomedical and Environmental Group, Ifakara Health Institute, PO Box 53, Off Mlabani Passage, Ifakara, Tanzania, <sup>2</sup>Laboratory of Entomology, Wageningen University & Research Centre, PO Box 8031, 6700 EH, Wageningen, the Netherlands, <sup>3</sup>Pest Management Center, Sokoine University of Agriculture, PO Box 3110, Morogoro, Tanzania, <sup>4</sup>Faculty of Biomedical and Life Sciences, University of Glasgow, 120 University Place, G12 8TA, Glasgow, UK, <sup>5</sup>Instituto de Patologia Tropical e Saúde Pública, Universidade Federal de Goiás, CP 131, 74001-970 Goiânia, GO, Brasil, <sup>6</sup>Department of Zoology and Marine Biology, University of Dar es Salaam, PO Box 35064, Dar es Salaam, Tanzania and <sup>7</sup>Vector Group, Liverpool School of Tropical Medicine, Liverpool, L3 5QA, UK

Email: Ladslaus L Mnyone\* - [llaurent@ihi.or.tz](mailto:llaurent@ihi.or.tz); Tanya L Russell - [trussell@ihi.or.tz](mailto:trussell@ihi.or.tz); Issa N Lyimo - [ilyimo@ihi.or.tz](mailto:ilyimo@ihi.or.tz); Dickson W Lwetoijera - [dwilson@ihi.or.tz](mailto:dwilson@ihi.or.tz); Matthew J Kirby - [mkirby@ihi.or.tz](mailto:mkirby@ihi.or.tz); Christian Luz - [wolf@iptsp.ufg.br](mailto:wolf@iptsp.ufg.br)

\* Corresponding author

Published: 1 December 2009

Received: 12 November 2009

*Parasites & Vectors* 2009, **2**:59 doi:10.1186/1756-3305-2-59

Accepted: 1 December 2009

This article is available from: <http://www.parasitesandvectors.com/content/2/1/59>

© 2009 Mnyone et al; licensee BioMed Central Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/2.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

### Abstract

The entomopathogenic fungus *Metarhizium anisopliae* isolate IP 46, originating from a soil sample collected in 2001 in the Cerrado of Central Brazil, was tested for its ability to reduce the survival of adult male and female *Anopheles gambiae* s.s. and *An. arabiensis* mosquitoes. A 6-h exposure to the fungus coated on test paper at a concentration of  $3.3 \times 10^6$  conidia  $\text{cm}^{-2}$  reduced the daily survival of both mosquito species (HR = 3.14,  $p < 0.001$ ), with higher risk of dying in *An. gambiae* s.s. relative to *An. arabiensis* (HR = 1.38,  $p < 0.001$ ). Fungal sporulation was observed in >95% of mosquito cadavers in the treatment groups. The results indicate that *M. anisopliae* IP 46 has the potential to be a bio-control agent for African malaria vector species, and is a suitable candidate for further research and development.

### Findings

*Metarhizium anisopliae* IP 46 has shown ovicidal effects against the eggs of *Aedes* spp in Brazil [1-3]. However, its pathogenicity against adult malaria vectors has never been explored. As such, we examined the effect of this strain against laboratory-reared adult *Anopheles gambiae sensu stricto* and *An. arabiensis*, with the aim to include IP 46 in the spectrum of fungal candidates available for use as bio-control agents.

The fungus was imported as conidia from the Institute of Tropical Pathology and Public Health, Federal University of Goiás, Goiânia, Brazil (Tropical Pesticides Research

Institute Import Permit No. 2471). Before conducting bioassays, the IP 46 isolate was host-passaged through laboratory-reared *An. gambiae* s.s. adults in order to maintain its virulence. Conidia were harvested from cultures grown on autoclaved rice substrate (200 g per bag) in nylon bags at 25 °C and 12 h photophase, after 15 d incubation. They were then dried in silica gel at 4 °C for 4 d. Preparation of stock- and working-solution concentrations formulated in Enerpar oil (Enerpar M002®, BP South Africa Ltd) followed standard protocols [4]. Before each experiment, conidia viability (>95% germination on Sabouraud Dextrose Agar) was confirmed. 1200  $\mu\text{l}$  of the working-solution was applied evenly to 15 × 25 cm proof-

ing paper using a metal bar (0.31 mm diameter; paper and applicator bar from RK Print Coat Instruments, London), giving a uniform concentration of  $3.3 \times 10^6$  conidia  $\text{cm}^{-2}$ . The treated paper was left to dry for 12 h at  $26 \pm 1^\circ\text{C}$  and  $80 \pm 5\%$  RH, and then used to line the inside of plastic exposure tubes (8.2 cm diameter  $\times$  12.5 cm height). Untreated control replicates used paper treated with Enerpar oil only.

A total of 30-40 unfed 3-7 d old adult *An. gambiae* s.s. (colony established in 1996, Njage village, Tanzania) or *An. arabiensis* (colony established in 2007, Sagamaganga village, Tanzania) were introduced to the exposure tubes. Four separate bioassays were run (both sexes for each species) and three replicates were carried out for each bioassay. Mosquitoes were held in the tubes for 6 h, after which they were transferred to 9  $\text{cm}^3$  holding cages at  $26 \pm 1^\circ\text{C}$  and  $90 \pm 5\%$  RH, and provided with 9% glucose/water (w/v) solution. The survival and fungus infection status of mosquitoes were monitored daily for up to 28 d, following procedures described elsewhere [4]. Mosquito survival was analysed by Kaplan-Meier pair-wise comparison and Cox regression analysis, using SPSS version 16. Cox regression generated hazard ratios (HR) indicating the daily risk of dying for a mosquito in each bioassay group.

*Metarhizium anisopliae* IP 46 was capable of infecting males and females of both mosquito species: >95% of *An. gambiae* s.s. and *An. arabiensis* cadavers showed fungus sporulation after incubation for 5-6 d. The fungus significantly reduced the survival of all exposed mosquitoes compared to controls ( $p < 0.001$ , Table 1, Fig. 1); >90% of mosquitoes in the exposure groups had died by day 14 while >25% of control mosquitoes were still alive by this time. All of the control mosquitoes in all bioassays had died by day 28. For *An. gambiae* s.s. the daily risk of dying was over three-fold greater in exposed females (HR = 3.18,  $p < 0.001$ ) and males (HR = 3.81,  $p < 0.001$ ) relative to their controls. A similar trend was observed in exposed females (HR = 2.28,  $p < 0.001$ ) and males (HR = 3.31,  $p < 0.001$ ) of *An. arabiensis*. The daily risk for males was higher than for females in both species (*An. gambiae* s.s. HR = 1.11,  $p = 0.001$  and *An. arabiensis* HR = 1.13,  $p =$


0.004). Overall, daily risk of dying was higher for exposed *An. gambiae* than *An. arabiensis* (HR = 1.38,  $p < 0.001$ ). The controls for *An. gambiae* survived relatively longer (males MST = 14 d; females MST = 16 d) than those of *An. arabiensis* (male MST = 12 d; female MST = 12 d, Table 1), but this difference was accounted for by Cox regression model which compares relative risks rather than fixed survival time values.

For effective malaria control, entomopathogenic fungi do not need to kill vector mosquitoes instantly [5]. If mosquitoes are able to reproduce and pass genes to the next generation before they are killed by an insecticide the selection pressure for the development of resistance is significantly reduced [6,7]. Here we have shown that the isolate *M. anisopliae* IP 46 kills females of *An. gambiae* s.s. and *An. arabiensis* on average 8-9 d after exposure. By day 14 the majority (>90%) of exposed mosquitoes had been killed. Given that the *Plasmodium* parasite requires approximately 9 to 14 d to infect the mosquito salivary glands, the risk of malaria transmission by fungus-infected mosquitoes is minimal [8]. Similar rates of mortality have been recorded for other entomopathogenic fungi against mosquitoes [4,9-11]. Perhaps most importantly, *M. anisopliae* IP 46 was effective against both *An. arabiensis* and *An. gambiae* s.s. suggesting that it could be used to target both indoor and outdoor resting anophelines. This is the first study demonstrating the susceptibility of adult *An. arabiensis* to *Metarhizium anisopliae*.

Ultimately the success of entomopathogenic fungi against malaria-carrying mosquitoes in any situation may depend on the choice of fungal isolate. This is because of the inter-isolate variation in virulence, spore production and persistence in relation to their ability to withstand sub-optimal environmental conditions [12-16]. The long-standing barriers that have prevented the widespread uptake of biological control agents include low virulence and short-term residual activity. In order to overcome such barriers it is necessary to screen an array of fungal strains to identify those with the greatest potential for development. We found that the isolate *M. anisopliae* IP 46 is able to reduce the survival of adult anophelines within the same time frame as other strains, *M. anisopliae* ICIPE-30 and *B. bassi*

**Table 1: Pair-wise Kaplan-Meier median survival times (MST) for adult *Anopheles gambiae* s.s. and *An. arabiensis* exposed to oil-formulated *M. anisopliae* IP 46 (treatment) or oil only (control).**

Species	Sex	MST $\pm$ I S.E.		$\chi^2$ value	p value
		Control	Treatment		
<i>An. gambiae</i> s.s.	Female	16 $\pm$ 0.51	9 $\pm$ 0.23	94.58	<0.001
	Male	14 $\pm$ 0.76	8 $\pm$ 0.30	133.07	<0.001
<i>An. arabiensis</i>	Female	12 $\pm$ 0.79	8 $\pm$ 0.38	63.04	<0.001
	Male	12 $\pm$ 0.45	6 $\pm$ 0.31	113.13	<0.001



**Figure 1**  
Survival of adult female and male a) *Anopheles gambiae* s.s. and b) *An. arabiensis* mosquitoes after 6 h exposure to *Metarhizium anisopliae* IP 46 conidia.

ana IMI 391510 [4,10,17]. We anticipate that our findings will encourage research into other strains and further investigation and development of IP 46.

### Competing interests

The authors declare that they have no competing interests.

### Authors' contributions

Conceived and designed the experiments: LLM TLR CL. Performed the experiments: LLM DWL INL. Analyzed the data: LLM TLR MJK INL. Wrote the paper: LLM MJK. Reviewed the paper: CL TLR.

### Acknowledgements

We wish to acknowledge Monika Mpingwa, Kassian Mbina, Ally Daraja, Paulina Kasanga and Emmanuel Simfukwe for rearing mosquitoes and providing technical assistance.

### References

- Luz C, Tai MHH, Santos AH, Silva HHG: **Impact of moisture on survival of *Aedes aegypti* eggs and ovicidal activity of *Metarhizium anisopliae* under laboratory conditions.** *Memórias do Instituto Oswaldo Cruz* 2008, **103**:214-215.
- Albernaz DAS, Tai MHH, Luz C: **Enhanced ovicidal activity of an oil formulation of the fungus *Metarhizium anisopliae* on the mosquito *Aedes aegypti*.** *Medical and Veterinary Entomology* 2009, **23**:141-147.
- Santos AH, Tai MH, Rocha LF, Silva HH, Luz C: **Dependence of *Metarhizium anisopliae* on high humidity for ovicidal activity on *Aedes aegypti*.** *Biological Control* 2009, **50**:37-42.
- Blanford S, Chan BHK, Jenkins N, Sim D, Turner RJ, Read AF, Thomas MB: **Fungal pathogen reduces potential for malaria transmission.** *Science* 2005, **308**:1638-1641.
- Hancock PA, Thomas MB, Godfray HC: **An age-structured model to evaluate the potential of novel malaria-control interventions: a case study of fungal biopesticide sprays.** *Proceedings of the Royal Society of London* 2008:1-10.
- Thomas MB, Read AF: **Can fungal biopesticides control malaria?** *Nature Reviews Microbiology* 2007, **5**:377-383.
- Ondiaka S, Bukhari T, Farenhorst M, Takken W, Knols BG: **Effects of fungal infection on the host-seeking behaviour and fecundity of the malaria mosquito *Anopheles gambiae* Giles.** *Proceedings of Netherlands Entomological Society Meeting* 2008, **9**:121-128.
- Talman AM, Domarle O, McKenzie FE, Arley F, Robert V: **Gametocytogenesis: the puberty of *Plasmodium falciparum*.** *Malaria Journal* 2004, **3**:24-38.
- Farenhorst M, Farina D, Scholte EJ, Takken W, Hunt RH, Coetzee M, Knols BGJ: **African water storage pots for the delivery of the entomopathogenic fungus *Metarhizium anisopliae* to the malaria vectors *Anopheles gambiae* s.s. and *Anopheles funestus*.** *American Journal of Tropical Medicine and Hygiene* 2008, **78**:910-916.
- Stevenson JC: **The use of fungi against adult malaria mosquitoes, PhD Thesis.** *London School of Hygiene and Tropical Medicine, London* 2008.
- Scholte EJ, Ng'habi K, Kihonda J, Takken W, Paaijmans K, Abdulla S, Killeen GF, Knols BGJ: **An entomopathogenic fungus for control of adult African malaria mosquitoes.** *Science* 2005, **308**:1641-1642.
- Ihara F, Toyama M, Higaki M, Mishiro K, Yaginuma K: **Comparison of pathogenicities of *Beauveria bassiana* and *Metarhizium anisopliae* to chestnut pests.** *Applied Entomology and Zoology* 2009, **44**:127-132.
- De La Rosa W, Alatorre R, Barrera JF, Toriello C: **Effect of *Beauveria bassiana* and *Metarhizium anisopliae* (Deuteromycetes) upon the coffee berry borer (Coleoptera: Scolytidae) under field conditions.** *Journal of Economic Entomology* 2000, **93**:1409-1414.
- Sun J, Fuxa JR, Henderson G: **Effects of virulence, sporulation, and temperature on *Metarhizium anisopliae* and *Beauveria***

**bassiana laboratory transmission in *Coptotermes formosanus*.** *Journal of Invertebrate Pathology* 2003, **84**:38-46.

15. Bugeme DM, Knapp M, Boga HI, Wanjoya AK, Maniania NK: **Influence of temperature on virulence of fungal isolates of *Metarhizium anisopliae* and *Beauveria bassiana* to the two-spotted spider mite *Tetranychus urticae*.** *Mycopathologia* 2009, **167**:221-227.
16. Ansari MA, Vestergaard S, Tirry L, Moens M: **Selection of a highly virulent fungal isolate, *Metarhizium anisopliae* CLO 53, for controlling *Hoplia philanthus*.** *Journal of Invertebrate Pathology* 2004, **85**:89-96.
17. Scholte EJ, Njiru BN, Smallegange RC, Takken W, Knols BG: **Infection of malaria (*Anopheles gambiae* s.s.) and filariasis (*Culex quinquefasciatus*) vectors with the entomopathogenic fungus *Metarhizium anisopliae*.** *Malaria Journal* 2003, **2**:29.

Publish with **BioMed Central** and every scientist can read your work free of charge

*"BioMed Central will be the most significant development for disseminating the results of biomedical research in our lifetime."*

Sir Paul Nurse, Cancer Research UK

Your research papers will be:

- available free of charge to the entire biomedical community
- peer reviewed and published immediately upon acceptance
- cited in PubMed and archived on PubMed Central
- yours — you keep the copyright

Submit your manuscript here:  
[http://www.biomedcentral.com/info/publishing\\_adv.asp](http://www.biomedcentral.com/info/publishing_adv.asp)

