

# Hip Structure Associated With Hip Fracture in Women: Data from the Geelong Osteoporosis Study (Gos) Data Analysis-Geelong, Australia

Najmah<sup>1</sup>, L. Gurrin<sup>2</sup>, M.Henry<sup>3</sup>, J.Pasco<sup>3</sup>

<sup>1</sup> Faculty of Public Health Sriwijaya University, Kampus Unsri Indralaya, Ogan Ilir, Sumatera Selatan, Indonesia.

<sup>2</sup> School of Population Health, The University of Melbourne, Australia.

<sup>3</sup> Department of Clinical and Biomedical Sciences, The University of Melbourne, Australia.

For reprint and all correspondence: Najma, Faculty of Public Health, Sriwijaya University, Kampus Unsri Indralaya, Ogan Ilir, Sumatera Selatan, Indonesia, Hp: +6285267412242

E-mail: najem240783@yahoo.com

## ABSTRACT

<b>Backgrounds</b>	Aging leads to changes in bones to be highly fragile causing fractures. In this research, changes in the dimensions of the hip structure can be measured by using a computer program called 'Hip Structural Analysis (HSA)'. The objective of this study is to estimate the association between hip geometries in Femoral Neck (FN) and the risk of hip fracture in older women.
<b>Methods</b>	A case control study was performed to explore the objective respectively using the data of participants from population cohort and fracture cohort of the Geelong Osteoporosis Cohort Geelong, Southern Victoria, Australia. Simple and multiple logistic regressions were performed.
<b>Results and Discussion</b>	Of total of 598, comparing Fracture group (44 subjects) and non-fracture group (454 subjects) aged over 63 years, the odds of hip fracture increased by approximately 2 fold for each 1 SD increase in width (OR=1.70(1.18-2.45,p 0.005), endocortical diameter (OR=1.80 (1,23-2.62, p=0.002), and buckling ratio (OR=1.85(1.32- 2.61, p < 0.0001) and for each 1 SD decrease in BMD (OR=1.98(1.21-3.23, p.0.006) and average cortical thickness (OR=2.02(1.23-3.34), p.0.006) controlling for age, height, weight and menopausal status. Findings suggest that not only is BMD associated with hip fractures, but also other hip geometry dimensions, including WID, ENDO, AVCO and AVBR, independent of age, height, weight and physical activity.
<b>Conclusions</b>	These results provide additional insights that the geometries of FN is associated with fracture neck of femur in older women and strongly suggest its potential value, not only BMD, as clinical predictors for assessing the risk of hip fracture in older women. In addition to this, utilization of some combined parameters of bone geometries in FN might be a more effective method in screening than case findings to reduce the burden of hip fracture in the future. Further statistical methods is needed to analyze the combined hip structure to predict hip fracture.

## INTRODUCTION

### *Background and Rationale*

Hip fracture is a major public health problem and a leading cause of morbidity, hospitalisation and mortality in particular cases (Deng et al., 2002, Hannan, 2001, Wehren and Magaziner, 2003, Khasraghi, 2003). Hip fractures are also associated with certain medical complications, including electrolyte imbalance, urinary tract infection, respiratory failure and delirium in both men and women (Khasraghi, 2003). In the early twenty-first century, the estimated number of incident osteoporotic hip fractures was 1.6 million worldwide; with approximately 70 % of these occurred in women. The Disability Adjusted Life Years (DALYs) for hip fracture was 0.82 million in men and 1.53 million in women within the same year. Moreover, hip fracture contributed for 41 % of the global burden of osteoporosis in 2000 (Johnell and Kanis, 2006).

In Australia, hip fractures were predicted to increase for 36 % between 1996 and 2006 (from 15000 to 21000 cases) (Sanders et al., 1999a). This significant increase was due to the increased number of elderly aged 85 years and over as Australia faces the challenges of an aging population (Sanders et al., 1999a). Furthermore, hip fractures are projected to increase by twofold in 2026 and fourfold in 2051 (Sanders et al., 1999a) with hospitalization rate of almost 100 % (Pasco et al., 2005). A number of studies showed that women have a higher incidence of hip fractures than men (Sanders et al., 1999b, Seeman, 2002, Wehren and Magaziner, 2003).

Bone mineral density (BMD), a surrogate marker of strength of bone (Bonnick, 2007, Rivadeneira et al., 2007), has also been widely used in clinical bone research as the main diagnostic factors to estimate fracture risk (Heaney, 2005, Pulkkinen, 2004). Measurement of BMD using DXA only generate length and breadth (two dimensional), acting as a surrogate for three dimensional volume (length, breadth and depth of bone structure) (Heaney, 2005). Thus, other important elements, for instance the size, shape, geometries, and relative amounts of bone in cortical and trabecular compartments that are related to bone strength is not measured directly by conventional DXA (Black et al., 2008). On the contrary, HSA has been able to generate 3 dimensional measurement from 2 dimensional measurement of DXA scans results using specific principles developed by Beck and colleagues (Beck et al., 1990).

Previous studies from the US (Kaptoge, 2008) and Netherland (Rivadeneira et al., 2007) population has shown that some parameters generated from HSA measurements, areal BMD, cortical thickness (AVCO) and buckling ratio

(AVBR), were associated with increased risk of hip fracture. These results support the strong potential of BMD and other hip geometries to be used as risk predictors for future hip fracture (Kaptoge et al., 2008, Rivadeneira, 2007). However, there are indications that variation in BMD and other geometrical dimension of bone is largely influenced by ethnicity (Wang, 2005, Nelson, 2004, Peacock et al., 2009). Therefore, the utility of this method needs to be confirmed in terms of its consistency in a different setting across different age. The Objective of the research was to examine the association between hip geometries in Femoral Neck and the risk of hip fracture in older women.

## METHODS

### **Study Design**

A case control design was used to determine risk factors of fracture, in particular, measured from FN geometries.

### **Data Collection**

This study was conducted using the data drawn from the large epidemiological project known as the Geelong Osteoporosis Study (GOS) with a population cohort (Henry et al., 2000) that commenced in 1994 and a fracture sample (Sanders, 1998) collected at the same time period. The date has been extracted and cleaned by Dr. Margaret Henry.

### **Study Area**

The GOS, a population-based cohort study was based in a region surrounding Geelong in Southern Victoria (Henry et al., 2000). The region named Barwon Statistical Divisions, was defined by the Australia Bureau of Statistics (Henry et al., 2000). The region consists of stable urban and rural populations that may represent the Australian populations (Henry et al., 2000). A small number of radiological centres indicates a complete ascertainment can be attained by accessing all radiological reports (Pasco et al., 1999). There are two cohort data in the GOS; fracture cohort (Sanders, 1998) and population cohort (Henry et al., 2000).

This study used the population and sample from the GOS. The detail of GOS has been published elsewhere (Henry et al., 2000, Sanders, 1998). Study Sample in this research is that the eligible case and control sample was made up of participants in fracture cohort and population cohort of the Geelong Osteoporosis Study respectively in older group aged  $\geq 64$  years old. The inclusion criterion for case group was also that participants have sustained in the low trauma group of hip fracture. Low trauma group was defined by the GOS researchers as fracture due to accidental fall from less than standing height (lying/sitting), accidental fall from standing height, a spontaneous

fracture and other fracture except due to transportation accidents. Moreover, the inclusion criterion for the control group was that participants have not sustained any fracture based on self-reported data in population cohort of GOS. Participants who had previous hip fractures were

excluded in fracture groups. The consideration of the exclusion was that hip structure in participants with previous fracture would be not similar with that in participants who never have hip fracture previously. Therefore, exclusion might reduce bias.




Figure 1 Hip Fracture picture and hip Anatomy (Matt, 2002, Matt, 2003)

### Statistical Analysis

All analysis procedures were performed using a statistical computer program, STATA version 10. Initially, case and control definitions were re-checked to eliminate the possibility of misclassification due to statistical properties. Missing values and wrong coding were also properly checked and corrected as necessary.

The outcome, fracture status, was measured dichotomously, coded into 1 for cases (fracture group) and 0 for control (non fracture group). At the beginning, basic descriptive statistics were performed to review the description of each variable. Means, standard deviation or standard errors were generated for continuous variables with normal distribution and median and range for those with asymmetric distribution, while proportion was used for categorical variables.

Exposures were hip geometries and outcome was risk of fracture. Univariable analysis was performed to find a difference between participants' characteristics and the outcome. Student t-test was used for the continuous exposure, while Chi-Square test was performed for

categorical exposure. Pearson's correlation was performed to find correlation value between two hip geometries. Subsequently, the association of hip geometries and risk of hip fracture was performed by using logistic regression. This estimated regression coefficients were expressed by odds ratio for the association between risk of fracture and hip geometries (dimensions). OR of hip fracture corresponds to 1 SD change (decrease or increase) in the Hip structural variables increasing risk of hip fracture, therefore the interpretation of possible hip predictors of fracture were able to be described clearly. Regression coefficient (the first objective), the Odds Ratio (the second objective), the 95 % confidence interval and *p* value were reported.

### RESULTS

**Association between hip geometries in Femoral Neck and risk of hip fracture Characteristics of sample**

**Table 1 Characteristics of Participants in Fracture and Non Fracture Group**

Characteristics	Fracture group N=44	Non fracture group N=454	P value*
Steroid/Corticosteroid Hormone Use , n, (%) Yes	3 (4%)	49 (10%)	0.383
Smoking Status, n, (%) Yes	18 (40%)	142 (31%)	0.232
Menopause Status , n, (%) Yes	45 (100%)	454 (100%)	-
Calcium/Multivitamin D Use , n, (%) Yes	4 (9 %)	92 (20%)	0.07
Hormone Replacement Therapy, n, (%) Yes	4 (9%)	82 (18 %)	0.303
Physical Activity n, (%)			
Active	8(18%)	179 (40%)	
Sedentary	15 (34 %)	180 (40%)	<0.0001
Limited	21 (8 %)	95 (82 %)	
Family History , n, (%) Yes	1 (2%)	29 (6%)	0.26
Drinking status, n, (%) Yes	35 (13 %)	10 (4 %)	0.001
	<b>Mean (SD)</b>		<b>P Value**</b>
Height ,cm	156.88 (6.28)	156 (6.17)	0.72
Current Age, year	79.1 (7.54)	75.65 (6.91)	0.001
Weight, kg	61.39 (11.17)	65.37 (12.64)	0.04
Bone mineral density, g/cm <sup>2</sup>	0.59 (0.14)	0.68 (0.14)	<0.0001
Cross sectional area, cm <sup>2</sup>	1.64 (0.40)	1.78 (0.36)	0.014
Cross sectional Moment of Inertia, cm <sup>4</sup>	1.52 (0.51)	1.43 (0.40)	0.016
Width, cm	2.90 (0.25)	2.73 (0.28)	0.0001
Section modulus, cm <sup>3</sup>	0.98 (0.30)	0.99 (0.23)	0.69
Endocortical diameter, cm	2.67 (0.26)	2.46 (0.30)	<0.0001
Average cortical thickness, cm	0.11 (0.03)	0.13 (0.03)	<0.0001
Buckling ratio	14.54 (4.35)	11.41 (3.33)	<0.0001
Centroid Position	0.48 (0.03)	0.49 (0.03)	0.04

\* Chi Square function    \*\* Student t-test function

Table 1 shows the basic characteristics of participants in this study according to fracture and non-fracture. There were 44 participants (8.8%) in fracture group and 454 participants (91.2%) in non-fracture group. Fracture group had lower weight, older age, and slightly higher height compared than non-fracture group. Furthermore, some hip geometry dimensions, including the average value of BMD, WID, ENDO, AVCO, and AVBR were considerably different among fracture and non fracture group ( $p \leq 0.0001$ ). Differences of characteristics of study participants between fracture and non fracture group were also found in physical activity ( $p < 0.0001$ ) (table 1)

#### **Association between hip structural analysis variables in Femoral Neck region and risk of hip fracture in older women**

Table 2 shows the unadjusted and adjusted model of the association between HSA Parameters and hip fracture. In model 1, BMD, WID, ENDO, AVCO, and AVBR were strongly associated (all  $p < 0.0001$ ) and CSMI and SectMod were weakly associated with risk of hip fracture. After adjusting for age, height, weight and physical activity, most of these associations remain, however the strength of this association was reduced (model 2, table 1).

**Table 2** Association of HSA parameters in NN region with hip fracture

Parameters (per 1 SD changes)	Direction of change <sup>^^</sup>	Model 1 <sup>†</sup>		Model 2 <sup>§</sup>	
		OR (95% CI)	P value	OR	P value
Bone mineral density(BMD)	(-)	2.18 (1.5, 3.18)	<0.0001	1.98(1.21, 3.23)	0.006
Cross sectional area(CSA)	(-)	1.53(1.09, 2.15)	0.015	1.26(0.81, 1.96)	0.311
Cross sectional Moment of Inertia (CSMI)	(+)	1.22(0.92, 1.62)	0.164	1.43(1.02, 2.02)	0.04
Width (WID)	(+)	1.94 (1.39, 2.70)	<0.0001	1.70 (1.18, 2.45)	0.005
Section modulus (SECMOD)	(+)	1.06(1.46, 0.78)	0.685	1.19 (1.75, 0.81)	0.368
Endocortical diameter (ENDO)	(+)	2.14(1.52, 3.00)	<0.0001	1.80(1.23, 2.62)	0.002
Average cortical thickness (AVCO)	(-)	2.24(1.52, 3.29)	<0.0001	2.02 (1.23, 3.34)	0.006
Buckling ratio	(+)	2.03(1.54, 2.67)	<0.0001	1.85(1.32, 2.61)	<0.0001
Centroid Position (CENPOS)	(-)	1.44(1.01, 2.03)	0.041	1.25(0.89,1.76)	0.188

† Unadjusted

§ Adjusted for dimension age, height, weight, and physical activity

<sup>^^</sup>The direction of change is indicated as increase (+) or decrease (-) 1 SD of hip geometries.

## DISCUSSION

This case control study also suggests that increased AVBR, WID, ENDO and a reduced BMD and AVCO in FN region are associated with an increased risk of hip fracture with independent effect of height, weight, age and physical activity. Findings suggest that not only is BMD associated with hip fractures, but also other hip geometry dimensions, including WID, ENDO, AVCO and AVBR, independent of age, height, weight and physical activity.

Association between hip geometries and hip fracture have been widely studied. Zebaze (2007) concluded in their study in Lebanon that the result of failure to adapt bone's architecture to loading may lead to bone fragility, not just low bone mass. However, Zebaze used a different method called micro Computed Tomography. Zebaze et al (2005) argues that DXA is not a suitable method to measure FN (FN) depth and volume BMD (vBMD) because DXA assume a circular and a square cross section. However, Beck et. al (1990) have validated that HSA using DXA as accurate method in measuring geometric measurements and strength estimates. Pulkkinen P (2004) and El-Kaissi et al (2005) found the similar conclusions to Zebaze. However, both studies used manufacture rulers to measure hip geometries using a single observer. Hence, the results might have been impacted by observer bias and measurement bias that might weaken the validity of these studies.

Two studies on Hip Structural Analysis parameters were found. Firstly, a research by Rivadeneira et al (2007) with 2740 women aged  $\geq 55$  years old (106 hip fracture) found that an decreased BMD in FN is a strong predictor of hip fracture in women. However, this result is not mechanically clear because this findings suggests that BMD is capturing a strength aspect (Kaptoge, 2008). SectMod as an indicator for bone strength is not present in this study (Rivadeneira et al., 2007). However, Kaptoge (2008), a study with 635 hip fracture cases among 7474 participants, found that each dimension in FN adjusted by age, weight and height was associated with hip fracture (Kaptoge, 2008).

Both studies used cohort design that could minimize some limitations in case control design, including bias and confounders. In this study, cohort study design was not performed, because the population cohort in the GOS consisted of only approximately 700 participants aged over 50 years at baseline. Only a small sample size could be obtained if cohort designs had been used. Moreover, Kaptoge's and Rivadeneira's research was conducted in the USA and the Netherlands respectively with different ethnics groups comparing to Australia. The results might not directly equate to the Australian population, because different races and ethnics have different bone structures (Wang, 2005, Nelson, 2004, Peacock et al., 2009). However, the writers also

suggest that hip geometries in FN indicates strong evidence associated with risk of fracture regardless of the different research methods, study design and ethnics based on this research and previous research above. Moreover, the utility of HSA method has been confirmed in terms of its consistency in a different setting across different age in this study.

Australia, like other western countries, has to cope with the health problem related to ageing as early as possible, particularly osteoporotic hip fractures. This condition is more likely to be major component of the expected increased demand for acute-care hospital services in the future (Pocock et al., 1999). Case findings of hip fracture related osteoporosis, using measurements based on only BMD in WHO criteria for this high risk group (prior fragility fracture, corticosteroid use, family history, and low body mass index), could be underestimating the scope of the problem (Kanis et al., 2002). Therefore, additional hip geometries as independent factor of hip fracture could be considered a clinical assessment method to evaluate risk of hip fractures in high risk group in the population.

There are also some limitations in this study. Firstly, investigating possible confounder factors in this study was based on self reporting; therefore recall bias might have also been present. However, those residual measurements are difficult to be controlled in every research. Secondly, limitation of DXA is that it only produces a two dimensional image of bone to predict a three dimensional image. Projection images from 3-dimensional can be performed using quantitative computed tomography (QCT) application. However, this costly measurement requires much higher effective dose (Genant, 1993). Thirdly, the small sample size in fracture group is a potential limitation. Moreover, cervical fracture and trochanteric fracture were combined as hip fracture. Hence, our study result should be interpreted cautiously to detect due to less statistical power. However, the results of this study could be as baseline of description that the importance of hip geometries in predicting hip fracture.

Some future research is proposed. Sensitivity and specificity can be considered to explore the predictive power of this model as the following study after this research. Moreover, a cohort study with a larger number of samples in the baseline could be performed to reduce the limitation of case control design. Another study that is proposed is meta-analysis of some studies by pooling other research with the same research questions and variables in order to obtain results with high statistical power.

## CONCLUSION

These results provide additional insights that the geometries of FN is associated with fracture neck of femur in older women and strongly suggest its potential value, as clinical predictors for assessing the risk of hip fracture in older women. In addition to this, utilization of some combined parameters of bone geometries in FN may be a more effective method in screening than case findings to reduce the burden of hip fracture in the future. However, more studies with stronger design are clearly needed to eliminate some limitations found in this study.

## ACKNOWLEDGEMENTS

I would like to say heap gratitude for Assoc prof Julie Pasco and Dr Margaret Henry for giving me a chance to analyze some part of GOS, and also for Assoc Prof Lyle Gurrin as my main supervisor during my study. I would like to extend special thanks to my friends, M. Bayu Sasongko, Mahmud Shahid, Aung Ko Win, Farhana Sultana and Roz Turton in School of Population Health, The University of Melbourne.

## REFERENCES

1. Beck, T. Hip Structural Analysis (HSA) Program,(BMD and Structural Geometry Methodology),As Used to Create NHANES III Dataset1.2002a.
2. Beck, T. J. Predicting Femoral-Neck strength from bone mineral data, A structural approach. *Investigative Radiology*, 1990. 25, 6-18.
3. Beck, T. J. *Hip Structural Analysis (HSA) Program (BMD and Structural Geometry Methodology), As Used to Create NHANES III Data*, Baltimore, Johns Hopkins University, School of Medicine. 2002b.
4. Beck, T. J., Oreskovic, T. L., Stone, K. L., Ruff, C. B., Ensrud, K., Nevitt, M. C., Genant, H. K. & Cummings, S. R. Structural Adaptation to Changing Skeletal Load in the Progression Toward Hip Fragility: The Study of Osteoporotic Fractures. *Journal of Bone and Mineral Research*, 2001. 16, 1108-1119.
5. Beck, T. J., Ruff, C. B., Warden, K. E., Mebe, Scott, W. S. & Rao, G. U. Predicting Femoral-Neck strength from bone mineral data, A structural approach. *Investigative Radiology*, 1990. 25, 6-18.
6. Black, D. M., Bouxsein, M. L., Marshall, L. M., Cummings, S. R., Lang, T. F., Cauley, J. A., Ensrud, K. E., Nielson, C. M. & Orwoll, E. S. Proximal Femoral Structure and the Prediction of Hip Fracture in Men: A Large Prospective

- Study Using QCT. *Journal of Bone and Mineral Research*, 2008. 23, 1326-1333.
7. Bonnicksen, S. L. HSA: Beyond BMD with DXA. *BONE*, 2007. 41, S9-S12.
  8. Bonura, F. Prevention, screening, and management of osteoporosis: An overview of the current strategies. *Postgraduate Medicine*, 2009. 121, 5-17.
  9. Deng, H.-W., Xu, F.-H., Davies, K. M., Heaney, R. & Recker, R. R. Differences in bone mineral density, bone mineral content, and bone areal size in fracturing and non-fracturing women, and their interrelationships at the spine and hip. *Journal of bone and mineral metabolism*, 2002. 20, 358-66.
  10. El-Kaissi, S., Pasco, J. A., Henry, M. J., Panahi, S., Nicholson, J. G., Nicholson, G. C. & Kotowicz, M. A. Femoral neck geometry and hip fracture risk: the Geelong osteoporosis study. (ORIGINAL ARTICLE). *Osteoporosis International*, 2005. 16, 1299(5).
  11. Fisher, A. A., O'Brien, E. D. & Davis, M. W. Trends in hip fracture epidemiology in Australia: Possible impact of bisphosphonates and hormone replacement therapy. *Bone*, 2009. 45, 246-253.
  12. Genant, H. K. Bone Densitometry - Current Assessment. *Osteoporosis International*, 1993. 3, S91-S97.
  13. Hannan, E. L. Mortality and locomotion 6 months after hospitalization for hip fracture - Risk factors and risk-adjusted hospital outcomes. *JAMA-Journal Of The American Medical Association*, 2001. 285, 2736-2742.
  14. Heaney, R. P. BMD: The problem. *Osteoporosis International*, 2005. 16, 1013-1015.
  15. Henry, M. J., Pasco, J. A., Nicholson, G. C., Seeman, E. & Kotowicz, M. A. Prevalence of osteoporosis in Australian women: Geelong osteoporosis study. *Journal of Clinical Densitometry*, 2000. 3, 261.
  16. Johnell, O. & Kanis, J. A. An estimate of the worldwide prevalence and disability associated with osteoporotic fractures. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*, 2006. 17, 1726-33.
  17. Kanis, J. A., Black, D., Cooper, C., Dargent, P., Dawson-Hughes, B., De Laet, C., Delmas, P., Eisman, J., Johnell, O., Jonsson, B., Melton, L., Oden, A., Papapoulos, S., Pols, H., Rizzoli, R., Silman, A. & Tenenhouse, A. A new approach to the development of assessment guidelines for osteoporosis. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA*, 2002. 13, 527-36.
  18. Kaptoge, S. Effects of gender, anthropometric variables, and aging on the evolution of hip strength in men and women aged over 65. *Bone*, 2003. 32, 561-70.
  19. Kaptoge, S. Effects of physical activity on evolution of proximal femur structure in a younger elderly population. *Bone*, 2007. 40, 506-15.
  20. Kaptoge, S. Prediction of Incident Hip Fracture Risk by Femur Geometry Variables Measured by Hip Structural Analysis in the Study of Osteoporotic Fractures. *Journal Of Bone And Mineral Research*, 2008. 23, 1892-1904.
  21. Kaptoge, S., Beck, T. J., Reeve, J., Stone, K. L., Hillier, T. A., Cauley, J. A. & Cummings, S. R. Prediction of incident hip fracture risk by femur geometry variables measured by hip structural analysis in the study of osteoporotic fractures. *Journal of bone and mineral research : the official journal of the American Society for Bone and Mineral Research*, 2008. 23, 1892-904.
  22. Khasraghi, F. A. The economic impact of medical complications in geriatric patients with hip fracture. *OrthopedicS*, 2003. 26, 49-53.
  23. Lim, L. S. Screening for osteoporosis in the adult U.S. population: ACPM position statement on preventive practice. *Am J Prev Med*, 2009. 36, 366-75.
  24. Matt. Hip Fractures. 2002.
  25. Matt. Hip Anatomy. 2003.
  26. Nelson, D. A. Sex and ethnic differences in bone architecture. *Curr Osteoporos Rep*, 2004. 2, 65-9.
  27. Pasco, J. A. The human cost of fracture. *Osteoporosis International*, 2005. 16, 2046-2052.
  28. Pasco, J. A., Henry, M. J., Gaudry, T. M., Nicholson, G. G. & Kotowicz, M. A. (1999) Identification of incident fractures: The Geelong osteoporosis study. *Australian and New Zealand Journal of Medicine*, 1999. 29, 203.
  29. Pasco, J. A., Sanders, K. M., Hoekstra, F. M., Henry, M. J., Nicholson, G. C. & Kotowicz, M. A. The human cost of



- fracture. *Osteoporosis International*, 2005.16, 2046-2052.
30. Peacock, M., Buckwalter, K. A., Persohn, S., Hangartner, T. N., Econs, M. J. & Hui, S. Race and sex differences in bone mineral density and geometry at the femur. *Bone*, 2009. 45, 218-225.
  31. Pocock, N. A., Culton, N. L. & Harris, N. D. The potential effect on hip fracture incidence of mass screening for osteoporosis. *The Medical journal of Australia*, 1999. 170, 486-8.
  32. Pulkkinen, P. Combination of bone mineral density and upper femur geometry improves the prediction of hip fracture. *Osteoporosis International*, 2004. 15, 274-280.
  33. Randell, A., Sambrook, P. N., Nguyen, T. V., Lapsley, H., Jones, G., Kelly, P. J. & Eisman, J. A. Direct clinical and welfare costs of osteoporotic fractures in elderly men and women. *Osteoporosis International*, 1995. 5, 427-432.
  34. Rivadeneira, F., Et Al. Femoral neck BMD is a strong predictor of hip fracture susceptibility in elderly men and women because it detects cortical bone instability: The Rotterdam study. *Journal of Bone and Mineral Research*, 2007. 22, 1781-1790.
  35. Rivadeneira, F., Zillikens, M. C., De Laet, C. E., Hofman, A., Uitterlinden, A. G., Beck, T. J. & Pols, H. A. P. Femoral neck BMD is a strong predictor of hip fracture susceptibility in elderly men and women because it detects cortical bone instability: The Rotterdam study. *Journal of Bone and Mineral Research*, 2007. 22, 1781-1790.
  36. Sanders, K. M. The exclusion of high trauma fractures may underestimate the prevalence of bone fragility fractures in the community: the Geelong Osteoporosis Study. *J Bone Miner Res*, 1998. 13, 1337-42.
  37. Sanders, K. M., Nicholson, G. C., Ugoni, A. C., Pasco, J. A., Seeman, E. & Kotowicz, M. A. Health burden of hip and other fractures in Australia beyond 2000 - Projections based on the Geelong Osteoporosis Study. *Medical Journal Of Australia*, 1999a. 170, 467-470.
  38. Sanders, K. M., Seeman, E., Ugoni, A. M., Pasco, J. A., Martin, T. J., Skoric, B., Nicholson, G. C. & Kotowicz, M. A. Age- and gender-specific rate of fractures in Australia: A population-based study. *Osteoporosis International*, 1999b.10, 240.
  39. Seeman, E. Osteoporosis: trials and tribulations. *The American journal of medicine*, 1997. 103, 74S-87S; d-89S.
  40. Seeman, E. Pathogenesis of bone fragility in women and men. *The Lancet*, 2002. 359, 1841-1850.
  41. Wang, X.-F. Varying contributions of growth and ageing to racial and sex differences in femoral neck structure and strength in old age. *Bone*, 2005. 36, 978-86.
  42. Wehren, L. E. & Magaziner, J. Hip fracture: Risk factors and outcomes. *Current Osteoporosis Reports*, 2003. 1, 78-85.
  43. Youm, T., Koval, K. J. & Zuckerman, J. D. The economic impact of geriatric hip fractures. *American journal of orthopedics (Belle Mead, N.J.)*, 1999. 28, 423-8.
  44. Zebaze, R. M. D. Construction of the femoral neck during growth determines its strength in old age. *Journal Of Bone And Mineral Research*, 2007. 22, 1055-1061.
  45. Zebaze, R. M. D., Jones, A., Weslh, F., Knackstedt, M. & Seeman, E. Femoral neck shape and the spatial distribution of its mineral mass varies with its size: Clinical and biomechanical implications. *BONE*, 2005. 37, 243-252.