

# CATENA

AN INTERDISCIPLINARY JOURNAL OF  
**PEDOLOGY - HYDROLOGY - GEOMORPHOLOGY**  
INTERDISZIPLINÄRE ZEITSCHRIFT FÜR  
**HYDROLOGIE - GEOMORPHOLOGIE - PEDOLOGIE**  
JOURNAL INTERDISCIPLINAIRE DE  
**GEOMORPHOLOGIE - PEDOLOGIE - HYDROLOGIE**

EDITORS:

F. Ahnert, Aachen  
G. Bartels, Köln  
L. Berry, Worcester, Mass.  
J. J. Bigarella, Curitiba  
B. Bunting, Hamilton, Ont.  
K. Butzer, Chicago  
G. Castary, Orléans  
J. Demek, Brno  
I. Douglas, Armidale, N. S. Wales  
G. H. Dury, Madison, Wisc.  
A. R. Eschner, Syracuse, N. Y.  
H. Fölster, Göttingen  
H. Faure, Marseille  
R. Herrmann, Bayreuth  
P. D. Jungerius, Amsterdam  
J. Kukla, Palisades  
L. B. Leopold, Berkeley  
J. A. Mabbutt, Kensington  
B. Messerli, Bern  
B. Meyer, Göttingen

P. Michel, Strasbourg  
R. B. Morrison, Denver  
D. Newson, Wallingford  
R. Paepe, Brussel  
H. Paloc, Orléans  
M. Pecsí, Budapest  
J. de Ploey, Leuven  
R. Pullan, Liverpool  
R. V. Ruhe, Bloomington, Ind.  
A. Ruellan, Rennes  
A. P. Schick, Jerusalem  
A. Semmel, Frankfurt  
O. Slaymaker, Vancouver  
L. Starkel, Krakow  
A. Velichko, Moskva  
J. Vogt, Orléans  
R. Webster, Oxford  
D. H. Yaalon, Jerusalem  
A. Young, Norwich

CHIEF EDITOR:  
H. Rohdenburg, Braunschweig

# CATENA

AN INTERDISCIPLINARY JOURNAL OF PEDOLOGY – HYDROLOGY – GEOMORPHOLOGY  
INTERDISZIPLINÄRE ZEITSCHRIFT FÜR HYDROLOGIE – GEOMORPHOLOGIE – PEDOLOGIE  
JOURNAL INTERDISCIPLINAIRE DE GEOMORPHOLOGIE – PEDOLOGIE – HYDROLOGIE

## GUIDE FOR AUTHORS

The title, CATENA, is chosen to represent the connections between different disciplines which the journal will seek to make. At the same time it symbolizes an attempt to unite scientists of different nations, modes of thought and, not least, different languages.

The editors of CATENA take the view that the emphasis on interdisciplinary aspects of PEDOLOGY, HYDROLOGY, GEOMORPHOLOGY and the related aspects of GEOLOGY, CLIMATOLOGY and the BIOLOGICAL SCIENCES can contribute to the advancement of these disciplines in their own right, as well as relating them to each other. Certainly there are at the present large gaps between specialisms, which it is worth attempting to fill.

CATENA publishes original research, reviews of research fields, and critical commentaries on particular articles or themes. Review sections will not seek complete coverage, but will indicate publications of particular importance in relation to its chosen interdisciplinary fields.

### Publication details

1. Suitable manuscripts will be published as rapidly as possible (4 - 6 months). The printing time will be reduced to the minimum required by technical necessities.
2. There is no general rule regarding length of manuscripts. However, general preference will be given to manuscripts of medium length (10 - 15000 words).
3. In order to meet these requirements, it is intended that there should be flexibility with respect to publication dates. Individual issues will be of varying length, depending on manuscripts offered. These will be grouped into continuously paginated volumes with a length of 400 - 450 pages.

In order to avoid delay in publication, manuscripts must conform with the directions outlined below. Manuscripts inadequately prepared will be returned to the author(s). Manuscripts will be reviewed by two editors (one of whom may be chosen by the author) regarding the following points: title, summary, length, arrangement and content of the text, adequate documentation, soundness of interpretations and conclusions as justified by the data, language standard, illustrations, bibliography.

Manuscripts should be sent to:

either a) one copy to an editor of free choice, and two copies to the Editorial Office (stating the name of editor already in possession of the third copy)

or b) directly to the Editorial Office in triplicate.

In case of several authors all correspondence will be made with the person submitting the manuscript.

The originals of manuscript, figures and photos should be sent to the Editorial Office; illustrations for the second and third manuscripts can be copies.

Manuscripts should be arranged in this sequence:

- a) Title
- b) Author(s)
- c) Summary
- d) Zusammenfassung (German) or Résumé (French)
- e) Contents
- f) List of illustrations } in more voluminous papers only
- g) Text
- h) Bibliography

On title page: title of article, author(s), affiliation(s), complete address(es).

All essays must have an English summary containing essential results.

English papers: Summary + Résumé or Zusammenfassung

French papers: Summary + Résumé

German papers: Summary + Zusammenfassung

Additional summaries in other languages can be printed. Translations of summaries into German can be provided by the Editorial Office.

RADIOCARBON CHRONOLOGY OF LATE QUATERNARY LAKES  
IN THE KALAHARI, SOUTHERN AFRICA

K. Heine

Geographisches Institut der Rheinischen Friedrich-Wilhelms-Universität  
Franziskanerstraße 2, D-5300 Bonn 1, Bundesrepublik Deutschland*(manuscript accepted for publication June 12, 1978)*SUMMARY

Some radiocarbon dates concerning the late Quaternary development of the southern Kalahari and the Ngami-Makgadikgadi area are given. The southern Kalahari is characterized by pluvial conditions during the last glacial maximum (ca. 19,000 to 12,000 yr b.p.) and relatively arid conditions during the Holocene and before ca. 19,000 yr b.p.. A very high lake level in the Makgadikgadi depression occurred during ca. 30,000 to 19,000 yr b.p.; here the last glacial maximum was arid; a second pluvial period with less high lake levels is dated ca. 12,000 yr b.p.. Minor fluctuations of the Makgadikgadi lake level eventually occurred during the Early Holocene. The preliminary chronostratigraphy is shown in table 1.

ZUSAMMENFASSUNG

Einige  $^{14}\text{C}$ -Daten zur spätquartären Entwicklung der südlichen Kalahari und des Ngami-Makarikari-Gebietes werden vorgestellt. Die südliche Kalahari wird durch pluviale Bedingungen während des letzten Hochglazials (ca. 19 000 bis 12 000 B.P.) und durch relativ aride Bedingungen während des Holozäns und vor ca. 19 000 B.P. charakterisiert. In der Makarikari-Depression bestand ein sehr hoher Seespiegel während ca. 30 000 bis 19 000 B.P.; hier war das letzte Hochglazial arid; eine zweite pluviale Phase mit weniger hohem Seespiegel wird auf ca. 12 000 B.P. datiert. Geringere Fluktuationen des Makarikari-Seespiegels gab es vermutlich im Frühholozän. Eine vorläufige Chronostratigraphie wird in Tabelle 1 wiedergegeben.

The recent increase in observational evidence for late Quaternary moist periods in North Africa, East Africa, Arabia, and Australia (ROGNON & WILLIAMS 1977, BUTZER et al. 1972, GASSE 1977, McCLURE 1976, BOWLER 1975), derived largely from dating and analysis of lake sediments, has been accompanied by little attention on fossil lake beds in the Kalahari of southern Africa, although fossil lakes have long been known to exist in the Kalahari region (PASSARGE 1904, JAEGER 1939, LANCASTER 1974). The results of a preliminary attempt to establish the age of a series of late Quaternary sediments and fluvial deposits in the Kalahari region are reported here.

The Kalaharian zone is situated on a plateau generally more than 1,000 m above sea level in Botswana, eastern South West Africa (Namibia), and north western Transvaal, stretching from the Okavango River in the north to close to the Orange River in the Republic of South Africa in the south (Fig. 1). It is flat or gently undulating, with sand dunes more frequently occurring in the south west. The true Kalahari is a huge sand-filled basin (DE VOS 1975). The zone is semi-arid. Rain-




Fig. 1: Location map of places mentioned in the text. 1, Swamp; 2, Lake; 3, Pan; 4, Perennial river; 5, Valley with episodic run-off; 6, Fossil lake shore during the most extensive late Quaternary transgressive phase; 7, Minor fossil lake shore; 8, International border.

main generations of lake beds in the Ngami-Okavango-Makgadikgadi region. The older lake may on one or more occasions have formed a very large and continuous body of water with the Okavango-Mababe-Ngami depression (GREY & COOKE 1977), estimated to have been about 60,000 km<sup>2</sup> in area. Aeolian sands antedate the sediments of the younger lake. The 'contouring' of the slopes of dunes of the Ntwetwe pan in the Makgadikgadi is believed to have been caused by the retreating waters of the younger lake that flooded the pan after the formation of the dunes (GROVE 1969).

In the southern Kalahari the fossil bearing fluvial deposits of the Molopo River are underlain by aeolian sands; modern dune sands of light colour cover the fluvial sediments of the valleys, whereas the sand ridges of the neighbouring areas show the bright red colour which is characteristic of the Kalahari dunes.

Radiocarbon dating of the lake beds from the Ngami-Makgadikgadi basins and of the fluvial sediments of the Molopo supports the hypothesis that two main periods of high water level occurred in the northern and middle Kalahari during the late Quaternary whereas the southern Kalahari is characterised by only one pluvial period (Table 1). In the tropical part of the Kalahari the dates place the first pluvial lake between 31,000 (or older) and 19,000 yr b.p., and the different stages of the second pluvial lake between ca. 12,000 and 9,000 yr b.p. (and younger). An arid period between ca. 19,000 and 12,000 yr b.p. separates the two pluvial lakes. In the subtropical southern part of the Kalahari as well as in the Cape province (VAN ZINDEREN BAKKER 1976) the period between ca. 19,000 and 12,000 yr b.p. shows fairly wet climatic conditions (pluvial).

fall is erratic, confined mainly to the period November to April and decreasing from about 500 mm annually in the north and east to about 200 mm in the south west. From 22° latitude southward the zone is dry grassland and savanna, receiving only little and unreliable rainfall (DE VOS 1975, DIEM 1977). The summer climate is dry and continental. Scarcity of surface water and poor soils are the main limiting factors to vegetative growth (COLE & BROWN 1976). Ground waters in the northern Kalahari are directly, and in some cases rapidly, recharged by rain (VERHAGEN et al. 1974). Surface water is available in only a few areas for a short time after the rains, when it collects in pans or shallow depressions. These pans play an important role in the ecology of the area (DE VOS 1975).

Lacustrine deposits are associated with spring tufas and other related deposits, paleosols, aeolian and alluvial deposits, and hold the key to the late Quaternary geomorphological and climatic evolution of the area. The lacustrine deposits consist of calcareous and sometimes fossiliferous marls, clays, and silts and sometimes contain snails and shells. Morphological, stratigraphical, and palaeontological criteria indicate the presence of two

Table 1: THE MAIN STAGES IN THE LATE QUATERNARY EVOLUTION OF THE NGAMI-MAKGADIKGADI BASINS AND THE MOLOPO RIVER VALLEY

Ngami-Makgadikgadi basins (19° - 21°S)			Molopo River valley (ca. 27°S)		
<sup>14</sup> C Age (yr b.p.)	Sedimentary facies	Humidity	<sup>14</sup> C Age (yr b.p.)	Sedimentary facies	Humidity
	aeolian sand and calcrete	alternating from wetter to drier		aeolian sand	very dry with wet intervals
C- 8,720±95 (Hv 8383) C- 9,390±80 (Hv 8378)	calcrete	wetter			?
M 11,920± <sup>1630</sup> <sub>1270</sub> (Hv 8367)	lacustrine sediments	wet/ very wet			?
C+ 13,275±110 (Hv 8380) C+ 14,300±190 (Hv 8386) C+ 14,620±90 (Hv 8381)	Dune sand, fluvial sand near the Okavango delta	dry	Mf 12,480±220 (Hv 8372)  Mf 15,580±350 (Hv 8368)	fluvial deposits (sand, silt, clay) gravel	wet
M 19,170±660 (Hv 8366) L 20,835±355 (Hv 8365) 20,990±1100 (+) L 24,330±270 (Hv 8364) M 25,910± <sup>1210</sup> <sub>1000</sub> (Hv 8371) L 27,050±450 (Hv 8379) M 27,350±550 (Hv 8370) L 30,250±520 (Hv 8382) C+ 31,750±500 (Hv 8387)	lacustrine sediments (chalk, marl, silt, clay)	very wet		aeolian sand	dry
	?	?			?

(+) High Makgadikgadi lake level dated by A.T. Grove, see ref. 17 and ref. 1, p. 306.

L Lacustrine chalk, indicating high lake level.

M Mollusca, indicating high lake level and/or fresh water supply.

Mf Mollusca, indicating a perennial river.

C- Calcrete without fossil calcite fragments.

C+ Calcrete with fossil calcite fragments; the <sup>14</sup>C determination does not represent the true age of the calcrete formation

The preliminary chronostratigraphy (Table 1) confirms the assumption that the major environmental changes along the temperate and tropical margins of the Kalahari seem to be in phase with the variations in precipitation and evaporation in North and East Africa as well as in Arabia and Australia. Variations in temperature seem to be synchronous worldwide; variations in humidity depend on a shifting of the climatic belts during the late Quaternary and therefore do not occur synchronously in various regions of southern Africa. In the Kalaharian zone, as in North and East Africa, Arabia, Australia, and Mexico (HEINE 1974, 1977), we have to distinguish between the following periods with different climatic conditions: (1) the high glacial period (19,000 to 12,000 yr b.p.) characterised by a cool and arid climate in the northern, western, middle, and eastern Kalahari and a cold and humid climate (winter rains) in the southern Kalahari, (2) the interstadial period ( $\geq 31,000 - 19,000$  yr b.p.) and the late Pleistocene/early Holocene (12,000 - 9,000 yr b.p.), both periods are characterised by pluvials in the Kalahari, and (3) the postglacial period ( $< 9,000$  yr b.p. = Holocene) with a warm and semihumid climate (tropical summer rains) in the northern, western, middle, and eastern Kalahari and a warm and semi-arid climate (subtropical anticyclone) in the southern Kalahari.

#### ACKNOWLEDGEMENTS

These investigations were sponsored by the DEUTSCHE FORSCHUNGSGEMEINSCHAFT. Special acknowledgement is due to Professor Dr. M.A. GEYH (Hannover) for radiocarbon dating and to Professor Dr. R. HUCKRIEDE (Marburg) for palaeontological determinations of shells and snails.

#### BIBLIOGRAPHY

- BOWLER, J.M. (1975): Deglacial Events in Southern Australia: Their Age, Nature, and Palaeoclimatic Significance. *Quaternary Studies*, The Royal Society of New Zealand, Bull. 13, 75 - 82, Wellington.
- BUTZER, K.W., ISAAC, G.L., RICHARDSON, J.L. & WASHBOURN-KAMAU, C. (1972): Radiocarbon dating of East African lake levels. *Science* 175, 1069 - 1076.
- COLE, M.M. & BROWN, R.C. (1976): The vegetation of the Ghanzi area of western Botswana. *Journal of Biogeography* 3, 169 - 196.
- DE VOS, A. (1975): Africa, the Devastated Continent? W. Junk, The Hague.
- DIEM, M. (1977): Ein weiträumiger extremer Regenfall in der Kalahari. *Meteorologische Rundschau* 30, 138 - 144.
- GASSE, F. (1977): Evolution of Lake Abhê (Ethiopia and TFAI), from 70,000 b.p. *Nature* 265, 42 - 45.
- GREY, D.R.C. & COOKE, H.J. (1977): Some Problems in the Quaternary Evolution of the Landforms of Northern Botswana. *Catena* 4, 123 - 133.
- GROVE, A.Z. (1969): Landforms and Climate in the Kalahari and Ngamiland. *The Geographical Journal* 135, 191 - 212.
- HEINE, K. (1974): Bemerkungen zu neueren chronostratigraphischen Daten zum Verhältnis glazialer und pluvialer Klimabedingungen. *Erdkunde* 28, 303 - 312.
- HEINE, K. (1977): Beobachtungen und Überlegungen zur eiszeitlichen Depression von Schneegrenze und Strukturbodengrenze in den Tropen und Subtropen. *Erdkunde* 31, 161 - 178.
- JAEGER, F. (1939): Trockenseen der Erde. *Petermanns Mitteilungen Erg.H.* 236, 1-159.
- LANCASTER, I.N. (1974): Pans of the Southern Kalahari. *Botswana Notes and Records* 6, 157 - 169.
- McCLURE, H.A. (1976): Radiocarbon chronology of late Quaternary lakes in the Arabian Desert. *Nature* 263, 755 - 756.
- PASSARGE, S. (1904): *Die Kalahari*. D. Reimer, Berlin.

- ROGNON, P. & WILLIAMS, M.A.J. (1977): Late Quaternary Climatic Changes in Australia and North Africa: A Preliminary Interpretation. *Palaeogeography, Palaeoclimatology, Palaeoecology* 21, 285 - 327.
- STREET, F.A. & GROVE, A.T. (1976): Environmental and climatic implications of late Quaternary lake-level fluctuations in Africa. *Nature* 261, 385 - 390.
- VAN ZINDEREN BAKKER, E.M. (1976): The Evolution of Late-Quaternary Palaeoclimates of Southern Africa. *Palaeoecology of Africa* 9, 160 - 202.
- VERHAGEN, B.Th., MAZOR, E. & SELLSCHOP, J.P.F. (1974): Radiocarbon and tritium evidence for direct rain recharge to ground waters in the northern Kalahari. *Nature* 249, 643 - 644.





---

E.W. ANDERSON & N.J. COX	
<i>A COMPARISON OF DIFFERENT INSTRUMENTS FOR MEASURING SOIL CREEP</i>	81
ANN R.M. YOUNG	
<i>THE INFLUENCE OF DEBRIS MANTLES AND LOCAL CLIMATIC VARIATIONS ON SLOPE STABILITY NEAR WOLLONGONG, AUSTRALIA</i>	95
A.C. IMESON	
<i>SLOPE DEPOSITS AND SEDIMENT SUPPLY IN A NEW ENGLAND DRAINAGE BASIN (AUSTRALIA)</i>	109
W. SYMADER & W. THOMAS	
<i>INTERPRETATION OF AVERAGE HEAVY METAL POLLUTION IN FLOWING WATERS AND SEDIMENT BY MEANS OF HIERARCHICAL GROUPING ANALYSIS USING TWO DIFFERENT ERROR INDICES</i>	131
K. HEINE	
<i>RADIOCARBON CHRONOLOGY OF LATE QUATERNARY LAKES IN THE KALAHARI, SOUTHERN AFRICA</i>	145
I.D.L. FOSTER	
<i>SEASONAL SOLUTE BEHAVIOUR OF STORMFLOW IN A SMALL AGRICULTURAL CATCHMENT</i>	151
R. HERRMANN	
<i>REGIONAL PATTERNS OF POLYCYCLIC AROMATIC HYDROCARBONS IN NE-BAVARIAN SNOW AND THEIR RELATIONSHIPS TO ANTHROPOGENIC INFLUENCE AND AIR FLOW</i>	165
A.M. HARVEY	
<i>DISSECTED ALLUVIAL FANS IN SOUTHEAST SPAIN</i>	177
P. MICHEL	
<i>LA VALLEE ALLUVIALE DU SENEGAL (AFRIQUE DE L'OUEST). RELATIONS GEOMORPHOLOGIE - SOLS - APTITUDES CULTURALES ET LEUR CARTOGRAPHIE AU 1/50 000</i>	213

The list of illustrations should contain titles (possibly abbreviated) pertaining to each illustration.

Manuscripts: typewritten, double-spaced,  
size of paper: 21 - 22 x 28 - 30 cm, written on one side only, margin: 4 cm.

Please avoid footnotes. If unavoidable they should be numbered in sequence.

Since page proofs will not be provided, cross-references within the text should refer to sections, not pages.

References in the text:

Every reference to publications in the text should appear in the list of references. References in the text: AUTHOR (year, page) or (AUTHOR year, page). Author's name in capital letters; page-numbers should be given whenever practicable; direct quotations should always be page-referenced.

List of references:

Arranged according to author's surname, and chronologically per author.

AUTHOR (year): Title of article. Journal or booktitle (for multi-author books give editor, volume number (please underline), pages; publisher and city (for books only).

Example:

ZEUNER, F.E. (1953): Das Problem der Pluvialzeiten. Geologische Rundschau 41, 241 - 253.

The journal's title should be written out in full rather than abbreviated.

Tables and illustrations should be enclosed separately (stating name of author, abbreviated title of essay and indication of top of illustration). Numbering should be continuous, and separate for Tables, drawings (marked as Figures) and photographs (marked as Photos). Legends of sufficient length, in the language of the paper and possibly in English, should be collected on a separate sheet (stating author's name and short title of paper).

For the rough draft of line drawings please observe the following guide-lines:

1. All figures and text will be reduced in size 23 %.
2. The maximum size possible for line drawings is 27 x 18.5 cm (after 23 % reduction: 20.8 x 14.2 cm). If larger formats are needed, please consult the Editorial Office.
3. Line drawings should be drafted with black India ink on white drawing paper; high quality prints are acceptable.
4. The final format for all line drawings will be decided by the Editorial Office. The authors may be asked to change their format. It is very important that the *letter size in the line drawings be no less than 2 mm.*
5. Use bar scales on all illustrations instead of numerical scales that must be changed with reduction.

Photographs are acceptable only when showing good contrast and intensity; sharp and glossy prints are required. Prints from existing half-tone illustrations cannot be accepted. Coloured illustrations cannot be included usually unless financed by the authors.

No manuscript or figures will be returned following publication. It is the author's responsibility to obtain written permission to reproduce material which has appeared in another publication. 100 reprints free of charge. Additional reprints may be ordered on the form accompanying the proofs.

## EDITORIAL OFFICE CATENA

c/o Lehrstuhl für Physische Geographie  
und Landschaftsökologie

Technische Universität Braunschweig

Langer Kamp 19 c

D 3300 BRAUNSCHWEIG / W. Germany

Telefon (0531) 391-2561

Fernschreiber: tu bewg 952526

Postfach der TU 3329

Individual issues will be of varying length and will be grouped into continuously paginated volumes of 400 - 450 pages.

Price of Vol. 5, 1978

U\$ S 43.50,

DM 98,-

Personal Subscribers

U\$ S 29.60,

DM 64,-

Copyright by CATENA VERLAG

Margot Rohdenburg M.A.

Brockenblick 8

D 3302 Cremlingen-Destedt/W. Germany.

Druck: Graphische Kunstanstalt W. Herr, 6300 Giessen