



**Geochemistry
Geophysics
Geosystems**

G³

AN ELECTRONIC JOURNAL OF THE EARTH SCIENCES

Published by AGU and the Geochemical Society

Data Brief

Volume 8, Number 10

18 October 2007

Q10009, doi:10.1029/2007GC001694

ISSN: 1525-2027



Bathymetry of the Amundsen Sea continental shelf: Implications for geology, oceanography, and glaciology

F. O. Nitsche and S. S. Jacobs

*Lamont-Doherty Earth Observatory of Columbia University, Route 9 W, Palisades, New York 10964, USA
(fnitsche@ldeo.columbia.edu)*

R. D. Larter

Geological Sciences Division, British Antarctic Survey, High Cross/Madingley Road, Cambridge CB3 0ET, UK

K. Gohl

Alfred Wegener Institute for Marine and Polar Research, P.O. Box 120161, D-27515 Bremerhaven, Germany

[1] The Amundsen Sea continental shelf is one of the most remote areas of coastal Antarctica and was relatively unexplored until the late 1980s. Over the last two decades, increased oceanographic and geological interest has led to several cruises that resulted in sufficient bathymetric data to compile a fairly detailed regional map of the Amundsen continental shelf. We have combined available multibeam and single-beam bathymetry data from various sources and created a new regional bathymetry of the Amundsen Sea continental shelf and margin. Deep trough systems that dominate the inner shelf are aligned with present glaciers and separated by shallower ridges. Shaped by paleo-ice streams, these features merge into a small number of broader troughs on the middle shelf and shoal seaward. They now serve as conduits and reservoirs for relatively warm Circumpolar Deep Water. This new compilation is a major improvement over previously available regional maps and should aid the numerical modeling of ocean circulation, the reconstructions of paleo-ice streams, and the refinement of ice sheet models.

Components: 5032 words, 5 figures, 1 table.

Keywords: West Antarctica; continental shelf; bathymetry; seafloor morphology.

Index Terms: 3002 Marine Geology and Geophysics: Continental shelf and slope processes (4219); 3045 Marine Geology and Geophysics: Seafloor morphology, geology, and geophysics; 4207 Oceanography: General: Arctic and Antarctic oceanography (9310, 9315).

Received 18 May 2007; **Revised** 6 August 2007; **Accepted** 15 August 2007; **Published** 18 October 2007.

Nitsche, F. O., S. S. Jacobs, R. D. Larter, and K. Gohl (2007), Bathymetry of the Amundsen Sea continental shelf: Implications for geology, oceanography, and glaciology, *Geochem. Geophys. Geosyst.*, 8, Q10009, doi:10.1029/2007GC001694.

1. Introduction

[2] The Amundsen Sea continental shelf is located between 100° and 135°W, south of 71°S, along the margin of the Marie Byrd Land sector of the West

Antarctic Ice Sheet (WAIS) (Figures 1 and 4). As one of the more remote Antarctic coastlines, with a nearly perennial sea ice cover and little if any bottom water formation, it remained largely unexplored until the mid 1980s. Interest in the region




Figure 1. Antarctica with the location of the Amundsen Sea. Outline is based on Antarctic Digital Database 4.1 [Thomson and Cooper, 1993] (<http://www.add.scar.org/>).

increased with the 1994 discovery that the ice shelf at the terminus of the Pine Island Glacier was melting orders of magnitude faster than the Ross and Filchner-Ronne Ice Shelves [Jacobs *et al.*, 1996; Jenkins *et al.*, 1997; Hellmer *et al.*, 1998], accompanied by increased velocity, thinning and grounding line retreat [Rignot, 1998; Rignot and Jacobs, 2002]. A melt rate exceeding 40 m/yr near its grounding line [Rignot and Jacobs, 2002] appears to be driven by “warm” Circumpolar Deep Water (CDW) that floods deeper areas of the Amundsen Sea continental shelf. Subsequent investigations have revealed high melt rates for other Amundsen Sea ice shelves, thinning and increased velocities of tributary glaciers, and decreasing ice surface elevations of the adjacent WAIS drainage basins [Wingham *et al.*, 1998; Rignot and Thomas, 2002; Shepherd *et al.*, 2004; Thomas *et al.*, 2004; Davis *et al.*, 2005]. Such findings have revived longstanding concerns about ice sheet stability in this sector [Hughes, 1973], particularly as the WAIS is mostly grounded below sea level [Drewry *et al.*, 1982] and the global ocean is warming [Levitus *et al.*, 2005].

[3] Much of the rapid melting occurs within deep troughs on the inner shelf, presumably cut when the ice streams of larger ice sheets were grounded on the present seafloor [Lowe and Anderson, 2002; Evans *et al.*, 2006]. As these features now extend beneath the small Amundsen Sea ice shelves, it is important to understand how the troughs facilitate CDW access to the ice, along with their character-

istics and connections to the shelf break. A better description of the continental shelf bathymetry could become a key element in improved models of ocean circulation, ocean-ice interactions and ice sheet behavior [Philippon *et al.*, 2006], and in reconstructions of paleo-ice flow [Kellogg and Kellogg, 1987; Lowe and Anderson, 2002; Evans *et al.*, 2006; Larter *et al.*, 2007].

[4] While several previous studies of Amundsen Sea bathymetry provided significant new insights, most have been restricted to local areas covered by single cruises. The aggregated tracks of several expeditions, including recent cruises on the RRS *James Clark Ross* and R/V *Polarstern* in 2006 and the *Nathaniel B. Palmer* in 2007, have now been compiled into a chart that provides an overview of the regional bathymetric setting and its geological, oceanographic and glaciological implications.

2. Setting of the Amundsen Sea Continental Shelf

[5] The tectonic structure of the Amundsen Sea continental shelf results mainly from the Late Cretaceous rifting and breakup between New Zealand and the Marie Byrd Land and Thurston Island blocks, and may also have been affected by the existence of an independent Bellingshausen Plate until the early Tertiary [Stock and Molnar, 1987; Mayes *et al.*, 1990; Larter *et al.*, 2002; Eagles *et al.*, 2004; Gohl *et al.*, 2007]. It has been suggested that the southern boundary of the Bellingshausen Plate crossed the continental shelf of the Amundsen Sea embayment [Larter *et al.*, 2002; Eagles *et al.*, 2004; Gohl *et al.*, 2007].

[6] The present continental shelf was sculpted during past glaciations, displaying surface and subsurface features that indicate grounded ice has reached the outer shelf several times [Kellogg and Kellogg, 1987; Nitsche *et al.*, 1997; Lowe and Anderson, 2002; Evans *et al.*, 2006]. Earlier cruises discovered several deep troughs on the inner shelf [SPRITE Group, 1992; T. B. Kellogg *et al.*, Former rock-floored ice-streams on the Amundsen Sea continental shelf, unpublished manuscript, 2000], and subsequent investigations of the trough originating in Pine Island Bay reported exposed crystalline bedrock on the inner shelf, changing to dipping strata with overlying recent sediment cover toward the mid-shelf [Lowe and Anderson, 2002; Uenzelmann-Neben *et al.*, 2007]. Mega-scale glacial lineations in this trough indicate that it was occupied by paleo-ice streams. The shallower outer

shelf areas (~ 500 m) show widespread iceberg furrows on the seafloor, and broad shelf break depressions that may be related to inner shelf troughs [Lowe and Anderson, 2002; Evans *et al.*, 2006; Walker *et al.*, 2007].

3. Data Compilation

[7] Multibeam swath-mapping on several recent expeditions has significantly improved our knowledge of bathymetric features on and along the Amundsen Sea continental shelf. Here we integrate those data and add other available bathymetric and geophysical measurements to create a new regional map of this sector of the continental margin. Our sources include single-beam and multibeam bathymetry available through the NSF-funded Antarctic Multibeam synthesis database [Carbotte *et al.*, 2007], soundings from earlier cruises in the GEODAS database [National Geophysical Data Center, 1996], and measurements from James Clark Ross cruises JR84 and JR141, Polarstern cruises ANT-XI/3, ANT-XII/4, ANT-XVIII/5a, and ANT-XXIII/4, and NB Palmer cruise 07-02 (Table 1 and Figure 2).

[8] Usually the shoreline boundary of bathymetry compilations is simply set to 0 m, whereas bathymetric features in West Antarctica often continue below the ice shelves and ice sheet. To accommodate these sub-ice features we integrated ice sheet basement data from the US Airborne Geophysical Survey of the Amundsen Sea Embayment, Antarctica (AGASEA) and the UK Basal Balance And Synthesis (BBAS) projects [Holt *et al.*, 2006; Vaughan *et al.*, 2006] and less detailed topography from the BEDMAP project [Lythe *et al.*, 2000] for areas west and north of the AGASEA/BBAS data sets. Our compilation extends from $\sim 100^\circ\text{W}$ to 135°W and 68°S to 76°S , reaching from the Abbot, Cosgrove and Pine Island ice fronts in the eastern Amundsen Sea to the western end of the Getz Ice Shelf, and covering the entire continental slope and most of the continental rise.

[9] Data processing and integration followed the scheme in Figure 3. All single-beam soundings were converted into a common format and then imported into ArcGIS software (ESRI). Most of the swath-bathymetry data sets were initially ping edited and processed aboard ship. We reduced the swath-bathymetry data to manageable size by creating grids of 200–250 m resolution using MB-system [Caress and Chayes, 1996] or Fledermaus IVS [Mayer *et al.*, 2000], depending on the original

format, and imported those grids into ArcGIS, re-converting raster to point data. These point data were the basis for generating new grids. Positional accuracy of the data sets varies, but most data used for this compilation (1988 and younger) were obtained with GPS systems that reached an accuracy of <100 m. Navigation data from earlier cruises, which represent a small fraction of the whole compilation and were used to fill gaps in sparsely covered parts of the continental rise, were collected using the older US Navy TRANSIT satellite navigation system, accurate to ~ 2 km or better, or celestial navigation. To accommodate positional inaccuracies and the varying density of available data we chose a 2 km grid resolution. After testing several interpolation schemes we settled on a “natural neighbor” algorithm [Sibson, 1981] as implemented by ArcGIS for grid formation. This algorithm exactly recovers reference points and can handle highly irregular data distributions [Sambridge *et al.*, 1995], suitable for the present Amundsen Sea where areas covered by high-density swath-bathymetry data are typically separated by large gaps (Figure 2). As data quality varies between individual surveys, several iterations were required to identify and remove disparate measurements. This included single or small groups of measurements that created obvious artifacts in the grid including local, unrealistically deep holes and edge effects resulting from less reliable outer beams of multibeam swaths.

4. New Amundsen Sea Bathymetric Map

[10] The resulting compilation outlines for the first time the full extent and shape of the Amundsen Sea continental shelf, defining its shelf break along most of its length (Figure 4). Seaward of the coastline, the continental shelf narrows westward from >400 km north of Pine Island Bay to 100–200 km west of Siple Island. The outstanding features on the continental shelf are its deep and rugged inner shelf troughs that typically merge and shoal seaward of the ice shelf fronts. Individual troughs often merge into single features by mid-shelf, with the eastern shelf dominated by two trough systems. The one in Pine Island Bay combines troughs that originate from the Pine Island, Thwaites, and Smith Glaciers. A second system includes troughs from the Dotson and eastern Getz ice shelves. Maximum depths in both systems exceed 1600 m, but are not always near the present-day ice fronts. These dendritic trough sys-

Table 1. Survey ID, Data Type, Navigation Equipment, Date, and Sources Used for Compilation, With Selected References Describing or Utilizing These Data

Survey ID	Data Type	Navigation	Year	Source	Reference
ELT11	single-beam PDR 12 kHz	sextant	1964	GEODAS ^a	
ELT17	single-beam PDR 12 kHz	sextant	1965	GEODAS ^a	
ELT33	single-beam PDR 12 kHz	TRANSIT/sextant	1968	GEODAS ^a	
DSDP35GC	single-beam 12 kHz/GIFFT	TRANSIT	1974	GEODAS ^a	<i>Tucholke and Houtz</i> [1976]
THB80	single-beam unknown	unknown	1981	GEODAS ^a	<i>Yamaguchi et al.</i> [1988]
DF85	single-beam SeaBeam 12 kHz	TRANSIT	1985	GEODAS ^a	<i>Kellogg and Kellogg</i> [1987]
PD190L02	single-beam EDO depth tracker	GPS	1990	GEODAS ^a	
NBP92-08	single-beam Simrad:EK500	GPS	1992	GEODAS ^a	
PS92	single-beam	GPS	1992	GEODAS ^a	<i>SPRITE Group</i> [1992]
RITS94B	multibeam SeaBeam	GPS	1993	NGDC ^b	
NBP94-02	single-beam Simrad:EK500	GPS	1994	GEODAS ^a	<i>Jacobs et al.</i> [1996]
ANT-XI/3	multibeam Hydrosweep DS-1	GPS	1994	AWI ^c	<i>Nitsche et al.</i> [2000]
ANT-XII/4	multibeam Hydrosweep DS-1	GPS	1995	AWI ^c	<i>Nitsche et al.</i> [2000]
NBP96-02	multibeam SeaBeam:2112	GPS	1996	AMBS ^d	
NBP99-02	multibeam SeaBeam:2112	GPS	1999	AMBS ^d	<i>Lowe and Anderson</i> [2002]
NBP99-09	multibeam SeaBeam:2112	GPS	1999	AMBS ^d	
NBP00-01	multibeam SeaBeam:2112	GPS	2000	AMBS ^d	
BEDMAP	various	various/GPS	2000	BEDMAP	<i>Lythe et al.</i> [2000]
ANT-XVIII/5a	multibeam Hydrosweep DS-2	GPS	2001	AWI ^c	<i>Scheuer et al.</i> [2006]
JR84	multibeam Simrad EM120	GPS	2003	SPRI ^e	<i>Dowdeswell et al.</i> [2006a]; <i>Evans et al.</i> [2006]
ANT-XXIII/4	multibeam Hydrosweep DS-2	GPS	2006	AWI ^c	<i>Larter et al.</i> [2007]; <i>Gohl et al.</i> [2007]; <i>Uenzelmann-Neben et al.</i> [2007]
JR141	multibeam Simrad EM120	GPS	2006	BAS ^f	<i>Larter et al.</i> [2007]
AGASEA/BBAS	radar	GPS	2006	AGASEA/BBAS	<i>Vaughan et al.</i> [2006]; <i>Holt et al.</i> [2006]
NBP07-02	multibeam Simrad EM120	GPS	2007	AMBS ^d	

^aGEODAS Web site: <http://www.ngdc.noaa.gov/mgg/geodas/geodas.html>.

^bNational Geophysical Data Center <http://www.ngdc.noaa.gov/>.

^cAlfred Wegener Institute for Polar and Marine Research.

^dAntarctic Multibeam Bathymetry and Geophysical Data Synthesis Web site: <http://www.marine-geo.org/antarctic/>.

^eScott Polar Research Institute.

^fBritish Antarctic Survey.




Figure 2. Distribution of the various data sources that form the basis for the new bathymetric grid. Red areas show the coverage of the multibeam bathymetry data, blue lines represent tracks of individual soundings, and green areas represent additional data sets that we used for areas covered by continental ice. Table 1 provides details.

tems are separated by a bank extending northward from the Bear Peninsula, with water depths of ~250–400 m, where deep-draft icebergs go aground. Recently collected swath bathymetry data that have been included in the new bathymetric compilation show that geomorphic features interpreted as subglacial meltwater features [Lowe and Anderson, 2003] are widespread throughout the inner shelf troughs.

[11] Several smaller troughs and trough systems are found near Cosgrove Ice Shelf, and off Carney Island and the central and western Getz Ice Shelf fronts. These inner shelf troughs commonly display rugged relief, to >200 m in Pine Island Bay, and a host of smaller features including striations, grooves, drumlins, and mega-scale lineations indicative of glacial erosion [Wellner et al., 2001; Lowe and Anderson, 2002]. Similar troughs also appear along other parts of the Antarctic continental margin [Pudsey et al., 1994; Canals et al., 2000, 2002; O Cofaigh et al., 2002, 2005] and previously glaciated margins in the northern hemisphere [e.g., Stokes and Clark, 2001; Ottesen et al., 2005; Dowdeswell et al., 2006b; Shaw et al., 2006].

[12] Depths along the continental shelf break average ~500 m, but vary from ~400 to >600 m (auxiliary material¹ Figure S2). An outer shelf depression near 114°W has been charted in some

detail [Dowdeswell et al., 2006a; Walker et al., 2007], but persistent sea ice has limited mapping of other shelf break depressions NW of Thurston and Siple Islands. Connections between these depressions and inner-shelf troughs remain to be confirmed by more detailed mapping of mid-shelf areas usually covered by perennial ice. West of Siple Island a trough near ~128°W can more clearly be followed from the ice edge to the shelf break.




Figure 3. Processing scheme used to integrate the different data sets and create the final grid.

¹Auxiliary materials are available in the HTML. doi:10.1029/2007GC001694.




Figure 4. Bathymetry of the Amundsen Sea based on data available through March 2007 (Table 1). Grid control is based on the ship tracks shown in Figure 2. The coastline, ice shelves, and grounding lines are based on the Antarctic Digital Database 4.1 but have been modified to fit more accurate MODIS satellite image [see also *Swithinbank et al.*, 2003, 2004]. The sub-ice topography is based on AGASEA/BBAS and BEDMAP data sets.




Figure 5. Other representations of Amundsen Sea bathymetry: (a) GEBCO [Mammerix and Cande, 1982], (b) ETOPO-2 [Smith and Sandwell, 1994, 1997], and (c) BEDMAP [Lythe et al., 2000], compared to (d) the new bathymetry grid. The largest differences appear in the outline of the shelf break and in the trough and ridge structures.

[13] The general morphology of the continental slope and rise is comparable to other parts of the Antarctic continental margins. The slope ranges from 2° – 5° with parts of the upper slope being dissected by gullies that in some areas converge downslope and feed into large channels that cross the continental rise [Dowdeswell et al., 2006a; Lowe and Anderson, 2002]. More detailed multibeam coverage of both the slope and shelf is needed to definitively relate gullies, depressions, lineations, troughs and paleo-ice streams. In addition to sedimentary features previously reported on the continental slope [Dowdeswell et al., 2006a] a series of mounds radiate away from the slope between 105°W and 110°W . These mounds connect to known features on the lower continental rise [Yamaguchi et al., 1988; Nitsche et al., 2000; Scheuer et al., 2006]. North of Siple Island (125°W – 127°W) the slope is interrupted by a large SE-NW oriented ridge at depths of 1–3 km, incised by steep canyons. Its shape and water depth preclude glacial links, but its proximity to the Mt. Siple volcano and Marie Byrd Seamounts suggests a tectonic origin.

[14] Earlier depictions of the Amundsen Sea region (BEDMAP/ETOPO/GEBCO in Figure 5) predated

the marine multibeam data utilized here, and so are deficient in several important respects. However, control for the current bathymetry is still uneven, with many large gaps between ship tracks (Figure 2), some resulting from the typical distribution of perennial sea ice. The depth accuracy varies accordingly, being highest in areas with modern multibeam cover, but containing significant uncertainties in the large interpolated areas. Seams between detailed multibeam bathymetry and interpolated areas can cause artifacts such as sudden changes in seafloor roughness and the outward bulge of the lower continental slope between 121°W and 125°W . The seafloor bathymetry is still unknown beneath the ice shelves, although recent airborne radar missions suggest the continuation of some troughs below the Smith, Thwaites and Pine Island Glaciers [Holt et al., 2006; Vaughan et al., 2006].

5. Implications

[15] Concerns about ice sheet mass balance and sea level change have recently focused on the Amundsen Sea because high basal melt rates of its small ice shelves are positively correlated with acceler-

ating ice streams and apparent draw-down of the adjacent ice sheet [Shepherd and Wingham, 2007]. The shelf bathymetry is thought to play a substantial role in this process, both as a gateway facilitating CDW intrusions near the shelf break and a link to deep ice shelf grounding lines. There are indications from observations and models that warm deep water may preferentially move onto continental shelves via shelf break curvatures and depressions [Dinniman *et al.*, 2003; Walker *et al.*, 2007]. However, CDW rises well above the Amundsen shelf break across a wide area [Jacobs *et al.*, 2002], and as yet little is known about the temporal variability of its penetration onto the shelf. Once in the troughs, the saltiest and therefore densest deep water sinks and flows into the ice shelf cavities, its melting potential enhanced by the seawater freezing point depression with increasing pressure [Jacobs *et al.*, 1996].

[16] One consequence of recent work in the Amundsen Sea is that models of ice sheet behavior must now be updated to include the effects of an evolving ocean and its ice shelves. Similarly, models of the ocean circulation must properly account for this remarkable continental shelf, its fringing ice shelves and persistent sea ice cover. Quite different results may be obtained from models based on the bathymetry reported here, compared to that shown on prior maps (Figure 5). The new bathymetry could impact estimates of ice shelf melting, WAIS mass balance, and ocean freshening. The locations and depths of the trough systems are also important for reconstructions of paleo-ice flow and deglaciation. This compilation of Amundsen Sea bathymetry represents a major improvement over previously available regional data sets, which involved far more interpolation between sparser, less accurate observations. A larger-scale version of the map is available in the auxiliary material (Figure S1), and the grid may be obtained from http://www.marine-geo.org/link/entry.php?id=Amundsen_Sea_Nitschke.

Acknowledgments

[17] We thank the personnel aboard the several ships, especially the *NB Palmer*, R/V *Polarstern*, and RRS *James Clark Ross*, who have been responsible for obtaining bathymetric data in the Amundsen Sea, often under difficult conditions. Multibeam data on cruises ANT-XXIII/4 and JR141 were collected and processed by S. Gauger and T. Deen, respectively. T. Kellogg digitized bathymetry from DF85, J. Dowdeswell made data available from JR84, H.-W. Schenke provided access to *Polarstern* data, and J. Anderson, S. Cande, and others contributed observations through the Antarctic Multibeam

Bathymetry Synthesis database. S. Carbotte, S. O'Hara, and the database team at Lamont provided key assistance. The manuscript benefited from the comments by two anonymous reviewers. This work was supported in part by U.S. National Science Foundation through a supplement to U.S. National Science Foundation ANT 04-40655 and by ANT 02-33303 and ANT 04-40775. It also forms part of the BAS Glacial Retreat in Antarctica and Deglaciation of the Earth System (GRADES) Program as well as the MARCOPOLI Work Packages MAR2 and POL6 of AWI. This is Lamont contribution 7068.

References

- Canals, M., R. Urgeles, and A. M. Calafat (2000), Deep seafloor evidence of past ice streams off the Antarctic Peninsula, *Geology*, *28*, 31–34.
- Canals, M., J. L. Casamor, R. Urgeles, A. M. Calafat, E. W. Domack, J. Baraza, M. Farran, and M. De Batist (2002), Seafloor evidence of a subglacial sedimentary system off the northern Antarctic Peninsula, *Geology*, *30*, 603–606.
- Carbotte, S. M., W. B. F. Ryan, S. O'Hara, R. Arko, A. Goodwillie, A. Melkonian, R. A. Weisell, and V. L. Ferrini (2007), Antarctic multibeam bathymetry and geophysical data synthesis: An on-line digital data resource for marine geoscience research in the Southern Ocean, in *Antarctica: A Keystone in a Changing World—Proceedings for the Tenth International Symposium on Antarctic Earth Sciences*, edited by A. K. Cooper, C. R. Raymond, and the ISAES Editorial Team, *U.S. Geol. Surv. Open File Rep.*, 2007-1047, *Short Res. Pap. 002*, doi:10.3133/of2007-1047.srp002, 4 pp.
- Caress, D. W., D. N. Chayes (1996), Improved processing of Hydrosweep DS multibeam data on the R/V Maurice Ewing, *Mar. Geophys. Res.*, *18*(6), 631–650.
- Davis, C. H., Y. Li, J. R. McConnell, M. M. Frey, and E. Hanna (2005), Snowfall-driven growth in East Antarctic ice sheet mitigates recent sea-level rise, *Science*, *308*, 1898–1901.
- Dinniman, M. S., J. M. Klinck, and W. O. Smith (2003), Cross-shelf exchange in a model of the Ross Sea circulation and biogeochemistry, *Deep Sea Res., Part II*, *50*, 3103–3120.
- Dowdeswell, J. A., J. Evans, C. Ó Cofaigh, and J. B. Anderson (2006a), Morphology and sedimentary processes on the continental slope off Pine Island Bay, Amundsen Sea, West Antarctica, *Geol. Soc. Am. Bull.*, *118*, 606–619.
- Dowdeswell, J. A., D. Ottesen, and L. Rise (2006b), Flow switching and large-scale deposition by ice streams draining former ice sheets, *Geology*, *34*, 313–316.
- Drewry, D. J., S. R. Jordan, and E. Jankowski (1982), Measured properties of the Antarctic ice sheet: Surface configuration, ice thickness, volume and bedrock characteristics, *Ann. Glaciol.*, *3*, 83–91.
- Eagles, G., K. Gohl, and R. D. Larer (2004), High-resolution animated tectonic reconstruction of the South Pacific and West Antarctic Margin, *Geochem. Geophys. Geosyst.*, *5*, Q07002, doi:10.1029/2003GC000657.
- Evans, J., J. A. Dowdeswell, C. Ó Cofaigh, T. J. Benham, and J. B. Anderson (2006), Extent and dynamics of the West Antarctic Ice Sheet on the outer continental shelf of Pine Island Bay during the last glaciation, *Mar. Geol.*, *230*, 53–72.
- Gohl, K., et al. (2007) Geophysical survey reveals tectonic structures in the Amundsen Sea embayment, West Antarctica, in *Antarctica: A Keystone in a Changing World—Proceedings for the Tenth International Symposium on Antarctic Earth Sciences*, edited by A. K. Cooper, C. R. Raymond, and the

- ISAES Editorial Team, *U.S. Geol. Surv. Open File Rep.*, 2007-1047, *Short Res. Pap.* 047, doi:10.3133/of2007-1047.srp047, 4 pp.
- Hellmer, H. H., S. S. Jacobs, and A. Jenkins (1998), Oceanic erosion of a floating Antarctic glacier in the Amundsen Sea, *Antarct. Res. Ser.*, 75, 83–99.
- Holt, J. W., D. D. Blankenship, D. L. Morse, D. A. Young, M. E. Peters, S. D. Kempf, T. G. Richter, D. G. Vaughan, and H. F. J. Corr (2006), New boundary conditions for the West Antarctic Ice Sheet: Subglacial topography of the Thwaites and Smith glacier catchments, *Geophys. Res. Lett.*, 33, L09502, doi:10.1029/2005GL025561.
- Hughes, T. (1973), Is West Antarctic Ice-Sheet disintegrating?, *J. Geophys. Res.*, 78, 7884–7910.
- Jacobs, S. S., H. H. Hellmer, and A. Jenkins (1996), Antarctic ice sheet melting in the southeast Pacific, *Geophys. Res. Lett.*, 23, 957–960.
- Jacobs, S. S., P. A. Mele, G. Krahnmann, and W. M. Smethie (2002), Coastal ocean measurements in the Amundsen and Ross Seas, 366 pp., NB Palmer Cruise 00-01, Feb–Mar 2000, *LDEO-2002–2a*, Lamont-Doherty Earth Observ., Palisades, N. Y.
- Jenkins, A., D. G. Vaughan, S. S. Jacobs, H. H. Hellmer, and J. R. Keys (1997), Glaciological and oceanographic evidence of high melt rates beneath Pine Island glacier, west Antarctica, *J. Glaciol.*, 43, 114–121.
- Kellogg, T. B., and D. E. Kellogg (1987), Recent glacial history and rapid ice stream retreat in the Amundsen Sea, *J. Geophys. Res.*, 92(B9), 8859–8864.
- Larter, R. D., A. P. Cunningham, P. F. Barker, K. Gohl, and F. O. Nitsche (2002), Tectonic evolution of the Pacific margin of Antarctica: 1. Late Cretaceous tectonic reconstructions, *J. Geophys. Res.*, 107(B12), 2345, doi:10.1029/2000JB000052.
- Larter, R. D., et al. (2007), West Antarctic Ice Sheet change since the Last Glacial Period, *Eos Trans. AGU*, 88, 189–196.
- Levitus, S., J. Antonov, and T. Boyer (2005), Warming of the world ocean, 1955–2003, *Geophys. Res. Lett.*, 32, L02604, doi:10.1029/2004GL021592.
- Lowe, A. L., and J. B. Anderson (2002), Reconstruction of the West Antarctic Ice Sheet in Pine Island Bay during the Last Glacial Maximum and its subsequent retreat history, *Quat. Sci. Rev.*, 21, 1879–1897.
- Lowe, A. L., and J. B. Anderson (2003), Evidence for abundant subglacial meltwater beneath the paleo-ice sheet in Pine Island Bay, Antarctica, *J. Glaciol.*, 49, 125–138.
- Lythe, M. B., D. G. Vaughan, and the BEDMAP Consortium (2000), BEDMAP—Bed topography of the Antarctic, 1:10,000,000 scale map, *Br. Antarct. Surv.*, Cambridge, U. K.
- Mammerix, J., and S. Cande (1982), *General Bathymetric Charts of the Oceans (GEBCO)*, Can. Hydrogr. Surv., Ottawa, Canada.
- Mayer, L. A., M. Paton, L. Gee, J. Gardner, and V. C. Ware (2000), Interactive 3D Visualization: A tool for seafloor navigation, exploration and engineering, paper presented at OCEANS 2000 MTS/IEEE Conference and Exhibition, Inst. of Electr. and Electron. Eng., Providence, 11–14 Sept.
- Mayes, C. L., L. A. Lawver, and D. T. Sandwell (1990), Tectonic history and new isochron chart of the South Pacific, *J. Geophys. Res.*, 95(B6), 8543–8567.
- National Geophysical Data Center (1996), Marine Geophysical trackline data (GEODAS/TRACKDAS), Boulder, Colo.
- Nitsche, F. O., K. Gohl, K. Vanneste, and H. Miller (1997), Seismic expression of glacially deposited sequences in the Bellingshausen and Amundsen seas, West Antarctica, in *Geology and Seismic Stratigraphy of the Antarctic Margin, Part 2, Antarct. Res. Ser.*, vol. 71, edited by P. F. Barker and A. K. Cooper, pp. 95–108, AGU, Washington, D. C.
- Nitsche, F. O., A. P. Cunningham, R. D. Larter, and K. Gohl (2000), Geometry and development of glacial continental margin depositional systems in the Bellingshausen Sea, *Mar. Geol.*, 162, 277–302.
- Ó Cofaigh, C., C. J. Pudsey, J. A. Dowdeswell, and P. Morris (2002), Evolution of subglacial bedforms along a paleo-ice stream, Antarctic Peninsula continental shelf, *Geophys. Res. Lett.*, 29(8), 1199, doi:10.1029/2001GL014488.
- Ó Cofaigh, C., R. D. Larter, J. A. Dowdeswell, C.-D. Hillenbrand, C. J. Pudsey, J. Evans, and P. Morris (2005), Flow of the West Antarctic Ice Sheet on the continental margin of the Bellingshausen Sea at the Last Glacial Maximum, *J. Geophys. Res.*, 110, B11103, doi:10.1029/2005JB003619.
- Ottesen, D., L. Rise, J. Knies, L. Olsen, and S. Henriksen (2005), The Vestfjorden-Trænadupet palaeo-ice stream drainage system, mid-Norwegian continental shelf, *Mar. Geol.*, 218, 175–189.
- Philippon, G., G. Ramstein, S. Charbit, M. Kageyama, C. Ritz, and C. Dumas (2006), Evolution of the Antarctic ice sheet throughout the last deglaciation: A study with a new coupled climate-north and south hemisphere ice sheet model, *Earth Planet. Sci. Lett.*, 248, 750–758.
- Pudsey, C. J., P. F. Barker, and R. D. Larter (1994), Ice sheet retreat from the Antarctic Peninsula shelf, *Cont. Shelf Res.*, 14, 1647–1675.
- Rignot, E., and S. S. Jacobs (2002), Rapid bottom melting widespread near Antarctic Ice Sheet grounding lines, *Science*, 296, 2020–2023.
- Rignot, E., and R. H. Thomas (2002), Mass balance of polar ice sheets, *Science*, 297, 1502–1506.
- Rignot, E. J. (1998), Fast recession of a West Antarctic glacier, *Science*, 281, 549–551.
- Sambridge, M., J. Braun, and H. McQueen (1995), Geophysical parameterization and interpolation of irregular data using natural neighbors, *Geophys. J. Int.*, 122, 837–857.
- Scheuer, C., K. Gohl, and G. Eagles (2006), Gridded isopach maps from the South Pacific and their use in interpreting the sedimentation history of the West Antarctic continental margin, *Geochem. Geophys. Geosyst.*, 7, Q11015, doi:10.1029/2006GC001315.
- Shaw, J., D. J. W. Piper, G. B. J. Fader, E. L. King, B. J. Todd, T. Bell, M. J. Batterson, and D. G. E. Liverman (2006), A conceptual model of the deglaciation of Atlantic Canada, *Quat. Sci. Rev.*, 25, 2059–2081.
- Shepherd, A., and D. Wingham (2007), Recent Sea-Level Contributions of the Antarctic and Greenland Ice Sheets, *Science*, 315, 1529–1532.
- Shepherd, A., D. Wingham, and E. Rignot (2004), Warm ocean is eroding West Antarctic Ice Sheet, *Geophys. Res. Lett.*, 31, L23402, doi:10.1029/2004GL021106.
- Sibson, R. (1981), A brief description of natural neighbor interpolation, in *Interpreting Multivariate Data*, edited by V. Barnett, pp. 21–36, John Wiley, New York.
- Smith, W. H. F., and D. T. Sandwell (1994), Bathymetric prediction from dense satellite altimetry and sparse shipboard bathymetry, *J. Geophys. Res.*, 99(B11), 21,803–21,824.
- Smith, W. H. F., and D. T. Sandwell (1997), Global seafloor topography from satellite altimetry and ship depth soundings, *Science*, 277, 1956–1961.
- SPRITE Group (1992), The Southern rim of the Pacific Ocean: Preliminary report of the Amundsen Sea-Bellingshausen Sea cruise of the Polar Sea, *Antarct. J. U.S.*, 27(1), 11–14.

- Stock, J., and P. Molnar (1987), Revised history of early Tertiary plate motion in the south-west Pacific, *Nature*, *325*, 495–499.
- Stokes, C. R., and C. D. Clark (2001), Palaeo-ice streams, *Quat. Sci. Rev.*, *20*, 1437–1457.
- Swithinbank, C., R. S. Williams Jr., J. G. Ferrigno, K. M. Foley, C. A. Hallam, and C. E. Rosanova (2003), Coastal-change and glaciological map of the Bakutis coast area, Antarctica: 1972–2002, in *Coastal-Change and Glaciological Maps of Antarctica*, 1 sheet, 1:1,000,000, *U.S. Geol. Surv. Geol. Invest. Ser., Map I-2600–F*.
- Swithinbank, C., R. S. Williams Jr., J. G. Ferrigno, K. M. Foley, C. E. Rosanova, and L. M. Dallide (2004), Coastal-change and glaciological map of the Eights coast area, Antarctica: 1972–2001, in *Coastal-Change and Glaciological Maps of Antarctica*, 1 sheet, 1:1,000,000, *U.S. Geol. Surv. Geol. Invest. Ser., Map I-2600–E*.
- Thomas, R., et al. (2004), Accelerated sea-level rise from West Antarctica, *Science*, *306*, 255–258.
- Thomson, J. W., and A. P. R. Cooper (1993), The Scar Antarctic Digital Topographic Database—Review, *Antarct. Sci.*, *5*(3), 239–244.
- Tucholke, B. E., and R. E. Houtz (1976), Sedimentary framework of the Bellingshausen basin from seismic profiler data, *Initial Rep. Deep Sea Drill. Proj.*, *35*, 197–227.
- Uenzelmann-Neben, G., K. Gohl, R. D. Larter, and P. Schlüter (2007), Differences in ice retreat across Pine Island Bay, West Antarctica, since the Last Glacial Maximum: Indications from multichannel seismic reflection data, in *Antarctica: A Keystone in a Changing World—Proceedings for the Tenth International Symposium on Antarctic Earth Sciences*, edited by A. K. Cooper, C. R. Raymond, and the ISAES Editorial Team, *U.S. Geol. Surv. Open File Rep., 2007-1047, Short Res. Pap. 084*, doi:10.3133/of2007-1047.srp084, 4 pp.
- Vaughan, D. G., H. F. J. Corr, F. Ferraccioli, N. Frearson, A. O'Hare, D. Mach, J. W. Holt, D. D. Blankenship, D. L. Morse, and D. A. Young (2006), New boundary conditions for the West Antarctic ice sheet: Subglacial topography beneath Pine Island Glacier, *Geophys. Res. Lett.*, *33*, L09501, doi:10.1029/2005GL025588.
- Walker, D. P., M. A. Brandon, A. Jenkins, J. T. Allen, J. A. Dowdeswell, and J. Evans (2007), Oceanic heat transport onto the Amundsen Sea shelf through a submarine glacial trough, *Geophys. Res. Lett.*, *34*, L02602, doi:10.1029/2006GL028154.
- Wellner, J. S., A. L. Lowe, S. S. Shipp, and J. B. Anderson (2001), Distribution of glacial geomorphic features on the Antarctic continental shelf and correlation with substrate: Implications for ice behavior, *J. Glaciol.*, *47*, 397–411.
- Wingham, D. J., A. J. Ridout, R. Scharroo, R. J. Arthern, and C. K. Shum (1998), Antarctic elevation change from 1992 to 1996, *Science*, *282*, 456–458.
- Yamaguchi, K., Y. Tamura, I. Mizukosho, and T. Tsuru (1988), Preliminary report of geophysical and geological surveys in the Amundsen Sea, West Antarctica, *Proc. NIRP Symp. Antarct. Geosci.*, *2*, 55–67.