



# 1 Meteorological observations from ship cruises during summer 2 to the central Arctic: A comparison with reanalysis data

3 C. Lüpkes,<sup>1</sup> T. Vihma,<sup>2</sup> E. Jakobson,<sup>3</sup> G. König-Langlo,<sup>1</sup> and A. Tetzlaff<sup>1</sup>

4 Received 29 January 2010; revised 31 March 2010; accepted 13 April 2010; published XX Month 2010.

5 [1] Near-surface meteorological observations and rawinsonde  
6 soundings from Arctic cruises with the German icebreaker RV  
7 Polarstern during August 1996, 2001, and 2007 are compared  
8 with each other and with ERA-Interim reanalyses. Although  
9 the observations are usually applied in the reanalysis, they  
10 differ considerably from ERA data. ERA overestimates the  
11 relative humidity and temperature in the atmospheric  
12 boundary layer and the base height of the capping inversion.  
13 Warm biases of ERA near-surface temperatures amount up  
14 to 2 K. The melting point of snow is the most frequent near-  
15 surface temperature in ERA, while the observed value is the  
16 sea water freezing temperature. Both observations and ERA  
17 show that above 400 m, in the North Atlantic sector 0–90 E,  
18 the warmest August occurred in 2001, and August 2007 had  
19 the highest humidity. In the Eastern Siberian and Beaufort  
20 Sea region ERA temperatures along 80 and 85 N were  
21 highest in 2007. **Citation:** Lüpkes, C., T. Vihma, E. Jakobson,  
22 G. König-Langlo, and A. Tetzlaff (2010), Meteorological observa-  
23 tions from ship cruises during summer to the central Arctic: A compar-  
24 ison with reanalysis data, *Geophys. Res. Lett.*, 37, LXXXXX,  
25 doi:10.1029/2010GL042724.

## 26 1. Introduction

27 [2] Projections of climate models for the 21st century  
28 show an especially pronounced warming in high latitudes  
29 usually called the arctic amplification [e.g., *Serreze et al.*,  
30 2009], and the presently available data already indicate a  
31 strong warming of the Arctic in the last decade. Arctic in-  
32 situ data are, however, rare, since longer time series of ob-  
33 servations result from a few coastal stations only, and in the  
34 inner Arctic mainly from buoys and rarely from drifting  
35 stations or from ship cruises. Especially, time series from  
36 Central Arctic in-situ observations above the surface layer  
37 are rare. They are available only from rawinsonde and te-  
38 thersonde soundings performed at drifting stations [e.g.,  
39 *Serreze et al.*, 1992; *Vihma et al.*, 2008] and during ship  
40 cruises. The analysis of climate trends on the basis of these  
41 data is a challenging task, because drifting stations and ship  
42 trajectories of different years are often separated from each  
43 other by hundreds of kilometers.

44 [3] In this work we will investigate routine meteorological  
45 observations and soundings from the German icebreaker RV  
46 Polarstern from three different years and consider as the first  
47 goal the differences between these years and to what extent

such data can contribute to the analysis of climate change, 55  
while paying attention to the spatial representativeness of the 56  
observations. We concentrate on the summer expeditions 57  
1996, 2001, and 2007 to the Central Arctic (Figure 1). The 58  
focus on these years is especially interesting, since the sea ice 59  
conditions differed strongly from each other and the identi- 60  
fication of differences in the meteorological conditions might 61  
help to better understand the reasons for the recent sea ice 62  
retreat. According to NSIDC (<http://nsidc.org/>) a large 63  
mean August sea ice extent was observed in 2001 (7.5 64  
mill. km<sup>2</sup>) and 1996 (8.2 mill. km<sup>2</sup>), but, as well known, in 65  
2007 it was extremely low (5.4 mill. km<sup>2</sup>) resulting finally in 66  
the historical September minimum. Also the sea ice surface 67  
characteristics differed considerably in these years with only 68  
very few or, north of 84°N, even no melt ponds along the 69  
cruise track in 1996 [*Augstein et al.*, 1997; *Haas and Eicken*, 70  
2001] and a large melt pond coverage in 2001 [*Thiede*, 2002] 71  
and 2007 [*Schauer*, 2008]. 72

[4] The spatial representativeness of the ship data will be 73  
investigated with the help of reanalysis data. However, since 74  
the accuracy of reanalysis in the Central Arctic regions is 75  
not well known, we concentrate also on the performance of 76  
the reanalysis by comparing it with the measurements along 77  
the ship tracks. For this task, ERA-Interim data are used 78  
representing the newest set of reanalysis data from the 79  
European Centre for Medium-Range Weather Forecasts 80  
(ECMWF). Although the routine meteorological Polarstern 81  
observations are always transmitted to the Global Tele- 82  
communication System (GTS) of the WMO and thus con- 83  
tribute to ERA, it cannot be expected that observations are 84  
perfectly reproduced by the reanalysis, and our study will 85  
help to identify the shortcomings of ERA-Interim over sea 86  
ice covered regions. 87

## 2. Data Used 88


[5] As in-situ data we use the routinely observed near- 89  
surface temperature from Polarstern and data from raw- 90  
insondes launched from the ship during the summer cruises 91  
ARK-XII (1996), ARK-XVII/2 (2001), and ARK-XXII/2 92  
(2007), whose meteorological data are compiled by *König- 93  
Langlo* [2005, 2008]. We concentrate on the August data 94  
and in case of soundings on data north of 80°N, since there 95  
the ship tracks (Figure 1) were always north of the ice edge 96  
and the overlapping times of the cruises were largest. The 97  
routine temperature measurements from Polarstern consid- 98  
ered here in a 10-minute resolution were carried out at 30 m 99  
height. 100

[6] The Polarstern temperature sensors (PT-100) are well 101  
protected against radiation and mounted at a well ventilated 102  
position at 30 m height [*König-Langlo et al.*, 2006]. They 103  
are calibrated during each cruise using an aspiration 104

<sup>1</sup>Alfred Wegener Institute for Polar and Marine Research, Bremerhaven, Germany.

<sup>2</sup>Finnish Meteorological Institute, Helsinki, Finland.

<sup>3</sup>Tartu Observatory, Toravere, Estonia.



**Figure 1.** Cruise tracks of RV Polarstern during August 1996 (blue), 2001 (green) and 2007 (red).

105 psychrometer (Assmann) as reference. Rawinsondes (Vaisala)  
 106 were usually launched twice a day, in 1996 at 02.30 and  
 107 10.30 UTC, in 2001 at 05.00 and 10.00 UTC, and in  
 108 2007 at 06.00 and 11.00 UTC. According to the above  
 109 position criteria we consider the soundings between 7 and  
 110 31 August of 2007 (44 soundings), between 4 and 31 August  
 111 (55 soundings) of 2001 and between 1 and 29 August of  
 112 1996 (58 soundings). All data are publicly available via  
 113 <http://www.pangaea.de/>.

114 [7] We compare the Polarstern near-surface data with  
 115 ERA-Interim data from the second-lowest model level  
 116 (about 38 m height) at the ship positions in 6-hour resolu-  
 117 tion. All reanalysis data represent means over  $0.72 \times 0.72^\circ$   
 118 in longitude and latitude. Furthermore, vertical profiles of  
 119 meteorological variables at the sounding locations are used,  
 120 interpolated to the above-mentioned ship times. This inter-  
 121 polation caused only slight changes in the monthly averages  
 122 considered in section 3.3. Sounding data are not distributed  
 123 regularly in height. Hence, before averaging, the data were  
 124 interpolated to a regular grid with 40 m vertical resolution.

### 125 3. Results

#### 126 3.1. August Conditions Along 80° and 85°N

127 [8] A hint on the spatial representativeness of temperature  
 128 and humidity measured in the considered years at an arbi-  
 129 trary position in the Arctic can be obtained from Figure 2  
 130 showing ERA-Interim results along 80° and 85°N.

131 [9] According to these results 2007 had the warmest and  
 132 most humid August along 85°N. Differences to other years  
 133 were especially large east of 150°E, outside of the sector  
 134 studied by the ship. In 1996 and 2001 temperatures and  
 135 specific humidities decreased slightly towards the east,  
 136 which was in contrast to 2007 with an almost constant near-  
 137 surface temperature and humidity along 85°N.

138 [10] A large difference exists between the sector 0–90°E  
 139 (North Atlantic sector, NA) and the remaining part. In the  
 140 first one, 2001 was by far the warmest year at 80°N, espe-  
 141 cially at the height of the 850 hPa pressure level, where  
 142 along 80°N the NA sector was in August 2001 up to 3.5 K  
 143 warmer than in both other years. Everywhere else August  
 144 2007 had the highest 850 hPa temperatures with nearly 6 K  
 145 difference to 1996 and 2001 at 80°N and 180°E. This was  
 146 similar for 85°N, except that 2001 was not that much

warmer in the NA sector and the warm 2007 anomaly was  
 most pronounced at 210–240°E.


[11] At 80°N the near-surface temperature and humidity  
 strongly decrease in all years towards east in the region at  
 about 270°E. This is due to the influence of the Greenland  
 ice sheet region, which we do not consider in the present  
 analysis.

[12] The above findings show that a comparison of data  
 obtained from the three cruises should be considered with  
 caution, since possible differences between ship observa-  
 tions from different years can result from different ship lo-  
 cations. On the other hand, it is important to note that all  
 ship tracks were in the sector 0–150°E and more close to the  
 85°N latitude, where the near-surface horizontal gradients  
 along the latitude were found to be relatively small.


#### 3.2. Near-Surface Data Along Ship Tracks

[13] Figure 3 shows the observed probability distributions  
 of the 30 m real air temperature as well as the distributions  
 obtained from ERA at the second-lowest model level  
 ( $\approx 38$  m height) along the ship tracks. Based on these  
 distributions the following two conclusions are possible.

[14] The first refers to an intercomparison of the data from  
 the different years. Obviously, the 1996 temperature distri-  
 bution has a larger width and shows lower temperatures than  
 in both other years. This holds for both observations and  
 ERA-Interim giving confidence to the ERA-based result  
 that 1996 was colder almost everywhere in the circumpolar  
 Arctic at 80° and 85°N (Figure 2). It may partly explain, why  
 the amount of observed open melt ponds was so small in  
 1996 compared with the other years.



**Figure 2.** August averages, as calculated from ERA Interim results, for the real air temperature and specific humidity in steps of 30 degrees longitude along 80°N and 85°N for 850 hPa and 38 m height. Polarstern operated mainly in the region left of the vertical dashed line. Values of 38 m at 270–330°E and 80°N are low, since locations are over mountains.



**Figure 3.** Probability density functions, mean values  $m$  and standard deviation  $\sigma$  of the 30 m real air temperature observed by Polarstern (red) and calculated from ERA-Interim along the ship tracks at  $\approx 38$  m height. The values are calculated for 0.5 K bins.

177 [15] The second conclusion is related to the large quali-  
 178 tative and quantitative differences between the distributions  
 179 based on ship and ERA data. The most frequently observed  
 180 temperature is in all years the freezing point of sea water  
 181 ( $\approx -1.8^\circ\text{C}$ ), whereas in the reanalysis the peak occurs  
 182 approximately at the melting point of snow ( $0^\circ\text{C}$ ). The latter  
 183 temperature appears in 1996 and 2007 also in the observed  
 184 distributions as a secondary peak. A related difference  
 185 between observation and ERA concerns the warm bias of  
 186 ERA in all the three August distributions, most pronounced  
 187 in 2007.

### 188 3.3. Soundings

189 [16] Profiles of temperature and humidity in Figure 4 are  
 190 approximately August averages (as explained in Section 2)  
 191 along the cruise tracks. Four results can be derived:

192 [17] 1. In heights above 400 m both the 2001 observations  
 193 and ERA data show clearly the highest temperatures  
 194 (Figure 4, top). Differences at 1200 m amount to roughly  
 195 3.5 K, while near the surface only small differences can be  
 196 found between 2001 and 2007, and slightly larger differences  
 197 between 2007 and 1996.

198 [18] 2. August 2007 differs from the other years [18] mainly by  
 199 its large humidity values, especially when compared with  
 200 2001.


201 [19] 3. Obviously, there is a large bias of the ERA results  
 202 in the height range below 800 m. This concerns both tem-  
 203 perature and specific humidity. The bias in the latter data is  
 204 especially pronounced in 2001 and 1996, where ERA  
 205 humidity is close to saturation in heights below 400 m, but  
 206 observed values are much lower.

207 [20] 4. ERA considerably overestimates the base height of  
 208 the boundary layer capping inversion, which can be seen in  
 209 the monthly averages, but it is especially obvious comparing  
 210 individual profiles at selected times (not shown). Differ-  
 211 ences were most pronounced in cases with observed surface  
 212 based inversions, while the ERA-Interim showed mostly  
 213 elevated inversions.

[21] We also found that ERA wind and observed wind 214  
 agreed well with almost identical mean values near the 215  
 surface and increasing differences in higher levels, which 216  
 were, however, smaller than  $1 \text{ ms}^{-1}$  and not significant on 217  
 the 95% confidence level. 218

## 4. Summary and Conclusions

[22] Both ERA Interim results and Polarstern observations 220  
 reveal large and consistent differences in the mean August 221  
 temperatures and humidities of 1996, 2001, and 2007. 222  
 Along 80 and 85°N and east of 90°E the highest August 223  
 temperatures occurred in 2007. However, in the North 224  
 Atlantic sector (0–90°E), where a large part of the Polarstern 225  
 cruises in 2007 and 2001 was carried out, the 2007 August 226  
 near-surface temperature close to 80°N was similar as in 227  
 2001. But further north 2001 was by far the warmest year 228  
 and the upper-level temperatures were up to 3.5 K higher 229  
 than in other years. In August 2007 the near-surface tem- 230  
 perature along 80 and 85°N peaked in the East Siberian 231  
 Arctic. *Serreze et al.* [2009] and *Overland and Wang* [2010] 232  
 suggest that the recent temperature increase in autumn is a 233  
 combined effect of sea ice loss and atmospheric circulation. 234  
 The latter is probably the main reason for the high upper- 235  
 level temperatures in the North Atlantic sector, since there 236  
 the 2001 sea ice cover was similar as in 1996 [see also 237  
*Graversen et al.*, 2008]. In 2007, however, also the exces- 238  
 sive sea ice loss may have affected air temperatures already 239  
 in August. The ship data show that in the second half of 240



**Figure 4.** Mean August profiles (dashed, rawinsondes; solid, ERA-Interim at the sounding positions; horizontal bars are the 95% confidence intervals). Figure 4 (top) shows the same profiles as below to allow a comparison between years.



241 August the air temperature often dropped well below  
 242  $-1.8^{\circ}\text{C}$ , i.e. an upward heat flux could develop over open  
 243 leads preventing the atmosphere from a stronger cooling.  
 244 ERA Interim near-surface data show a warm bias of 1.5–  
 245 2 K, although the Polarstern data were assimilated into  
 246 the reanalysis. This confirms similar findings of *Liu et al.*  
 247 [2005] on the basis of SHEBA data and demonstrates a  
 248 strong need to improve methods of near-surface data  
 249 assimilation for ERA. The differences in the most fre-  
 250 quent values of surface temperatures ( $-1.8^{\circ}\text{C}$  measured  
 251 and  $0^{\circ}\text{C}$  in ERA) might hint at a non-adequate treatment  
 252 of thermodynamic effects of open leads, whose surface  
 253 temperature is usually at the freezing point of sea water.  
 254 In ERA the sea ice concentration north of  $84^{\circ}\text{N}$  is always  
 255 100%, which is unrealistic, particularly for 2007. The  
 256 comparison of the whole time series (not shown here) of  
 257 ERA and ship data revealed that a similar warm bias of  
 258 ERA occurred also between  $-2$  and  $-10^{\circ}\text{C}$  air tempera-  
 259 ture so that also an overestimation by ERA of warm air  
 260 advection, cloud cover or turbulent mixing could explain  
 261 the differences. A possible measurement error can be  
 262 excluded due to the often calibrated sensors. Solar radi-  
 263 ation and heating by the ship could only contribute to a  
 264 warm bias of the observations.  
 265 [23] Large differences between ERA and observations  
 266 occur also in the boundary layer. ERA overestimates the  
 267 base height of the capping inversion sometimes by more  
 268 than a factor of two and the stratification is biased towards  
 269 neutral values. These biases may be attributed to too  
 270 excessive turbulent mixing. Relative humidity is over-  
 271 estimated by ERA in the boundary layer especially in 2001  
 272 by about 15% in the August mean value. This finding earns  
 273 much attention, since clouds play a critical role for varia-  
 274 tions in the Arctic sea ice cover [*Francis and Hunter, 2006*].  
 275 Note that errors in the cloud cover would cause a long chain  
 276 of other drawbacks in the modeling of meteorological  
 277 parameters.  
 278 [24] Finally, we stress the need for a regular validation of  
 279 reanalysis data against independent observations, since our  
 280 present knowledge of arctic climate change strongly relies  
 281 on high quality reanalyses. So the present comparison  
 282 should also be carried out for other reanalyses than for ERA.  
 283 Vice versa, we have shown that to some extent, data from  
 284 ship cruises in different years can be used to identify dif-  
 285 ferences in the climatic conditions. However, to account for  
 286 spatial variations, this should always be done in combina-

tion with reanalysis data keeping in mind, however, their 287  
 uncertainties. 288

[25] **Acknowledgments.** This study was supported by the EU project 289  
 DAMOCLES (grant 18509), which is part of the Sixth Framework 290  
 Programme. The ECMWF is acknowledged for providing us with the 291  
 ERA-Interim data. 292

## References

- Augstein, E., et al. (1997), The expedition ARCTIC '96 of RV "Polarstern" 294  
 (ARK XII) with the Arctic Climate System Study (ACSYS), *Rep. on* 295  
*Polar Res.* 234, 54 pp., Alfred Wegener Inst., Bremerhaven, Germany. 296  
 Francis, J. A., and E. Hunter (2006), New insight into the disappearing 297  
 Arctic sea ice, *Eos Trans. AGU*, 87(46), doi:10.1029/2006EO460001. 298  
 Graverson, R. G., T. Mauritsen, M. Tjernström, E. Källen, and G. Svensson 299  
 (2008), Vertical structure of recent Arctic warming, *Nature*, 451, 53–56. 300  
 Haas, C., and H. Eicken (2001), Interannual variability of summer sea ice 301  
 thickness in the Siberian and central Arctic under different atmospheric 302  
 circulation regimes, *J. Geophys. Res.*, 106, 4449–4462. 303  
 König-Langlo, G. (2005), Meteorological observations during Polarstern 304  
 cruises ARK-XII; ARK-XVII/2, Alfred Wegener Inst., Bremerhaven, 305  
 Germany. 306  
 König-Langlo, G. (2008), Meteorological observations during Polarstern 307  
 cruise ARK-XXII/2, Alfred Wegener Inst., Bremerhaven, Germany. 308  
 König-Langlo, G., B. Loose, and B. Bräuer (2006), 25 years of Polarstern 309  
 meteorology, *WDC-MARE Rep. 4*, World Data Cent. for Mar. Environ. 310  
 Sci., Bremerhaven, Germany. 311  
 Liu, J., J. A. Curry, W. B. Rossow, J. R. Key, and X. Wang (2005), 312  
 Comparison of surface radiative flux data sets over the Arctic Ocean, 313  
*J. Geophys. Res.*, 110, C02015, doi:10.1029/2004JC002381. 314  
 Overland, J., and M. Wang (2010), Large scale atmospheric circulation 315  
 changes are associated with the recent loss of Arctic sea ice, *Tellus*, 316  
*Ser. A*, 62, 1–9. 317  
 Schauer, U. (2008), The expedition ARKTIS-XXII/2 of the research vessel 318  
 "Polarstern" in 2007, *Rep. Polar Mar. Res.* 579, 271 pp., Alfred 319  
 Wegener Inst., Bremerhaven, Germany. 320  
 Serreze, M. C., J. D. Kahl, and R. C. Schnell (1992), Low-level tempera- 321  
 ture inversions of the Eurasian Arctic and comparisons with Soviet 322  
 drifting station data, *J. Clim.*, 5, 615–629. 323  
 Serreze, M. C., A. P. Barrett, D. N. Stroeve, and M. M. Holland (2009), The 324  
 emergence of surface-based Arctic amplification, *Cryosphere*, 3, 11–19. 325  
 Thiede, J. (2002), Polarstern ARKTIS XVII/2, cruise report: AMORE 2001 326  
 (Arctic Mid-Ocean Ridge Expedition), *Rep. Polar Mar. Res.* 421, 327  
 390 pp., Alfred Wegener Inst., Bremerhaven, Germany. 328  
 Vihma, T., J. Jaagus, E. Jakobson, and T. Palo (2008), Meteorological 329  
 conditions in the Arctic Ocean in spring and summer 2007 as recorded 330  
 on the drifting ice station Tara, *Geophys. Res. Lett.*, 35, L18706, 331  
 doi:10.1029/2008GL034681. 332  
 E. Jakobson, Tartu Observatory, 61602 Toravere, Tartumaa, Estonia. 333  
 G. König-Langlo, C. Lüpkes, and A. Tetzlaff, Alfred Wegener Institute 334  
 for Polar and Marine Research, Postfach 120161, D-27515 Bremerhaven, 335  
 Germany. (christof.luepk@awi.de) 336  
 T. Vihma, Finnish Meteorological Institute, PO Box 506, FIN-00101 337  
 Helsinki, Finland. 338