

Mathematical Journal of Okayama University

Volume 16, Issue 1

1973

Article 10

SEPTEMBER 1973

On torsion free modules over regular rings

Kiyoichi Oshiro*

*Yamaguchi University

Copyright ©1973 by the authors. *Mathematical Journal of Okayama University* is produced by
The Berkeley Electronic Press (bepress). <http://escholarship.lib.okayama-u.ac.jp/mjou>

ON TORSION FREE MODULES OVER REGULAR RINGS

KIYOICHI OSHIRO

Throughout this paper, all rings have identity and all modules are unital.

Let M be a left R -module. We denote its torsion submodule in the sense of Gentile [2] by $T(M)$, i. e.,

$$T(M) = \{x \in M \mid \text{Hom}_R(Rx, I(R)) = 0\}$$

where $I(R)$ is the injective hull of R as a left R -module.

A left R -module M is said to be torsion free if $T(M) = 0$. For the properties of torsion free modules, the reader is referred to [6].

In section 2, we show that a regular ring R is left continuous if and only if every cyclic torsion free R -module is projective.

In section 3, we obtain module theoretic characterizations of a commutative regular ring R whose maximal ring of quotients coincides with the Baer hull of R . In connection with this, Examples A and C are given which answer Pierce's questions (4) and (11) of [6] in the negative, respectively.

The author wishes to thank Prof. H. Tachikawa for his helpful suggestions and also Prof. Y. Kurata.

1. Preliminaries

Let Q be a ring of left quotients of R . It is well known that a torsion free injective left R -module can be turned into a left Q -module (see [8]). So, if M is a torsion free left R -module, M is embedded in the Q -submodule QM of the injective hull of M . Note that for any two elements $\sum_{i=1}^n p_i a_i$ and $\sum_{j=1}^m q_j b_j$ in QM , $\sum_{i=1}^n p_i a_i = \sum_{j=1}^m q_j b_j$ if and only if, for any $r \in \bigcap_{i=1}^n (R : p_i)^{1)}$ and $0 \neq r' \in R$, there exists $r'' \in \bigcap_{j=1}^m (R : r q_j)$ such that $r'' r' \neq 0$ and $\sum_{i=1}^n (r'' r p_i) a_i = \sum_{j=1}^m (r'' r q_j) b_j$.

In latter section 3, we need the following lemma, the proof of which is straightforward, and will be omitted.

¹⁾ $(R : p) = \{r \in R \mid r p \in R\}$.

Lemma 1.1. *Let M be a torsion free left R -module and let $\{M_i\}$ a collection of R -submodules of M indexed by A . If $M = \sum_{i \in A} \oplus M_i$, then $QM = \sum_{i \in A} \oplus QM_i$.*

2. Continuous regular rings

We recall that a regular ring R is said to be left continuous if the lattice $L(R)$ of principal left ideals of R is continuous (see [9]).

Theorem 2.1. *Let R be a regular ring and Q its maximal ring of left quotients. Then the following conditions are equivalent :*

- (a) *R is left continuous.*
- (b) *Every idempotent in Q is contained in R .*
- (c) *$A = Q(A \cap R)$ for any left ideal A of Q .*
- (d) *Every principal left ideal of Q is generated by an idempotent of R .*
- (e) *Every cyclic torsion free left R -module is projective.*

Proof. (a) \Rightarrow (b) By [10, Lemma 8].

(b) \Rightarrow (c) Since Q is a regular ring, this is evident.

(c) \Rightarrow (d) Let Qx be a principal left ideal of Q . Since $Qx = Q(Qx \cap R)$, we have $x = \sum_{i=1}^n q_i r_i$ for some $q_1, q_2, \dots, q_n \in Q$ and $r_1, r_2, \dots, r_n \in Qx \cap R$. Since R is a regular ring, there exists an idempotent e in $Qx \cap R$ such that $\sum_{i=1}^n Rr_i = Re$. Then we have $Qx = Qe$.

(d) \Rightarrow (a) By [9, Theorem 2.1], Q is left continuous. It is easily seen from (d) that $L(Q)$ is lattice isomorphic to $L(R)$. Hence R is left continuous.

(e) \Rightarrow (b) Let e be an idempotent of Q . Then, $R + Re$ is projective, since $R + Re = Re \oplus R(1 - e)$. Therefore, by [3, Lemma 4], R is a direct summand of $R + Re$. However, since R is an essential submodule of $R + Re$, this implies that $Re + R = R$ and $e \in R$.

(c) \Rightarrow (e) Let Rx be a cyclic torsion free left R -module. By [1, Theorem 2.1], Qx is Q -projective. Here we claim that $Q \otimes_R Rx \simeq Qx$ canonically. To this end, we consider the exact sequence

$$0 \longrightarrow \text{Ann}_R(x) \longrightarrow R \longrightarrow Rx \longrightarrow 0$$

where $\text{Ann}_R(x) = \{r \in R \mid rx = 0\}$. Since Q is flat as a right R -module, the induced sequence of left Q -modules

$$0 \longrightarrow Q\text{Ann}_R(x) \longrightarrow Q \longrightarrow Q \otimes_R Rx \longrightarrow 0$$

is exact. Now to show that $Q \otimes_R Rx \simeq Qx$, it is sufficient to show that $Q \text{Ann}_R(x) = \text{Ann}_Q(x)$. Clearly $Q \text{Ann}_R(x) \subseteq \text{Ann}_Q(x)$. Conversely if $q \in Q$ such that $qx=0$, then it is easily seen from $Q(Qq \cap R) = Qq$ that $q \in \text{Ann}_Q(x)$. Hence we have $Q \text{Ann}_R(x) = \text{Ann}_Q(x)$ and $Q \otimes_R Rx \simeq Qx$. So, $Q \otimes_R Rx$ is Q -projective and therefore, by [7, Theorem 2.8], Rx is R -projective.

3. Baer ring of quotients

Throughout this section, we assume that a ring R is commutative.

Let M be a torsion free R -module. In the following we shall consider the following conditions :

- (α) M is a direct sum of cyclic R -modules.
- (β) M is isomorphic to an essential submodule of a direct sum of cyclic torsion free R -modules.
- (γ) M is isomorphic to a submodule of a direct sum of cyclic torsion free R -modules.

Remark 1. The following question has been asked by Pierce [6, p. 109]: “Let M be an R -module (where R is a regular ring) which is a finite direct sum of cyclic R -modules. Let N be a finitely generated R -submodule of M . Is N a direct sum of cyclic R -modules?” However there exists an example of a module over a Boolean ring which satisfies the condition (β) but not the condition (α). Therefore Pierce’s question has a negative answer.

For constructing such an example, we need a lemma.

Lemma 3.1. *Let R be a Boolean ring, and Q its maximal ring of quotients. For e and f in Q , the following conditions are equivalent :*

- (a) $Re + Rf$ satisfies the condition (α).
- (b) ef is contained in $Re + Rf$.

Proof. (a) \Rightarrow (b) By the assumption, $Re + Rf$ is decomposed into a direct sum of cyclic R -submodules, say $Re + Rf = Rg_1 \oplus Rg_2 \oplus \dots \oplus Rg_n$. Then, by Lemma 1.1, $Qg_1 + Qg_2 + \dots + Qg_n = Qg_1 \oplus Qg_2 \oplus \dots \oplus Qg_n$, i. e., $\{g_i \mid i = 1, 2, \dots, n\}$ is a set of orthogonal elements. Hence, when we view Q as a partially ordered set, $g_1 + g_2 + \dots + g_n$ is the supremum of $Rg_1 + Rg_2 + \dots + Rg_n$. On the other hand, the supremum of $Re + Rf$ is $e + f - ef$. Hence we have $e + f - ef = g_1 + g_2 + \dots + g_n$ and $ef \in Re + Rf$.

(b) \Rightarrow (a) If $ef \in Re + Rf$, then, clearly, $Re + Rf = Re(f - 1) \oplus \dots$

$Rf(e - 1) \oplus Ref$.

Example A. Let S be an infinite set. Let Q be the set of all subsets of S , and R the set of all finite subsets and all cofinite subsets (i. e., complements of finite subsets) of S . Q become a Boolean ring by the following definition: for a, b in Q .

$$a + b = (a \cup b) \cap (a \cap b)^c$$

where $(a \cap b)^c$ denotes the complement of $a \cap b$ in S ,

$$ab = a \cap b.$$

Note that the empty set is the zero element and S is the identity. Moreover R is a subring of Q and its maximal ring of quotients coincides with Q (see [4, p. 45]).

Now we can select e and f in Q such that $e, f, e \cap f, e \cap (e \cap f)^c$ and $f \cap (e \cap f)^c$ are infinite sets. Put $e' = e \cap (e \cap f)^c$ and $f' = f \cap (e \cap f)^c$. Then $Re' + Rf' + Ref = Re' \oplus Rf' \oplus Ref$ and it contains $Re + Rf$ as an essential submodule. However $Re + Rf$ can not be decomposed into a direct sum of cyclic R -modules. For, if it is decomposed into a direct sum of cyclic R -modules, then, by Lemma 3.1, $ef = re + sf$ for some r, s in R . It is easily seen that $e' \subseteq r^c$ and $f' \subseteq s^c$. If both r and s are finite sets, then ef is a finite set. If r is a cofinite set, then e' is a finite set. Similarly if s is a cofinite set, then f' is a finite set. At any rate we have a contradiction. Therefore $Re + Rf$ can not be decomposed into a direct sum of cyclic R -modules.

Lemma 3.2. *Let R be a ring and Q its ring extension with the same identity. If $\{e_i \mid i = 1, 2, \dots, n\}$ is a set of idempotents of Q , then there exists a set $\{f_i \mid i = 1, 2, \dots, m\}$ of orthogonal idempotents in Q such that*

$$\sum_{i=1}^n Re_i \subseteq \sum_{i=1}^m Rf_i \text{ and } \sum_{i=1}^n Qe_i = \sum_{i=1}^m Qf_i.$$

Proof. We proceed by induction on the number n of elements of $\{e_i \mid i = 1, 2, \dots, n\}$.

$n = 1$. Obvious.

Assume $n > 1$ and the lemma is true on $n = k - 1$.

$n = k$. Let $\{e_i \mid i = 1, 2, \dots, k\}$ be a set of idempotents of Q . By the induction assumption, there exist orthogonal idempotents f_1, f_2, \dots, f_m such that

$$\sum_{i=1}^{k-1} Re_i \subseteq \sum_{i=1}^m Rf_i \text{ and } \sum_{i=1}^{k-1} Qe_i = \sum_{i=1}^m Qf_i.$$

Then we have

$$\begin{aligned} \sum_{i=1}^k Re_i &\subseteq \sum_{i=1}^m Rf_i + Re_k \\ &\subseteq Re_k(1 - \sum_{i=1}^m f_i) \oplus \sum_{i=1}^m Rf_i \oplus R(1 - e_k)f_i \oplus \sum_{i=1}^m Qe_k f_i \end{aligned}$$

and

$$\sum_{i=1}^k Qe_i = Qe_k(1 - \sum_{i=1}^m f_i) + \sum_{i=1}^m Q(1 - e_k)f_i + \sum_{i=1}^m Qe_k f_i.$$

Hence the lemma is true on $n=k$, and our proof is complete.

We recall that if R is a semi-prime ring, then R has a unique minimal Baer ring of quotients which is called by Mewborn the Baer hull of R . And it is known in [5, Proposition 2.5] that the Baer hull of R coincides with the ring generated by R and all idempotents of the maximal ring of quotients of R . Hence it follows from Theorem 2.1 that a commutative regular ring R is a Baer ring if and only if it is continuous.

Proposition 3.3. *If R is a regular ring, then its Baer hull is a continuous regular ring.*

Proof. It is easily seen from Lemma 3.2 that Baer hull of R is a regular ring. So, it is a continuous regular ring.

Lemma 3.4. *Let R be a regular ring and Q its maximal ring of quotients. If, for any x in Q , $R + Rx$ satisfies the condition (γ) , then Q coincides with the Baer hull of R .*

Proof. First we assume that R is a continuous regular ring and show that $Q = R$. Let x be in Q . Since $R + Rx$ satisfies the condition (γ) , by (e) of Theorem 2.1, $R + Rx$ is isomorphic to a submodule of a projective module. Hence, by [3, Lemma 4], R is a direct summand of $R + Rx$ and it follows that $R + Rx = R$ and $Q = R$.

Next let R be an arbitrary regular ring. Then, by Proposition 3.3, the Baer hull of R , say P , is a continuous regular ring. We claim that $Q = P$. If $Q \neq P$, then by the above observation, there exists x in Q such that $P + Px$ does not satisfy the condition (γ) . But this provides that $R + Rx$ does not satisfy the condition (γ) , a contradiction. Thus Q must coincide with P as desired.

Lemma 3.5. *Let R be a semi-prime ring and Q its maximal ring of quotients. Then every finitely generated torsion free R -module can be*

embedded in an external direct sum of finitely generated R -submodules of Q as an essential submodule.

Proof. See [11, Corollary 5].

Lemma 3.6. *Let R be a semi-prime ring, and Q its maximal ring of quotients. If Q coincides with the Baer hull of R , then every finitely generated torsion free R -module satisfies the condition (β).*

Proof. In view of Lemma 3.5, it is sufficient to show the statement for finitely generated R -submodules of Q .

First, for a cyclic R -submodule $Rq \subseteq Q$, we show that there exist idempotents e_i , $i = 1, 2, \dots, m$, in Q such that $\sum_{i=1}^m Re_i$ contains Rq as an essential submodule. Since Q coincides with the Baer hull of R , by Lemma 3.2, there exist r_i , $i = 1, 2, \dots, m$, in R and orthogonal idempotents g_i , $i = 1, 2, \dots, m$, in Q such that $q = \sum_{i=1}^m r_i g_i$. Then Rq is an essential submodule of $\sum_{i=1}^m Rr_i g_i$. Since Q is a regular ring, it is easily seen that we can find idempotents e_i , $i = 1, 2, \dots, m$, in Q such that $Rr_i g_i \subseteq Re_i$ and $Qr_i g_i = Qe_i$, $i = 1, 2, \dots, m$. Hence Rq is an essential submodule of $\sum_{i=1}^m Re_i$.

Now let $M = \sum_{i=1}^n Rq_i$ be any finitely generated R -submodule of Q . For each i , $i = 1, 2, \dots, n$, there exist idempotents e_{ij} , $j = 1, 2, \dots, n_i$, in Q such that $\sum_{j=1}^{n_i} Re_{ij}$ contains Rq_i as an essential submodule. Hence $\sum_{i=1}^n \sum_{j=1}^{n_i} Re_{ij}$ contains $\sum_{i=1}^n Rq_i$ as an essential submodule. On the other hand, by Lemma 3.2, there exist orthogonal idempotents f_i , $i = 1, 2, \dots, s$, in Q such that $\sum_{i=1}^s \oplus Rf_i$ contains $\sum_{i=1}^n \sum_{j=1}^{n_i} Re_{ij}$ as an essential submodule. Consequently $\sum_{i=1}^s \oplus Rf_i$ contains $\sum_{i=1}^n Rq_i$ as an essential submodule.

By Lemmas 3.4 and 3.6, we have

Theorem 3.7. *Let R be a commutative regular ring, and Q its maximal ring of quotients. Then the following conditions are equivalent:*

- (a) *Every finitely generated torsion free R -module satisfies the condition (β).*
- (b) *Every finitely generated torsion free R -module satisfies the condition (γ).*
- (c) *Q coincides with the Baer hull of R .*

Remark 2. (i) Reviewing our observation, the following conditions are also equivalent to the conditions in Theorem 3.7:

(a') For any x in Q , $R + Rx$ satisfies the condition (β).

(b') For any x in Q , $R + Rx$ satisfies the condition (γ).

However, by Example A, the following condition is not equivalent to the above conditions:

(d) Every finitely generated torsion free R -module satisfies the condition (α).

(ii) As is seen from the following example, in order that a regular ring satisfies the conditions of Theorem 3.7, it need not be self-injective or Boolean.

Example B. Let D be a finite field. Take an infinite index set A , and let

$$Q = \prod_{i \in A} D_i, \quad D_i = D \quad \text{for all } i \in A$$

and

$$R = \sum_{i \in A} \oplus D_i + 1 \cdot D,$$

where 1 is the identity of Q . Then R is a regular ring and its maximal ring of quotients is Q . Since D is a finite field, it is easily seen that Q coincides with the ring generated by R and all idempotents of Q , i. e., Q is the Baer hull of R .

The proof of the following lemma is easy.

Lemma 3.9. Let R be a ring and Q its maximal ring of quotients. For any idempotent e in Q , Re is R -injective if and only if Re is a self-injective ring.

By Lemmas 3.6 and 3.9, we have

Theorem 3.10. Let R be a commutative semi-prime ring such that its maximal ring of quotients coincides with its Baer hull. Then a torsion free R -module M is finitely generated and injective if and only if $M \simeq R/J_1 \oplus R/J_2 \oplus \cdots \oplus R/J_n$ (as a module), where J_i is an ideal of R such that R/J_i is self-injective for $i = 1, 2, \dots, n$.

Remark 3. In case R is a Boolean ring, Theorem 3.10 was shown by Pierce [6, p.104] without the assumption that M is torsion free, and he asked if this is valid for an arbitrary regular ring ([6, p.110]). However we can answer this question in the negative by giving the following example.

Example C. Let D be a finite field, say $D = \{a_1, a_2, \dots, a_n\}$, and let D' be a proper subfield of D . Take an infinite index set A and let

$$Q = \prod_{i \in A} D_i, \quad D_i = D \text{ for all } i \in A.$$

Here we denote by R the subring of Q consisting of all elements such that all but a finite number of whose components belong to D' . Then R is a continuous regular ring and its maximal ring of quotients coincides with Q (cf. [10, Example 3]). Moreover if we denote by x_i the element of Q such that all components are a_i , $i = 1, 2, \dots, n$, then we have $Q = Rx_1 + Rx_2 + \dots + Rx_n$. Hence Q is a finitely generated injective R -module, but not a direct sum of cyclic R -modules. Because, if it is a direct sum of cyclic R -modules, then it is projective, since by Theorem 2.1 each component is so. By [3, Lemma 4], this implies that $Q = R$, a contradiction.

REFERENCES

- [1] V.C. CATEFORIS: On regular self-injective rings, Pacific J. Math. **30** (1969), 39—45.
- [2] E. GENTILE: Singular submodule and injective hull, Indag. Math. **24** (1962), 426—433.
- [3] I. KAPLANEKY: Projective modules, Ann. of Math. **68** (1958), 372—377.
- [4] J. LAMBEK: Lectures on rings and modules. Blaisdell Publishing Co., Waltham, Mass., 1966.
- [5] A.C. MEWBORN: Regular rings and Baer rings, Math. Z. **12** (1971), 211—219.
- [6] R.S. PIERCE: Modules over commutative regular rings, Memoirs Amer. Math. **128** (1967),
- [7] F.L. SANDOMIERSKI: Non-singular rings, Proc. Amer. Math. Soc. **19** (1968), 225—230.
- [8] B. STENSTRÖM: Rings and modules of quotients, Springer Lecture Notes 237, 1971.
- [9] Y. UTUMI: On continuous regular rings, Canad. Math. Bull. **4** (1961) 63—69.
- [10] —————: On continuous regular rings and semisimple self-injective rings, Canad. J. Math. **12** (1960), 597—605.
- [11] J. ZELMANOWITZ: Injective hulls of torsion free modules, Canad. J. Math. **23** (1971), 1094—1101.

DEPARTMENT OF MATHEMATICS,
YAMAGUCHI UNIVERSITY

(Received November 18, 1972)