

Expression systems for industrial Gram-positive bacteria with low guanine and cytosine content

Willem M de Vos*[†], Michiel Kleerebezem* and Oscar P Kuipers*

Recent years have seen an increase in the development of gene expression systems for industrial Gram-positive bacteria with low guanine and cytosine content that belong to the genera *Bacillus*, *Clostridium*, *Lactococcus*, *Lactobacillus*, *Staphylococcus* and *Streptococcus*. In particular, considerable advances have been made in the construction of inducible gene expression systems based on the capacity of these bacteria to utilize specific sugars or to secrete autoinducing peptides that are involved in quorum sensing. These controlled expression systems allow for present and future exploitation of these bacteria as cell factories in medical, agricultural, and food biotechnology.

Addresses

*Department of Biophysical Chemistry, NIZO, PO Box 20, Kernhemseweg 2, 6710 BA Ede, The Netherlands

[†]Laboratory for Microbiology, Department of Biomolecular Sciences, Wageningen Agricultural University, Wageningen, The Netherlands; e-mail: willem.devos@algemeen.micr.wau.nl
Correspondence: Willem M de Vos

Current Opinion in Biotechnology 1997, 8:547–553

<http://biomednet.com/elecref/0958166900800547>

© Current Biology Ltd ISSN 0958-1669

Abbreviations


ABE	acetone-butanol-ethanol
CcpA	catabolite control protein
GC	guanine and cytosine
LAB	lactic acid bacteria
LGB	low GC Gram-positive bacteria
RCR	rolling circle replication
RR	response regulator

Introduction

Gram-positive bacteria are well known for their important contributions to medical, agricultural and food biotechnology and as production organisms for industrial enzymes. The latter capacity is largely based on the common architecture of their cell envelope, which allows for direct secretion of proteins into the fermentation medium, which in *Bacilli* has reached a high level of efficiency [1]. Based on the guanine and cytosine (GC) content of their DNA, Gram-positive bacteria can be divided into two major groups, the low and high GC groups. This dichotomy is confirmed by phylogenetic analysis based on 16S rRNA sequence comparisons that emphasize the separated evolutionary positions of the two groups [2]. As a consequence, the industrially relevant but highly diverse actinomycetes, which belong to the high GC group, are not discussed here, although *Streptomyces* spp. have been developed into efficient hosts for heterologous protein production, in spite of their GC content of

more than 70% [3]. This review will focus on the low GC Gram-positive bacteria (LGB) that have established biotechnological applications and belong to the genera *Bacillus*, *Clostridium*, *Lactococcus*, *Lactobacillus*, *Staphylococcus* and *Streptococcus*. Phylogenetic analysis indicates that these genera, in particular those belonging to the lactic acid bacteria (LAB) that include *Lactococcus*, *Lactobacillus* and *Streptococcus* spp., form a homogeneous group and are relatively closely related in evolutionary terms (Figure 1). This is in line with the widely spread presence, or functionality upon introduction, of highly conserved plasmids and transposons in LGB that are excellent tools for the development of genetic systems. Furthermore, recent findings, discussed below, indicate that common mechanisms of initiation and density-dependent control of gene expression are operating in the LGB. These common genetic characteristics are highly relevant for the further development of LGB to become expression hosts. As a consequence, these genetic characteristics are briefly summarized below before recent progress on controlled and efficient gene expression in the individual genera is discussed.

Figure 1



Unrooted phylogenetic tree based on 16S rRNA sequences showing the relative positions of the genera *Bacillus*, *Clostridium*, *Staphylococcus*, *Streptococcus*, *Lactobacillus*, and *Lactococcus*.

Common characteristics of LGB: mechanisms and applications

Conserved plasmids and transposons

Plasmids with a highly conserved unit of replication, or 'replicon', are found in all the genera of LGB discussed in the introduction and replicate by two well-studied mechanisms, theta replication or rolling circle replication (RCR) [4,5]. Large plasmids usually replicate via theta replication and belong either to the pAM β 1 family of broad host-range conjugative plasmids or to the pCI305

family, the members of which have a limited host-range. A variety of useful vectors have been based on the replicon of pAM β 1, which is functional in all studied genera of LGB [5,6]. These vectors have been equipped with *Escherichia coli* replicons to allow efficient shuttling between Gram-positive and Gram-negative bacteria [7]. The detailed analysis of its replicon allowed for the development of plasmid vectors for LGB which can switch from high to low copy number, and hence are useful for modulating the expression level of cloned genes [8*]. This was achieved by introducing an inactivating linker cassette in the repressor gene, *copF*, that regulates the *repE* gene coding for the rate-limiting replication enzyme. Removal of the cassette by a simple digestion with the infrequently-cutting restriction enzyme *KpnI* reconstitutes a functional *copF* gene, resulting in a change from high to low copy number.

Small plasmids of several kilobases in general use RCR, are ubiquitous in all studied genera, and also belong to a limited number of families [4,5,9]. The replicons of these plasmids have been developed in a series of highly useful cloning vectors that replicate in a broad range of genera, notably those based on the related *Lactococcus lactis* plasmids pWV01/pSH71, which also replicate in *E. coli* [5]. A thermosensitive derivative of pWV01, termed pG+host, has been generated and has shown its potential as a delivery system in LGB [10]. By introduction of the lactococcal insertion sequence *ISS1* element in pG+host, a random transposition system has been constructed that is functional in a broad range of genera [11*] and may be used to generate mutants or introduce foreign genes. This system may also be used as an alternative to the naturally occurring transposons that are found in LGB [12]. These include the transposons related to Tn917 that have been used for insertional mutagenesis in *Bacillus* and several LAB [12,13]. Recently, a Tn917 derivative was optimized for promoter screening in *Lactococcus* [14]. In addition, conjugative transposons have been found in several genera [15]; however, most use has been made of the well-characterized Tn916, which is able to generate insertions in all LGB [12,15]. The application of the described tools into suitable cloning, integration and expression vectors has been the subject of several recent reviews, with specific attention being paid to LAB [5,8*,9,12].

Initiation of transcription and translation

The composition of core RNA polymerases in LGB resembles that of *E. coli* [16]. In addition, while the number of identified sigma factors may vary between the different genera, the main sigma factor, sigma A, is also conserved [5,16]. This is in line with the observation that most sigma A-dependent promoters in LGB contain the canonical -35 and -10 sequences [5,9,16]. Recently, a vital -16 region was identified that affects promoter efficiency in *Bacillus subtilis* and *E. coli* [17]. In addition, sequence analysis of *B. subtilis* sigma A-

dependent promoters indicated that there are extended contacts in this region and also upstream as far as -70 [18]. This confirms previous analysis of LGB other than *B. subtilis* that show conservation of a thymine-guanine dinucleotide in the -16 region and an excess of adenine and thymine sequences upstream of the -35 consensus [5,9]. Furthermore, it may explain why some *E. coli* promoters are not efficiently used in LGB; however, within LGB the efficiency of transcription initiation may also vary in a species-dependent way, as was observed when transcriptional *gusA* fusions were studied in different LAB, suggesting that optimization for each expression host is advisable [19].

Translation initiation in LGB shows remarkable features that discriminate it from that in other Gram-positive and Gram-negative bacteria, confirming its phylogenetic position [20]. Only LGB have been found to lack the ribosomal protein S1, which is coupled to a high translational selectivity not observed with other bacteria, including the high GC Gram-positive ones. This is reflected in a more-extensive complementarity between the Shine-Dalgarno sequences and the 3' end of the 16S rRNA than found in other bacteria [5,9,20]. The requirement for a stringent Shine-Dalgarno region for high translational efficiency has been observed in many cases and explains previous observations that *E. coli* genes, which do not generally have Shine-Dalgarno sequences with such complementarity, are not well expressed in several LGB [21].

Catabolite repression

While catabolite repression in *E. coli* and other Gram-negative bacteria has been well characterized and involves the catabolite repressor protein and its interaction with cyclic AMP, these compounds have not been found to a similar extent in LGB [22]. In contrast, it has been shown that catabolite repression in LGB is mediated via a catabolite control protein (CcpA) and the *cis*-acting sequence *cre* [22,23]. CcpA interacts with the serine-phosphorylated form of the heat stable protein (HPr), a phospho-carrier involved in the phosphotransferase system mediating the active uptake of sugars [23,24]. The serine-phosphorylation of HPr, which seems exclusive to LGB, is part of the signalling pathway, since it is catalyzed by a serine kinase when the intracellular fructose-1,6-diphosphate concentration is high. Immunological studies have shown the presence of a CcpA-like protein in all genera of LGB discussed here [25]. Recently, this has been supported by genetic data on the sequence and functionality of the *ccpA* genes [25,26]. While some *ccpA* genes seem constitutively expressed, the *ccpA* gene in *Staphylococcus xylosus* is transcribed from two promoters, one of which contains a *cre* sequence and is subject to CcpA-mediated autoregulation [27]. Further insight in the global control system exerted by CcpA will allow for optimal exploitation of sugar inducible expression systems that are often used to control gene expression in LGB. Well-established


examples are the xylose inducible systems studied in *Bacillus*, *Staphylococcus*, and *Lactobacillus*, which are subject to specific repression by a homologous repressor XylR and subject to negative regulation via CcpA [28–30]. Finally, analysis of the CcpA mediated control may contribute to eliminating catabolite repression that is often observed during industrial mixed substrate fermentations.

Quorum sensing

Cell density-dependent gene expression, also known as quorum sensing, appears to be widely spread in microbes. In Gram-negative bacteria, this form of cell-cell communication has been studied extensively and involves diffusible *N*-acetyl homoserine lactones as autoinducing signals, which interact with LuxR-like transcriptional activators [31]. Recent studies have indicated that LGB also communicate with each other but use peptide pheromones as signal molecules [32•] (Figure 2). These peptide pheromones are made as precursors that are post-translationally processed and exported via dedicated ATP binding cassette transporters. In addition, the extracellular peptide pheromones function as input signals for a two component signal transduction system consisting of a membrane located sensor histidine protein kinase and a cytoplasmatic response regulator (RR), which activates transcription. It is assumed that binding of the peptide pheromone to histidine protein kinase results in a phosphorylation cascade resulting in formation of a phosphorylated RR, which subsequently binds and transcriptionally activates one or more dedicated promoters, including the promoter of an operon of the peptide pheromone precursor gene and the *hprk* and *rr* genes, encoding HPR and RR, respectively (Figure 2).

Examples of this novel quorum sensing involving autoinduction of peptide pheromones have been found in all members of LGB described here and include the development of competence in *Bacillus* [33,34] and *Streptococcus spp.* [35,36], the development of virulence in *Staphylococcus spp.* [37,38], and the production of antimicrobial peptides by *Lactobacillus spp.* [39,40] or *Lactococcus* [41]. The latter system concerns the production of the lantibiotic nisin, an extensively post-translationally modified peptide that is used in the food industry because of its activity against a wide range of LGB, including pathogens and food spoilage bacteria [42]. The nisin autoinduction system was the first example of quorum sensing involving antimicrobial peptide production. It has now been developed into an efficient and industrially applicable controlled gene expression system by introduction of a gene of interest (*gene X*) under control of a nisin-inducible promoter either in a nisin producing strain (Figure 2a) or in a strain harboring only the genes for the signal transduction pathway (Figure 2b) [43••]. In the first case, expression of *gene X* is dependent on the cell density and occurs at the end of the growth (Figure 2a). In the

Figure 2



General outline of the quorum-sensing systems in LGB. In cell-density-dependent expression (a) the peptide pheromone is the signalling molecule, the concentration of which accumulates during growth in a fermentor and triggers its own production and that of the gene of interest (*gene X*) via a cascade response. Alternatively, for induced gene expression (b) the structural gene for the peptide pheromone precursor is not present and induction of the gene(s) of interest depends on the exogenous addition of the pheromone. ABC, ATP-binding cassette; HPK, histidine protein kinase; P, promoter; P_i, inorganic phosphate; ppp, peptide pheromone precursor gene.

second configuration, *gene X* expression can be induced at will by the addition of the inducing peptide pheromone (Figure 2b). Examples of these systems in *L. lactis* have been described [43••]. Similar approaches could be applied to the quorum sensing modules that are presently known in LGB (see above) or may be found in future studies.

Controlled and efficient expression systems in LGB: new developments and applications

Bacilli

One of the first controlled expression systems in *B. subtilis* was based on the *E. coli lacI* repressor gene, equipped with an appropriate promoter and ribosome binding site in conjunction with the inducible *Spac-1* promoter, consisting of the promoter of phage SPO-1 coupled to the *lac* operator [44]. This system, however, could only be used for specific, small-scale applications because of the need to add the gratuitous inducer isopropyl- β -D-thiogalactosidase. A variety of other systems have been reported in the meantime but it was only recently that another successful *E. coli* system, the T7 system, was implemented in *B. subtilis* [45].

The use of the T7 system was achieved in *B. subtilis* by inserting the T7 RNA polymerase gene *rpoT7* into the chromosome under control of a xylose-inducible promoter (see above). In conjunction with a plasmid carrying the T7 promoter driving expression of the gene of interest, this heterologous expression system resulted in high induction and production levels of both intracellular and extracellular enzymes upon addition of xylose and may be suitable for larger scale applications. The xylose-inducible system was also used recently—without the T7 amplification step—in a set of cassettes that could be integrated in the *amyE* locus of the *B. subtilis* chromosome [46]. High induction levels were also obtained here upon xylose addition and only repression by glucose, not by other rapidly metabolizable compounds, was observed. This versatile system is not only suitable for fundamental studies aimed at comparing expression efficiencies but may also have potential for further applications, since it may be used in other, industrially more relevant *Bacilli* and eliminate the need for unstable plasmid vectors and their selection. These may include hosts such as *Bacillus licheniformis* and *Bacillus amyloliquefaciens* [1]. In addition, in the past year significant progress has been reported with *Bacillus brevis*, which produces large amounts of surface layer proteins. Based on the appropriate signals of these surface layer proteins, expression-secretion vectors were constructed that allowed for the extracellular production of large amounts (more than 1 g/l) of functional human epidermal growth factor [47*]. Similar, although less efficient, production was obtained in another study with animal erythropoietin and the extracellular domain of its receptor [48]. Finally, hyperproduction of the *Clostridium* alpha-toxin was observed in a similar approach in *B. brevis* which showed ten times more production than *B. subtilis* [49].

Clostridia

In spite of their considerable medical and economic importance, genetic systems for *Clostridium* spp. are not very well developed, hampering their use as expression hosts [50]. Considerable attention has been given to the medical aspects of *Clostridia*, which have been reviewed

recently together with the present state of its molecular biology [51]. Next to their role in pathogenesis and as producers of exoenzymes, some *Clostridia*, notably *Clostridium acetobutylicum* are well known for the production of acetone–butanol–ethanol (ABE). In an attempt to increase the ABE fermentation, the well-known *C. acetobutylicum* strain ATCC 824 was transformed with plasmids carrying homologous sporulation genes. Transformants were reported not to survive, however, and so an integration approach was followed that resulted in mutants with scrambled chromosomal inserts, some of which showed increased stability and efficiency of ABE fermentation [52]. In addition to its established application areas, a novel potential application for *Clostridia* has been recently described that is based on its potential to target hypoxic regions of tumour cells and specifically germinate at these locations [53*]. This would create a specific delivery system for toxic compounds that could inactivate the cancer cells. This interesting possibility was tested with *Clostridium beijerinckii* expressing the *E. coli* gene for cytosine deaminase, which may convert the nontoxic prodrug 5-fluorocytosine into an active chemotherapeutic compound. *In vitro* conversion of the prodrug was achieved with a clostridial supernatant [53*], illustrating the feasibility of this approach.

Staphylococci

Most studies on staphylococcal expression systems have concentrated on *Staphylococcus carnosus* or *Staphylococcus xylosus*, that are non-pathogenic and have a long tradition of safe use as starters for meat fermentations. An improvement of the xylose induction system (see above) from *S. xylosus* has been developed [29]. Using a previously constructed lipase reporter system it was shown that the new expression vector pTX15 carrying the xylose inducible *xylA* promoter showed better induction and lower background activity than a previously described plasmid [54]. The utility of the vector pTX15 was demonstrated in *S. carnosus* with the *Staphylococcus epidermidis* *epiB* gene, which is involved in the production of lanthionine rings in the lantibiotic epidermin. Finally, the T7 system also has been applied to *S. carnosus*, again using the xylose-inducible *xylA* promoter [55]. *S. carnosus* was also further developed for the display of recombinant proteins at its cell surface, exploiting the anchor domain of the staphylococcal protein A [56]. This approach is aimed at developing non-pathogenic *Staphylococci* with a long tradition of safe use as novel vaccine carriers. Similarly, strains of both *S. carnosus* and *S. xylosus* appeared to be excellent hosts for the surface production of functional single-chain Fv antibodies [57]. It is speculated that this approach may lead to the development of these LGB as whole-cell diagnostic devices or as alternatives to filamentous phages for surface display of single chain Fv libraries.

Streptococci

Interest in the genetics of a number of different streptococcal species is in many cases supported by their role

as pathogens or their contribution to food-fermentations. Although a variety of plasmid vectors is available for *Streptococci*, integrative gene expression systems were evaluated in two different *Streptococcus* spp., representing each of these application groups. A promoterless chloramphenicol acetyltransferase gene was inserted in between the chromosomally located *lacS* and *lacZ* genes of the yoghurt bacterium *Streptococcus thermophilus*, the expression of which is induced during growth on lactose [58]. Although LacZ activity was reduced to about 10 %, the chloramphenicol acetyltransferase activities paralleled those of β -galactosidase, indicating that both genes were expressed and regulated by the same control elements. In addition, a host-vector system for heterologous gene expression was developed for the pathogenic *Streptococcus gordonii* (formerly *Streptococcus sanguis*), which included the streptococcal M6 protein to allow for surface expression [59].

Lactobacilli

The development of convenient cloning and controlled expression systems in *Lactobacilli* has been recently reviewed [9,60,61]. Increasing attention is being given to *Lactobacillus plantarum*, which has a reasonable capacity to secrete proteins. This was exploited by the secretion of α -amylase and levanase using homologous expression-secretion signals both located on the chromosome and on plasmids [62]. In addition, using a similar approach as described above for *B. brevis*, high level production of heterologous proteins was obtained in both *Lactobacillus* and *Lactococcus* with a secretion system based on the signals of the surface layer protein discovered in *Lactobacillus brevis* [63].

Lactococci

By far the best studied LAB are the *Lactococci*, specifically *L. lactis*, an important starter in cheese production. In the past few years several expression systems, both constitutive and regulated, have been developed [5,6,61]. Recent advances include the isolation and characterization of more than 10 different promoter elements from the chromosome of *L. lactis*, exhibiting a 70-fold range of activity [64]. The strongest promoter, P15, shows unusual spacing between 35 and -10 regions, which could indicate a regulation mechanism acting at the transcriptional level. In addition, several reports have appeared describing the use of the T7 expression system in *L. lactis*, which was based on the inducible *lac* promoter [65]. High, inducible extracellular production of interleukin-2 was achieved in *Lactococci* [66]. The potential of *L. lactis* as a vaccine carrier was further developed and a protective immune response was obtained with strains secreting the tetanus toxin fragment C [67]. Recently, several different systems for inducible gene expression were developed that are briefly discussed below.

Based on the complete nucleotide sequence of the temperate *L. lactis* bacteriophage ϕ r1t, repressor-operator

sequences were identified and exploited by developing a system in which gene expression could be activated by the addition of mitomycin C, resulting in an induction factor of about 70 [68]. Recently, this system was further improved by development of a thermosensitive repressor, which facilitates a wider applicability of this system (J Kok, personal communication).

A system for explosive gene expression was developed based on the ϕ 31 middle promoter and origin of replication (*ori*) [69••]. Essentially, any gene of interest can be cloned under control of the middle promoter combined with a low copy number vector containing the phage *ori*, which will be induced by infection by ϕ 31, resulting in an increase in plasmid copy number and considerable overproduction of the protein of interest to >1000-fold. The protein(s) of interest will be released into the growth medium since phage infection will lyse the cells within one hour.

A system has been developed for controlled overproduction of proteins, employing the two-component regulatory system for the biosynthesis of the antimicrobial peptide nisin (see above, Figure 2) [42]. It was shown that the *nisA* promoter could be employed in a series of vectors for expression of several homologous and heterologous genes, which can be induced by the addition of subinhibitory amounts of the signalling molecule nisin [43,70]. A variety of genes have been expressed in *L. lactis* using this system and the hyperproduction (more than 50% protein) of the intracellular aminopeptidase N was reported [42]. Food-grade expression vectors were also developed based on the dominant *lacF* selection system of *L. lactis*, allowing the overproduction of proteins to be used in the food industry [43]. This study also showed the inducible (Figure 2a) and cell density dependent expression of a reporter gene under control of the *nisA* promoter (Figure 2b). Recently, two compatible, broad host-range plasmids were constructed, one based on a theta replicon and expressing the *nisRK* genes and the other based on a high copy number RCR plasmid carrying the *nisA* promoter controlling the expression of a reporter gene. This dual plasmid system could be introduced and maintained in *Lactococcus*, *Lactobacillus* and *Streptococcus* spp., and resulted in nisin controlled gene expression (Kleerebezem M, Vaughan EE, de Vos WM, Kuipers OP, unpublished data). As a consequence, the nisin inducible systems hold a high potential for a variety of applications in different LGB. Major advantages are the food-grade nature of the plasmids, strains and inducing agent, the high expression levels that can be reached and the low background activity of the *nisA* promoter, which enables induction factors of over 1000.

Conclusion

A great variety of inducible expression systems have now been developed for LGB. Most of these have been based on the advanced knowledge of gene expression in these hosts and naturally occurring regulatory systems that seem

to have a wide host range in LGB. Various species of LGB, including most of the genera discussed here, are now being analyzed by genomic sequencing. It may be expected that this will, on the one hand, promote the development of new or improved expression systems and, on the other hand, lead to an increased use of these systems in order to exploit LGB as cell factories in medical, agricultural and food biotechnology.

Acknowledgements

We are grateful to Arthur Wolterink for help with Figure 1 and Luis Quadri for handling the Figures. Part of the work in the authors' laboratory was supported by BIOT contracts from the European Union.

References and recommended reading

Papers of particular interest, published within the annual period of review have been highlighted as:

- of special interest
 - of outstanding interest
1. Ferrari E, Jarnagin AS, Schmidt BF: **Commercial production of extracellular enzymes.** In *Bacillus Subtilis and Other Gram-Positive Bacteria. Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:917-937.
 2. Woese CR: **Bacterial evolution.** *Microbiol Rev* 1987, **51**:221-271.
 3. Binnie C, Cossar JD, Stewart DIH: **Heterologous biopharmaceutical protein expression in Streptomyces.** *Trends Biotechnol* 1997, **15**:315-320.
 4. Janniere L, Gruss A, Ehrlich SD: **Plasmids.** In *Bacillus Subtilis and Other Gram-Positive Bacteria. Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:625-645.
 5. De Vos WM, Simons G: **Gene cloning and expression systems in lactococci.** In *Genetics and Biotechnology of Lactic Acid Bacteria*. Edited by Gasson MJ, de Vos WM. Oxford: Chapman and Hall; 1994:52-105.
 6. Renault P: **Genetic engineering strategies.** In *Lactic Acid Bacteria: Current Advances in Metabolism, Genetics and Applications*. Edited by Bozoglu TF, Ray B. Berlin: Springer-Verlag; 1996:1-37. [NATO Symposium Series.]
 7. O'Sullivan DJ, Klaenhammer: **High- and low-copy-number Lactococcus shuttle cloning vectors with features for clone screening.** *Gene* 1993, **137**:227-231.
 8. Renault P, Corthier G, Goupil N, Delorme C, Ehrlich SD: **Plasmid vectors for gram-positive bacteria switching from high to low copy number.** *Gene* 1996, **183**:175-182.
- Illustrates the rational design of useful tools based on insight in the copy number control system of the broad host-range plasmid pAM β 1. Described vectors will be useful for many low guanine and cytosine Gram-positive bacteria.
9. Pouwels PH, Leer RJ: **Genetics of lactobacilli: plasmids and gene expression.** *Antonie van Leeuwenhoek* 1993, **64**:85-107.
 10. Maguin E, Duwat P, Hege T, Ehrlich SD, Gruss A: **New thermosensitive plasmid for gram-positive bacteria.** *J Bacteriol* 1992, **174**:5633-5638.
 11. Maguin E, Prevost H, Ehrlich SD, Gruss A: **Efficient insertional mutagenesis on lactococci and other Gram-positive bacteria.** *J Bacteriol* 1996, **178**:931-935.
- Delivery system for a broad host-range transposon eliminating the need for efficient transformation. Useful tool that will allow genetic analysis of wide range of low guanine and cytosine Gram-positive bacteria.
12. Gasson MJ, Fitzgerald GF: **Gene transfer systems and transposition.** In *Genetics and Biotechnology of Lactic Acid Bacteria*. Edited by Gasson MJ, de Vos WM. Oxford: Chapman and Hall; 1994:1-51.
 13. Youngman P: **Transposons and their applications.** In *Bacillus Subtilis and Other Gram-positive Bacteria. Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:585-596.
 14. Israelson H, Madsen SM, Vrang A, Hansen EB, Johansen E: **Cloning and partial characterization of regulated promoters from Lactococcus lactis Tn917-lacZ integrants with the new promoter probe vector, pAK80.** *Appl Environ Microbiol* 1995, **61**:2540-2547.
 15. Scott JR: **Conjugative transposons.** In *Bacillus Subtilis and Other Gram-Positive Bacteria. Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:597-613.
 16. Moran CP: **RNA polymerase and transcription factors.** In *Bacillus Subtilis and Other Gram-Positive Bacteria. Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:653-668.
 17. Voskuil MI, Voepel K, Chambliss GH: **The -16 region, a vital sequence for the utilization of a promoter in Bacillus subtilis and Escherichia coli.** *Mol Microbiol* 1995, **17**:271-279.
 18. Helmann JD: **Compilation and analysis of Bacillus subtilis sigma A-dependent promoter sequences: evidence for extended contact between RNA polymerase and upstream promoter DNA.** *Nucleic Acids Res* 1995, **23**:2351-2360.
 19. Platteeuw C, Simons G, de Vos WM: **Use of the Escherichia coli β -glucuronidase (gusA) gene as a reported gene for analyzing promoters in lactic acid bacteria.** *Appl Environ Microbiol* 1995, **60**:587-593.
 20. Vellanoweth RL: **Translation and its regulation.** In *Bacillus Subtilis and Other Gram-Positive Bacteria. Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:699-711.
 21. Rabinowitz JC, Roberts M: **Translational barriers limiting expression of E. coli genes in Bacillus subtilis and other Gram-positive organisms.** In *Antibiotic Resistance Genes: Ecology, Transfer, and Expression*. Edited by Levy SB, Novick RP. Cold Spring Harbor: Cold Spring Harbor Laboratories; 1986:297-312.
 22. Chambliss GH: **Carbon source mediated catabolite repression.** In *Bacillus Subtilis and Other Gram-Positive Bacteria. Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:213-219.
 23. Hueck C, Hillen W: **Catabolite repression in Bacillus subtilis: a global regulatory mechanism for the Gram-positive bacteria?** *Mol Microbiol* 1995, **15**:395-401.
 24. Deutscher J, Kuster E, Bergstedt U, Charrier V, Hillen W: **Protein-kinase dependent HPr/CcpA interaction links glycolytic activity to carbon catabolite repression in Gram-positive bacteria.** *Mol Microbiol* 1995, **15**:1049-1053.
 25. Kuster E, Luesink EJ, de Vos WM, Hillen W: **Immunological cross-reactivity to catabolite control protein CcpA from B. megaterium is found in many Gram-positive bacteria.** *FEMS Microbiol Lett* 1996, **139**:109-115.
 26. Luesink EJ, Vaughan EE, Kuipers OP, de Vos WM: **Catabolite repression in lactic acid bacteria.** *Le Lait* 1997, in press.
 27. Egeter O, Brückner R: **Catabolite repression mediated by the catabolite control protein CcpA in Staphylococcus xylosum.** *Mol Microbiol* 1997, **21**:739-749.
 28. Jacob S, Allmansberger R, Gartner D, Hillen W: **Catabolite repression of the operon for xylose utilization from Bacillus subtilis W23 is mediated at the level of transcription and depends on a cis site in the xylA reading frame.** *Mol Gen Genet* 1991, **229**:189-196.
 29. Sizemore C, Wieland B, Gotz F, Hillen W: **Regulation of the Staphylococcus xylosum xylose utilization genes at the molecular level.** *J Bacteriol* 1991, **174**:3042-3048.
 30. Lokman BC, Heerikshuizen M, van den Broek A, Borsboom Y, Chaillou S, Postma PW, Pouwels PH: **Regulation of the Lactobacillus pentosus xylAB operon.** *J Bacteriol* 1997, **179**:in press.
 31. Salmond GPC, Bycroft BW, Stewart GSAB, Williams P: **The bacterial enigma: cracking the code of cell-cell communication.** *Mol Microbiol* 1995, **16**:615-624.
 32. Kleerebezem M, Quadri LEN, Kuipers OP, de Vos WM: **Quorum sensing by peptide pheromones and two-component signal transduction systems in Gram-positive bacteria.** *Mol Microbiol* 1997, **24**:895-905.
- Overview of the peptide pheromone-induced quorum-sensing modules that are operating in low guanine and cytosine Gram-positive bacteria and function via a two-component regulatory system.

33. Magnusson R, Solomon J, Grossman AD: **Biochemical and genetic characterization of a competence pheromone from *Bacillus subtilis***. *Cell* 1994, **77**:207-216.
34. Grossman DA: **Genetic networks controlling initiation of sporulation and development of genetic competence in *Bacillus subtilis***. *Ann Rev Genet* 1995, **29**:477-508.
35. Håvarstein LS, Coomaraswamy G, Morrison DA: **An unmodified heptadecapeptide pheromone induces competence for genetic transformation in *Streptococcus pneumoniae***. *Proc Natl Acad Sci USA* 1995, **92**:11140-11144.
36. Pestova EV, Håvarstein LS, Morrison DA: **Regulation of competence for genetic transformation in *Streptococcus pneumoniae* by an auto-induced peptide pheromone and a two-component regulatory system**. *Mol Microbiol* 1996, **21**:853-862.
37. Ji GY, Beavis RC, Novick RP: **Cell density control of staphylococcal virulence mediated by an octapeptide pheromone**. *Proc Natl Acad Sci USA* 1995, **92**:12055-12059.
38. Guangyong Ji, Beavis R, Novick RP: **Bacterial interference caused by autoinducing peptide variants**. *Science* 1997, **276**:2027-2030.
39. Diep DB, Havarstein LS, Nes IF: **A bacteriocin-like peptide induces bacteriocin synthesis in *Lactobacillus plantarum* C11**. *Mol Microbiol* 1995, **18**:631-639.
40. Eijsink VGH, Brurberg MB, Middelhoven PH, Nes IF: **Induction of bacteriocin production in *Lactobacillus sake* by a secreted peptide**. *J Bacteriol* 1996, **178**:2232-2237.
41. Kuipers OP, Beerthuyzen MM, de Ruyter PGG, Luesink EJ, de Vos WM: **Autoregulation of nisin biosynthesis in *Lactococcus lactis* by signal transduction**. *J Biol Chem* 1995, **270**:27299-27304.
42. De Vos WM, Kuipers OP, van der Meer JR, Siezen RJ: **Maturation pathway of nisin and other lantibiotics: post-translationally modified antimicrobial peptides exported by Gram-positive bacteria**. *Mol Microbiol* 1995, **17**:427-437.
43. Kuipers OP, de Ruyter PGG, Kleerebezem M, de Vos WM: **Controlled overproduction of proteins by lactic acid bacteria**. *Trends Biotechnol* 1997, **15**:135-140.
- This report discusses the different systems used for enzyme overproduction in lactic acid bacteria and describes the nisin-inducible gene expression system resulting in controlled hyperproduction of intracellular enzymes in *L. lactis*.
44. Yansura DG, Henner DJ: **Use of the *Escherichia coli* lac repressor and operator to control gene expression in *Bacillus subtilis***. *Proc Natl Acad Sci USA* 1984, **81**:439-443.
45. Conrad B, Savchenko RS, Breves R, Hofeweister J: **A T7 promoter-specific, inducible protein expression system for *Bacillus subtilis***. *Mol Gen Genet* 1996, **250**:230-236.
46. Kim L, Mogk A, Schumann W: **A xylose-inducible *Bacillus subtilis* integration vector and its application**. *Gene* 1996, **181**:71-76.
47. Ebisu S, Takagi H, Kadowaki K, Yamagata H, Udaka S: **The efficient production of human epidermal growth factor by *Bacillus brevis***. *Ann N Y Acad Sci* 1996, **782**:115-122.
- Describes the capacity of *B. brevis* for the extracellular production of proteins and illustrates the potential of an alternative *Bacillus* host.
48. Nagao M, Inoue K, Moon SK, Masuda S, Takagi H, Udaka S, Sasaki R: **Secretory production of erythropoietin and the extracellular domain of the erythropoietin receptor by *Bacillus brevis*: affinity purification and characterization**. *Biosci Biotechnol Biochem* 1997, **61**:670-674.
49. Nagaham M, Michiue K, Sakurai J: **Production and purification of *Clostridium perfringens* alpha-toxin using a protein-hyperproducing strain, *Bacillus brevis* 47**. *FEMS Microbiol Lett* 1996, **145**:239-243.
50. Young M, Cole ST: ***Clostridia***. In *Bacillus Subtilis and Other Gram-Positive Bacteria*. *Biochemistry, Physiology and Molecular Biology*. Edited by Sonensheim AL, Hoch JA, Losick R. Washington: American Society for Microbiology; 1993:35-53.
51. Rood JI, McClane BA, Songer JG, Titball RW: *The Clostridia: Molecular Biology and Pathogenesis*. New York: Academic Press; 1997.
52. Wong J, Bennet GN: **Recombination-induced variants of *Clostridium acetobutylicum* ATCC 824 with increased solvent production**. *Curr Microbiol* 1996, **32**:349-356.
53. Fox ME, Lemmon MJ, Mauchline ML, Davis TO, Giacca AJ, Minton NP, Brown JM: **Anaerobic bacteria as a delivery system for cancer therapy: *in vitro* activation of 5-fluorocytosine by genetically engineered clostridia**. *Gene Ther* 1996, **3**:173-178.
- The authors describes an original approach that may allow for the therapeutic use of *Clostridia*, due to their potential to reach and kill cancer cells.
54. Peschel A, Ottenwalder B, Gotz F: **Inducible production and cellular location of the epidermin biosynthetic enzyme EpiB and improved staphylococcal expression system**. *FEMS Microbiol Lett* 1996, **137**:279-284.
55. Weiland K-P, Weiland B, Gotz F: **A promoter-screening plasmid and xylose-inducible, glucose-repressible expression vectors for *Staphylococcus carnosus***. *Gene* 1996, **158**:91-96.
56. Samuelsson P, Hannson M, Ahlborg N, Androni C, Gotz F, Bach T, Nguyen TN, Binz H, Uhlen W, Stahl S: **Cell surface display of recombinant protein on *Staphylococcus carnosus***. *J Bacteriol* 1995, **177**:1470-1476.
57. Gunneriusson E, Samelson P, Uhlen M, Nygren PA, Stahl S: **Surface display of a functional single chain Fv antibody on staphylococci**. *J Bacteriol* 1996, **178**:1341-1346.
58. Mollet B, Knol J, Poolman B, Marciset O, Delley M: **Directed genomic integration, gene replacement, and integrative gene expression in *Streptococcus thermophilus***. *J Bacteriol* 1993, **175**:4315-4324.
59. Oggioni MR, Pozzi G: **A host-vector system for heterologous gene expression in *Streptococcus gordonii***. *Gene* 1996, **169**:85-90.
60. Klaenhammer TR: **Genetics of intestinal lactobacilli**. *Int Dairy J* 1995, **5**:39-86.
61. Kok J: **Inducible gene expression and environmentally regulated genes in lactic acid bacteria**. *Antoine van Leeuwenhoek* 1996, **70**:129-145.
62. Hols P, Ferain T, Garmyn D, Bernard N, Delcour J: **Use of homologous expression-secretion signals and vector-free stable chromosomal integration in engineering of *Lactobacillus plantarum* for α -amylase and levanase expression**. *Appl Environ Microbiol* 1994, **60**:1401-1413.
63. Savijoki K, Kahala M, Palva A: **High level heterologous protein production in *Lactococcus* and *Lactobacillus* using a new secretion system based on the *Lactobacillus brevis* S-layer signals**. *Gene* 1997, **186**:255-262.
64. Waterfield NR, LePage RWF, Wilson PW, Wells JM: **The Isolation of Lactococcal promoters and their use in investigating bacterial luciferase synthesis in *Lactococcus lactis***. *Gene* 1995, **165**:9-15.
65. Wells JM, Wilson PW, Norton PM, Gasson MJ, Le Page RWP: ***Lactococcus lactis*: high level expression of tetanus toxin fragment C and protection against lethal challenge**. *Mol Microbiol* 1995, **8**:1155-1162.
66. Steidler L, Wells JM, Raemakers A, Vanderkerckhove J, Fiers W, Remaut E: **Secretion of biologically active murine interleukin-2 by *Lactococcus lactis* subsp. *lactis***. *Appl Environ Microbiol* 1995, **61**:1627-1629.
67. Robinson K, Chamberlain LM, Schofield KM, Wells JM, LePage RWF: **Oral vaccination of mice against tetanus with recombinant *Lactococcus lactis***. *Nat Biotechnol* 1997, **15**:653-657.
68. Nauta A, van Sinderen D, Karsens H, Smit E, Venema G, Kok J: **Inducible gene expression mediated by a repressor-operator system isolated from *Lactococcus lactis* bacteriophage-R1T**. *Mol Microbiol* 1996, **19**:1331-1341.
69. O'Sullivan DJ, Walker SA, West SG, Klaenhammer TR: **Development of an expression strategy using a lytic phage to trigger explosive plasmid amplification and gene expression**. *Bio-Technology* 1996, **14**:82-87.
- The authors describe an elegant system describing coupling of overproduction and lysis of the expression host *L. lactis*. Induction is not triggered by the addition of a chemical but of a bacteriophage particle.
70. De Ruyter PGG, Kuipers OP, de Vos WM: **Controlled gene expression systems for *Lactococcus lactis* with the food-grade inducer nisin**. *Appl Environ Microbiol* 1996, **62**:3662-3667.