

Unravelling the secret of the resistance of desert strains of *Chroococcidiopsis* to desiccation and radiation

Willemotte Annick³, Daniela Billi¹, Claudia Faglierone¹, Cyprien Verseux¹, Claudia Mosca¹ and Mickael Baqué^{1,2}

¹ University of Rome Tor Vergata, Dept Biology, Rome Italy

² Astrobiological Laboratories Research Group, Institute of Planetary Research, Management and Infrastructure, German Aerospace Center (DLR), Berlin, Germany

³ InBioS-Centre for Protein Engineering, University of Liège, Liège, Belgium
E-mail: awillemotte@ulg.ac.be

Chroococcidiopsis is a unicellular cyanobacterial genus that is growing in extreme dry conditions, either in low or high temperatures. At the lower end of the spectrum, they live as cryptoendoliths in rocks of the Mc Murdo Dry Valleys in Antarctica where they were discovered by Imre Friedmann, while at the higher end, they grow as hypoliths/endoliths in hot deserts, e.g. Negev, Gobi, Atacama (Friedman, 1980).

The capacity of desert strains of *Chroococcidiopsis* to stabilize their sub-cellular organization is so efficient that, when dried, they can cope with simulated space and Martian conditions (Billi et al 2011 ; Baqué et al. 2013a) as well as with high doses of ionizing and UV radiations (Verseux et al. 2017 ; Baqué et al. 2013b).

Since it is known for radiation/desiccation tolerant bacteria that the capability to avoid protein oxidation is critical to cope with such stressors (Frederickson et al. 2008; Daly et al. 2007), the present study investigates the protein oxidation after prolonged desiccation, irradiation with gamma-rays up to 25kGy and treatment with hydrogen peroxide in a selection of desert *Chroococcidiopsis* isolates, including 2 Antarctic strains: CCME134 and CME171 isolated from Beacon Valley and University Valley, respectively (Mc Murdo Dry Valleys).

A tight correlation was observed between the desiccation and radiation tolerance of the investigated desert strains and the absence of oxidative damage to proteins. The efficiency of the antioxidant systems of the desert strains of *Chroococcidiopsis* was highlighted also by the lack of protein carbonylation until treatment with 1M of oxygen peroxide. The phylogenetic analysis of the investigated 11 desert strains of *Chroococcidiopsis* is reported.

This work was supported by the Italian National Antarctic Research Program

This work is dedicated to the memory of Roseli Ocampo-Friedmann and E. Imre Friedmann who pioneered the research on *Chroococcidiopsis* and life in extreme environments.

References

- Baqué, M. et al. 2013a. Biofilm and planktonic lifestyles differently support the resistance of the desert cyanobacterium *Chroococcidiopsis* under space and Martian simulations. *Origin of Life and Evolution of Biospheres* 3,377-89.
- Baqué, M. et al. 2013b. Endurance of the endolithic desert cyanobacterium *Chroococcidiopsis* under UVC radiation. *Extremophiles* 17,161-169.
- Billi, D. et 2011. Damage escape and repair in dried *Chroococcidiopsis* spp. from hot and cold deserts exposed to simulated space and Martian conditions. *Astrobiology* 11,65-73.
- Daly, M.J. et al., 2007. Protein oxidation implicated as the primary determinant of bacterial radioresistance. *PLoS Biology*, 5(4), p.e92.
- Frederickson, J.K. et al. (2008) Protein oxidation: key to bacterial desiccation resistance? *Int J Syst Evol Microbiol* 2:393-403
- Friedmann, E.I. 1980. Endolithic microbial life in hot and cold deserts. *Origins of Life and Evolution of Biospheres*10, 223-235.
- Verseux, C. et al. 2017. Evaluation of the resistance of *Chroococcidiopsis* spp. to sparsely and densely ionizing irradiation. *Astrobiology* 17,118-125