

## Development of a regional glycerol dialkyl glycerol tetraether (GDGT) temperature calibration for Antarctic and sub-Antarctic lakes

Roberts Stephen<sup>1</sup>, Louise Foster<sup>1,2</sup>, Emma Pearson<sup>2</sup>, Steve Juggins<sup>2</sup>, Dominic Hodgson<sup>1</sup>, Krystyna Saunders<sup>3,4</sup> and Elie Verleyen<sup>5</sup>

<sup>1</sup> British Antarctic Survey, Natural Environment Research Council, High Cross, Madingley Road, Cambridge, CB3 0ET, UK  
E-mail: [sjro@bas.ac.uk](mailto:sjro@bas.ac.uk)

<sup>2</sup> School of Geography, Politics and Sociology, Newcastle University, Newcastle-upon-Tyne, NE1 7RU, UK

<sup>3</sup> Institute of Geography and the Oeschger Centre for Climate Change Research, University of Bern, 3012, Bern, Switzerland

<sup>4</sup> Australian Nuclear Science and Technology Organisation, New Illawarra Road, Lucas Heights, New South Wales 2234, Australia

<sup>5</sup> Gent University, Protistology and Aquatic Ecology, Krijgslaan 281 S8, 9000 Gent, Belgium

Modern-day temperature calibration models based on the relative abundances of Glycerol dialkyl glycerol tetraethers (GDGTs – temperature-sensitive membrane lipids of Archaea and bacteria), in surface sediments have been used to reconstruct past temperatures from marine and terrestrial sedimentary records, but have not been widely applied in high latitude terrestrial lacustrine environments. This is mainly because the presence, type and provenance of GDGT compounds in remains uncertain or unknown in many lacustrine settings, while the performance of GDGT-temperature calibrations at lower temperatures is comparatively poorly understood. To address these issues, we examined surface sediments from 38 Antarctic, sub-Antarctic and Southern Chilean lakes (Foster et al., 2016). First, we quantified which GDGT compounds were present and then investigated modern-day environmental controls on GDGT composition. GDGTs were found in 37 of 38 lakes studied. Branched GDGTs (brGDGTs) were dominant in 36 lakes and multivariate statistical analyses showed that their composition was strongly correlated with mean summer air temperature (MSAT) rather than pH, conductivity or water depth. Second, we developed the first regional brGDGT-temperature calibration for Antarctic and sub-Antarctic lakes based on four brGDGT compounds (GDGT-Ib, GDGT-II, GDGT-III and GDGT-IIIb). The GDGT-IIIb compound had not been included in previous global calibrations, but proved particularly important in cold lacustrine environments. Compared to previous global calibrations (Pearson et al., 2011), our new brGDGT-Antarctic temperature calibration dataset exhibited an improved statistical performance at low temperatures ( $r^2=0.83$ , RMSE=1.45°C, RMSEP-LOO=1.68°C, n=36 samples). Third, as a proof of concept test for the new Antarctic calibration model, we applied the new Antarctic brGDGT-temperature calibration to strategically-located and well-dated lake sediment records from the Antarctic Peninsula and South Georgia. Downcore temperature reconstructions using the Antarctic brGDGT-temperature calibration proved to be more sensitive to even relatively minor temperature variations compared to existing global calibrations, and each record reproduced known periods of mid-late Holocene warmth. In conclusion, our study highlights the importance of basing palaeotemperature reconstructions on regional GDGT-temperature calibrations for the Polar Regions, where the inclusion of additional location-specific compounds leads to improved model performance.

### References

- Foster, L.C., Pearson, E.J., Juggins, S., Hodgson, D.A., Saunders, K.M., Verleyen, E., Roberts, S.J. 2016. Development of a regional glycerol dialkyl glycerol tetraether (GDGT)-temperature calibration for Antarctic and sub-Antarctic lakes. *Earth and Planetary Science Letters* 433: 370–379. <http://dx.doi.org/10.1016/j.epsl.2015.11.018>
- Pearson, E.J., Juggins, S., Talbot, H.M., Weckstrom, J., Rosen, P., Ryves, D.B., Roberts, S.J., Schmidt, R., 2011. A lacustrine GDGT-temperature calibration from the Scandinavian Arctic to Antarctic: renewed potential for the application of GDGT-paleothermometry in lakes. *Geochim. Cosmochim. Acta* 75: 6225–6238. <http://dx.doi.org/10.1016/j.gca.2011.07.042>