



# Mycochemical study of polysaccharides from edible mushroom *Cortinarius caperatus* (Gypsy mushroom)

Zacchigna M.<sup>a</sup>, Altieri T.<sup>a</sup>, Beltrame G.<sup>a</sup>, Cateni F.<sup>a</sup>, Procida G.<sup>a</sup>

<sup>a</sup>Department of Chemical and Pharmaceutical Sciences, University of Trieste, P.zle Europa 1, 34127 Trieste, Italy  
E-mail: zacchigna@units.it



## INTRODUCTION

Among basidiomycete molecules, cell wall polysaccharides have been recognized as a major class of bioactive constituents. [1] They are safe molecules and they have a wide spectrum of biological activities, such as immunostimulatory and antioxidant, therefore they possess a prominent role in health benefits coming from mushroom consumption. These properties make mushroom polysaccharides potential candidates for nutraceutical applications and bioactive ingredients production. [2]

[1] - Zhang M. et al. (2007), Trends in Food Science & Technology 18, 4-19  
[2] - Zhu F. et al (2015), Journal of Food Composition and Analysis 41, 165-173

## STRUCTURAL ELUCIDATION OF ISOLATED POLYSACCHARIDES

Structural elucidation of isolated polysaccharides has been performed using mono- and bi-dimensional NMR spectroscopy (<sup>1</sup>H-NMR, <sup>13</sup>C-NMR, DEPT, <sup>1</sup>H-<sup>1</sup>H COSY, DQCOSY, TOCSY, HSQC, and HMBC), together with GC analysis of silylated monosaccharides and methylation analysis [4] to determine the monosaccharide composition and the linkage type, respectively.

[4] - Ciucanu I. & Kerek F. (1984), Carbohydrate Research, 131(2): 209-217




Fig. 4 HSQC of the branched  $\alpha,\beta$  glucan





Fig. 5 TOCSY of the branched  $\alpha,\beta$  glucan

## PURIFICATION PROCEDURES

### LIPIDIC FRACTION



### Soluble Polysaccharide




Fig. 1 <sup>1</sup>H-NMR spectrum of the soluble polysaccharide  $\beta$  1,6 glucan




Fig. 2 HMBC spectrum of the soluble polysaccharide  $\beta$  1,6 glucan

### Insoluble Polysaccharide





Fig. 3 <sup>1</sup>H-NMR of the branched  $\alpha,\beta$  glucan

Sugar linkage	Residue	C1	C2	C3	C4	C5	C6
6)- $\beta$ -(1-	A	103.40	73.64	76.06	70.01	75.40	69.33
6)- $\alpha$ -(1-	A'	100.70	73.55	--	70.10	--	68.76
$\alpha$ -(1-	B	98.37	--	--	69.11	--	61.25
4,6)- $\beta$ -(1-	C	103.40	--	--	76.14	--	67.03

## ANTIOXIDANT ACTIVITY




The antioxidant activity of the soluble polysaccharide fraction has been evaluated as radical-scavenging activity by DPPH assay and the  $\beta$ -(1 $\rightarrow$ 6)-D-glucan showed significant antioxidant activity.

## RESULTS

Fractionation of the hot aqueous extract of *Cortinarius caperatus* led to isolation of two polysaccharidic fractions characterized by spectroscopic analyses (<sup>1</sup>H-NMR, <sup>13</sup>C-NMR, DEPT, <sup>1</sup>H-<sup>1</sup>H COSY, DQCOSY, TOCSY, HSQC, HMBC and HMQC), mass spectrometry (EI-MS, ESI-MS), infrared spectroscopy (FT-IR), chemical reactions of hydrolysis and derivatization followed by GC and HPLC analyses. [5] The mycochemical study revealed a water-soluble fraction characterized as  $\beta$ -(1 $\rightarrow$ 6)-D-glucan, whose presence inside *C. caperatus* has never, to the best of our knowledge, been reported before. Moreover, a purified water insoluble fraction has been characterized as a branched  $\alpha,\beta$  glucan (Figures 3-5).

[5] - Ruthes A. C. et al. (2015), Carbohydrate Polymers 117, 753-761

## POLYSACCHARIDIC FRACTION<sup>[3]</sup>



[3] Smiderle F.R. et al. (2013), Carbohydrate Polymers, 94(1):91-99