

Journal of Environmental Monitoring. 2010. Vol 12, issue 5, 1159-1169.

Assessing multiple novel tracers to improve the understanding of the contribution of agricultural farm waste to diffuse water pollution

Granger, S. J.¹, Bol, R.¹, Dixon, L.¹, Naden, P.², Old, G.², Marsh, J. K.³, Bilotta, G.⁴, Brazier, R.⁵, White, S. M.⁶ and Haygarth, P. M.⁷

¹North Wyke Research, North Wyke, Okehampton, Devon EX20 2SN, UK.

²Centre for Ecology and Hydrology, Benson Lane, Crowmarsh Gifford, Wallingford, Oxfordshire OX10 8BB, UK.

³Environmental Tracing Systems Ltd, The Coach House, Bannachra, Helensburgh G84 9EF, UK.

⁴School of Environment and Technology, University of Brighton, Cockcroft Building, Brighton, East Sussex BN2 4GJ, UK.

⁵School of Geography, University of Exeter, Amory Building, Rennes Drive, Exeter, Devon EX4 4QJ, UK.

⁶Natural Resources Department, Cranfield University, Cranfield, Bedfordshire MK43 0AL, UK.

⁷Centre for Sustainable Water Management, University of Lancaster, Lancaster LA1 4YQ, UK.

Summary

A study was undertaken on drained and undrained 1ha grassland lysimeters to assess the effectiveness of multiple novel tracing techniques in understanding how agricultural slurry waste moves from land to water. Artificial fluorescent particles designed to mimic the size and density of organic slurry particles were found to move off the grassland via inter-flow (surface + lateral through flow) and drain-flow. Where both pathways were present the drains carried the greater number of particles. The results of the natural fluorescence and $\delta^{13}\text{C}$ of water samples were inconclusive. Natural fluorescence was higher from slurry amended lysimeters than from zero slurry lysimeters, however, a fluorescence decay experiment suggested that no slurry signal should be present given the time between slurry application and the onset of drainage. The $\delta^{13}\text{C}$ values of $>0.7\mu\text{m}$ and $<0.7\mu\text{m}$ material in drainage were varied and unrelated to Q. The mean value of $>0.7\mu\text{m}$ $\delta^{13}\text{C}$ in

water from the drain-flow pathways were higher from the lysimeter which had received naturally enriched maize slurry compared to the lysimeter which received grass slurry indicating a contribution of slurry derived material. Values of $<0.7\mu\text{m } \delta^{13}\text{C}$ from the same pathway however produced counter intuitive trends and may indicate that different fractions of the slurry have different $\delta^{13}\text{C}$ values.

Introduction

With the recent introduction of legislation and policy aimed at improving aquatic ecology and water quality in river basins (*e.g.* EU Water Framework Directive), attention is increasingly focusing on efforts to decrease inputs of pollutants into surface waters (*i.e.* rivers and lakes) and groundwater. Agricultural areas typically represent a large proportion of the land surface in many countries, and in Europe and North America this land often receives large quantities of agricultural amendments such as inorganic fertilizers or managed animal manure (*e.g.* farmyard manure (FYM) and slurry) which are applied to improve plant production. However with the intensification of farming, and subsequent increases in stocking densities, managed manures are often applied to land in excess. These organic amendments can have a significant impact if released into river systems as they contain high levels of nutrients, particularly nitrogen (N) and phosphorus (P) and have a high biological oxygen demand which can lead to a decrease in river oxygen levels (Hooda et al., 2000). Increases in plant nutrients in aquatic systems can lead to the eutrophication of an ecosystem with a host of associated detrimental effects (Conley et al., 2009). The application of these amendments to land has long been implicated by studies as a source of aquatic diffuse N and P pollution (Heathwaite, 1997; Smith et al., 1998; McDowell and Sharpley, 2001; Smith et al., 2002; McDowell et al., 2007). However, few studies actually provide evidence of a direct link between the organic sources of pollution applied to land and the occurrence of pollutants in water bodies (Haygarth et al., 2006). The potential to apply tracing techniques – particularly the simultaneous use of multiple tracers – to provide substantive evidence for the nature of this link at the field scale has previously been highlighted (Granger et al., 2007). To provide proof of concept, a tracer study was undertaken on 1ha intensively managed, drained and undrained, grassland lysimeters in the south west U.K. Three tracing techniques were chosen: i) natural abundance carbon (C) isotopes, ii) labelled synthetic particles, and iii) natural fluorescence.

It is believed that the use of stable C isotopes at natural abundance should be able to provide evidence for the movement of slurry derived C from land into drainage waters. A previous unreported study undertaken at the small plot scale had shown that the amount of slurry-derived organic C lost in drainage varied from 10 to 75%, with the maximum loss occurring shortly after slurry application during a rainfall event. Furthermore, slurry derived C continued to be detectable in drainage waters for over 2 months, long after pollutants had returned to pre-slurry levels (Bol, personal communication). The principles behind the stable C isotope tracing technique have been described in detail elsewhere (Amelung et al., 1999; Bol et al., 2000; Dungait et al., 2005). In short, all grasslands in the U.K. are C₃ systems ($\delta^{13}\text{C}$ about -26‰). Cattle over-wintered on silage produced from this herbage produce a slurry of similar enrichment while cattle over-wintered on a C₄ plant-based silage produce an isotopically higher slurry ($\delta^{13}\text{C}$ about -13‰) (Glaser et al., 2001). Thus, any increase in the $\delta^{13}\text{C}$ signal of C present in the drainage water leaving a C₃ lysimeter which receives C₄ slurry, when compared to a C₃ lysimeter which receives C₃ slurry, can be attributed to the presence of the C₄ slurry. A lysimeter receiving C₃ slurry is used as a control rather than a zero-slurry lysimeter to account for any differences in the soil biogeochemical processes due to the presence of slurry and it is assumed that the processes that affect both C₃ and C₄ slurries are the same. By examining the $\delta^{13}\text{C}$ of both particulates and filtrates in the drainage waters, this technique may be extended to provide evidence for differing loss processes.

The use of labelled synthetic particles has been previously used in maritime and terrestrial aquatic environments (Marsh et al., 1991; Marsh et al., 1993; McComb and Black, 2005). These particles typically comprise a polymer combined with a fluorescent dye manufactured with a density matched to that of the particles which are being mimicked. To an extent, similar size distributions may also be achieved by initial sieving, with further control over particle size being applied at the sample analysis phase. Thus particles of roughly equivalent size and density to particles typically present in slurry can be created and subsequently added to and mixed with slurry before its application to the

grassland lysimeters. The presence, or absence, of these particles in water draining the lysimeters is a direct indication that particulate material applied with the slurry is able to move, and has moved from the system.

Recent advances in fluorescence spectroscopy have enabled rapid automated collection of fluorescence intensity data across a wide range of excitation and emission wavelengths. The observed centres of fluorescence in river water samples correspond to those produced by known fluorophores, predominantly the amino acids tryptophan and tyrosine, and high molecular weight organic molecules, such as humic and fulvic acids. Hudson *et al.* (2007) provide an excellent review of both the phenomenon of natural fluorescence and its measurement. Farm wastes have also been analysed for their fluorescent properties and have been found to have very high intensities of tyrosine-like and tryptophan-like fluorescence and high ratios of these to fulvic/humic-like fluorescence compared to stream waters (Baker, 2002). This suggests that pollution of waters by these agricultural wastes could leave a distinct fluorescence signature reflecting both the source of the dissolved organic matter and its concentration.

This combination of tracers may therefore be able to provide both evidence of a direct connection between pollutant losses and organic amendments applied to agricultural grassland but also to discriminate between dissolved and particulate slurry fractions. The aim of this paper is to present the results from the field experiment in terms of both the evidence provided by the tracers for direct slurry losses into drainage waters and to provide an assessment of the tracing techniques used.

Experimental

Site

The study was sited on the field-scale lysimeters (1ha) of the Rowden experimental research platform (RERP) in Devon, Southwest England (National Grid Reference: SX 650995). The experimental site is maintained as permanent grassland and is grazed by

beef heifers at stocking densities of approximately four livestock units per hectare in order to manage sward height during the months of June to September. The soil is a clayey non-calcareous pelostagnogley of the Hallsworth series (Dystic Gleysol, FAO; HOST class 24 (Boorman et al., 1995)), overlying clay shales of the Crackington Formation (Culm measures) and the site slopes between 5 and 10% westwards. For details of soil properties see Scholefield et al (1993) and Armstrong and Garwood (1991). The annual rainfall at this site averages 1055mm, where the majority falls between October and March. As a consequence of the virtually impermeable clay layer below 30cm the soil remains waterlogged for much of the winter period. Half of the lysimeters at the RERP have been agriculturally drained while the other half has no drainage installed. Lysimeters in which no drainage has been installed are dependent upon natural drainage via surface-flow and lateral through-flow pathways, and are termed 'undrained' (Figure 5.1a). Lysimeters which have drainage installed are termed 'drained' and have mole drains at a depth of 55cm which are intercepted by gravel filled trenches with 85cm deep permanent pipe drains (Figure 5.1b). All the lysimeters have perimeter gravel filled ditches to a depth of 30cm which collect any surface runoff plus any lateral through-flow (combined referred to as inter-flow). The flow is channelled through 45° V-notch weirs. The drained lysimeters have additional and separate V-notch weirs for measuring water that flows through the drainage system (termed drain-flow) (Figure 5.1b). The stage height (h) of both inter-flow and drain-flow were measured using solar powered Starlevel flow sensors on a 5 minute time-step with data recorded by Campbell radio loggers and subsequently transmitted via radio modem to a central computer. To convert h to discharge (Q) stage-discharge relationships were produced for each weir, including estimates of the errors involved in the calibration (Bilotta et al., 2008). These were used to produce hydrographs with associated uncertainty bands (Q_{\max} and Q_{\min}) through a technique developed by Kreuger *et al* (2008). Rainfall at the site was measured using a tipping-bucket rain gauge (Rainwise, Bar Harbor, ME), which recorded the total number of tips min^{-1} (each tip equivalent to 0.254mm rainfall).




Figure 5.1. The hydrological pathways of the undrained (a) and drained (b) lysimeters at the Rowden Experimental Research Platform (adapted from Granger et al.(2008)). The location of the lysimeters used for each tracer application is presented in (c).

Experimental Design

The experiment began on the 18th April 2006. Prior to this, slurry was sourced from two local farms where cattle were over-wintered on a diet of either ryegrass (*Lolium perenne* L, a C₃ plant with a $\delta^{13}\text{C}$ value of about -26‰) or maize (*Zea mays* L, a C₄ plant with a $\delta^{13}\text{C}$ value of about -13‰) based silage. After collection and prior to application the slurries were stored in two slurry mini-stores so that they could be well mixed. Each store was used exclusively for the storage and mixing of either the C₃ or the C₄ slurry to avoid cross contamination.

Prior to mixing and application to the lysimeters, the C₄ slurry received a dose of fluorescent tracer particles. These were manufactured by Environmental Tracing Systems Ltd, Helensburgh, U.K. with a particle size distribution (PSD) based on the PSD of an initial test slurry (Figure 5.2a). This data showed a bimodal distribution with one peak between 27-37µm. The artificial fluorescent particles were manufactured to match this peak within this size range (Figure 5.2b) and a density of 1.15g cm⁻³ so as to be comparable with the organic particles within the slurry. The PSD data for the raw tracer particles (Figure 5.2b) show a relatively large proportion of very fine particles below 2µm. Given that these were not present in the slurry used as an initial guide, all particles below 2µm were discounted during the analysis of any samples. Excluding the <2µm particles, the D₅₀ of the tracer was about 12µm with the main peak between 16-19µm. Analysis of the two stock solutions showed that 4.4 x 10¹³ particles were applied per release. Mixing was accomplished through the pump and return of slurry out of, and back into the slurry towers using a conventional vacuum tanker. Tracer was then added to the slurry before further mixing was undertaken.

The experimental design comprised six hydrologically-isolated lysimeters at the RERP consisting of three drained and undrained pairs. Two pairs received an application of either C₃ or C₄ slurry while the third pair received nothing (Figure 5.1c). The maximum recommended application rate for UK grassland systems is 50m³ ha⁻¹, however, due to environmental constraints at the RERP at the time of application only 21m³ ha⁻¹ was applied. The C₃ and C₄ slurries were applied to the drained lysimeters on the 18th April, while application to the undrained lysimeters occurred one week later on the 25th April. Slurry was applied using a conventional vacuum tanker fitted with a splash plate in three applications starting with the third of the lysimeter furthest away from the access point. Slurry was remixed before each tanker load was removed from the slurry mini-stores. A 10m margin was left around the edge of each lysimeter in order to prevent contamination of the surface drains in accordance with the 'Code of Good Agricultural Practice' (Defra, 2002).




Figure 5.2. Particle size distributions of a) initial test dairy slurry, b) the raw artificial fluorescent tracer particles and c) the maize slurries applied to drained and undrained lysimeters after the addition of the tracer particles.

Sampling

Soil cores were collected from across the RERP to ascertain the $\delta^{13}\text{C}$ of the total soil C prior to slurry application. Five points within four of the lysimeters at the RERP were used which included two of the lysimeters used in this study. At each point 10 cores (2.5cm diameter) were taken to a depth of 7.5cm and these cores were then bulked.

Sub-samples of slurry were collected from each batch of slurry that was applied to each of the four lysimeters and stored in a refrigerator at 4°C until analysis.

Samples of drainage water were taken as follows: all V-notch weir systems were fitted with an internal stainless steel mixing plate below the drainage input to the weir. The drainage flowing through the mixing plate was then sampled on hourly or sub-hourly time-steps either using an automated sampler or through the collection of manual grab samples. Sampling of drainage water commenced after the first rainfall event to initiate

drainage from the lysimeters. This occurred on the 19th May 2006, some 31 days after slurry application (Fig. 3). A further two rainfall events were sampled, occurring on the 21st and 24th of May, and these three events are referred to as events 1, 2 and 3 respectively. Events 1 and 3 were sampled using automated samplers while event 2 was sampled manually by taking hourly grab samples. This was undertaken as during event 1 it was found that at low flows the auto-samplers were unable to collect sufficient sample for analysis. Event 3 was sufficiently large for resumption of automated sampling. Auto-samplers were set to sample using the same atomic clock time as the flow loggers. Samples were stored in a refrigerated environment in 1000ml polyethylene bottles.

In addition, temporal changes in the natural fluorescence signal of the slurry were monitored by applying slurry to a compartmented tray of a sterile medium (vermiculite) at the same rate (*i.e.* depth) as to the lysimeters. The use of a sterile medium was chosen so as not to introduce additional sources of fluorescence. There was no attempt to simulate soil processes. Four replicate sub-samples and two controls were collected each week and extracted using de-ionised water for analysis.

Analysis

Chemistry. Slurry samples were analysed for total solids by drying a pre-weighed sample in an oven at 100°C and subsequently re-weighing until a constant weight is obtained. Organic matter was determined through weight loss on ignition at 500°C of the dried slurry.

Within 24 hours of arrival at the laboratory, drainage waters had sub-samples for total phosphorus (TP) analysis transferred to polypropylene autoclavable bottles. Concentration of TP was then determined through an acid persulphate digestion using a method described by Eisenreich *et al.* (1975). Absorbance was measured colourimetrically on a spectrophotometer (Cecil Instruments, Cambridge, UK) calibrated by standards prepared fresh on the day of analysis. Concentrations of suspended solids (SS) and volatile solids (VS) were determined through vacuum filtration, and subsequent drying at 105°C

for at least 1 hour, of a known sample volume through a pre-weighed GF/F filter paper (Whatman), with particle size retention of 0.7 μ m. Filter papers were then heated to 500°C for at least 30 minutes to determine the total residual ash. The difference between SS and total ash is considered to represent VS (UK Standing Committee of Analysts, 1980).

Artificial fluorescent particles. Particle size distributions for both the raw tracer particles and tracer particles mixed with slurry were measured using a LISST 100 laser diffraction particle size analyser (Sequoia Scientific, USA). In both cases, ten analyses were undertaken and a mean result generated. Water samples collected from the lysimeters were agitated and sonified to disaggregate and suspend any solids and tracer particles. A measured volume, between 5 and 25ml depending on turbidity, of the sample was then filtered through a 0.45 μ m cellulose nitrate filter paper (Whatman). The filter paper was carefully transferred using forceps from the filter housing to a labelled microscope slide. Each slide was then analysed manually using a specially modified scanning fluorescent microscope (Leica, Germany). The total counts for the measured volume were then converted to tracer counts per ml (cts ml⁻¹). Replicate analyses were carried out on 15% of the samples analysed with the results indicating less than 9% error on all samples.

Prior to the release of the tracer particles some raw slurry samples and background water samples were collected from the RERP to determine whether there was any background fluorescence. While some weakly fluorescent particles were present in these samples, likely to be either inorganic minerals and/or biological material, they did not interfere with the very much brighter fluorescent yellow particles applied due to the different excitation and emission wavelengths.

Natural fluorescence. Sub-samples of drainage water were taken from the ISCO sample bottles in acid-washed brown glass bottles and kept in cool, dark conditions. Samples were analysed at room temperature and before analysis, samples were filtered using pre-ashed GF/C glass fibre filters (Whatman) with particle size retention of 1.2 μ m.

The fluorescence measurements were carried out on a Varian Cary Eclipse fluorescence spectrophotometer fitted with a Xenon flash lamp using slit widths of 5nm, an integration time of 12.5ms and voltage of 725V. Excitation wavelengths from 240 to 400nm (in steps of 5nm) and emission wavelengths from 280 to 500nm (in steps of 2nm) were used to generate excitation/emission matrices (EEM). The absorbance of each sample was measured in a 1cm cuvette on an Agilent 8453 spectrophotometer at 1nm intervals from 200 to 800nm. Samples with an absorbance value of $>0.02\text{cm}^{-1}$ at 650nm were omitted from the analysis as this was indicative of inefficient filtering. Samples which had an absorbance of $0-0.02\text{cm}^{-1}$ had their absorbance corrected using the long-wavelength scatter-correction method of Blough *et al* (1993). Raw fluorescence data was then corrected for lamp output and for instrument sensitivity following manufacturer's instructions (Haines, personal communication). Inner-filtering corrections, calculated from the corrected absorbance data using the method of Lackowicz (1983) were then applied. The fluorescence data were finally converted to Raman units using the method of Lawaetz and Stedmon (2009).

Following investigations using known mixtures of test solutions prepared with ultra-pure water and DL-Tyrosine, L-Tryptophan, IHSS Suwannee River Fulvic Acid Standard II and Humic Acid Standard II, indices representing tryptophan-like and fulvic/humic-like fluorescence were calculated as the mean fluorescence intensity over the following ranges:

Tryptophan Index (TI): Excitation 265-285nm/Emission 346-354nm

Fulvic/Humic Index (FI): Excitation 325-375nm/Emission 450-500nm.

This provides a ratio of TI:FI. This ratio is typically >2 for slurry and <1 in drainage waters in the absence of slurry (Naden *et al*, 2010). Thus, slurry losses should be identifiable in the enrichment of the TI:FI ratio. Where results are presented as means, two standard errors are presented in parenthesis.

Isotopic. Bulk soil cores were sieved through a 2mm sieve to remove large stones and organic fragments before being dried overnight at 30°C to a constant weight and ground into a fine powder. Slurry sub-samples were dried to a constant weight in an oven at 100°C before being milled to a fine powder. Drainage samples were collected and delivered to the laboratory within 24 hours of being sampled. They were then refrigerated at 5°C until ready for preparation. Samples were subject to vacuum filtration through pre-combusted (500°C), washed and dried 0.7µm glass fibre filters (GF/F, Whatman). The resultant filtrate was then frozen and freeze dried prior to analysis. Filter papers upon which >0.7µm particulate material was collected were dried in an oven at 105°C and stored individually in airtight bags and placed in a desiccator until ready for analysis.

Total C (TC) and ¹³C content was determined by dry combustion on a Carlo Erba NA 1500 elemental analyser connected to an automated continuous flow ANCA 20/20SL system (Europa, Crewe, UK). Prepared soil and slurry samples were weighed into 8 x 5mm tin capsules and combusted using 25ml of O₂. Samples of particulate material on glass fibre filter papers were cut to fit inside 10 x 10mm tin capsules and combusted in the elemental analyser via a large bore auto-sampler (AS128) using a 50ml oxygen loop. Despite the 10 x 10mm capsule being able to hold significantly more sample than an 8 x 5mm capsule, the actual amounts of sample C were still found to be low. To test the precision of the mass spectrometer with small amounts of C, a series of 63 flours (42.2% C; -26.4‰ δ¹³C) from 5mg down to 0.25mg (*i.e.* 5, 4, 3, 2, 1, 0.5, and 0.25) were used where each weight was weighed to within ± 5% of the target weight. Sample data was statistically adjusted based on the exponential curves generated by results up to 5mg flour. Freeze dried <0.7µm filtrates were also weighed into 10 x 10mm tin capsules and combusted using 50ml of O₂, however in this case sufficient sample C was supplied to not warrant statistical adjustment of the data.

Ground wheat flour referenced against CaCO₃ (NBS-19, Gaithersburg, MD, USA) was used as a calibration standard. The natural abundance values were expressed as δ values, which represents the ratios of ¹³C/¹²C relative to the international VPDB standard. The $\delta^{13}\text{C}$ values (per mil) are defined as:

$$\delta^{13}\text{C} = [(\text{atom } \% \text{ } ^{13}\text{C}_{\text{sample}} - \text{atom } \% \text{ } ^{13}\text{C}_{\text{VPDB}}) / \text{atom } \% \text{ } ^{13}\text{C}_{\text{VPDB}}] \times 1000$$

Internal quality controls were included with samples during analysis comprising no less than 10% of the total number of samples and drift correction was made against internal standards during the run. Replication of samples was typically better than 0.8‰ for GF/F particulate samples and 1‰ for freeze-dried filtrates with mean values of 0.1 (±0.2) and 0.1 (±0.5) respectively (two standard errors are presented in parenthesis).

Results and discussion

Hydrological response

Three storms were monitored during the period 19th- 26th May 2006 which generated hydrographs from the nine hydrological pathways contained within the six paired lysimeters. Due to a logger failure, discharge data from the zero slurry and maize slurry lysimeters was not recorded from 22nd May until the end of the study. The Q record from the grass slurry lysimeters was unaffected. To estimate the missing Q record for this time period Q has been taken from other lysimeters at the RERP site which have been shown to behave in a hydrologically similar fashion. These lysimeters were selected by visual comparison of the hydrographs and assessment of the monthly flow statistics recorded from all lysimeters compared to those with missing data over the period March – May 2006.

The amounts of rainfall leading to the Q events sampled were 13, 7 and 14mm occurring on the 19th, 21st and 24th May respectively. The previous month, during which

slurry had been applied to the lysimeters, had been unusually dry having received just 18mm rainfall compared to the mean April rainfall of 68mm. For the month of May, up to the 19th, 30mm of rainfall had fallen which was still lower than average for the time of year. By the time of drainage caused by event 1 on the 19th May occurred the soil moisture deficit (SMD) for the site, as calculated by the Penman equation (Monteith, 1980), was approximately 25mm. This deficit explains why a rainfall event of 13mm over 6 hours did not produce significant Q for event 1. By event 2 on the 21st May the SMD had been reduced to about 4mm, however the relatively small amount of rainfall that caused it also meant that Q was low from the lysimeters. However, by event 3 on the 24th May the soil had no SMD and the 14mm of rainfall that fell at the site caused Q to be significantly greater than event 1.

The hydrological response of the lysimeters is typical of this site and of other clay soil sites both with and without mole drainage (Trafford and Rycroft, 1973; Harris et al., 1984; Armstrong and Garwood, 1991). Discharge responds rapidly to the onset of rainfall forming characteristically peaky hydrographs, especially from the undrained lysimeters. Where mole drains have been installed, one could expect the drainage pathway to carry the greater proportion of storm water (Harris et al., 1984; Armstrong and Garwood, 1991). However, should the drainage system not operate efficiently through collapse of the moles the peaky drain-flow hydrograph would be attenuated while the inter-flow pathway would carry a greater percentage of the total drainage (Armstrong and Garwood, 1991). During this study, while drain-flow from the lysimeters remained peaky, the hydrographs are more smoothed in comparison to the inter-flow hydrographs. Furthermore, the volume of water carried by the drain-flow pathway was approximately equal to that of the inter-flow (Bilotta et al., 2008).

Chemical response

A summary of the chemical data for each lysimeter and hydrological pathway for the three monitored drainage events is presented in Table 5.1. This data shows that there was

considerable variation in TP concentrations between events, slurry treatments and drainage status of the lysimeters with no consistent treatment effect evident. The mean concentrations of TP from the zero slurry undrained lysimeters were 146, 540 and 247 $\mu\text{g l}^{-1}$ for events 1, 2 and 3 respectively. Mean concentrations from slurry amended undrained lysimeters were higher during events 1 and 3, but lower during event 2. Mean concentrations of TP from the drained lysimeters were always higher in the drain-flow than the inter-flow. Concentrations in both pathways were also found to be higher from the zero slurry lysimeter compared to both the slurry amended lysimeters during events 1 and 2 although during event 3 concentrations were higher in the drain-flow of both the slurry amended lysimeters.

Suspended sediment concentrations also showed cross-site variation with no distinct event or slurry treatment effects. Mean event concentrations from the zero slurry undrained lysimeters were 78, 185 and 51 mg l^{-1} while from the grass slurry amended lysimeter they were 44, 38 and 90 mg l^{-1} and from the maize slurry amended lysimeter 116, 78 and 45 mg l^{-1} . Mean concentrations from the drained lysimeters showed a similar lack of event or slurry effect although concentrations were typically higher in the inter-flow than in the drain-flow for any given lysimeter. Concentrations of VS also showed no distinct event or slurry treatment effect although as with SS mean concentrations tended to be slightly higher in the inter-flow than the drain-flow. The composition of the SS is typically mineral with maximum mean VS from the zero slurry lysimeter of 24% from the undrained and 26/24% from the drain-flow/inter-flow pathways of the drained lysimeters. The proportion of VS was generally higher in lysimeters which received slurry compared to those which did not for any given event, with the drain-flow of the drained lysimeters having a larger percentage of VS to SS compared to the inter-flow.

The highest concentrations of SS and TP typically coincide with or occur just before peak Q and then decrease with decreasing Q as previously described by Bilotta *et al.* (2008). The concentrations reported here are comparable with those reported previously during a more comprehensive study of SS and TP from zero slurry lysimeters at this site

(Bilotta et al., 2008). The number of significant correlations between SS and TP increase through events 1 to 3 but show no link to drainage status of the lysimeter, the hydrological pathway or the application of slurry (Table 5.1). During event 1 there were virtually no significant relationships while all pathways in event 3 show highly significant correlations. This perhaps indicates that the size of the drainage event maybe more important in SS and TP mobilisation rather than the drainage status of the grassland and that the soil, not the slurry, was the dominant source of SS and TP.

Artificial fluorescent particles

The PSD data for the slurries mixed with tracer is presented in Figure 5.2c and given the much smaller volume of tracer added to a large volume of slurry the PSD data essentially reflects the slurry used for each study. The data indicate the maize slurry used to mix with the fluorescent tracer was not the same as the slurry used as an initial guide. The maize slurry used for the undrained and drained plots show a mainly unimodal distribution with a peak around 60-80 μm and not a bi-modal distribution as the initial guide slurry had shown. The PSD data for the slurries applied to drained and undrained lysimeters show differences although given they were different batches and not collected at the same time this is perhaps not surprising. The fluorescent tracer particles used tended to represent the initial 40-50% of the slurry based on PSD data with more than 50% of the slurry consisting of particles larger than the tracer used. The PSD data of the slurry tracer mixture also shows that both have an increase in the number of particles $<2\mu\text{m}$, a feature not present in the initial test slurry. This is interpreted as purely a function of the addition of the tracer which has a high number of $<2\mu\text{m}$ particles, which were excluded from analysis.

The number of tracer particles lost from each pathway increased through events 1 to 3 (Table 5.1); the mean number of particles lost during each event from the inter-flow pathway of the undrained lysimeter was 1, 8 and 84cts ml^{-1} , while a comparable response is also present in the drain-flow of the drained lysimeter (2, 23 and 90cts ml^{-1}). While

there are no data for event 1 from the inter-flow pathway of the drained lysimeter, there is a marked increase in tracer particle numbers in event 3 compared with event 2, however the numbers of particles lost in events 2 and 3 are notably lower from this pathway than from the other two pathways (0 and 58cts ml⁻¹), particularly during event 2. During event 1 no relationship between tracer particles and Q was observed in the undrained lysimeter, however a significant positive relationship ($r_7 = 0.83$, $p < 0.01$) occurs in the drain-flow from the drained lysimeter. Similarly during event 2 no relationship was present between tracer particles and Q in the inter-flow from either lysimeter; although tracer particle numbers had increased compared to event 1 from the undrained lysimeter and were in turn greater than those from the drained lysimeter inter-flow pathway. No significant relationship was found between particle tracer number and Q from the drain-flow pathway during event 2; however the number of particles lost were greatest from this pathway during this event. As event 3 Q from these two lysimeters is interpolated Q and not true Q from these lysimeters no statistical analysis of Q v tracer particles has been undertaken, however a clear trend is apparent. From all three pathways particle cts ml⁻¹ appear to exhibit a positive relationship with Q with the maximum cts ml⁻¹ occurring at, or shortly after, the maximum Q (Figure 5.3). The increase in numbers is rapid on the rising limb of the hydrograph, while numbers decline more slowly on the falling limb, mirroring the nature of the Q itself.




Figure 5.3. The response of the fluorescent tracer particles to Q during the three monitored rainfall events.

Table 5.1. Summary of data obtained from RERP lysimeters during the three monitored events. Drainage status: UD = undrained, D = drained. Pathway: IF = inter-flow, DF = drain-flow. Correlations: ns = not significant, * = p<0.05, ** = p<0.01, * = p<0.001.**

	Event ID	Event 1									Event 2									Event 3										
		Treatment	Zero Slurry			Grass Slurry			Maize Slurry			Zero Slurry			Grass Slurry			Maize Slurry			Zero Slurry			Grass Slurry			Maize Slurry			
			Drainage	UD	D		UD	D		UD	D		UD	D		UD	D		UD	D		UD	D		UD	D		UD	D	
				Pathway	IF	DF	IF	IF	DF	IF	IF	DF	IF	IF	DF	IF	IF	DF	IF	IF	DF	IF	IF	DF	IF	IF	DF	IF	IF	DF
TP ($\mu\text{g l}^{-1}$)	Min	83	168	85	182	197	88	85	107		444	287	234	203	197	138	253	113	61	202	154	220	210	180		139	221	107		
	Mean	146	346	161	360	321	109	288	171		540	313	303	267	214	178	307	216	114	247	206	261	359	204		183	281	142		
	Max	269	516	359	680	615	140	439	283		707	353	355	305	234	207	348	316	236	398	255	326	818	237		307	601	185		
	n=	6	8	8	4	5	4	8	7		6	6	6	4	6	6	6	6	5	20	20	20	7	8		13	13	13		
SS (mg l^{-1})	Min	37	36	40	20	6	28	49	3		104	43	58	23	17	44	60	20	30	28	30	48	15	11		21	35	17		
	Mean	78	60	81	44	28	33	116	22		185	50	73	38	21	56	78	33	43	51	44	66	90	22		45	45	23		
	Max	303	108	292	75	95	45	154	41		361	72	104	54	26	71	97	49	68	121	86	121	271	38		128	66	33		
	n=	8	8	8	4	10	4	10	9		6	6	6	4	6	6	6	6	4	20	20	20	7	8		13	13	13		
Correlation SS v. TP	r=	-0.08	0.23	-0.32	0.96	0.04	0.82	-0.07	0.64		0.96	0.87	0.22	-0.04	0.22	0.89	0.94	0.94	1.00	0.97	0.86	0.82	0.96	0.88		0.99	0.75	0.71		
	p=	ns	ns	ns	*	ns	ns	ns	ns		**	*	ns	ns	ns	*	**	**	***	***	***	***	***	**		***	**	***		
VS (mg l^{-1})	Min	6	8	10	10	1	9	9	3		20	12	11	8	8	9	13	7	9	5	6	10	7	4		5	14	8		
	Mean	12	14	16	17	8	10	19	7		30	13	15	10	8	10	14	9	11	11	10	14	21	6		10	17	9		
	Max	34	19	48	31	25	14	26	13		61	14	20	15	9	15	15	15	19	15	20	54	9		20	27	12			
	n=	8	8	8	4	9	4	10	9		6	6	6	4	6	6	6	6	4	20	20	20	7	8		13	13	13		
% VS	Min	11	16	16	30	4	26	14	21		14	19	16	21	29	15	14	23	23	16	18	14	20	21		16	29	33		
	Mean	18	24	22	39	31	29	17	41		16	26	21	28	40	19	18	30	28	24	24	22	34	31		25	38	42		
	Max	25	32	26	49	43	31	21	100		19	29	25	42	51	21	21	33	32	40	31	29	47	40		33	47	67		
	n=	8	8	8	4	9	4	10	9		6	6	6	4	6	6	6	6	4	20	20	20	7	8		13	13	13		
Correlation SS v. VS	r=	0.99	0.59	1.00	0.95	0.86	0.96	0.86	0.84		0.97	0.62	0.65	0.68	-0.46	0.82	0.45	0.93	0.97	0.90	0.89	0.84	0.99	0.76		0.97	0.78	0.64		
	p=	***	ns	***	*	**	*	**	**		**	ns	ns	ns	ns	*	ns	**	*	***	***	***	***	*		***	**	*		
$\delta^{13}\text{C}$ ($>0.7\mu\text{m}$)	Min				-28.2	-27.8	-27.8	-28.9	-27.1					-27.3	-27.8	-27.2	-27.8	-26.8	-28.1				-28.8	-29.4		-29.1	-28.7	-28.3		
	Mean				-27.3	-27.0	-27.4	-28.3	-26.3					-27.1	-27.1	-26.6	-27.6	-26.5	-27.4				-28.2	-28.4		-27.7	-27.5	-27.6		
	Max				-27.0	-26.0	-27.1	-27.6	-25.5					-26.9	-26.7	-26.3	-27.2	-26.2	-26.9				-27.7	-27.8		-27.2	-27.1	-27.0		
	n=				4	9	4	8	6					4	6	6	6	6	6	4			7	8		12	13	13		
$\delta^{13}\text{C}$ ($<0.7\mu\text{m}$)	Min				-28.9	-26.1	-28.9	-30.5	-29.6					-28.7	-26.9	-28.4	-29.0	-28.0	-30.0				-29.5	-26.9		-28.7	-29.3	-28.8		
	Mean				-27.6	-21.9	-26.9	-27.7	-27.1					-25.6	-25.4	-28.0	-28.4	-27.5	-29.1				-29.0	-25.6		-27.3	-28.8	-27.9		
	Max				-24.9	-4.8	-24.4	-20.1	-20.9					-17.3	-22.4	-27.6	-27.6	-26.8	-28.2				-28.4	-21.7		-24.5	-28.1	-25.0		
	n=				4	9	3	8	6					4	6	6	6	6	4				7	8		12	13	13		
Tracer Particles (cts ml^{-1})	Min							0	0																	28	21	28		
	Mean							1	2																	84	90	58		
	Max							4	7																	135	129	92		
	n=							8	9																	13	13	13		
TI:FI	Min										0.36	0.40	0.45	0.70	0.48	0.39	0.40	0.57	0.50											
	Mean										0.37	0.41	0.49	0.80	0.53	0.47	0.41	0.61	0.59											
	Max										0.38	0.43	0.54	0.94	0.56	0.54	0.44	0.67	0.68											
	n=										5	6	6	4	6	6	6	5	5											

Natural fluorescence

It is important to analyse samples for natural fluorescence as quickly as possible after sampling as it is known that fluorescence may change over time and this may be related to many factors (*e.g.* microbial activity, oxygen availability, temperature in sampler) which makes the magnitude of change for any given sample very difficult to predict. Therefore the data presented here is restricted to storm 2 (Table 5.1.) for which the period between sampling to analysis was <3days.

Spatial sampling in a nearby first order catchment has shown that drainage waters not directly contaminated by farm waste have TI:FI ratios of 0.3 to 0.4; at the RERP subsequent sampling of non-slurry affected lysimeters gave mean TI:FI ratios of 0.45 (± 0.01). For the samples discussed here, the mean value of TI:FI in inter-flow from the zero slurry lysimeter was 0.37 (± 0.00) while mean values from the two lysimeters which received slurry were higher, one marginally so at 0.41 (± 0.01) while the other was much higher at 0.80 (± 0.06). Similarly mean values recorded in the inter-flow of the drained lysimeters were the same (0.47 (± 0.02)) or higher (0.59 (± 0.03)) from slurry-amended lysimeters compared to the zero slurry lysimeter (0.49 (± 0.1)). Mean values of TI:FI from the drain-flow appear to be more consistently affected by the presence of slurry; the value from the zero slurry lysimeter was 0.41 (± 0.00), while values from the two slurry-amended lysimeters were 0.53 (± 0.01) and 0.61 (± 0.02). This suggests some evidence for the loss of slurry-derived material in a dissolved phase during event 2. However, results from the tray experiment indicate that within 4-5 weeks much of the natural fluorescence signal from the slurry (TI and FI) has been lost and the results from the TI:FI are inconclusive.


Isotopic response

No significant difference in the $\delta^{13}\text{C}$ of the soil was found between lysimeters at the RERP and therefore a grand mean of -28.7‰ (± 0.2) is taken to represent the $\delta^{13}\text{C}$ of the soil at the site. Slurry was applied to the lysimeters at two distinct times, with the drained lysimeters receiving an application of grass and maize slurry one week before the undrained lysimeters. The $\delta^{13}\text{C}$ of the grass slurries were -28.2‰ (± 0.3) and -27.3‰ (± 0.1) applied to drained and undrained lysimeters respectively, while the maize slurries had $\delta^{13}\text{C}$ values of -18.9‰ (± 0.2) and -18.0‰ (± 0.2) having a higher $\delta^{13}\text{C}$ value of $\sim 9\text{‰}$ which was less than reported elsewhere (Dungait et al., 2005).

Summary values of $\delta^{13}\text{C}$ measured in material >0.7 and $<0.7\mu\text{m}$ leaving the lysimeters are presented in Table 5.1. There was no overall significant relationships between the $\delta^{13}\text{C}$ of the <0.7 and $>0.7\mu\text{m}$ fractions. Variation in $\delta^{13}\text{C}$ values are greatest in the $<0.7\mu\text{m}$ fraction compared to the $>0.7\mu\text{m}$ fraction and both fractions show no clear trends occurring within individual storm events. There were also no clear trends in the $\delta^{13}\text{C}$ of >0.7 and $<0.7\mu\text{m}$ fractions across the three events in the inter-flow of both the undrained and drained lysimeters. However, due to auto-sampler failure no data from event one on the maize slurry amended, and event 3 on the grass slurry amended drained lysimeters was collected from this pathway on the drained lysimeters which makes it impossible delineate trends. Values of $\delta^{13}\text{C}$ from the drain-flow pathway of the drained lysimeters do however show a consistent trend across all three events in both the grass and maize slurry amended lysimeters. Here values of $\delta^{13}\text{C}$ in both >0.7 and $<0.7\mu\text{m}$ material become lower with each subsequent event (Figure 5.4). The trend is clearest in the $>0.7\mu\text{m}$ material and less so in the $<0.7\mu\text{m}$ fraction, which are strongly influenced by a few samples which have a far higher $\delta^{13}\text{C}$ than the bulk of the data especially during event 1. This trend of lower values across subsequent storms does not appear to be affected by the slurry treatment.

To establish what slurry derived organic material is leaving the lysimeters a comparison between the $\delta^{13}\text{C}$ of lysimeters that received grass slurry and naturally enriched maize slurry should be made. To do this the mean value of $\delta^{13}\text{C}$ (± 2 standard errors) from the maize-slurry amended lysimeters during each event was taken and compared to the mean obtained from the grass-slurry amended lysimeters from the corresponding event and pathway (Figure 5.4). The >0.7 and $<0.7\mu\text{m}$ material obtained from the undrained lysimeter were similar in that although there were apparent differences between grass and maize slurry treatments, there was no specific direction to the trends. A more pronounced trend was observed in the $\delta^{13}\text{C}$ of material derived from the drain-flow of the drained lysimeter. The $\delta^{13}\text{C}$ values of the $>0.7\mu\text{m}$ fraction from the maize slurry lysimeter were always higher relative to that from the grass slurry lysimeter. This increase was only markedly different during events 2 and 3 with increases of 0.6‰ (± 0.20 Standard Error of the Difference (SED)) and 0.9‰ (± 0.16 SED) respectively. The reverse was true of the $\delta^{13}\text{C}$ values of the $<0.7\mu\text{m}$ material which were higher from the grass slurry lysimeter relative to the maize slurry lysimeter; 2.1‰ (± 0.68 SED) and 3.2‰ (± 0.37 SED) for events 2 and 3 respectively. Assuming that the increase in $\delta^{13}\text{C}$ of the $>0.7\mu\text{m}$ particles was caused as a result of maize

slurry material becoming entrained in the drain-flow it is estimated that between 6% (± 2)



and 10% (± 2) of this material originated from the slurry during events 2 and 3 respectively.

What is less clear is why an unexpected and counter intuitive difference in $\delta^{13}\text{C}$ values was present within the $<0.7\mu\text{m}$ fraction derived during events 2 and 3. We have no simple explanation for this behaviour. The above observation may point to the fact that: (i) the maize slurry was not homogeneously naturally enriched with the $<0.7\mu\text{m}$ fraction being less labelled than the particulate $<0.7\mu\text{m}$ fraction, (ii) the finer material of the surface applied $<0.7\mu\text{m}$ fraction is predominantly held in the soil matrix at the soil surface, hence what is found in the drains is soil derived material which might have a different $\delta^{13}\text{C}$ value between the two plots, (iii) the $<0.7\mu\text{m}$ material contained forms of C that were significantly different in their isotopic composition from that of C_3 and C_4 plant matter in the slurry; this material may have been sourced from either, or both, of the slurries themselves or from the soil/soil-water system, or (iv) the differential fractionation in the soil of the <0.7 and $>0.7\mu\text{m}$ material differs between maize and grass slurry (Bol et al., 1999). All of these tentative suggestions partly counter the assumption on which the natural abundance labelling technique is based (Balesdent and Mariotti, 1987) and require further investigation.

Figure 5.4. Means $\delta^{13}\text{C}$ (± 2 standard errors) of the <0.7 and $>0.7\mu\text{m}$ material in waters from the undrained and drained lysimeters which received an application of either grass or maize based slurry.

Analysis of the tracing approaches

The effectiveness of any tracer is wholly dependent on the degree to which it represents the material that is being traced. All the tracers being applied to this study are novel to some extent; natural fluorescence signals at the field lysimeter scale have not been widely studied and reported previously, artificial fluorescent particles have been used previously but not in the terrestrial environment, while natural abundance C isotopes have been used previously to assess the contribution of dung derived C to soil systems but not to aquatic ones nor at this scale. To evaluate fully the effectiveness of the tracers being used we need to reconsider the strengths and limitations of the individual approaches.

The artificial fluorescent particles appear to have been a success in terms of improving our understanding of the processes and pathways that organic material sourced from agricultural amendments follow. However, while these particles may mimic the size and density of a large fraction of slurry particles there are other aspects which remain uncertain such as the shapes of slurry particles and the nature of their surface charge (the particles being manufactured with a positive charge). This obviously has implications for the way such particles may flocculate and move (Droppo, 2001), however unpublished data indicates that the particles do adsorb natural charge and therefore behave similarly when looking at hydraulic and flocculation processes (J.K. Marsh, personal communication). Furthermore, the distribution and number of tracer particles mixed with the slurry may not represent the distribution and number of particles within the slurry, and counts of particles in drainage water were made irrespective of particle size. Therefore it is uncertain whether the sizes of particles yielded in drainage are representative of the initial tracer as a whole or whether one particular size range dominates. Additionally, to understand better the consequences of slurry particle movement more information is needed on what pollutants are associated with these particles.

While natural abundance C isotopes have been used as tracers with a great deal of success elsewhere (Amelung et al., 2008), results from this application of the technique were inconclusive. This may have resulted from the low initial slurry application reducing the strength of the maize slurry $\delta^{13}\text{C}$ signal, or may be a consequence of the long dry period between slurry application and drainage causing slurry-derived material to be (partly) decomposed or sorbed onto soil surfaces. Furthermore, the nature of the label within the C_4 slurry was uncertain, appearing to be only partly labelled with a δ value of -18‰ compared

to other C₄ slurries with δ values of about. -12‰ (Dungait et al., 2005). Unlike direct excreta, slurry is a complex composite of potentially many materials and while the 'particulates' in the slurry might be considered to represent pieces of undigested or partially digested C₄ plant material and therefore fully labelled, the colloidal and soluble components may represent material not sourced from the C₄ substrate and therefore not labelled to the same extent. These different size fractions are also subject to different processes and rates of decomposition (Fangueiro et al., 2008), with smaller organic particles being metabolised differently than larger ones. Other issues were methodological such as the separation of the >0.7 and $<0.7\mu\text{m}$ material. Collecting and analysing $>0.7\mu\text{m}$ material collected on GF/C papers severely restricts the mass of actual particulate material available to the MS and although increasing the volume of sample filtered would have increased the amount of particulate C on the filters, they would become rapidly clogged leading to large sample preparation times and the retention of material $<0.7\mu\text{m}$. Furthermore, the freeze dried $<0.7\mu\text{m}$ material represented a composite of all colloids and solutes $<0.7\mu\text{m}$, thus may have contained a component of inorganic C, which if present may have increased the $\delta^{13}\text{C}$ values.

Natural fluorescence has potential to act as an indicator for slurry-derived material however the nature of the signal is time dependent and probably only persists for 3-4 weeks after slurry application. Thus, only direct losses may be traced using natural fluorescence rather than any longer term impacts. There is also a requirement for the samples to be analysed quickly after sampling which may also limit the potential applications of the technique.

Conclusions

The artificial fluorescent particles show that organic slurry particles have the potential to move readily from land to water and numbers increase and decrease with Q. Particles were lost via the inter-flow pathway of both the undrained and drain lysimeters and where both inter-flow and drain-flow pathways were present the drain-flow carried the greater number.

Natural fluorescence TI:FI ratios were inconclusive. Although ratios were higher in slurry amended lysimeters when compared to the zero slurry lysimeter, given the time interval between the application of slurry and the monitored runoff it is likely that most of the natural fluorescence signal from the slurry would have been lost.

The data obtained from the stable C isotope tracing approach were also inconclusive. The $\delta^{13}\text{C}$ of both particulate and dissolved phases were highly variable over time and only the particulate material in the drain-flow appeared to carry a slurry signature.

While many questions have been raised concerning the interpretation of the evidence from individual tracers, the simultaneous use of multiple tracers has been useful in helping to corroborate the available evidence. One of the main limitations in the experiment was the low application rate of slurry and further work needs to be carried out using a higher rate of application. Another limitation was the length of time between the applied slurry and the first rainfall event which meant that the use of some of the tracers was limited. There is a need to follow this up by more controlled experiments looking specifically at the mechanisms behind the tracers and the environmental processes through which they are changed.

Acknowledgments

The work was supported through a Department of Environment, Fisheries and Rural Affairs (Defra) grant (PE0120). The first author acknowledges the support of Cranfield University and North Wyke Research through his PhD studentship award as well as advice from Dr. Phil Owen of the University of Northern British Columbia. North Wyke Research is grant aided by the Biotechnology and Biological Sciences Research Council (BBSRC). We acknowledge the work of Tobias Krueger, Jim Freer, John Quinton (University of Lancaster), and Kit Macleod (N. Wyke) in establishing the flow measurements and uncertainties for the RERP.

References

- AMELUNG, W., BOL, R. & FRIEDRICH, C. 1999. Natural ^{13}C abundance: a tool to trace the incorporation of dung-derived carbon into soil particle-size fractions. *Rapid Communications in Mass Spectrometry*, 13, 1291-1294.
- AMELUNG, W., BRODOWSKI, S., SANDHAGE-HOFMANN, A. & BOL, R. 2008. Combining biomarker with stable isotope analysis for assessing the transformation and turnover of soil organic matter. *Advances in Agronomy*, 100, 155-250.
- ARMSTRONG, A. C. & GARWOOD, E. A. 1991. Hydrological consequences of artificial drainage of grassland. *Hydrological Processes*, 5, 157-174.

- BAKER, A. 2002. Fluorescence properties of some farm wastes: implications for water quality monitoring. *Water Research*, 36, 189-195.
- BALESDENT, J. & MARIOTTI, A. 1987. Natural ^{13}C abundance as a tracer for studies of soil organic matter dynamics. *Soil Biology and Biochemistry*, 19, 25-30.
- BILOTTA, G. S., BRAZIER, R. E., HAYGARTH, P. M., MACLEOD, C. J. A., BUTLER, P., GRANGER, S., KRUEGER, T., FREER, J. & QUINTON, J. 2008. Rethinking the contribution of drained and undrained grasslands to sediment-related water quality problems. *Journal of Environmental Quality*, 37, 906-914.
- BLOUGH, N. V., ZAFIRIOU, O. C. & BONILLA, J. 1993. Optical absorption spectra of waters from the Orinoco River outflow: terrestrial input of coloured organic matter to the Caribbean. *Journal of Geophysical Research - Oceans*, 98, 2271-2278.
- BOL, R., AMELUNG, W., FRIEDRICH, C., LAUF, J. & SHEPHARD, M. Year. The isotopic fractionation of dung-derived C in water extracts from a temperate grassland soil. *In: Arbeitsgemeinschaft Stabile Isotope*, 4-6 October 1999 1999 Göttingen, Germany. 70.
- BOL, R., AMELUNG, W., FRIEDRICH, C. & OSTLE, N. 2000. Tracing dung-derived carbon in temperate grassland using ^{13}C natural abundance measurements. *Soil Biology and Biochemistry*, 32, 1337-1343.
- BOORMAN, D. B., HOLLIS, J. M. & LILLY, A. 1995. Hydrology of soil types: a hydrologically based classification of the soils of the United Kingdom. Wallingford, Oxfordshire: Institute of Hydrology report No.126.
- CONLEY, D. J., PAERL, H. W., HOWARTH, R. W., BOESCH, D. F., SEITZINGER, S. P., HAVENS, K. E., LANCELOT, C. & LIKENS, G. E. 2009. Controlling eutrophication: nitrogen and phosphorus. *Ecology*, 323, 1014-1015.
- DEFRA 2002. *Code of good agricultural practice for the protection of water*, London, Defra Publications.
- DROPPO, I. G. 2001. Rethinking what constitutes suspended sediment. *Hydrological Processes*, 15, 1551-1564.
- DUNGAIT, J. A. J., BOL, R. & EVERSLED, R. P. 2005. Quantification of dung carbon incorporation in a temperate grassland soil following spring application using bulk stable carbon isotope determinations. *Isotopes in Environmental and Health Studies*, 41, 3-11.
- EISENREICH, S. J., BANNERMAN, R. T. & ARMSTRONG, D. E. 1975. A simplified phosphorus analysis technique. *Environmental Research Letters*, 9, 43-53.

- FANGUEIRO, D., CHADWICK, D. & BOL, R. 2008. Assessment of the potential N mineralization of different particle-size fractions in two dairy cattle slurries. *Journal of Plant Nutrition and Soil Science*, 171, 313-315.
- GLASER, B., BOL, R., PREEDY, N., MCTIERNAN, K., CLARK, M. & AMELUNG, W. 2001. Short-term sequestration of slurry-derived carbon and nitrogen in temperate grassland soil as assessed by ^{13}C and ^{15}N natural abundance measurements. *Journal of Plant Nutrition and Soil Science*, 164, 467-474.
- GRANGER, S. J., BOL, R., BULTER, P. J., HAYGARTH, P. M., NADEN, P., OLD, G., OWENS, P. N. & SMITH, B. P. G. 2007. Processes affecting transfer of sediment and colloids, with associated phosphorus, from intensively farmed grasslands: tracing sediment and organic matter. *Hydrological Processes*, 21, 417-422.
- HARRIS, G. L., GOSS, M. J., DOWDELL, R. J., HOWSE, K. R. & MORGAN, P. 1984. A study of mole drainage with simplified cultivation for autumn-sown crop on a clay soil. *Journal of Agricultural Science, Cambridge*, 102, 561-581.
- HAYGARTH, P. M., BILOTTA, G. S., BOL, R., BRAZIER, R. E., BUTLER, P. J., FREER, J., GIMBERT, L. J., GRANGER, S. J., KRUEGER, T., MACLEOD, C. J. A., NADEN, P., OLD, G., QUINTON, J. N., SMITH, B. & WORSFOLD, P. 2006. Processes affecting transfer of sediment and colloids, with associated phosphorus, from intensively farmed grasslands: an overview of key issues. *Hydrological Processes*, 20, 4407-4413.
- HEATHWAITE, A. L. 1997. Sources and pathways of phosphorus loss from agriculture. In: TUNNEY, H., CARTON, O. T., BROOKES, P. C. & JOHNSON, A. E. (eds.) *Phosphorus Loss from Soil to Water*.: CAB International. Wallingford, UK.
- HOODA, P. S., EDWARDS, A. C., ANDERSON, H. A. & MILLER, A. 2000. A review of water quality concerns in livestock farming areas. *The Science of the Total Environment*, 250, 143-167.
- HUDSON, N., BAKER, A. & REYNOLDS, D. 2007. Fluorescence analysis of dissolved organic matter in natural, waste and polluted waters - a review. *River Research and Applications*, 23, 631-649.
- KRUEGER, T., QUINTON, J. N., FREER, J., MACLEOD, C. J. A., BILOTTA, G. S., BRAZIER, R. E., BUTLER, P. & HAYGARTH, P. M. 2008. Uncertainties in data and models to describe event dynamics of agricultural sediment and phosphorus transfer. *Journal of Environmental Quality*, 38, 1137-1148.
- LAKOWICZ, J. R. 1983. *Principles of Fluorescence Spectroscopy*, New York, Plenum Press.
- LAWAETZ, A. J. & STEDMON, C. A. 2009. Fluorescence intensity calibration using the Raman scatter peak of water. *Applied Spectroscopy*, 63, 936-940.

- NADEN, P. S., OLD, G. H., ELIOT-LIAZE, C., GRANGER, S. J., HAWKINS, J. M. B., BOL, R., HAYGARTH, P. M. 2010. Assessment of natural fluorescence as a tracer of diffuse agricultural pollution from slurry spreading on intensely-farmed grasslands. *Water Research*, 44, 1701-1712.
- MARSH, J. K., BALE, A. J., UNCLES, R. J. & DYER, K. R. 1991. A tracer technique for the study of suspended sediment dynamics in aquatic environments. *Proceedings of the International Symposium on the Transport of Suspended Sediments and its Mathematical Modelling*. Florence, Italy, 2-5 September: International Association of Hydraulic Research.
- MARSH, J. K., BALE, A. J., UNCLES, R. J. & DYER, K. R. 1993. Particle tracing experiment in a small shallow lake: Loe Pool, UK. In: MCMANUS, J. & DUCK, R. W. (eds.) *Geomorphology and Sedimentology of Lakes and Reservoirs*. Chichester: John Wiley and Sons Ltd.
- MCCOMB, P. & BLACK, K. 2005. Detailed observations of littoral transport using artificial sediment tracer, in a high energy, rocky reef and iron sand environment. *Journal of Coastal Research*, 21, 358-373.
- MCDOWELL, R. W., NASH, D. M. & ROBERTSON, F. 2007. Sources of phosphorus lost from a grazed pasture receiving simulated rainfall. *Journal of Environmental Quality*, 36, 1281-1288.
- MCDOWELL, R. W. & SHARPLEY, A. N. 2001. Phosphorus losses in subsurface flow before and after manure application to intensively farmed land. *The Science of the Total Environment*, 278, 113-125.
- MONTEITH, J. L. 1980. The development and extension of Penman's Evaporation formula. In: HILLEL, D. (ed.) *Applications of Soil Physics II*. New York: Academic Press Incorporated.
- SCHOLEFIELD, D., TYSON, K. C., GARWOOD, E. A., ARMSTRONG, A. C., HAWKINS, J. & STONE, A. C. 1993. Nitrate leaching from grazed lysimeters: effects of fertilizer input, field drainage, age of sward and patterns of weather. *Journal of Soil Science*, 44, 601-613.
- SMITH, K. A., BECKWITH, C. P., CHALMERS, A. G. & JACKSON, D. R. 2002. Nitrate leaching following autumn and winter application of animal manures to grassland. *Soil Use and Management*, 18, 428-434.
- SMITH, K. A., CHALMERS, A. G., CHAMBERS, B. J. & CHRISTIE, P. 1998. Organic manure phosphorus accumulation, mobility and management. *Soil Use and Management*, 14, 154-159.
- TRAFFORD, B. D. & RYCROFT, D. W. 1973. Observations on the soil-water regimes in a drained clay soil. *Journal of Soil Science*, 24, 380-391.
- UK STANDING COMMITTEE OF ANALYSTS 1980. *Suspended, settleable and total dissolved solids in waters and effluents*, London, HMSO.

