

Physical processes controlling the spatial distributions of relative humidity in the tropical tropopause layer over the Pacific

Eric J. Jensen, NASA Ames Research Center
Troy D. Thornberry, NOAA Earth System Research Laboratory
Andrew W. Rollins, NOAA Earth System Research Laboratory
Rei Ueyama, NASA Ames Research Center
Leonhard Pfister, NASA Ames Research Center
Thaopaul Bui, NASA Ames Research Center
Glenn S. Diskin, NASA Langley Research Center
Joshua P. DiGangi, NASA Langley Research Center
Eric Hints, NOAA Earth System Research Laboratory
Ru-Shan Gao, NOAA Earth System Research Laboratory
Sarah Woods, SPEC, Inc.
R. Paul Lawson, SPEC, Inc.
Jasna Pittman, Harvard University

Abstract

The spatial distribution of relative humidity with respect to ice (RHI) in the boreal wintertime tropical tropopause layer (TTL, $\approx 14\text{--}18$ km) over the Pacific is examined with the measurements provided by the NASA Airborne Tropical Tropopause EXperiment. We also compare the measured RHI distributions with results from a transport and microphysical model driven by meteorological analysis fields. Notable features in the distribution of RHI versus temperature and longitude include (1) the common occurrence of RHI values near ice saturation over the western Pacific in the lower to middle TTL; (2) low RHI values in the lower TTL over the central and eastern Pacific; (3) common occurrence of RHI values following a constant mixing ratio in the middle to upper TTL (temperatures between 190 and 200 K); (4) RHI values typically near ice saturation in the coldest airmasses sampled; and (5) RHI values typically near 100% across the TTL temperature range in air parcels with ozone mixing ratios less than 50 ppbv. We suggest that the typically saturated air in the lower TTL over the western Pacific is likely driven by a combination of the frequent occurrence of deep convection and the predominance of rising motion in this region. The nearly constant water vapor mixing ratios in the middle to upper TTL likely result from the combination of slow ascent (resulting in long residence times) and wave-driven temperature variability. The numerical simulations generally reproduce the observed RHI distribution features, and sensitivity tests further emphasize the strong influence of convective input and vertical motions on TTL relative humidity.

Journal Geophysical Research Atmosphere