

Flux Cancellation: The Key to Solar Eruptions

Abstract


Solar coronal jets are magnetically channeled eruptions that occur in all types of solar environments (e.g. active regions, quiet-Sun regions and coronal holes). Recent studies show that coronal jets are driven by the eruption of small-scale filaments (*minifilaments*). Once the eruption is underway magnetic reconnection evidently makes the jet spire and the bright emission in the jet base. However, the triggering mechanism of these eruptions and the formation mechanism of the pre-jet minifilaments are still open questions. In this talk, mainly using SDO/AIA and SDO/HMI data, first I will address the question: what triggers the jet-driving minifilament eruptions in different solar environments (coronal holes, quiet regions, active regions)? Then I will talk about the magnetic field evolution that produces the pre-jet minifilaments. By examining pre-jet evolutionary changes in line-of-sight HMI magnetograms while examining concurrent EUV images of coronal and transition-region emission, we find clear evidence that flux cancellation is the main process that builds pre-jet minifilaments, and is also the main process that triggers the eruptions. I will also present results from our ongoing work indicating that jet-driving minifilament eruptions are analogous to larger-scale filament eruptions that make flares and CMEs. We find that persistent flux cancellation at the neutral line of large-scale filaments often triggers their eruptions. From our observations we infer that flux cancellation is the fundamental process for the buildup and triggering of solar eruptions of all sizes.

Flux Cancellation: The Key to Solar Eruptions

Navdeep K. Panesar

(NASA, Marshall Space Flight Center, Huntsville, AL)



Alphonse Sterling (NASA), Ronald Moore (NASA), Prithi Chakrapani (Hunter College),
Davina Innes (MPS), Don Schmit (LMSAL), Sanjiv Tiwari (LMSAL)



OUTLINE


- Background
- Triggering of Jet-Driving Minifilament Eruptions
- Formation of Pre-Jet Minifilaments
- Triggering of CME-producing Filament Eruptions
- Summary

Background




- Coronal jets are frequent magnetically channeled narrow eruptions. They occur in various solar environments: quiet regions, coronal holes and active regions.
- They are relatively short-lived features (of about 10 minutes; *Shimojo et al 1996*, *Savcheva et al 2007*) occur at a rate of ~60 per day in polar coronal holes (*Savcheva et al 2007*).

standard jet



blowout jet




Moore et al 2010

- All coronal jets observed in EUV and X-ray images show a bright spire with a base brightening, also known as jet bright point (JBP). X-ray jets were first detailed study with Yohkoh satellite (*Shibata et al 1992*), later they were studied with Hinode satellite (*Savcheva et al 2007, Cirtain et al 2007*).
- There are two types of jets (*Moore et al 2010, 2013*): standard jets and blowout jets.

Solar Jet Models


- Some studies suggested that flux emergence may lead to the jet eruptions (e.g. *Shibata et al. 1992, 2007, Moreno-Insertis et al 2008*).
- Recent studies show that coronal jets are driven by small-scale filament eruptions (e.g. *Hong et al. 2011, Shen et al. 2012, Adams et al. 2014, Sterling et al 2015*).
- Sterling et al. 2015* did extensive study of 20 polar coronal hole jets and found that X-ray jets are mainly driven by the eruption of *minifilaments*.
- What leads to these minifilament eruptions?**
- How and when are minifilaments formed?**

Emerging-flux jet model



Shibata et al 1992, 2007

Minifilament-eruption jet model



Sterling et al 2015

(I) Triggering of Pre-Jet Minifilament Eruptions

Quiet region jets

Coronal hole jets

Active region jets


Quiet region jets

- We examined the magnetic cause of 10 random on-disk quiet region jet eruptions by using SDO/HMI magnetograms and SDO/AIA images.

Measured parameters for the observed quiet-region jets:



Event No.	Date	Time ^a (UT)	Location ^b x,y (arcsec)	Jet Speed ^c (km s ⁻¹)	Jet Dur. ^d min.	Jet-base ^e Width (km)	Minifil. length ^f (± 1700 km)	Φ values ^g 10 ¹⁹ Mx	% of Φ^h reduction
J1	2012 Mar 22	04:46	-470,-100	100 \pm 30	15 \pm 5	10500 \pm 500	9800	1.6	52 \pm 5.8
J2	2012 Jul 01	08:32	-44, 285	100 \pm 10	10 \pm 2	27000 \pm 500	25000	4.0	18 \pm 6.8
J3	2012 Jul 07	21:31	-192,-180	120 \pm 15	14 \pm 3	16500 \pm 400	10500	- ^j	-
J4	2012 Aug 05 ^l	02:20	-485, 190	140 \pm 35	10 \pm 3	22000 \pm 1000	31000	1.5	21 \pm 6.0
J5	2012 Aug 10	23:03	-168,-443	125 \pm 15	15 \pm 2	16000 \pm 400	10000	0.9	57 \pm 5.4
J6	2012 Sept 20	22:56	-158,-486	35 \pm 5	9 \pm 2	20000 \pm 500	36000	2.0	23 \pm 4.6
J7	2012 Sept 21	03:33	-115, -485	135 \pm 30	12 \pm 1	17500 \pm 500	15000	1.0	36 \pm 7.2
J8	2012 Sept 22	01:25	-338, 103	110 \pm 45	11 \pm 1	13000 \pm 600	5700	0.9	50 \pm 5.1
J9	2012 Nov 13	04:21	-28,-307	55 \pm 5	9 \pm 3	18000 \pm 1000	25000	1.7	34 \pm 3.2
J10	2012 Dec 13	10:36	26, 50	65 \pm 20	10 \pm 2	9500 \pm 500	12500	1.2	38 \pm 5.0

Quiet region jet (J7)






- A minifilament (length ~ 15000 km) is present in the jet-base region prior to jet eruption.
- It resides over the neutral line between the opposite-polarity flux patches.
- The JBP occurs at the pre-eruption location of the minifilament.
- The jet spire extends upward with an average speed of 135 ± 30 kms^{-1} .

Quiet region jet (J7)





- The minifilament was present at the neutral line for 34 hours before the jet eruption.
- The jet-producing eruptions and JBPs are similar to typical solar flare eruption, in which a flare arcade grows over the neutral line in the wake of the filament.

Flux cancellation leading to minifilament eruption



- Both polarities approach towards the neutral line, and eventually cancel with each other just before the eruption. Flux cancellation continued until the minority-polarity flux patch completely disappeared.
- We find in each of the ten jets that opposite polarity magnetic flux patches converge and cancel, with a flux reduction of 20-60 % until jet erupts.

Quiet region jet (J9)



- The minifilaments show a slow-rise, followed by a fast-rise as they erupt, analogous to larger-scale filament eruptions.

(I) Triggering of Pre-Jet Minifilament Eruptions

Quiet region jets


Coronal hole jets

Active region jets

Coronal Hole Jets



Event No.	Date	Time ^a (UT)	Location ^b Helio. Cord.	Jet Speed ^c (km s ⁻¹)	Jet Dur. ^d (minutes)	Jet-Base ^e Width (km)	XRT ^f Coverage	Φ values ^g 10 ¹⁹ Mx	% of Φ^h Reduction
J1	2012 Jul 02	02:11	N21, W07	65±1.5	10±2	22500±1500	No	1.5	33±4.5
J2	2015 Aug 18	13:07	N27, E02	40±20	12±3	9700±1000	Yes	0.9	31±10
J3	2015 Dec 28	09:54	N36, E19	110±40	7±1	13000±900	No	0.9	48 ±8.8
J4	2015 Dec 28	16:02	N37, E03	35±7	11±1	12000±2500	No	1.2	52±4.8
J5	2015 Dec 30	15:14	N36, W23	70±30	7±1	6700±1000	No	0.7	43 ±10
J6	2015 Dec 31	19:04	N43, W34	27±4	6±1	6600±500	No	0.6	41 ±8.3
J7	2016 Jan 01	11:45	N08, E30	30±5	8±1	18500±3500	No	0.7	52 ±6.6
J8	2016 Jan 01	18:11	N41, E39	204±70	4±1	1200±500	No	0.5	37 ±9.8
J9	2016 Apr 21	06:15	S01, E12	240±70	8±1 ⁱ	10500±700	Yes	- ^j	-
J10	2016 Sep 15	23:36	S06, E00	- ^k	6±1 ^l	19000±2000	Yes	- ^m	-
J11	2017 Jan 03	14:59	N20, E02	105±30	6±1	18700±6000	Yes	1.6	33±5.3
J12	2017 Jan 04	09:26	N24, W03	92±30	15±3	9500±1000	Yes	2.0	73 ±5.0
J13	2017 Jan 04	17:08	N19, W10	103±30	4±1	7000±500	Yes	0.9	21±8.0

Coronal hole jet (J11)






- A minifilament resides (1 hour before the eruption) over the neutral line between the opposite-polarity flux patches.
- The JBP occurs at the pre-eruption location of the minifilament.
- The jet spire extends upward with an average speed of $105 \pm 30 \text{ km s}^{-1}$.

Coronal hole jet (J11)





Flux cancellation leading to minifilament eruption




- The positive flux continuously decreases with time, which is clear evidence of flux cancellation at the neutral line of the minifilament.
- HMI time-distance map shows the convergence and cancellation of the jet-base polarities.
- We find in each of the 13 jets that opposite polarity magnetic flux patches converge and cancel, with a flux reduction of 20-75 % until jet erupts.

Coronal hole jet (J1)



Schematic Illustration of Observations



- The minifilament (blue) initially resides in sheared/twisted field between patches of majority (positive) and minority (negative) flux.
- These two flux patches converge and cancel with each other. Continuous flux cancellation at the neutral line eventually destabilizes the filament field to erupt outwards and undergo external reconnection with the surrounding coronal field.
- The external reconnection opens the erupting closed field, allowing hot reconnection-heated material and cool minifilament material to escape along the far-reaching field as the jet spire.


(I) Triggering of Pre-Jet Minifilament Eruptions

Quiet region jets

Coronal hole jets


Active region jets

Active Region Jets




Active Region Jet

2015-01-14 14:03:11




AIA 171 movie

2015-01-14 13:02:39





HMI movie


Flux cancellation leading to the jet eruption



Box (b1) Positive Flux



AIA 94 Intensity




(II) Minifilament Formation


Quiet Region Minifilament Formation

Event No.	Minifil. formation ^a time (UT)	Minifil. eruption ^b time (UT)	Location ^c helio. cord.	Duration of ^c minifil. (hrs)	Width of ^d minifil. (km)	No. of ^e Jets	Φ values ^f 10^{19} Mx	% of Φ ^g reduction
J1	2012 Mar 21 22:46	2012 Mar 22 04:46	S09, E29	6	2000±500	1	1.6	20 ± 6.8
J2	2012 Jul 01 05:58	2012 Jul 01 08:29	N12, E02	2.5	1500±200	1	1.9 ^h	20 ± 7.3
J3	2012 Jul 07 — ⁱ	2012 Jul 07 21:31	S15, E12	–	2200±200	1	–	–
J4	2012 Aug 04 05:14	2012 Aug 05 01:58 ^j , 2012 Aug 05 02:20	N07, E30	21	2500±500	2	5.8	14 ± 4.6
J5	2012 Aug 10 19:43	2012 Aug 10 23:03	S31, E11	3.2	1500±200	1	0.9	27 ± 6.1
J6	2012 Sept 19 17:15	2012 Sept 20 22:52	S34, E11	34	2500±500	2	3.0	9 ± 5.3
J7	2012 Sept 21 00:51	2012 Sept 21 03:33	S34, E08	3.5	2500±500	1	1.7	38 ± 2.6
J8	2012 Sept 21 23:55	2012 Sept 22 01:25	N01, E20	1.5	1500±500	1	0.9	38 ± 5.5
J9	2012 Nov 11 02:56	2012 Nov 11 13:08, 2012 Nov 12 17:06, 2012 Nov 12 21:34, 2012 Nov 13 04:20	S23, E01	49.5	2500±500	4	– ^k	–
J10	2012 Dec 13 08:06	2012 Dec 13 10:11, 2012 Dec 13 10:36	S01, W01	2.5	1600±200	2	1.2	7.0 ± 8.3

Minifilament Formation (J2)









Minifilament Formation (J2)




- Duration of minifilament ~ 2.5 hours.
- Brightenings appear at the location where the minifilament subsequently forms.

Minifilament Formation and Flux Cancellation



Homologous Jet Eruptions




- We also observe more than a single jet from the same neutral line. A minifilament erupts and drives a jet, reforms/reappears at the same location, and then again erupts, driving the next jet.
- This process occurs as flux cancelation is ongoing and continues until all the minority-polarity flux vanishes. Eventually, the neutral line disappears, no more minifilaments and homologous jets are produced.

Origin of Minority-Polarity Flux Patch

We found the three scenarios:

- In some events tiny grains of flux coalesce to make a minority-polarity flux clump.
- The minority-polarity foot of a newly-emerged bipole became the minority-polarity flux patch.
- In some cases the minority-polarity clump preexisted as it rooted onto the Earthward side of the Sun 2-3 days before our observations began.







Schematic Illustration of Observations





Continuous flux cancelation between a minority-polarity flux clump and a majority-polarity flux clump builds a highly sheared minifilament field.

Flux rope model



- Footpoints get displaced along the PIL due to the differential rotation of the Sun.
- Footpoints come closer, due to the flux convergence at the PIL.
- Reconnection at the footpoints lead to the formation of longer helical field lines which carried the prominence plasma.




Polar Crown Prominence Observed on 13 Feb 2011



Magnetic Field Structure




Flux Linkage Model




- The interaction between two unconnected bi-poles, one is older and diffused which is at higher latitude, the other polarity is at lower latitude.
- This results in the highly sheared field at the PIL. Thus, after the many repetitions of this process, it forms the long helical structures that can partly or fully cover the polar region of the Sun.

(III) CME-producing Filament Eruptions

Filament Observed on 04 May 2013





AIA 193 movie




HMI Magnetogram movie

Flux Cancellation at the Neutral line




Filament Observed on 27 March 2012

2012-03-27 00:00:55



AIA 193 movie

2012-03-26 19:59:47



HMI Magnetogram movie

- Continuous flux cancellation at the neutral line of the filaments often triggers their eruptions. This corresponds to the finding that persistent flux cancellation at the neutral is the cause of jet-producing minifilament eruptions.
- Thus our observations support coronal jets being miniature versions of CMEs.

Summary

- We examined in detail random on-disk quiet-region jets and coronal hole jets. In each event a cool-transition-region material, a minifilament, initially resides at a neutral line inside the jet-base region.
- Our observations suggest that flux cancelation is usually the trigger of coronal jets.
- We found that flux cancelation is the key agent responsible for building a highly sheared minifilament field, leading to the formation of minifilaments.
- All the jet-producing eruptions are similar to typical solar flare eruptions, where a solar flare arcade forms during the filament eruption along the neutral line along which the filament resided prior to its eruption.
- Thus our observations support coronal jets being miniature versions of CMEs.