



Short-term root and leaf decomposition of two dominant plant species in a Siberian tundra

Wang, P., van Ruijven, J., Heijmans, M. M. P. D., Berendse, F., Maksimov, A., Maximov, T. C., & Mommer, L.

This is a "Post-Print" accepted manuscript, which has been published in "Pedobiologia"

This version is distributed under a non-commercial no derivatives Creative Commons



(CC-BY-NC-ND) user license, which permits use, distribution, and reproduction in any medium, provided the original work is properly cited and not used for commercial purposes. Further, the restriction applies that if you remix, transform, or build upon the material, you may not distribute the modified material.

Please cite this publication as follows:

Wang, P., van Ruijven, J., Heijmans, M. M. P. D., Berendse, F., Maksimov, A., Maximov, T. C., & Mommer, L. (2017). Short-term root and leaf decomposition of two dominant plant species in a Siberian tundra. *Pedobiologia*. DOI: 10.1016/j.pedobi.2017.08.002

You can download the published version at:

<https://doi.org/10.1016/j.pedobi.2017.08.002>

16 **Abstract**

17 In tundra ecosystems, global warming is expected to accelerate litter decomposition and to lead to
18 shifts in vegetation composition. To understand these shifts, it is important to understand the
19 interactions between global warming, vegetation composition, litter quality and decomposition in the
20 tundra. In addition, it is important to consider root litter since roots are the major part of plant biomass
21 in the tundra. In order to increase our understanding of decomposition, and root decomposition in
22 particular, we performed a litter transplant experiment in northeastern Siberia, in which we measured
23 mass loss for leaf and root litter (live and dead material) of the two dominant plant species, graminoid
24 *Eriophorum vaginatum* and shrub *Betula nana*, in three vegetation types (*E. vaginatum* or *B. nana*
25 dominated and mixed vegetation) during the growing season.

26 Our results show that although leaf decomposition did not differ between the two species, root
27 decomposition showed significant differences. Mass loss of live roots was higher for *E. vaginatum*
28 than for *B. nana*, but mass loss of *E. vaginatum* dead roots was lowest. In addition, we found evidence
29 for home-field advantage in litter decomposition: litter of a plant decomposed faster in vegetation
30 where it was dominant. Mass loss rates of the litter types were significantly correlated with
31 phosphorus content, rather than nitrogen content. This indicates that phosphorus limits decomposition
32 in this tundra site.

33 The low decomposition rate of *B. nana* live roots compared to *E. vaginatum* live roots suggests that
34 the acceleration of decomposition in the Arctic may be partly counteracted by the expected expansion
35 of shrubs. However, more information on litter input rates and direct effects of climate change on
36 decomposition rates are needed to accurately predict the effects of climate change on carbon dynamics
37 in tundra ecosystems.

38 *Keywords:* Arctic tundra, mass loss, *Betula nana*, *Eriophorum vaginatum*, home-field advantage, leaf
39 litter, root litter

40 **1. Introduction**

41 Arctic soils are an important global carbon reservoir, as half of the terrestrial belowground organic
42 carbon pool is sequestered in the northern circumpolar soil (Tarnocai et al., 2009). One of the key
43 processes in the global carbon cycle is the decomposition of organic plant litter (Bonan et al., 2013;
44 Wieder et al., 2013). It has been estimated that decomposition of plant litter accounts for half of the
45 terrestrial carbon release into the atmosphere (Houghton, 2007). Therefore, changes in decomposition
46 rates will greatly affect the soil carbon stocks of the Arctic ecosystems.

47 Important abiotic factors controlling decomposition rates include soil moisture, temperature and
48 nutrient availability (Swift et al., 1979). In the Arctic, temperature arguably is the most important
49 driver of decomposition (Hobbie, 1996; Robinson, 2002), as the soil is frozen for most of the year,
50 strongly limiting decomposition of plant litter. However, due to climate change, temperature has
51 already increased by about 1 °C in the last century and is predicted to further increase by 2 – 8 °C this
52 century (IPCC, 2013; Jones et al., 2012). Consequently, Arctic tundra soils will be warmer, permafrost
53 will thaw and decomposition of organic carbon will be accelerated (Cornelissen et al., 2007; Davidson
54 and Janssens, 2006; Schuur et al., 2009). Ultimately, the Arctic tundra may shift from a net carbon
55 sink to a net carbon source (Belshe et al., 2013; Webb et al., 2016).

56 In addition to abiotic factors, litter quality is an important driver of decomposition (Cornwell et al.,
57 2008). In general, plant litter with high nutrients and low lignin content decays faster than litter with
58 low nutrients and high lignin content (Freschet et al., 2012; Zhang et al., 2008). In most studies,
59 nitrogen appears to be the most important nutrient limiting decomposition, but phosphorus content has
60 also been found to be related to decomposition (Cornwell et al., 2008; Enriquez et al., 1993). Litter
61 decomposition can differ substantially between plant species or plant functional types (PFTs) within
62 the same ecosystem. In the tundra, the main PFTs are dwarf shrubs and graminoids. Generally, it is
63 thought that shrub litter is less decomposable than graminoid litter, as the former has a higher lignin
64 concentration (Cornelissen et al., 2007; Hobbie, 1996; Zhang et al., 2008).

65 Decomposition rates may also differ between different plant tissues of the same species. For
66 example, root litter generally decays slower than leaf litter (Birouste et al., 2011; Bryant et al., 1998;

67 Fujii and Takeda, 2010; Ma et al., 2016; Robinson et al., 1997; Thormann et al., 2001). In Arctic
68 tundra, up to 70% of plant biomass is allocated belowground (Iversen et al., 2015; Poorter et al., 2012;
69 Wang et al., 2016a). This high fraction suggests that root litter is a major source of carbon input and
70 root litter decomposition is likely to be an important component of the carbon cycle in this ecosystem.
71 However, detailed information about differences in root litter decomposition rates among species or
72 PFTs in the field is scarce. Hobbie (1996) and Robinson et al. (1999) showed that root and/or leaf
73 litters of graminoid species decomposed faster than those of shrub species, but these experiments were
74 performed under controlled conditions.

75 Understanding the differences in decomposition rates between PFTs is particularly relevant in the
76 Arctic as both recent observations and experimental researches show that climate warming affects the
77 distribution and abundance of the different PFTs (Elmendorf et al., 2012; Hill and Henry, 2011; Tape
78 et al., 2006). Shrub expansion has been observed in many tundra ecosystems (Callaghan et al., 2011;
79 Myers-Smith et al., 2011a; Myers-Smith et al., 2011b; Tape et al., 2006; Wookey et al., 2009). Such
80 changes in plant distribution and abundance will likely lead to differences in the quantity and quality
81 of litter input into the soil, which may affect decomposition rates and thus carbon cycling (Aerts,
82 2006; Berendse et al., 1989; Berendse et al., 1987; Cornelissen et al., 2007).

83 Moreover, there are feedbacks among vegetation type and decomposition rates (Ward et al., 2015).
84 Decomposition of plant litter can be up to 70% faster in the species' own habitat compared to a
85 different environment, a phenomenon referred to as "home-field advantage" (Gholz et al., 2000;
86 Strickland et al., 2009; Veen et al., 2015a). Home-field advantage effects on decomposition have been
87 observed worldwide (Ayres et al., 2009; Veen et al., 2015a), but have rarely been studied in tundra
88 ecosystems. One study we know determined home-field advantage in a sub-arctic tundra ecosystem
89 but found no support for it (Veen et al., 2015b).

90 Here, we determined the decomposition rates of leaf and root litter for the two dominant species,
91 representing the main PFTs in Siberian tundra, and tested for home-field advantage effects. We
92 performed a litter transplant experiment, in which leaf and root litters of both the graminoid

93 *Eriophorum vaginatum* L. and the deciduous shrub *Betula nana* L. were incubated for five weeks
94 during the growing season in three different vegetation types: *E. vaginatum*-dominated, *B. nana*-
95 dominated, and mixed vegetation. We hypothesized that:

- 96 1. The decomposition rate of root litter is lower than that of leaves,
- 97 2. The decomposition rate of *E. vaginatum* litter is lower than that of *B. nana*,
- 98 3. Differences in decomposition rates between litter types and species are related to litter quality,
- 99 4. Litter of a species decomposes faster in its own vegetation (i.e. home-field advantage occurs).

100

101 **2. Material and methods**

102 *2.1. Study site*

103 The study site is at the Chokurdakh Scientific Tundra Station (70°49'28'' N, 147°29'23'' E;
104 elevation 11 m a.s.l.) in Kytalyk Wildlife Reserve, which is located in the lowlands of the Indigirka
105 River in northeastern Siberia, Russia. The mean annual air temperature at the nearest climate station
106 (Chokurdakh, WMO station code 21946, 27 km away from the study site) is -13.4 °C (1981 – 2010),
107 with 10.3 °C as the mean July temperature. Annual precipitation is 196 mm (1981 – 2010), of which
108 on average 76 mm falls in the summer (June – August). The study area is the former lake bed of a
109 drained thermokarst lake, which has a shallow (20 – 45 cm) active layer (the soil layer that thaws in
110 the summer) underlain by thick continuous permafrost (Blok et al., 2010; Nauta et al., 2015).

111 The vegetation surrounding the Chokurdakh Scientific Tundra Station is classified as G4,
112 consisting of tussock-sedges (i.e. graminoids), dwarf-shrubs and moss on the Circumpolar Arctic
113 Vegetation Map (Walker et al., 2005). In the lake bed we distinguished 3 vegetation types: vegetation
114 dominated by the tussock-forming sedge *E. vaginatum* (> 70% cover); vegetation dominated by the
115 deciduous shrub *B. nana* (> 70% cover) and a mixed vegetation of both species (Wang et al., 2016b).
116 Other co-existing species with minor abundances include the grasses *Arctagrostis latifolia* (R. Br.)
117 Griseb and *Calamagrostis holmii* Lange, the sedge *Carex aquatilis* Wahlenberg, the deciduous shrub
118 *Salix pulchra* Cham, the evergreen shrubs *Vaccinium vitis-idaea* L and *Rhododendron subarcticum*

119 Harmaja. A moss layer with some lichen species is present throughout the study area (Blok et al.,
120 2010).

121

122 2.2. Experimental design

123 We focused on the two dominant plant species from different PFTs, the graminoid *E. vaginatum*
124 and the deciduous shrub *B. nana*. For *E. vaginatum* it was possible to collect dead roots as its roots are
125 mostly annual (Chapin, 1974; Sullivan et al., 2007; but see Iversen et al., 2015) and white-colored
126 when alive, and become black after senescence. Roots of *E. vaginatum* grow from the base of the tiller
127 and are unbranched with a uniform diameter of ~ 1 mm along the length, so all *E. vaginatum* roots
128 were considered fine roots. In addition, *B. nana* roots are usually ectomycorrhizal while *E. vaginatum*
129 roots are non-mycorrhizal (Iversen et al., 2015). We also included senesced leaves of *E. vaginatum*
130 (referred to as dead leaves here) to compare the decomposition leaf and root litter of this species. It
131 was impossible to collect shrub root litter (*B. nana* dead roots) from the soil, because roots are not
132 likely to shed discretely like leaves, but rather gradually lose functions and become colonized by
133 decomposers as they age (Hobbie et al., 2010). Hence, for *B. nana* we only could collect live fine roots
134 (< 1 mm in diameter) (referred to as live roots in the text). To reliably compare root and leaf
135 decomposition for *B. nana*, we also included leaves that were alive at the time of sampling (e.g. green;
136 referred to as live leaves here). To reliably compare decomposition of the two species, we also
137 included live leaves and live roots of *E. vaginatum*, and dead leaves of *B. nana*. In total we included
138 seven litter types in this experiment: live and dead leaves of *E. vaginatum* and *B. nana*, live and dead
139 roots of the *E. vaginatum*, and live roots of *B. nana*.

140 Due to logistic constraints, the litters were collected in different years and dried differently (see
141 Table S1 for an overview). Three litter (live leaves of *E. vaginatum* and *B. nana*, and live roots of *B.*
142 *nana*) were collected in July 2013 from the 8 blocks where the litter bags would be buried in (see
143 below). *E. vaginatum* live leaves were collected in the *E. vaginatum* dominated vegetation, and *B.*
144 *nana* live leaves and live roots were collected in *B. nana* dominated vegetation. Live leaves of the two
145 species were collected by clipping leaves from leaf bases of *E. vaginatum* and *B. nana* shoots. Live

146 fine roots of *B. nana* were collected by taking soil cores from the *B. nana* dominated vegetation and
147 picking out roots manually with forceps (Wang et al., 2016b). As mentioned above, *B. nana* roots are
148 not likely to shed discretely, so it is possible that the roots from *B. nana* classified as ‘alive’ contained
149 some dead roots. However, when collecting *B. nana* roots, roots that were darker in color and easily
150 torn apart were excluded. Thus, dead roots should only account for a very minor part of the samples.
151 The samples collected in 2013 were oven dried at 65 °C for 72 h after collection and stored in dry
152 conditions until the start of the decomposition experiment.

153 The other four litter types (live leaves, dead leaves and dead roots of *E. vaginatum*, and dead leaves
154 of *B. nana*) were collected in July 2015 at a location close to the experimental plots. Dead leaves of
155 the two species were collected from the ground underneath the *E. vaginatum* dominated or *B. nana*
156 dominated vegetation, respectively. As dead leaves of *B. nana* on the soil surface were probably
157 recently shed, we collected dead *E. vaginatum* leaves that were still standing up-right (assuming that
158 older dead leaves would be lying close to the ground) to minimize the age difference of the two
159 species’ leaf litter we collected. Live and dead roots of *E. vaginatum* were collected in the center of
160 tussocks formed by *E. vaginatum*. Roots of *E. vaginatum* were either white (live) or black (dead), non-
161 woody, unbranched, and densely clustered underneath the tussock. The samples collected in July 2015
162 were air-dried around 10 °C for 24 hours prior to filling the litter bags.

163 These differences in collection time and processing could potentially affect our results. To take this
164 into account, we focused on pairwise comparisons of litter collected in the same year (e.g. dead leaves
165 of *B. nana* and *E. vaginatum*) as much as possible. In the discussion, we will critically evaluate the
166 potentially confounding effects of differences in collection time and processing on our conclusions.

167 *E. vaginatum* leaves and roots, both live and dead, were cut to pieces of 3 – 5 cm long to fit in the
168 litter bags. Samples of each litter type were mixed and then carefully placed into litterbags, which
169 were 10×10 cm made from nylon mesh with a 0.5 mm mesh size (Top Zeven B.V., the Netherlands).
170 Each litter bag contained one type of litter with approximately 0.4 g dry weight. We recorded the
171 initial weight of each sample before putting it to a litter bag. To close the litter bags, they were folded
172 and staple-sealed with stainless steel staples.

173 Litter bags were buried in the three vegetation types. The three vegetation types differ in abiotic
174 factors, with *E. vaginatum* vegetation higher in soil moisture (volumetric moisture content 51%, 39%,
175 24% in the late growing season, for *E. vaginatum*, mixture, and *B. nana* vegetation respectively) and
176 exchangeable nutrients (total inorganic N 55, 30, 26 $\mu\text{g g}^{-1}$ resin, available P 5, 3, 3 $\mu\text{g g}^{-1}$ resin, K
177 124, 90, 50 $\mu\text{g g}^{-1}$ resin, for *E. vaginatum*, mixture, and *B. nana* vegetation respectively; see Wang et
178 al., 2016b). In the study area, eight blocks were selected in which all three vegetation types were close
179 to each other (3 – 10 m distance). Each block was 40 – 140 m away from the next block. In each of the
180 24 plots, seven litter bags (representing the different litter types) were buried. In total, we buried 168
181 litter bags (7 litter types \times 8 blocks \times 3 vegetation types) on 6 July 2015. Before being buried into the
182 soil, the litter bags were moderately moisturized for 10 minutes. A spade was used to cut a gap in the
183 soil with a 45° angle to the moss surface, and then one litter bag was placed at a depth of 5 cm from
184 the moss surface to the upper edge of the litter bag.

185 After 38 days, on 13 August 2015, the litter bags were harvested. After the litter bags were gently
186 removed from the soil, organic matter and soil on the surface of the litter bags was carefully brushed
187 off. The litter bags were stored in paper envelopes and air-dried in the field, then they were transported
188 to the Netherlands, where they were oven-dried at 60 °C for at least 48 hours and weighed. Mass loss
189 was calculated as the difference between the initial dry weight and the final dry weight, divided by the
190 initial dry weight. To determine the water content of each litter type in the initial weight, additional six
191 samples of each litter type (four samples for *B. nana* live roots, see section 2.3), with known initial
192 weight at the time of filling the litter bags, were oven-dried and weighed at Wageningen University,
193 the Netherlands, and then average water contents were calculated. Initial weights were corrected for
194 their water content when calculating mass loss.

195

196 2.3. Litter quality

197 To determine the relationship between decomposition and litter quality, the chemical composition
198 of the seven litter types was determined. For each litter type, six samples (four for *B. nana* live roots
199 due to a limited amount of samples) were taken to measure the initial tissue moisture content (see

200 section 2.2) and litter quality. Three samples of each litter type were then used to analyze the initial
201 carbon, nitrogen, and phosphorus concentration. The other three samples (one sample for *B. nana* live
202 roots) were used for lignin analysis. Carbon and nitrogen concentrations were determined with an
203 elemental analyzer (Fisons EA 1108 CHN-O). Phosphorus concentration was determined with a
204 segmented flow analyzer (SKALAR SAN Plus System, Breda, The Netherlands) after digestion with
205 H₂SO₄-salicylic acid-H₂O₂ and selenium (Novozamsky et al., 1983). Acid detergent lignin was
206 determined with Ankom 220 Fiber Analyzer (Ankom Technology, USA). C:N, C:P, lignin:N, lignin:P
207 ratios were calculated. Because lignin and N and P concentrations were measured in separate samples,
208 lignin:N and lignin:P ratios were calculated using mean values of lignin and N and P concentrations in
209 each litter type.

210

211 2.4. Calculations and statistical analysis

212 Mass loss data of the tissue types which we had for both species, i.e., live leaves, live roots, and dead
213 leaves of the two species, were used to test HFA. Home-field advantage (HFA) was calculated
214 following the method described in Ayres et al. (2009):

$$222 A_{RMLa} = \frac{A_a}{A_a + B_a} \times 100$$

$$223 HFAI = \left[\frac{A_{RMLa} + B_{RMLb}}{2} / \frac{A_{RMLb} + B_{RMLa}}{2} \right] \times 100 - 100$$

215 in which A_{RMLa} is the relative mass loss of species *A* at site *a*, A_a and B_a are the percent mass loss of
216 species *A* and *B* at site *a*. HFAIs were calculated separately for live and dead leaf, and live root litter
217 for each block in the field. The formula controls for inherent habitat differences in decomposition, i.e.,
218 in one habitat the decomposition of most litter may be faster than in other habitats. Note that this
219 formula only tests for the presence of a general HFA at the site and it does not quantify the HFA for an
220 individual species. To calculate the HFA for individual species requires three or more reciprocally
221 transplanted species (Ayres et al., 2009), which is beyond the scope of this study.

224 We used linear mixed effects models (LMM) to take into account that mass loss of samples in the
225 same plot or block are not fully independent. As the experimental design in terms of litter species and

226 dead vs live plant material was not fully balanced (because we did not include *B. nana* dead roots), we
227 tested live and dead plant material separately. In the model for live leaves and roots, vegetation type,
228 species and tissue type (leaf, root) were included as fixed effects. In the model for dead leaves and
229 roots, vegetation type and litter type (*E. vaginatum* leaf, *E. vaginatum* root, *B. nana* leaf) were
230 included as fixed effects. In both models block and plot were included as random effects with a nested
231 structure (plot within block). Mass loss data were ln transformed. Least significance difference (LSD)
232 method was used for post hoc tests when an effect was significant in one of the models.

233 To test if the HFAI for each litter type is significantly larger than zero and if it differed
234 significantly between tissue types, we ran a linear mixed model with litter type as fixed effect and
235 block as random effect.

236 Litter quality was compared among the seven litter types using a model with litter type as fixed
237 factor, block and plot as random factors with nested structure for each chemical characteristic. To
238 determine relationships between litter quality and litter mass loss, linear regression models were fitted
239 to the average mass loss of each litter type, using important chemical characteristics, including
240 nitrogen, phosphorus, lignin concentration, and C:N, C:P, lignin:N, lignin:P ratios as predictors. We
241 also calculated the AIC (Akaike information criterion) values and Akaike weight of each model to
242 evaluate which chemical characteristics best explained mass loss. The lowest AIC indicates the most
243 preferable among a set of models based on the goodness of fit and the number of parameters (Burnham
244 and Anderson, 2004), and an Akaike weight is the probability that a model is the actual best model
245 among a set of models (Wagenmakers and Farrell, 2004).

246

247 **3. Results**

248 *3.1. Mass loss of leaf and root litter and home-field advantage*

249 When focusing on live leaves and roots, differences in mass loss between the two species depended
250 on tissue type (roots vs. leaves; significant interaction of species \times tissue; Table 1). Leaves of the two
251 species showed a similar mass loss ($F_{1,42} = 0.7$, $P = 0.424$), but mass loss of *E. vaginatum* live roots

252 was significantly higher than that of *B. nana* live roots ($F_{1,21} = 747$, $P < 0.001$; Fig. 1a). Vegetation
 253 type effects on mass loss significantly differed between the two species (significant interaction of
 254 species \times vegetation; Table 1): live leaves and roots of *E. vaginatum* had similar mass losses in the
 255 three types of vegetation ($F_{2,30} = 0.4$, $P = 0.657$), whereas live roots (but not leaves) of *B. nana* had
 256 significantly larger mass losses in *B. nana* vegetation than in *E. vaginatum* vegetation ($F_{2,21} = 4.2$, $P =$
 257 0.03; Fig. 1a).

258

259 **Table 1**

260 Effects of vegetation, species and tissue (leaf/root) on live litter mass loss.

Source	df	F value	P value
Vegetation	2	1.4	0.259
Species	1	264.3	< 0.001 *
Tissue	1	40.9	< 0.001 *
Vegetation \times species	2	4.2	0.019 *
Vegetation \times tissue	2	0.8	0.441
PFT \times tissue	1	310.4	< 0.001 *
Vegetation \times species \times tissue	2	0.2	0.829

261

262 When focusing on dead leaves and roots, decomposition of different litter types was significantly
 263 different (Table 2): mass loss of *B. nana* and *E. vaginatum* leaves was similar, but both were
 264 significantly higher than *E. vaginatum* roots ($P = 0.002$ and $P < 0.001$ respectively; Fig. 1b). Similar to
 265 live plant tissues, vegetation type effects on mass loss depended on species (significant interaction of
 266 litter type \times vegetation; Table 2). Dead roots of *E. vaginatum* decayed significantly faster in *E.*
 267 *vaginatum* vegetation than in *B. nana* vegetation ($F_{2,30} = 5.1$, $P = 0.013$), while live roots and dead
 268 leaves of *B. nana* had significantly larger mass loss in *B. nana* vegetation than in *E. vaginatum*
 269 vegetation ($F_{2,21} = 4.5$, $P = 0.023$). As indicated by these significant interactions between vegetation
 270 type and species, we found a clear home-field advantage for our litter types. This effect was
 271 significantly greater than zero for live roots and dead leaves (see Fig. S1).


272

273 **Table 2**

274 Effects of vegetation and dead litter type (*E. vaginatum* and *B. nana* leaves, *E. vaginatum* roots) on
 275 mass loss.

Source	df	F value	P value
Vegetation	2	0.4	0.697
Litter type	2	12.4	< 0.001 *
Vegetation × litter type	4	4.2	0.005 *

276



277

278 **Fig. 1.** Mass loss of live (a) and dead (b) litter types in the three vegetation types: *E. vaginatum* (Ev),
 279 mixed (mix) and *B. nana* (Bn). Bars are means ± SE, n = 8. Scales of y-axes in (a) and (b) are different
 280 as mass loss of dead litter was much smaller. Capital letters represent significant ($P < 0.05$) pairwise
 281 differences in mass loss between litter types for live and dead litter respectively; lowercase letters
 282 represent significant ($P < 0.05$) pairwise differences in mass loss between vegetation types within litter
 283 types.

284

285 *3.2. Relations between mass loss and litter quality*

286 The seven litter types differed significantly in nitrogen, phosphorus and lignin content and related
 287 ratios (Table 3). In general, *B. nana* litter was characterized by higher lignin concentrations than litter
 288 of *E. vaginatum*. Not surprisingly, dead plant material showed lower nitrogen and phosphorus
 289 concentrations than live plant material, but dead roots and leaves of *E. vaginatum* had particularly low
 290 nitrogen and phosphorus contents (Table 3). Across all litter types, nitrogen content differed up to 5
 291 fold (between *B. nana* live leaves and *E. vaginatum* dead roots), whereas phosphorus content differed
 292 up to 17 fold (between *B. nana* live leaves and *E. vaginatum* dead leaves; Table 3).

293 The average mass loss of a litter type was significantly related to litter characteristics involving
 294 phosphorus. Mass loss significantly increased with increasing P concentration, and decreased with the
 295 C:P and lignin:P ratios (Fig. 2). Mass loss also significantly decreased, albeit weaker, with the
 296 lignin:N ratio. No significant relationships between mass loss and nitrogen concentration, C:N and
 297 lignin concentration were found (Fig. 2). Model comparisons revealed that phosphorus concentration
 298 and lignin:P were the best predictors for mass loss of the different litters in the tundra (Table S2).

299


300 **Table 3**

301 Initial chemical characteristics of the different litter types. Different letters indicate differences
 302 between litter types using Tukey HSD post hoc method. N= 3 except for lignin concentration in *B.*
 303 *nana* roots. Lignin:N and lignin:P were calculated with mean values of lignin and N/P concentration as
 304 they were from separate samples.

Chemical characteristics	<i>E. vaginatum</i> live leaves	<i>E. vaginatum</i> dead leaves	<i>B. nana</i> live leaves	<i>B. nana</i> dead leaves	<i>E. vaginatum</i> live roots	<i>E. vaginatum</i> dead roots	<i>B. nana</i> live roots
C (%)	45.9 ± 0.2 ^{ab}	45.7 ± 0.2 ^{ab}	47.0 ± 2.6 ^{ab}	50.2 ± 0.3 ^a	43.2 ± 0.1 ^b	46.8 ± 0.2 ^{ab}	47.4 ± 1.0 ^{ab}
Lignin (%)	3.4 ± 0.02 ^d	4.7 ± 0.1 ^d	18.9 ± 0.5 ^b	35.3 ± 0.2 ^a	3.2 ± 0.3 ^d	11.4 ± 0.8 ^c	34.6
N (mg g ⁻¹)	16.5 ± 0.6 ^b	6.5 ± 0.8 ^c	27.2 ± 1.6 ^a	12.8 ± 0.5 ^b	13.3 ± 1.0 ^b	5.9 ± 0.9 ^c	15.1 ± 1.4 ^b

P (mg g ⁻¹)	2.77 ± 0.04 ^c	0.20 ± 0.02 ^f	3.37 ± 0.01 ^a	0.69 ± 0.04 ^e	3.05 ± 0.03 ^b	0.21 ± 0.01 ^f	1.15 ± 0.04 ^d
N:P	6.0 ± 0.3 ^a	32.0 ± 2.7 ^d	8.1 ± 0.5 ^a	18.8 ± 1.8 ^{bc}	4.4 ± 0.4 ^a	28.2 ± 4.7 ^{cd}	13.2 ± 1.2 ^{ab}
C:N	27.9 ± 1.0 ^b	72.4 ± 9.4 ^a	17.3 ± 0.4 ^c	39.4 ± 1.8 ^b	32.8 ± 2.7 ^b	82.0 ± 10.9 ^a	31.9 ± 3.5 ^b
C:P	166 ± 2 ^d	2277 ± 181 ^a	139 ± 8 ^d	735 ± 44 ^b	142 ± 1 ^d	2212 ± 132 ^a	413 ± 18 ^c
Lignin:N	2.1	7.2	7.0	27.6	2.4	19.2	22.9
Lignin:P	12.5	229.7	56.1	513.5	10.4	535.9	301.1

305



306

307 **Fig. 2.** Relationships between mass loss of the seven litter types and their initial chemical
308 characteristics. Relationships between mass loss and litter quality were particularly strong for P-related
309 characteristics (right column). Symbols show average mass loss (n = 24) and chemical characteristic
310 values (n = 1 for lignin content of *B. nana* dead roots and lignin:N, lignin:P; n = 3 for other
311 characteristics; see Table 3).

312 **4. Discussion**

313 The decomposition of root litter was different between the two species, in contrast to the
314 decomposition of leaf litter. The differences in mass loss rate among the litter types were significantly
315 correlated to phosphorus content, not to nitrogen content. In addition, we found evidence for home-
316 field advantage in litter decomposition: litter of the two species decomposed faster in vegetation in
317 which they were dominant. As root litter constitutes a considerable fraction of the organic matter input
318 in this system (Freschet et al., 2013), the difference in root decomposition rates between the two
319 dominant species at the research site suggests that changes in carbon dynamics with vegetation shifts
320 in tundra ecosystems will largely depend on root decomposition.

321

322 *4.1. Decomposition of leaves and roots of the two species*

323 The decomposition of leaf litter did not differ between the two species. This was true for both dead
324 and live leaves. However, the decomposition of live roots was significantly different between the two
325 species, although we cannot exclude a potential small effect of the inclusion of some *B. nana* dead
326 roots in the 'live' root samples, which might have led to an overestimation of the difference between
327 the decomposition of *E. vaginatum* and *B. nana* roots. These results only partly confirm our
328 hypotheses that root decomposition is slower than leaf decomposition and that the decomposition of *B.*
329 *nana* litter is slower than that of *E. vaginatum* litter, and they suggest that we need to consider the
330 differences between tissue types and species or PFTs, as PFTs are a reasonable classification of major
331 plant species in the tundra (Chapin et al., 1996), at the same time.

332 Shrub litter is generally thought to be less decomposable than graminoid litter, as the former has a
333 higher lignin concentration and C:N ratio (Cornelissen et al., 2007; Hobbie, 1996; Zhang et al., 2008).
334 In our study, this is only the case for root litter, as leaf litter of the two species exhibited similar mass
335 loss rates. *B. nana* leaves had higher nitrogen and phosphorus concentrations than *E. vaginatum*
336 leaves, however, the former also had higher lignin concentration, which could offset the positive
337 effects of higher nutrient concentrations on the litter decomposition.

338 Mass loss of live roots of the two species showed large difference. *B. nana* fine roots in our
339 samples are mainly first- and second-order roots, which fact could make the actual decomposition of
340 the whole root system a bit higher, as some studies found that lowest order roots decay slower than
341 higher order roots (Fan and Guo, 2010; Goebel et al., 2011). However, the lower turnover rates of
342 higher order roots (McCormack et al., 2015) make it unlikely to offset or reverse this difference we
343 found. Therefore, we suggest that root decomposition in the tundra can differ between shrubs and
344 graminoids, at least in the early stage of decomposition for the two dominant species. However, in the
345 later stages of decomposition the differences may change, as the decomposition of *E. vaginatum* dead
346 roots decreased drastically. Our results emphasize that it is important to consider root litter separately
347 from leaf litter when comparing different species or PFTs, even when the decomposition of leaf litter
348 does not show differences.

349 Many studies have shown that root litter is less decomposable than leaf litter as roots contain more
350 chemically recalcitrant substances (Freschet et al., 2012; Freschet et al., 2013; Gorissen and Cotrufo,
351 2000; Ma et al., 2016). In our study this is only partly true, as *E. vaginatum* live leaves were
352 decomposed slower than its live roots. However, the dead leaves of the *E. vaginatum* decomposed
353 faster than its dead roots. This discrepancy between live and dead plant material when comparing leaf
354 and root decomposition illustrates that live roots may not necessarily be a good predictor of dead root
355 decomposition, particularly for *E. vaginatum* roots, and that conclusions about decomposition based
356 on live tissues of different plants should be treated with caution, especially if species differ in
357 characteristics such as nutrient resorption efficiency (Scheffer and Aerts, 2000; Snyder and
358 Rejmánková, 2015).

359 Litter quality (e.g., nitrogen and lignin concentration, C:N ratio) is one of the most important
360 factors in decomposition from grassland to forest ecosystems (Cornwell et al., 2008; Freschet et al.,
361 2012; Zhang et al., 2008). It is well known that nutrient concentrations, particularly nitrogen, are
362 among the best predictors during the early stage of decomposition, while lignin is the best predictor
363 during later stages (Berg and McClaugherty, 2014). In our study, traits related to phosphorus content
364 were identified as the main drivers of litter decomposition, suggesting that at least the early phases of

365 decomposition are phosphorus-limited. This finding is consistent with another study (Beermann et al.,
366 2014) which suggested that at the research site nitrogen mineralization and fixation by bacteria are
367 limited by phosphorus availability. On the other hand, according to Koerselman and Meuleman
368 (1996), an N:P ratio below 14 indicates nitrogen limitation for plant growth. The low N:P ratio of
369 green leaves in our study (6.0 – 8.1; Table 3) suggests that plant growth was limited by nitrogen
370 availability. The explanation for the limitation of microbial growth and plants growth by different
371 elements could be that the accessibility to phosphorus is different for them. While microbes in the
372 shallow soil are limited by phosphorus, graminoids can exploit the deep soil, where larger amounts of
373 phosphorus are available (Beermann et al., 2014; Chapin et al., 1978). Shrubs can depend on
374 mycorrhizal fungi to absorb phosphorus from the deep soil (Bolan, 1991; Landeweert et al., 2001).
375 Efficient phosphorus resorption by plants from senescing plant parts could be another explanation of
376 the phosphorus limitation of microbes. However, we could not examine this hypothesis with our data
377 as the dead leaves/roots were not freshly senesced and we do not know to what extent the massive
378 differences in phosphorus concentration and N:P can be attributed to plant resorption and microbial
379 mobilization, respectively.

380

381 4.2. Home-field advantage effects on decomposition

382 Decomposition of both species tended to be faster in the vegetation in which they were dominant,
383 i.e. in their 'own' vegetation, suggesting home-field advantage effects in our study. Home-field
384 advantage was significant for *B. nana* (for roots and dead leaves) and *E. vaginatum* (dead leaves), even
385 though there are big differences in decomposability between these litter types. In fact, in our study the
386 site effects are stronger for dead than for live tissue (i.e. stronger for tissues with lower
387 decomposability; see Fig S1), consistent with the idea that litter with low decomposability requires
388 more specialized decomposers (Ayres et al., 2009; Milcu and Manning, 2011). However, in a subarctic
389 tundra in northern Sweden, Veen et al. (2015b) found no significant home-field advantage in the
390 decomposition of leaf litter. The reason could be that in their experiment they used a mixture of litters

391 from different plant species, therefore the HFA effect at the species level, as shown in our study, could
392 be masked by the community level measurement.

393 The *E. vaginatum* and *B. nana* vegetation in our study differ in abiotic factors. *E. vaginatum*
394 vegetation is wetter and more nutrient-rich than the *B. nana* vegetation (Nauta et al., 2015; Wang et
395 al., 2016b). However, the lack of overall vegetation effects on mass loss suggests that abiotic factors
396 are not decisive in our experiment. Instead, the different environmental conditions may have shaped
397 different microbial communities, that also determine the home-field advantage effects in litter
398 decomposition (Wallenstein et al., 2007).

399

400 4.3. Influences of the collection years and drying of litter

401 As mentioned in the methods, there were differences in litter collection and processing between
402 live and dead plant material. All live materials, except *E. vaginatum* live roots, were collected in 2013,
403 dried at 65°C and stored until the decomposition experiment, whereas all dead materials were
404 collected in 2015 and air-dried. *E. vaginatum* live roots were collected in 2015, together with the
405 collection of dead materials, and air-dried. This could potentially affect our results, but we avoided
406 this problem by drawing conclusions based on comparisons of litter types collected in the same year
407 (i.e., with the same drying process). The only exception are the live roots. *E. vaginatum* live roots were
408 collected in 2015, whereas those of *B. nana* were collected in 2013. This could potentially affect our
409 first conclusion that root decomposition differs between *E. vaginatum* and *B. nana*. To check this, we
410 compared the initial P and N concentrations in *E. vaginatum* live roots of both years (a small amount
411 of *E. vaginatum* live roots was also collected in 2013 and dried at 65°C in another experiment at the
412 same site), and found no significant differences between years for P (2.97 ± 0.04 vs 3.05 ± 0.03
413 mg/g, $P = 0.85$), and a marginally significant difference for N (17.0 ± 1.6 vs 13.3 ± 1.0 mg/g, $P =$
414 0.09). Given these small differences, it is unlikely that the large (3-fold) difference in mass loss we
415 report for *B. nana* and *E. vaginatum* roots (Fig. 1) is affected by differences in year of collection or
416 drying.

417 Across all seven litter types, P concentration was the main driver of differences in decomposition
418 (Fig. 2). However, as can also be seen in this figure, the three dead litters collected in 2015 (except *E.*
419 *vaginatum* live root) did show lower mass loss than the ones collected in 2013, suggesting that the P
420 concentration effect could also be driven by year of litter collection. We checked this by comparing
421 our litter quality model to one that explains mass loss by year of collection. The results show that year
422 did not have a significant effect on decomposition ($F_{1,5} = 0.55$, $P = 0.49$), which is not surprising given
423 that litters collected in 2015 included both *E. vaginatum* live roots and three dead materials and thus
424 the variance of mass loss was large. In addition, when comparing the AIC values of both models it
425 was clear that litter quality (P and lignin:P) performed better (AIC = -20.3 and -19.9, respectively)
426 than the year model (AIC = -5.8). Thus, although we cannot completely rule out that year of
427 collection affected mass loss, it is more likely that decomposition is indeed driven by litter quality.
428 This is supported by the lack of difference in P content between *E. vaginatum* live roots collected in
429 2013 and 2015, as explained above.

430

431 4.4. Implications for carbon dynamics in the tundra

432 There is major concern that tundra ecosystems might shift from a carbon sink to a carbon source
433 with warmer climates (Belshe et al., 2013; Oechel et al., 1993; Webb et al., 2016). A warmer climate
434 will increase primary productivity of tundra vegetation and thus increase carbon uptake by the
435 ecosystem (Epstein et al., 2012; Hill and Henry, 2011; Verbyla, 2008). On the other hand, higher
436 temperatures also accelerate decomposition and thus increase carbon emission from the soil (Davidson
437 and Janssens, 2006; Hobbie, 1996). The balance between these two changes will determine whether
438 tundra ecosystems will continue to act as a carbon sink or will shift to a carbon source. These changes
439 in carbon dynamics can be modified by shifts in tundra vegetation composition due to climate
440 warming. The home-field advantage in litter decomposition in our study suggests that litter
441 decomposition rates may be temporarily reduced when vegetation shifts occur. However, whether this
442 reduction in decomposition due to home-field advantage can at least temporarily offset the increase in
443 decomposition due to climate warming needs further investigation.

444 Shifts in vegetation composition also affect decomposition via changes in litter quality (Cornelissen
445 et al., 2007). Focusing on the aboveground tissues, our study does not provide evidence for vegetation
446 induced changes in decomposition rates, as the decomposability of leaf litter did not differ between the
447 two species. However, root litter decomposition was lower for *B. nana* than for *E. vaginatum*, at least
448 in the early stage. Also, it is known that root turnover rates of shrubs are lower than that of graminoids
449 in the tundra (Mack et al., 2004; Shaver and Chapin, 1991; Sullivan et al., 2007). When extrapolating
450 this finding, it would suggest that shrub expansion with increasing temperatures could reduce
451 decomposition and increase carbon storage. However, it may be important to consider that graminoids
452 roots typically grow deeper than shrubs (Miller et al., 1982; Shaver and Chapin, 1991; Wang et al.,
453 2016b). As deeper soil layers will be colder, decomposition of graminoid roots may be slower than
454 observed in litter bag studies in the upper soil layer. This is illustrated by the observation that yedoma
455 (windblown dust, deposited during the glacial age) permafrost contains almost intact graminoid roots
456 (Zimov et al., 2006). To accurately predict the long-term effects of vegetation shifts on decomposition
457 and the carbon balance, detailed knowledge of the temporal dynamics in root turnover and
458 decomposition at different soil depths in relation to soil temperature are needed.

459

460 **5. Conclusion**

461 Our study shows that although leaf litter decomposition did not differ between the two dominant
462 species, root litter decomposition was significantly higher for the graminoid *E. vaginatum* than for the
463 shrub *B. nana*. The differences we found in decomposability could be mainly attributed to litter traits
464 related to phosphorus. In addition, home-field advantage effects were found, which may lead to
465 temporary reductions in litter decomposition when vegetation shifts occur. Our results indicate that
466 root decomposition can be an important driver of changes in carbon dynamics when vegetation shifts
467 and consequently plant litter changes in the tundra. The large difference between the mass loss of the
468 live and dead plant materials, particularly between the live and dead *E. vaginatum* roots, suggests that
469 only looking at the initial phase of decomposition does not give a clear indication of decomposition
470 rate over time.

471

472 **Acknowledgements**

473 We thank staff of IBPC, Yakutsk, and staff of the Regional Inspection of Nature Protection of
474 Allaikhovsky Region, Chokurdakh, for logistic support and assistance. We thank Hongmei Chen for
475 providing litter bags used in the experiment. PW acknowledges financial support from China
476 Scholarship Council (CSC, No.201206040062), MH and LM acknowledge support from The
477 Netherlands Organisation for Scientific Research (NWO-ALW, VIDI grants 864.09.014 and
478 864.15.006).

479

480 **References**

- 481 Aerts, R., 2006. The freezer defrosting: global warming and litter decomposition rates in cold biomes. *Journal of*
482 *Ecology* 94, 713-724.
- 483 Ayres, E., Steltzer, H., Simmons, B.L., Simpson, R.T., Steinweg, J.M., Wallenstein, M.D., Mellor, N., Parton,
484 W.J., Moore, J.C., Wall, D.H., 2009. Home-field advantage accelerates leaf litter decomposition in forests. *Soil*
485 *Biology and Biochemistry* 41, 606-610.
- 486 Beermann, F., Teltewskoi, A., Fiencke, C., Pfeiffer, E.-M., Kutzbach, L., 2014. Stoichiometric analysis of
487 nutrient availability (N, P, K) within soils of polygonal tundra. *Biogeochemistry* 122, 211-227.
- 488 Belshe, E.F., Schuur, E.A.G., Bolker, B.M., 2013. Tundra ecosystems observed to be CO₂ sources due to
489 differential amplification of the carbon cycle. *Ecology Letters* 16, 1307-1315.
- 490 Berendse, F., Bobbink, R., Rouwenhorst, G., 1989. A comparative study on nutrient cycling in wet heathland
491 ecosystems. II. Litter decomposition and nutrient mineralization. *Oecologia* 78, 338-348.
- 492 Berendse, F., Oudhof, H., Bol, J., 1987. A comparative study on nutrient cycling in wet heathland ecosystems. I.
493 Litter production and nutrient losses from the plant. *Oecologia* 74, 174-184.
- 494 Berg, B., McLaugherty, C., 2014. *Plant Litter. Decomposition, Humus Formation, Carbon Sequestration*, Third
495 ed. Springer-Verlag Berlin Heidelberg.
- 496 Birouste, M., Kazakou, E., Blanchard, A., Roumet, C., 2011. Plant traits and decomposition: are the relationships
497 for roots comparable to those for leaves? *Annals of Botany*.
- 498 Blok, D., Heijmans, M.M.P.D., Schaepman-Strub, G., Kononov, A.V., Maximov, T.C., Berendse, F., 2010.
499 Shrub expansion may reduce summer permafrost thaw in Siberian tundra. *Global Change Biology* 16, 1296-
500 1305.
- 501 Bolan, N.S., 1991. A critical review on the role of mycorrhizal fungi in the uptake of phosphorus by plants. *Plant*
502 *and Soil* 134, 189-207.
- 503 Bonan, G.B., Hartman, M.D., Parton, W.J., Wieder, W.R., 2013. Evaluating litter decomposition in earth system
504 models with long-term litterbag experiments: an example using the Community Land Model version 4 (CLM4).
505 *Global Change Biology* 19, 957-974.

- 506 Bryant, D.M., Holland, E.A., Seastedt, T.R., Walker, M.D., 1998. Analysis of litter decomposition in an alpine
507 tundra. *Canadian Journal of Botany* 76, 1295-1304.
- 508 Burnham, K.P., Anderson, D.R., 2004. Multimodel inference understanding AIC and BIC in model selection.
509 *Sociological Methods & Research* 33, 261-304.
- 510 Callaghan, T., Tweedie, C., Åkerman, J., Andrews, C., Bergstedt, J., Butler, M., Christensen, T., Cooley, D.,
511 Dahlberg, U., Danby, R., Daniëls, F.A., de Molenaar, J., Dick, J., Mortensen, C., Ebert-May, D., Emanuelsson,
512 U., Eriksson, H., Hedenås, H., Henry, G.H.R., Hik, D., Hobbie, J., Jantze, E., Jaspers, C., Johansson, C.,
513 Johansson, M., Johnson, D., Johnstone, J., Jonasson, C., Kennedy, C., Kenney, A., Keuper, F., Koh, S., Krebs,
514 C., Lantuit, H., Lara, M., Lin, D., Lougheed, V., Madsen, J., Matveyeva, N., McEwen, D., Myers-Smith, I.,
515 Narozhniy, Y., Olsson, H., Pohjola, V., Price, L., Rigét, F., Rundqvist, S., Sandström, A., Tamstorf, M., Van
516 Bogaert, R., Villarreal, S., Webber, P., Zemtsov, V., 2011. Multi-decadal changes in tundra environments and
517 ecosystems: synthesis of the International Polar Year-Back to the Future Project (IPY-BTF). *AMBIO* 40, 705-
518 716.
- 519 Chapin, F.S., III, 1974. Morphological and physiological mechanisms of temperature compensation in phosphate
520 absorption along a latitudinal gradient. *Ecology* 55, 1180-1198.
- 521 Chapin, F.S., III, Barsdate, R.J., Barèl, D., 1978. Phosphorus cycling in alaskan coastal tundra: A hypothesis for
522 the regulation of nutrient cycling. *Oikos* 31, 189-199.
- 523 Chapin, F.S., III, Bret-Harte, M.S., Hobbie, S.E., Zhong, H., 1996. Plant functional types as predictors of
524 transient responses of arctic vegetation to global change. *Journal of Vegetation Science* 7, 347-358.
- 525 Cornelissen, J.H., Van Bodegom, P.M., Aerts, R., Callaghan, T.V., Van Logtestijn, R.S., Alatalo, J., Chapin,
526 F.S., III, Gerdol, R., Gudmundsson, J., Gwynn - Jones, D., 2007. Global negative vegetation feedback to climate
527 warming responses of leaf litter decomposition rates in cold biomes. *Ecology letters* 10, 619-627.
- 528 Cornwell, W.K., Cornelissen, J.H.C., Amatangelo, K., Dorrepaal, E., Eviner, V.T., Godoy, O., Hobbie, S.E.,
529 Hoorens, B., Kurokawa, H., Pérez-Harguindeguy, N., Quested, H.M., Santiago, L.S., Wardle, D.A., Wright, I.J.,
530 Aerts, R., Allison, S.D., Van Bodegom, P., Brovkin, V., Chatain, A., Callaghan, T.V., Díaz, S., Garnier, E.,
531 Gurvich, D.E., Kazakou, E., Klein, J.A., Read, J., Reich, P.B., Soudzilovskaia, N.A., Vaieretti, M.V., Westoby,
532 M., 2008. Plant species traits are the predominant control on litter decomposition rates within biomes worldwide.
533 *Ecology Letters* 11, 1065-1071.
- 534 Davidson, E.A., Janssens, I.A., 2006. Temperature sensitivity of soil carbon decomposition and feedbacks to
535 climate change. *Nature* 440, 165-173.
- 536 Elmendorf, S.C., Henry, G.H.R., Hollister, R.D., Björk, R.G., Bjorkman, A.D., Callaghan, T.V., Collier, L.S.,
537 Cooper, E.J., Cornelissen, J.H.C., Day, T.A., Fosaa, A.M., Gould, W.A., Grétarsdóttir, J., Harte, J., Hermanutz,
538 L., Hik, D.S., Hofgaard, A., Jarrad, F., Jónsdóttir, I.S., Keuper, F., Klanderud, K., Klein, J.A., Koh, S., Kudo, G.,
539 Lang, S.I., Loewen, V., May, J.L., Mercado, J., Michelsen, A., Molau, U., Myers-Smith, I.H., Oberbauer, S.F.,
540 Pieper, S., Post, E., Rixen, C., Robinson, C.H., Schmidt, N.M., Shaver, G.R., Stenström, A., Tolvanen, A.,
541 Totland, Ø., Troxler, T., Wahren, C.-H., Webber, P.J., Welker, J.M., Wookey, P.A., 2012. Global assessment of
542 experimental climate warming on tundra vegetation: heterogeneity over space and time. *Ecology Letters* 15, 164-
543 175.
- 544 Enriquez, S., Duarte, C.M., Sand-Jensen, K., 1993. Patterns in decomposition rates among photosynthetic
545 organisms: the importance of detritus C: N: P content. *Oecologia* 94, 457-471.
- 546 Epstein, H.E., Reynolds, M.K., Walker, D.A., Bhatt, U.S., Tucker, C.J., Pinzon, J.E., 2012. Dynamics of
547 aboveground phytomass of the circumpolar Arctic tundra during the past three decades. *Environmental Research*
548 *Letters* 7, 015506.
- 549 Fan, P., Guo, D., 2010. Slow decomposition of lower order roots: a key mechanism of root carbon and nutrient
550 retention in the soil. *Oecologia* 163, 509-515.
- 551 Freschet, G.T., Aerts, R., Cornelissen, J.H.C., 2012. A plant economics spectrum of litter decomposability.
552 *Functional Ecology* 26, 56-65.


- 553 Freschet, G.T., Cornwell, W.K., Wardle, D.A., Elumeeva, T.G., Liu, W., Jackson, B.G., Onipchenko, V.G.,
554 Soudzilovskaia, N.A., Tao, J., Cornelissen, J.H.C., 2013. Linking litter decomposition of above- and below-
555 ground organs to plant–soil feedbacks worldwide. *Journal of Ecology* 101, 943-952.
- 556 Fujii, S., Takeda, H., 2010. Dominant effects of litter substrate quality on the difference between leaf and root
557 decomposition process above- and belowground. *Soil Biology and Biochemistry* 42, 2224-2230.
- 558 Gholz, H.L., Wedin, D.A., Smitherman, S.M., Harmon, M.E., Parton, W.J., 2000. Long - term dynamics of pine
559 and hardwood litter in contrasting environments: toward a global model of decomposition. *Global Change*
560 *Biology* 6, 751-765.
- 561 Goebel, M., Hobbie, S.E., Bulaj, B., Zadworny, M., Archibald, D.D., Oleksyn, J., Reich, P.B., Eissenstat, D.M.,
562 2011. Decomposition of the finest root branching orders: linking belowground dynamics to fine - root function
563 and structure. *Ecological Monographs* 81, 89-102.
- 564 Gorissen, A., Cotrufo, M.F., 2000. Decomposition of leaf and root tissue of three perennial grass species grown
565 at two levels of atmospheric CO₂ and N supply. *Plant and Soil* 224, 75-84.
- 566 Hill, G.B., Henry, G.H.R., 2011. Responses of High Arctic wet sedge tundra to climate warming since 1980.
567 *Global Change Biology* 17, 276-287.
- 568 Hobbie, S.E., 1996. Temperature and plant species control over litter decomposition in Alaskan tundra.
569 *Ecological Monographs* 66, 503-522.
- 570 Hobbie, S.E., Oleksyn, J., Eissenstat, D.M., Reich, P.B., 2010. Fine root decomposition rates do not mirror those
571 of leaf litter among temperate tree species. *Oecologia* 162, 505-513.
- 572 Houghton, R.A., 2007. Balancing the global carbon budget. *Annu. Rev. Earth Planet. Sci.* 35, 313-347.
- 573 IPCC, 2013. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth*
574 *Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge University Press, Cambridge,
575 United Kingdom and New York, NY, USA.
- 576 Iversen, C.M., Sloan, V.L., Sullivan, P.F., Euskirchen, E.S., McGuire, A.D., Norby, R.J., Walker, A.P., Warren,
577 J.M., Wullschlegel, S.D., 2015. The unseen iceberg: plant roots in arctic tundra. *New Phytologist* 205, 34-58.
- 578 Jones, P.D., Lister, D.H., Osborn, T.J., Harpham, C., Salmon, M., Morice, C.P.C.D., 2012. Hemispheric and
579 large-scale land-surface air temperature variations: An extensive revision and an update to 2010. *J. Geophys.*
580 *Res.* 117, D05127.
- 581 Koerselman, W., Meuleman, A.F.M., 1996. The vegetation N:P ratio: A new tool to detect the nature of nutrient
582 limitation. *Journal of Applied Ecology* 33, 1441-1450.
- 583 Landeweert, R., Hoffland, E., Finlay, R.D., Kuyper, T.W., van Breemen, N., 2001. Linking plants to rocks:
584 ectomycorrhizal fungi mobilize nutrients from minerals. *Trends in Ecology & Evolution* 16, 248-254.
- 585 Ma, C., Xiong, Y., Li, L., Guo, D., 2016. Root and leaf decomposition become decoupled over time:
586 implications for below- and above-ground relationships. *Functional Ecology* 30, 1239-1246.
- 587 Mack, M.C., Schuur, E.A.G., Bret-Harte, M.S., Shaver, G.R., Chapin, F.S., III, 2004. Ecosystem carbon storage
588 in arctic tundra reduced by long-term nutrient fertilization. *Nature* 431, 440-443.
- 589 McCormack, M.L., Dickie, I.A., Eissenstat, D.M., Fahey, T.J., Fernandez, C.W., Guo, D., Helmisaari, H.S.,
590 Hobbie, E.A., Iversen, C.M., Jackson, R.B., 2015. Redefining fine roots improves understanding of below -
591 ground contributions to terrestrial biosphere processes. *New Phytologist* 207, 505-518.
- 592 Milcu, A., Manning, P., 2011. All size classes of soil fauna and litter quality control the acceleration of litter
593 decay in its home environment. *Oikos* 120, 1366-1370.

- 594 Miller, P.C., Mangan, R., Kummerow, J., 1982. Vertical distribution of organic matter in eight vegetation types
595 near Eagle Summit, Alaska. *Ecography* 5, 117-124.
- 596 Myers-Smith, I., Hik, D., Kennedy, C., Cooley, D., Johnstone, J., Kenney, A., Krebs, C., 2011a. Expansion of
597 canopy-forming willows over the twentieth century on Herschel Island, Yukon Territory, Canada. *AMBIO* 40,
598 610-623.
- 599 Myers-Smith, I.H., Forbes, B.C., Wilmking, M., Hallinger, M., Lantz, T., Blok, D., Tape, K.D., Macias-Fauria,
600 M., Sass-Klaassen, U., Lévesque, E., Boudreau, S., Ropars, P., Hermanutz, L., Trant, A., Collier, L.S., Weijers,
601 S., Rozema, J., Rayback, S.A., Schmidt, N.M., Schaepman-Strub, G., Wipf, S., Rixen, C., Ménard, C.B., Venn,
602 S., Goetz, S., Andreu-Hayles, L., Elmendorf, S., Ravolainen, V., Welker, J., Grogan, P., Epstein, H.E., Hik, D.S.,
603 2011b. Shrub expansion in tundra ecosystems: dynamics, impacts and research priorities. *Environmental*
604 *Research Letters* 6, 045509.
- 605 Nauta, A.L., Heijmans, M.M.P.D., Blok, D., Limpens, J., Elberling, B., Gallagher, A., Li, B.X., Petrov, R.E.,
606 Maximov, T.C., van Huissteden, J., Berendse, F., 2015. Permafrost collapse after shrub removal shifts tundra
607 ecosystem to a methane source. *Nature Climate Change* 5, 67-70.
- 608 Novozamsky, I., Houba, V.J.G., van Eck, R., van Vark, W., 1983. A novel digestion technique for multi -
609 element plant analysis. *Communications in Soil Science and Plant Analysis* 14, 239-248.
- 610 Oechel, W.C., Hastings, S.J., Vourlitis, G., Jenkins, M., Riechers, G., Grulke, N., 1993. Recent change of Arctic
611 tundra ecosystems from a net carbon dioxide sink to a source. *Nature* 361, 520-523.
- 612 Poorter, H., Niklas, K.J., Reich, P.B., Oleksyn, J., Poot, P., Mommer, L., 2012. Biomass allocation to leaves,
613 stems and roots: meta-analyses of interspecific variation and environmental control. *New Phytologist* 193, 30-50.
- 614 Robinson, C., Kirkham, J., Littlewood, R., 1999. Decomposition of root mixtures from high arctic plants: a
615 microcosm study. *Soil Biology and Biochemistry* 31, 1101-1108.
- 616 Robinson, C.H., 2002. Controls on decomposition and soil nitrogen availability at high latitudes. *Plant and Soil*
617 242, 65-81.
- 618 Robinson, C.H., Michelsen, A., Lee, J., Whitehead, S., Callaghan, T.V., Press, M., Jonasson, S., 1997. Elevated
619 atmospheric CO₂ affects decomposition of *Festuca vivipara* (L.) Sm. litter and roots in experiments simulating
620 environmental change in two contrasting arctic ecosystems. *Global Change Biology* 3, 37-49.
- 621 Scheffer, R.A., Aerts, R., 2000. Root decomposition and soil nutrient and carbon cycling in two temperate fen
622 ecosystems. *Oikos* 91, 541-549.
- 623 Schuur, E.A.G., Vogel, J.G., Crummer, K.G., Lee, H., Sickman, J.O., Osterkamp, T.E., 2009. The effect of
624 permafrost thaw on old carbon release and net carbon exchange from tundra. *Nature* 459, 556-559.
- 625 Shaver, G.R., Chapin, F.S., III, 1991. Production: biomass relationships and element cycling in contrasting arctic
626 vegetation types. *Ecological Monographs* 61, 1-31.
- 627 Snyder, J.M., Rejmánková, E., 2015. Macrophyte root and rhizome decay: the impact of nutrient enrichment and
628 the use of live versus dead tissue in decomposition studies. *Biogeochemistry* 124, 45-59.
- 629 Strickland, M.S., Lauber, C., Fierer, N., Bradford, M.A., 2009. Testing the functional significance of microbial
630 community composition. *Ecology* 90, 441-451.
- 631 Sullivan, P., Sommerkorn, M., Rueth, H., Nadelhoffer, K., Shaver, G., Welker, J., 2007. Climate and species
632 affect fine root production with long-term fertilization in acidic tussock tundra near Toolik Lake, Alaska.
633 *Oecologia* 153, 643-652.
- 634 Swift, M.J., Heal, O.W., Anderson, J.M., 1979. *Decomposition in Terrestrial Ecosystems*. University of
635 California Press, Berkeley and Los Angeles.

- 636 Tape, K., Sturm, M., Racine, C., 2006. The evidence for shrub expansion in northern Alaska and the Pan -
637 Arctic. *Global Change Biology* 12, 686-702.
- 638 Tarnocai, C., Canadell, J., Schuur, E., Kuhry, P., Mazhitova, G., Zimov, S., 2009. Soil organic carbon pools in
639 the northern circumpolar permafrost region. *Global Biogeochemical Cycles* 23, GB2023.
- 640 Thormann, M.N., Bayley, S.E., Currah, R.S., 2001. Comparison of decomposition of belowground and
641 aboveground plant litters in peatlands of boreal Alberta, Canada. *Canadian Journal of Botany* 79, 9-22.
- 642 Veen, G.F., Freschet, G.T., Ordonez, A., Wardle, D.A., 2015a. Litter quality and environmental controls of
643 home-field advantage effects on litter decomposition. *Oikos* 124, 187-195.
- 644 Veen, G.F., Sundqvist, M.K., Wardle, D.A., 2015b. Environmental factors and traits that drive plant litter
645 decomposition do not determine home-field advantage effects. *Functional Ecology* 29, 981-991.
- 646 Verbyla, D., 2008. The greening and browning of Alaska based on 1982–2003 satellite data. *Global Ecology and*
647 *Biogeography* 17, 547-555.
- 648 Wagenmakers, E.-J., Farrell, S., 2004. AIC model selection using Akaike weights. *Psychonomic Bulletin &*
649 *Review* 11, 192-196.
- 650 Walker, D.A., Raynolds, M.K., Daniëls, F.J., Einarsson, E., Elvebakk, A., Gould, W.A., Katenin, A.E., Kholod,
651 S.S., Markon, C.J., Melnikov, E.S., 2005. The circumpolar Arctic vegetation map. *Journal of Vegetation Science*
652 16, 267-282.
- 653 Wallenstein, M.D., McMahon, S., Schimel, J., 2007. Bacterial and fungal community structure in Arctic tundra
654 tussock and shrub soils. *FEMS Microbiology Ecology* 59, 428-435.
- 655 Wang, P., Heijmans, M.M.P.D., Mommer, L., van Ruijven, J., Maximov, T.C., Berendse, F., 2016a.
656 Belowground plant biomass allocation in tundra ecosystems and its relationship with temperature.
657 *Environmental Research Letters* 11, 055003.
- 658 Wang, P., Mommer, L., van Ruijven, J., Berendse, F., Maximov, T.C., Heijmans, M.M.P.D., 2016b. Seasonal
659 changes and vertical distribution of root standing biomass of graminoids and shrubs at a Siberian tundra site.
660 *Plant and Soil* 407, 55-65.
- 661 Ward, S.E., Orwin, K.H., Ostle, N.J., Briones, M.J.I., Thomson, B.C., Griffiths, R.I., Oakley, S., Quirk, H.,
662 Bardgett, R.D., 2015. Vegetation exerts a greater control on litter decomposition than climate warming in
663 peatlands. *Ecology* 96, 113-123.
- 664 Webb, E.E., Schuur, E.A.G., Natali, S.M., Oken, K.L., Bracho, R., Kraepke, J.P., Risk, D., Nickerson, N.R.,
665 2016. Increased wintertime CO₂ loss as a result of sustained tundra warming. *Journal of Geophysical Research:*
666 *Biogeosciences* 121, 249-265.
- 667 Wieder, W.R., Bonan, G.B., Allison, S.D., 2013. Global soil carbon projections are improved by modelling
668 microbial processes. *Nature Climate Change* 3, 909-912.
- 669 Wookey, P.A., Aerts, R., Bardgett, R.D., Baptist, F., Bråthen, K.A., Cornelissen, J.H.C., Gough, L., Hartley, I.P.,
670 Hopkins, D.W., Lavorel, S., Shaver, G.R., 2009. Ecosystem feedbacks and cascade processes: understanding
671 their role in the responses of Arctic and alpine ecosystems to environmental change. *Global Change Biology* 15,
672 1153-1172.
- 673 Zhang, D., Hui, D., Luo, Y., Zhou, G., 2008. Rates of litter decomposition in terrestrial ecosystems: global
674 patterns and controlling factors. *Journal of Plant Ecology* 1, 85-93.
- 675 Zimov, S.A., Schuur, E.A.G., Chapin, F.S., III, 2006. Permafrost and the global carbon budget. *Science* 312,
676 1612-1613.
- 677

678 **Supplementary materials**

679



680

681 Fig. S1. Home-field advantage index (HFAI) of litter from different tissue types. Letters above error

682 bars indicate significant difference between groups. Asterisks show that HFAI was significantly

683 different from zero for live root and dead leaf litter. Dotted line represents zero level of HFAI.

684 Symbols with error bars show mean \pm SE, n = 8 blocks.

685

686 Table S1. Overview of the collection and preparation of samples used in this experiment

	Litter type	Time of collection	Location of collection	Drying method	Drying time
Live materials	<i>E. vaginatum</i> live leaves	July 2013	<i>E. vaginatum</i> dominated vegetation	65 °C oven dried	72 h
	<i>B. nana</i> live leaves	July 2013	<i>B. nana</i> dominated vegetation	65 °C oven dried	72 h
	<i>B. nana</i> live roots	July 2013	<i>B. nana</i> dominated vegetation	65 °C oven dried	72 h
	<i>E. vaginatum</i> live roots	July 2015	Nearby <i>E. vaginatum</i> dominated vegetation	~ 10 °C air dried	24 h
Dead materials	<i>E. vaginatum</i> dead leaves	July 2015	Nearby <i>E. vaginatum</i> dominated vegetation	~ 10 °C air dried	24 h
	<i>E. vaginatum</i> dead roots	July 2015	Nearby <i>E. vaginatum</i> dominated vegetation	~ 10 °C air dried	24 h
	<i>B. nana</i> dead leaves	July 2015	Nearby <i>B. nana</i> dominated vegetation	~ 10 °C air dried	24 h

687

688

689 Table S2. Comparison of the regression models of mass loss and chemical characteristics using AIC
690 values and Akaike weights.

Model parameter	Log-likelihood	AIC	Δ AIC	Akaike weight
N	6.95	-7.91	12.38	0.001
P	13.15	-20.29	0.00	0.479
Lignin	6.26	-6.52	13.77	0.001
C:N	7.69	-9.38	10.91	0.002
C:P (ln transformed)	11.68	-17.36	2.93	0.111
Lignin:N (ln transformed)	9.31	-12.63	7.66	0.010
Lignin:P (ln transformed)	12.95	-19.91	0.38	0.396

691