

1 **Local diversity of native arbuscular mycorrhizal symbionts differentially affects growth and nutrition of three**
2 **crop plant species**

3 Alessandra Turrini^{1*a}, Alberico Bedini^{1,2a}, Mario Bonilla Loo¹, Gaia Santini¹, Cristiana Sbrana³, Manuela
4 Giovannetti¹, Luciano Avio^{3*}

5 ¹ Department of Agriculture, Food and Environment, University of Pisa, Italy

6 ² INOQ GmbH, Schnega, and Humboldt-Universität zu Berlin, Department of Biology, Germany

7 ³ Institute of Agricultural Biology and Agrobiotechnology, CNR, UOS Pisa, Italy

8

9

10 ^aThese two authors contributed equally to the work

11

12 *Authors for correspondence:

13 Alessandra Turrini, Department of Agriculture, Food and Environment, University of Pisa, Via del Borghetto 80,
14 56124, Phone: +39 0502216646, Fax: +39 0502210606, e-mail address: alessandra.turrini@unipi.it;

15 Luciano Avio, Institute of Agricultural Biology and Agrobiotechnology, CNR, UOS Pisa, Via Moruzzi 1, Phone: +39
16 0502216646, e-mail address: avio@ibba.cnr.it

17

18 ORCID of authors:

19 Alessandra Turrini: 0000-0002-7186-4418

20 Cristiana Sbrana: 0000-0002-8058-8566

21 Manuela Giovannetti: 0000-0002-0716-7837

22 Luciano Avio: 0000-0003-2468-3400

23

24 **Abstract**

25 Intact whole native AMF communities occurring across a 100 m long field were used for the evaluation of plant
26 performance, as determined by the actual fungal species colonizing host roots. The soil from distinct plots within a
27 “hot spot” field was collected to set up 54 experimental units where three different plant species were grown, in order
28 to test whether the whole native AMF communities were able to differentially affect plant growth, to assess the
29 genetic identity of the AMF actually colonizing the tested plants and to analyze their community composition in the
30 different hosts. Molecular analyses revealed that plant growth and nutrition of the crop plants was differentially
31 affected by the diverse native arbuscular mycorrhizal communities colonizing the roots of the three plants, whose

32 performance varied depending on the identity of plant hosts and fungal symbionts, more than on a rich and
33 diversified AMF community. Such results, improving our understanding of AMF distribution at the local scale,
34 represent a starting point allowing the selection, isolation and characterization of the most efficient AMF
35 assemblages to be used as inoculants in sustainable food production systems.

36
37 **Keywords:** root AMF communities; small scale AMF diversity; small ribosomal subunit (SSU rDNA); plant
38 performance; functional diversity.

39
40 **Introduction**

41 Arbuscular mycorrhizal fungi (AMF, Glomeromycotina, Spatafora et al. 2016) represent a group of beneficial soil
42 biota which establish root symbioses with the majority of land plants, including the most important food crops. They
43 provide essential ecosystem services, improving plant nutrient availability, soil structure formation, and tolerance to
44 biotic and abiotic stresses, and promote plant diversity (Gianinazzi et al. 2010; Smith and Read 2008). In addition,
45 they enhance the biosynthesis of beneficial phytochemicals in food plants, representing a valuable and innovative
46 tool for the healthy food production chain (Rouphael et al. 2015; Sbrana et al. 2014). After establishing the
47 symbiosis, AMF produce extensive underground extraradical mycelia that spread from the roots into the
48 surrounding soil, and uptake and transport soil nutrients, mainly P and N, to the host plants (Battini et al. 2017; Pepe
49 et al. 2017); in exchange, AMF receive C compounds, which they are unable to synthesize, or to feed off as
50 saprotrophic organisms (Smith and Read 2008). The availability of soil mineral nutrients is also synergistically
51 promoted by a diversified community of beneficial bacteria thriving in association to AMF spores and extraradical
52 mycelium (Agnolucci et al. 2015; Battini et al. 2016; Hildebrandt et al. 2006; Morrison et al. 2017).

53 AMF are globally distributed, and have been found in the most diverse plant biomes, from grasslands to
54 desert, forest, shrublands, agriculture and wetlands, wherever their host plants were reported to occur (Allen et al.
55 1995; Oehl et al. 2017; Öpik et al. 2006; Read 1991; Treseder and Cross 2006). Investigations on the distribution of
56 AMF at the global scale revealed its relationship with plant community composition, environmental variables,
57 dispersal ability and geographical distance (Davison et al. 2015; Kivlin et al. 2011; Öpik et al. 2006; Treseder and
58 Cross 2006). Although spatial factors on a broad global scale were considered the main elements affecting AMF
59 distribution (Kivlin et al. 2011; Turrini and Giovannetti 2012), a recent work reported a low level of AMF
60 endemism, with 93% of taxa occurring in multiple continents, suggesting that AMF distribution among sites may be
61 driven by very efficient dispersal vectors, including wind, birds, water and human activities (Davison et al. 2015).

62 A few studies performed at the regional scale reported that AMF community composition and diversity
63 were not affected by geographical distance (max. 100 km) in apple orchards across Belgium (Van Geel et al. 2015),
64 in coffee plantations across Ethiopia (max. 82 km) (De Beenhouwer et al. 2015) and in *Trifolium repens* and *Lolium*
65 *perenne* across Ireland (max. 392 km) (Hazard et al. 2013). By contrast, the distribution of AMF species was
66 strongly affected by geographical distance in agricultural soils across England (max. 250 km) and Switzerland (max.
67 294 km) (Jansa et al. 2014; Van der Gast et al. 2011). A recent molecular study, carried out at a more local scale
68 (max. 27 km), confirmed such findings, as different AMF communities were detected in two geographical locations
69 in South Tyrol (Turrini et al. 2017).

70 Works focused on AMF genetic diversity and distribution at the local scale showed high differences among
71 AMF genotypes and communities occurring in the same field (Croll et al. 2008; Koch et al. 2004), when separated
72 by a few meters (Stukenbrock and Rosendahl 2005) or even at the sub-meter scale (Mummey and Rillig 2008).
73 Moreover, a high functional diversity was found within AMF species, even among genotypes originating from a
74 single field (Angelard et al. 2010; Croll et al. 2008; Koch et al. 2004; Munkvold et al. 2004).

75 Knowledge of AMF diversity and community composition at a small scale is relevant not only for
76 improving our understanding of AMF distribution, but also for functional characterization studies of the native AMF
77 communities, in order to isolate and reproduce the most efficient ones to be used as inoculants in sustainable and
78 organic agriculture. Actually, while it has been long known that distinct assemblages of AMF species may occur in
79 different plants (host preference) (Bever et al. 2009; Johnson et al. 2004; Koorem et al. 2017; Rodríguez-Echeverría
80 et al. 2017; Vandenkoornhuysen et al. 2003), and determine plant productivity, after isolation and inoculation in soil
81 microcosms (van der Heijden et al. 1998), only a few studies analyzed the symbiotic performance of whole native
82 AMF communities, utilized as inoculum (Burrows and Pflieger 2002; Moora et al. 2004; Uibopuu et al. 2012).

83 A previous study identified a site with a very high AMF richness, within the UNESCO Man and Biosphere
84 Reserve, named 'Selva Pisana'
85 (<http://www.unesco.org/mabdb/br/brdir/directory/biores.asp?code=ITA+08&mode=all>). Such a site, represented by
86 a small field, used periodically for crop production since 1974, but often left uncultivated, is relevant for the study of
87 local AMF diversity, as it has been described as a global "hot spot" of AMF species richness, encompassing 58
88 different AMF species belonging to 14 genera, the highest number reported so far from a single site (Njeru et al.
89 2015), and for studies on the symbiotic performance of whole native AMF communities. In the present study, we
90 hypothesized that native AMF communities may show differential symbiotic performance, as determined by the
91 actual fungal species composition occurring in the roots of host plants and by the identity of the plant host. We used
92 the soil from six distinct plots within the "hot spot" field to set up 54 experimental units where three different plant

93 species were grown, *Allium cepa*, *Capsicum annuum* and *Lactuca sativa*. The main aims of this multimodal study
94 were: i) to test whether the whole native AMF communities originating from the six distinct plots were able to
95 differentially affect the growth of the three plant species, ii) to assess the genetic identity of the AMF actually
96 colonizing the tested plants and iii) to analyse their community composition in the different hosts. The results
97 obtained will improve our understanding of AMF distribution at the local scale, that is relevant for functional
98 characterization studies of native AMF communities to be used as inoculants in sustainable food production
99 systems.

100

101 **Materials and Methods**

102 **Field experimental site and soil sampling**

103 The experimental site was located at Interdepartmental Centre for Agri-environmental Research ‘Enrico Avanzi’
104 (CIRAA), University of Pisa, S. Piero a Grado, Pisa, Italy (latitude 43° 40’ N, longitude 10° 19’ E), within the
105 UNESCO Man and Biosphere Reserve named ‘Selva Pisana’. Since 1974 the field was periodically used for the
106 production of various crops (maize, different horticultural crops, durum wheat, perennial alfalfa) and was also
107 uncultivated for several years (from 1988 to 1997, 2006 to 2011, and 2014 to 2015), allowing the development of
108 spontaneous flora. Since October 2011 to September 2012 the field site was used for cover crops and tomato crop
109 production under the EU-RTD FP7-funded project Strategies for Organic and Low input Integrated Breeding and
110 Management (SOLIBAM 2010-2014) during which the soil was evaluated for AMF occurrence and diversity. In an
111 area of about 2600 m², 58 AMF species, belonging to 14 genera were identified by classical morphological spore
112 identification (Njeru et al. 2015).

113 Soil samples were collected from 12 field plots (3x4m each), distributed in an area of 13x114m in spring 2015 (Fig.
114 1). In each plot a “X” shaped sampling pattern was used, in which 5 points were selected, one at the centre of the
115 plot and the other four 1 m far from the first. Soil cores from each point were taken at a depth of 5 to 15 cm, then
116 mixed and labelled as a single sample. Finally soil samples were air dried and then sieved through a 5 mm mesh
117 sized sieve.

118

119 **Soil characterization**

120 ***Soil physical and chemical analyses***

121 Soil samples were analysed for: total N, available P, soil organic matter, pH_(H2O), cation exchange capacity, and
122 texture. Total N was evaluated by the macro-Kjeldahl digestion procedure (Bremner and Mulvaney 1982) and
123 available P was determined by colorimetric analysis using the Olsen method (Olsen and Sommers 1982). Organic

124 matter was determined using the modified Walkley-Black wet combustion method (Nelson and Sommers 1982).
125 Soil pH was measured potentiometrically in a 1:2:5 soil water suspension, the other soil parameters were determined
126 according to the standard methods (Gee and Bauder 1986; Sumner and Miller 1996).

127
128 ***Mycorrhizal inoculum potential of the field soil***
129 Mycorrhizal inoculum potential (MIP) bioassay was performed to verify the activity of AMF propagules occurring
130 in the soil of each sample and was assessed using *Cichorium intybus* L. as host plant. Five 50 ml replicate tubes
131 were filled with 40 ml of sampled field soil and sown using the biotest plant. Then they were put in sun-transparent
132 bags and maintained in a growth chamber at 25 °C under a 16/8 h light/dark daily cycle. Four days after
133 germination, plants were thinned to three per tube and harvested 28 days after sowing, by removing roots from soil
134 and washing them with tap water. Roots were then cleared in 10% KOH in a 80 °C water bath for 15 min and
135 stained with Trypan blue in lactic acid (0.05 %) after 10 min in 2 % aqueous HCl. The percentage of AMF
136 colonization was calculated using a dissecting microscope at x25 or x40 magnification and the gridline intersect
137 method (Giovannetti and Mosse 1980).

138
139 **Evaluation of the symbiotic performance of native AMF communities**
140 On the basis of similarity indices of their soil properties, plots 2, 5, 9, 10, 14 and 22 were selected to set up the
141 microcosm experiments aimed at evaluating the performance of three host plants: *A. cepa* cv. Rossa dolce di Tropea
142 (onion), *C. annuum* cv. Rosso quadrato d'Asti (pepper) and *L. sativa* cv. Foglia di quercia (lettuce) (Fig. 1). Onion,
143 pepper and lettuce were seeded in a sterile calcinated clay (OILDRI Chicago, IL, USA) and grown in a growth
144 chamber at 25 °C under a 16/8 h light/dark daily cycle. Ten days after sowing plantlets were transferred to 8-cell
145 trays (5 cm diameter), which were filled with soil from the 6 field plots selected as described above. Soil from each
146 plot represented a different native AMF treatment, hereafter named with the number of the original soil plot. Three
147 replicate trays of each host plant species were set up for the 6 native AMF treatments. Onion, pepper and lettuce
148 plants were grown in a glasshouse for 8, 5 and 6 weeks, respectively, under ambient natural light and temperature
149 conditions (min T 15-20°C , max T 27-33°C), relative humidity from 50 to 80 % and supplied with tap water as
150 needed. The differential harvest times were chosen on the basis of the length of host life cycles.

151 At harvest, plants were analysed for shoot dry matter and shoot N and P concentrations (Jones et al. 1991). Roots
152 from the 8 plants of each replicate tray were collected and pooled to assess mycorrhizal root colonization, using the
153 protocol described above for MIP analysis. Aliquots (100 mg) of the pooled roots were collected for DNA extraction
154 and stored at -80 °C.

155
156 **Molecular analyses**
157 The root samples of three replicates of onion, pepper and lettuce from native AMF treatments 2 and 22 were used
158 for DNA extraction (Fig. 1). Genomic DNA was obtained from roots ground in liquid nitrogen by using DNeasy
159 Plant Mini Kit (Qiagen Milan, Italy), and 1 µl of extracted DNA was used as template in PCR reactions. Partial
160 small subunit (SSU) of ribosomal RNA gene fragments were amplified in volumes of 25 µl with 0.125 U of GoTaq
161 G2 Flexi DNA Polymerase (Promega, Milan, Italy), 0.4 µM of each primer (AML1/AML2, Lee et al. 2008), 0.2
162 mM of each dNTP, 1.5 mM of MgCl₂ and 1× the manufacturer's reaction buffer. The thermal cycler (Eppendorf
163 Mastercycler personal, Eppendorf, Milan, Italy) was programmed as follows: a manual hot start at 94 °C for 3 min,
164 30 cycles at 94 °C for 30 s, 58 °C for 40 s, 72 °C for 55 s and a final extension step at 72 °C for 10 min. Reactions
165 yields were estimated by using a 1 % agarose gel containing ethidium bromide (0.5 µg ml⁻¹). Wizard® SV Gel and
166 PCR Clean-up system (Promega) was used to purify amplicons from onion, pepper and lettuce roots, which were
167 then ligated into pGem-T Easy vector (Promega) to transform XL10-Gold Ultracompetent *Escherichia coli* cells
168 (Stratagene, La Jolla, CA, USA). The composition of the AM fungal communities was determined using PCR-RFLP
169 screening of clone libraries (AML1/AML2 primers and HinfI and AluI restriction enzymes). Forty clones per clone
170 library were screened by PCR-RFLP analysis. Plasmids of representative clones of each RFLP pattern in each
171 library were purified by Wizard® Plus SV Minipreps (Promega) and sequenced using T7 vector primers at GATC
172 Biotech (Köln, Germany). The new unique cloned sequences reported in this study have been deposited in the
173 'European Nucleotide Archive' (<http://www.ebi.ac.uk/ena>) under the accession numbers LT856601-LT856682
174 (study_id PRJEB21051).

175
176 **Bioinformatics**
177 Sequences from *E. coli* libraries were edited in MEGA 6.0 and their similarities were determined using the Basic
178 Local Alignment Search Tool (BLASTn) provided by NCBI. The detection of chimeric sequences was performed
179 using USEARCH 6.0 (http://fungene.cme.msu.edu/FunGenePipeline/chimera_check/form.spr). Sequences were
180 aligned with those corresponding to the closest matches from GenBank as well as with sequences from major clades
181 of Glomeromycotina using MUSCLE as implemented in MEGA6. Phylogenetic trees were inferred by Neighbour-
182 joining analysis. The evolutionary distances were computed using the Maximum Composite Likelihood method.
183 The confidence of branching was assessed using 1000 bootstrap resamplings.

184
185 **Statistical analyses**

186 Chemical and physical data of soil plots were used to compute Euclidean similarities indices in PAST 3.0 after
187 standardization. Root colonization (after arcsin transformation) and shoot dry weight were analysed by two way
188 ANOVA and means were separated by simple main effect test with Sidak adjustment. Nutrient content data did not
189 fulfill two way ANOVA assumptions and were analyzed by one way ANOVA followed by Tukey HSD for mean
190 separation. Analyses were carried out in IBM SPSS statistics version 24 software (IBM Corporation, Armonk, NY,
191 USA).

192 Estimates of community diversity were determined as richness (S), bias-corrected Chao1 richness, Shannon
193 (H) and Simpson diversity (1-D) index and evenness (e^H/S). The indices were calculated using PAST 3.0 and 1000
194 bootstraps were used to determine confidence intervals. Non parametric Kruskal-Wallis analysis was used to
195 determine differences in the diversity indices among AMF communities colonizing the roots of the three different
196 plant species grown in treatments 2 and 22. We determined the rarefaction curves with PAST 3.0 software to
197 estimate whether the number of screened sequenced were sufficient to capture AMF diversity of each host. AMF
198 communities were also evaluated by two ways permutational multivariate analysis of variance (PERMANOVA), to
199 test the effects of soil microcosm and the host plant species, performed in PAST 3.0. Multivariate analyses (PCA
200 and RDA) were performed using Canoco 5.0.

201

202 **Results**

203 **Analyses of soil samples**

204 *Soil physical and chemical characteristics*

205 Soil physical and chemical analyses showed that the experimental site had a sandy loam soil texture with low
206 available P (Olsen) values (Table 1). A low degree of variability was detected among the different plots; for
207 example, N concentration ranged in most plots from 1.55‰ to 1.91‰, showing the lowest (1.32‰) and highest
208 (2.23‰) values in plots 17 and 21, respectively. Organic matter ranged from 2.5% to 2.8% in the majority of plots,
209 with minimum and maximum of 1.63% and 3.01%, in plots 17 and 5, respectively. However, Euclidean similarity
210 indices of soil properties ranged from 0.84, in the case of the plot pair showing the highest similarity (9 vs. 14), to
211 7.07, in the case of the plot pair with the lowest similarity (17 vs. 21). In order to select the most similar plots, in
212 terms of physical and chemical soil properties, we sorted all the pairwise comparison by similarity, and selected five
213 pairs (2/5, 9/10, 9/14, 10/14, 2/22) with similarity values ranging from 0.84 to 1.82, involving 6 out of 12 field plots.
214 The soil from the 6 selected field plots was used to set up the native AMF treatments for the plant performance
215 experiment.

216 *MIP bioassay*

217 The analysis of the activity of AMF soil propagules, as assessed by the MIP bioassay, showed no significant
218 differences among the soil originating from all 12 plots ($F_{11,33}$, $P=0.81$). The percentage of mycorrhizal root length
219 of biotest plants (*C. intybus*) varied from 37.7 ± 2.9 to 55.7 ± 3.7 , showing a good level of AMF activity in all plots.

220

221 **Plant performance in microcosm experiment**

222 The percentage of colonized root length was affected by host plant species and native AMF treatments and their
223 interaction (Table 2). Colonization levels in onion, the most mycotrophic plant of the experiment, were significantly
224 higher than those detected in lettuce and pepper (Table 3).

225 Significantly different growth responses among native AMF treatments were observed within each of the
226 three plant species (Table 2). Lettuce plants grown in treatment 22 showed the highest shoot dry weight (SDW),
227 while the lowest SDW value was observed in treatment 2 (Table 3). Consistently with the previous results, the
228 highest and lowest SDW values of pepper plants were in treatments 22 and 2, respectively. Statistically significant
229 differences were observed also in the growth of onion plants, with the highest and lowest biomass in treatment 5 and
230 2 (Tables 2, 3).

231 Native AMF treatments produced significant differences in N shoot content of lettuce and pepper (Table 2).
232 The highest N contents were detected in all host plants growing in treatment 22, while the lowest contents were
233 found in treatment 2 (lettuce and onion), and in treatment 14 (pepper) (Table 3).

234 Statistically significant differences in P content, among native AMF treatments, were detected in lettuce
235 and pepper (Table 2). The lowest values were shown by plants grown in treatment 2, while the highest occurred in
236 treatments 22, and 5, depending on the plant species (Table3).

237 In order to select the root samples for molecular analyses, we evaluated plant responses, in terms of plant
238 growth and nutrient uptake, obtained in the different treatments. To this aim, the selected five treatment pairs
239 associated with the most similar soil in terms of physical and chemical properties, were compared for their plant
240 performance. A radar graph, obtained by adding the relative increases in SDW, N and P content values of each pair
241 for the three plant species, allowed the detection of treatments 2 and 22 as the pair showing the largest plant
242 differential performance (Fig. 2a). The principal component analysis (PCA) performed using the same plant
243 response variables supported the selection of treatments 2 and 22 for molecular analyses (Fig. 2b).

244

245

246 **Molecular identification of native AMF communities**

247 The DNA extracted from onion, lettuce and pepper roots of treatments 2 and 22 was successfully amplified using
248 the primer pair AML1/AML2, obtaining a fragment of the expected size (~800 bp). A total of 720 clones from the
249 18 clone libraries were examined for the presence of the insert (40 clones/library) and 707 positive clones were
250 screened by RFLP analysis, obtaining 19 RFLP groups. For each RFLP group, clones of the different libraries were
251 sequenced accounting for a total of 214 sequences. BLAST analyses showed that all the sequences had a high
252 similarity (97-100% identity) to glomeromycotan sequences. No chimeric sequences were found. All non-redundant
253 sequences from the 18 clone libraries (82 out of 214) and 25 references from GenBank were used for neighbour-
254 joining phylogenetic analyses (Fig. 3). After RFLPs, BLASTn and phylogenetic analyses, the sequences were
255 grouped into 14 OTUs supported by a bootstrap value >84%. Of these, 10 belonged to the family Glomeraceae, one
256 to Gigasporaceae, one to Claroideoglomeraceae and two to Paraglomeraceae. Among Glomeraceae, five genera
257 (*Dominikia*, *Funneliformis*, *Rhizoglopus*, *Sclerocystis*, *Septoglopus*) were retrieved. *Racocetra*, *Claroideoglopus*
258 and *Paraglopus* were the genera found for Gigasporaceae, Claroideoglomeraceae and Paraglomeraceae,
259 respectively. The genus *Rhizoglopus* was the most abundant, accounting for 84% of the total sequences, followed
260 by *Septoglopus* (6.1%), *Claroideoglopus* (3.5%), *Funneliformis* (2.8%) and *Paraglopus* (2.0%). *Dominikia*,
261 *Sclerocystis* and *Racocetra* sequences corresponded to less than 1 % of total sequences.

262 Within the genus *Rhizoglopus*, one OTU (Rh3, 11.2% of the total sequences) was identified as
263 *Rhizoglopus irregulare* (synonym *Rhizophagus irregularis*, basionym *Glomus irregulare*); the remaining three
264 OTUs (Rh1, Rh2, Rh4) represented sequences of uncultured species (Table 4). Rh1 and Rh2 were the most abundant
265 sequences, accounting for 54.5% and 16.1% of the total, respectively. One OTU (Fun1, 2.4% of total sequences) in
266 the genus *Funneliformis* was identified as *Funneliformis mosseae* (Table 4), while sequences grouped in the OTU
267 Fun2 (0.4% of total sequences) were close to *Funneliformis geosporus* (99% similarity) (Fig. 3). The remaining
268 OTUs within Glomeraceae (Glo, Scle, Sept1 and Sept2) matched to sequences of either known (*Dom*, *Dominikia*
269 *iranica* basionym *Glomus iranicum*), or unknown species (*Sclerocystis* sp., *Septoglopus* sp.). Sequences grouped in
270 the OTU Clar showed a high similarity with *Claroideoglopus etunicatum* (Table 4), forming a unique clade together
271 with the sequences of such species in the phylogenetic analysis (Fig. 3). OTUs Rac and Par2 matched sequences of
272 *Racocetra fulgida* and *Paraglopus laccatum* (Table 4), and Par1 sequences showed high homology with sequences
273 of uncultured *Paraglopus* species. MaarjAM database (<http://maarjam.botany.ut.ee/>, accessed on February 2017)
274 was used to confirm the assignment of detected OTUs to sequences of Glomeromycotina. The number of analysed
275 sequences was generally sufficient to capture the AMF diversity in the roots of most host plants as shown by
276 rarefaction analyses, since the curves almost reached the asymptote (Online Resource 1).

277

278 **AMF root community composition as affected by native AMF treatment and host plant species**

279 The composition of AMF root communities differed significantly, when considering both the two native AMF
280 treatments (2 and 22, $P < 0.001$), the host plants (*A. cepa*, *C. annuus* and *L. sativa*, $P < 0.001$), and the interaction
281 between native AMF and host plants ($P = 0.024$) as revealed by two-ways PERMANOVA analysis. We detected 6
282 and 13 OTUs in the roots of plants from treatments 2 and 22, respectively (Fig. 4). All the OTUs in treatment 2 were
283 found also in 22, except one (Rac), matching *R. fulgida* sequences, but their frequencies were very different between
284 the two treatments. This trend was confirmed by diversity indices values, which showed significant differences in
285 richness (S), Chao and Shannon (H) diversity values related to the AMF communities occurring in the roots of all
286 the host plants from treatment 2, compared with 22 (Table 5). Interestingly, in onion and lettuce, Simpson (1-D)
287 index was significantly higher in AMF treatment 22 than in 2. Actually, root AMF communities found in treatment
288 2 were similar for all diversity parameters, while those found in treatment 22 showed a clear difference between
289 pepper and onion and lettuce.

290 Overall, in treatment 2, Rh1 was highly dominant, (83% of all sequences), while the OTUs Rh2, Sept2,
291 Fun1 (*F. mosseae*), Clar (*C. etunicatum*) and Rac (*R. fulgida*) were found at very low percentages (8.7%, 3.7%,
292 3.7%, 0.6%, 0.3%, respectively). Interestingly, lettuce hosted all six OTUs, while onion missed Clar and Rac, and
293 pepper harbored only Rh2 in association with Rh1 (Fig. 4).

294 On the contrary, in treatment 22, we found a more homogeneous distribution of OTUs, as Rh1, Rh2, Rh3
295 occurred each in the range 22.5% - 25.4% of total sequence number. In treatment 22, onion and lettuce hosted the
296 highest amounts of OTUs (12 and 11, respectively), while only 7 OTUs were detected in pepper roots, confirming
297 the low species richness obtained in treatment 2 in this host plant (Fig. 4). Rh1 and Rh2 sequences were highly
298 dominant in pepper (37.9% and 55.2%, respectively), while their frequencies were 17.8 and 10.2%, and 20.5 and
299 6%, in lettuce and onion, respectively. Other OTUs found in lettuce (Rh3, Rh4 and Clar) and in onion (Rh3, Sept2
300 and Clar) occurred with higher frequencies than in pepper. Moreover the AMF community composition change can
301 be ascribed to the occurrence of further OTUs in lettuce and onion (Fun1, Fun2, Glo, Par1, Par2, Scl and Sept1),
302 together representing 8.8% and 18.8% of the total sequences (Fig. 4).

303 Multivariate RDA analysis showed that 51.2% of the total variance (I and II axes) of AMF communities
304 occurring in roots of onion, pepper and lettuce from treatments 2 and 22 could be explained by native AMF
305 treatments and host species (Fig. 5). Moreover, Monte Carlo permutation tests on RDA confirmed that such
306 variables significantly affected the composition of root AMF communities ($P = 0.002$).

307

308 **Discussion**

309 This work shows that diverse whole native arbuscular mycorrhizal communities, occurring across a 100 m long
310 field, differentially affected growth and nutrition of three crop plant species, *A. cepa*, *C. annuum* and *L. sativa*. Our
311 data revealed that plant performance varied depending on the identity of host plants and fungal symbionts, more
312 than on a rich and diversified native AMF community.

313

314 **Plant performance in the different whole native AMF community treatments**

315 The six different native AMF treatments showed variable plant performances in the three host species. Treatment 22
316 produced, in most cases, the highest plant SDW, N and P content, compared with the other five native AMF
317 treatments. In particular, it significantly enhanced pepper and lettuce SDW, by 57 and 119%, respectively, compared
318 with treatment 2. Treatment 22 was the most efficient in N shoot uptake, although the increases in N levels varied in
319 the different plant species (18, 37 and 98% in onion, pepper and lettuce, respectively), compared with treatment 2. P
320 contents followed the same trend, with variable increases depending on the identity of the host plant. Differential
321 plant responses to AMF (functional host preference, see Walder and van der Heijden 2015) may be explained by the
322 fact that during plant/fungal interactions plant gene expression is modulated depending on the plant and fungal
323 genotypes involved in the symbiosis (Feddermann et al. 2008; Hohnjec et al. 2005; Massoumou et al. 2007).

324 Here, intact whole native AMF communities were used for the evaluation of plant performance. Previous
325 works carried out experiments with diluted or modified native communities sampled from natural soils and utilized
326 as inocula (Burrows and Pflieger 2002; Ji et al. 2010; Moora et al. 2004; Uibopuu et al. 2012), while most studies
327 employed single isolated species or artificially assembled AMF communities (Gustafson and Casper 2006; Jansa et
328 al. 2008; Munkvold et al. 2004; van der Heijden et al. 1998). In this work, six selected native AMF treatments
329 showed variable effects on growth and nutrition of the diverse host plants, in agreement with other studies, where
330 different native AMF communities collected from grassland and forest or from young and old forest in Estonia,
331 produced differential plant performance in rare and common *Pulsatilla* species or in different plant species typical of
332 the Koeru boreonemoral forest (Moora et al. 2004; Uibopuu et al. 2012). Other authors showed that spores
333 originating from native AMF communities collected from serpentine and prairie grassland and used as inoculum
334 differed functionally, depending on their local host-soil environment (Ji et al. 2010). Interestingly our results were
335 obtained using native AMF originating from sampling points within a single field, in a very short distance, whereas
336 the communities analysed in the previous papers belonged to distant sites and to different biomes. Here, in order to
337 assess the relationship between plant performance and the native AMF communities originating from the diverse
338 plots and actually colonizing the three host plants, further molecular analyses were performed.

339

340 **Molecular identification of native AMF communities**

341 Overall, we detected 14 OTUs in the roots of host plants, a number similar to that found in maize roots growing in
342 the same area (Turrini et al. 2016) and in other Mediterranean agroecosystems (Brito et al. 2012; Cesaro et al. 2008;
343 Pivato et al. 2007). The detection of sequences of *C. etunicatum*, *F. geosporus*, *F. mosseae*, *P. laccatum*, *R. fulgida*
344 and *R. irregulare*, whose spores were identified also by Njeru et al. (2015) in the same field site using a
345 morphological approach, confirmed their occurrence in this AMF hot spot within the Biosphere Reserve “Selva
346 Pisana”. Although molecular analyses of whole roots showed a lower number of fungal phylotypes compared with
347 laser microdissected arbusculated cells (Berruti et al. 2013), our work allowed the detection of additional ribotypes
348 corresponding to uncultured species, confirming that studies exclusively based on spore morphological
349 characterization are not sufficient to describe the whole biodiversity of AMF (Kivlin et al. 2011). Indeed, the
350 number and diversity of spores occurring in soil are not always correlated with AMF actively colonizing roots (Oehl
351 et al. 2005), as some AMF may occur in the soil only in the form of hyphae and colonized roots rather than spores
352 (Abbott and Gazey 1994). Moreover, morphological identification of spores may be either difficult for their partial
353 degradation or parasitization, or impossible as sporulation is seasonal-dependent in some AMF species (Helgason et
354 al. 2002, Oehl et al. 2005). Actually, spore production was reported to depend on fungal physiological parameters
355 and environmental conditions (Redecker 2002; Redecker et al. 2003). Our data suggest that a multidisciplinary
356 approach is the best strategy for a complete assessment of AMF biodiversity in natural and agro-ecosystems (Oehl et
357 al. 2010; Redecker et al. 2003).

358 Here, we retrieved sequences of *C. etunicatum*, *F. mosseae* and *R. irregulare*, considered generalist fast
359 root and soil colonizers, frequently found in arable soils (Oehl et al. 2003, 2005). Interestingly, we detected also
360 sequences of the rare species *R. fulgida*, which was reported to occur in coastal sand dunes not far from our
361 experimental field (Błaszowski et al 2004; Turrini et al. 2008). The finding of both generalist and rare species and
362 of both cultured and uncultured taxa may be related to the climate of the Mediterranean Basin (where temperate and
363 sub-tropical species co-occur), and also to the diversified land use history of the site during the past 40 years (Njeru
364 et al. 2015). The area was occasionally used for crop production, but it was also left unploughed, allowing the
365 development of a rich and diversified AMF community. Such findings agree with the results of a recent meta-
366 analysis (Ohsowski et al. 2014) on the occurrence of cultured and uncultured Glomeromycotina in different habitats
367 worldwide, reporting a greater association of cultured AM fungal taxa with human-impacted habitat and cultivated
368 plants when compared to uncultured AMF, usually occurring in natural habitats and wild plants. In this perspective,
369 the study site within the Mediterranean Basin, which is considered a hyper-hotspot of biodiversity (Myers et al.
370 2000), assumes a great importance for AMF biodiversity conservation.

371
372 **AMF root community composition as affected by native AMF treatment and host plant species**
373 The roots of plants growing in treatment 22 showed a more diversified AMF community, compared with treatment
374 2, where only one dominant species was detected, OTU Rh1, which represented 81-86% of the analyzed clones from
375 the three plants of treatment 2. Thus, assuming the high frequency of such sequences as prognostic of a main role of
376 this OTU, the lower plant performance reported in treatment 2, compared with treatment 22, could be ascribed to its
377 low efficiency, irrespective of host plant identity. On the other hand, it can be speculated that in pepper the better
378 performance in treatment 22 could be the result of the activity of a single distinct AMF OTU, as Rh2 dominance on
379 Rh1 may have changed the final outcome of the symbiotic relationship. In lettuce and onion, hosting a richer AMF
380 community, plant performance cannot be clearly ascribed to a single OTU, as an interaction among the different
381 OTUs may have occurred. However, such diverse AMF assemblages in treatment 22 produced a better plant
382 performance, compared with treatment 2, only in lettuce, confirming that AMF species identity may be more
383 important than community diversity (Njeru et al. 2017; Vogelsang et al. 2006). Indeed, the occurrence of a
384 diversified and rich AMF community has long been known to enhance plant growth and nutrition by different
385 mechanisms, i.e by functional complementarity (Koide 2000; Maherali and Klironomos 2007), by buffering
386 negative effects of either unfavorable abiotic and biotic stresses (Pringle and Bever 2002) or inefficient AMF
387 species, which may persist in host roots within a heterogeneous AMF community (Hart et al. 2013).

388 Interestingly, in all plant roots from treatment 22, we found phylogenetically highly related sequences,
389 affiliated to the genus *Rhizoglossus*, although with different frequencies, confirming the highly infective behaviour
390 of *Rhizoglossus* spp. (Alkan et al. 2006; Jansa et al. 2008). Previous works reported that phylogenetic relatedness
391 was positively associated with coexistence in the same root system and, interestingly, it was also positively
392 associated with plant growth (Roger et al. 2013). Though, almost nothing is known about the factors that control
393 AMF coexistence and competition in roots, which are processes depending on spatial scales, ecosystem types, host
394 plant quality and identity (Davison et al. 2016; Knecht et al. 2016).

395 Our data show that root AMF community composition varied with the identity of the three host plants. Host
396 plant identity has long been known to be one of the most important elements shaping AMF community composition
397 (Gollotte et al. 2004; Helgason et al. 2002; Mummey and Rillig 2008; Scheublin et al. 2004; Sýkorová et al. 2007;
398 Vandenkoornhuyse et al. 2003), more than habitat (Becklin et al. 2012), seasonality (Davison et al. 2011) or
399 agricultural practices (Vályi et al. 2015). Some authors suggested that AMF communities are not random
400 assemblages, but can differentially colonize the roots of ecologically diverse groups of plant species, depending on
401 AMF specific characteristics, i.e. habitat generalist vs. forest specialist AMF (Davison et al. 2011). Recently,

402 distinct AMF communities were found associated with different plant species within the same mixed cover crops (*V.*
403 *villosa* and *Trifolium* spp. vs *Avena* sp. and *P. tanacetifolia* and in the successive maize), revealing a strong effect of
404 the host on AMF communities actually occurring in the roots (Turrini et al. 2016).

405

406 **Conclusions**

407 Molecular analyses of host plant roots revealed that the best plant performance was not necessarily associated with a
408 richer and more diversified whole native AMF community. Indeed, AMF species belonging to Glomeraceae,
409 Acaulosporaceae, Gigasporaceae and Diversisporaceae may differentially activate important core sets of symbiosis-
410 associated genes, i.e. transport, defense, metabolic process (Fedderman et al. 2008; Massoumou et al. 2007), which
411 can be related to the functionality of the symbiosis. It is tempting to speculate that the three host plants assessed in
412 this experimental work may have benefited from the most efficient AMF species combinations in agreement with
413 new insights into AMF/plant interaction and cooperation, suggesting that host plant reward the best fungal
414 symbionts with a larger nutrient transfer (Kiers et al. 2011). Our results on the relationships between plant
415 performance and whole native AMF communities diversity represent a starting point allowing the isolation,
416 characterization and selection of single native AMF species from the “hot spot” site. Such genotypes could be
417 further studied in order to identify the most efficient ones, to be used as single inoculants and/or as assemblages in
418 sustainable food production systems.

419

420 **Acknowledgments**

421 This work was funded by the University of Pisa through the project “Molecular and functional biodiversity of plant
422 associated microorganisms” and through Fondi di Ateneo and by the National Research Council of Italy. The authors
423 wish to thank Prof. Marco Mazzoncini and Dr. Rosalba Risaliti for their precious help in preserving the field site
424 and determining soil physical and chemical properties.

425 The authors declare that they have no conflict of interest.

426

427 **References**

428 Abbott LK, Gazey C (1994) An ecological view of the formation of VA mycorrhizas. *Plant Soil* 159:69
429 Agnolucci M, Battini F, Cristani C, Giovannetti M (2015) Diverse bacterial communities are recruited on spores of
430 different arbuscular mycorrhizal fungal isolates. *Biol Fertil Soils* 51:379-389

431 Alkan N, Gadkar V, Yarden O, Kapulnik Y (2006) Analysis of quantitative interactions between two species of
432 arbuscular mycorrhizal fungi, *Glomus mosseae* and *G. intraradices*, by real-time PCR. Appl Environ Microbiol
433 72:4192-4199

434 Allen EB, Allen MF, Helm DJ, Trappe JM, Molina R, Rincon E. (1995) Patterns and regulation of mycorrhizal plant
435 and fungal diversity. Plant Soil 170:47-62

436 Angelard C, Colard A, Niculita-Hirzel H, Croll D, Sanders IR (2010) Segregation in a mycorrhizal fungus alters rice
437 growth and symbiosis-specific gene transcription. Curr Biol 20:1216-1221

438 Battini F, Cristani C, Giovannetti M, Agnolucci M (2016) Multifunctionality and diversity of culturable bacterial
439 communities strictly associated with spores of the plant beneficial symbiont *Rhizophagus intraradices*.
440 Microbiol Res 183:68-79

441 Battini F, Grønlund M, Agnolucci M, Giovannetti M, Jakobsen I (2017) Facilitation of phosphorus uptake in maize
442 plants by mycorrhizosphere bacteria. Sci Rep 7:4686 DOI:10.1038/s41598-017-04959-0

443 Becklin KM, Hertweck KL, Jumpponen A (2012) Host identity impacts rhizosphere fungal communities associated
444 with three alpine plant species. Microb Ecol 63:682-693

445 Berruti A, Borriello R, Lumini E, Scariot V, Bianciotto V, Balestrini R (2013) Application of laser microdissection
446 to identify the mycorrhizal fungi that establish arbuscules inside root cells. Front Plant Sci 4

447 Bever JD, Richardson SC, Lawrence BM, Holmes J, Watson M. (2009) Preferential allocation to beneficial
448 symbiont with spatial structure maintains mycorrhizal mutualism. Ecol Lett 12:13-21

449 Błaszczowski J, Blanke V, Renker C, Buscot F (2004) *Glomus aurantium* and *G. xanthium*, new species in
450 Glomeromycota. Mycotaxon 90:447-467

451 Bremner JM, Mulvaney CS (1982) Nitrogen total. In: Page AL, Miller RH, Keeney DR (Eds) Methods of Soil
452 Analysis. Part 2. American Society of Agronomy, Madison, WI, pp 595-624

453 Brito I, Goss MJ, de Carvalho M, Chatagnier O, van Tuinen D (2012) Impact of tillage system on arbuscular
454 mycorrhiza fungal communities in the soil under Mediterranean conditions. Soil Till Res 121:63-67

455 Burrows RL, Pflieger FL (2002) Host responses to AMF from plots differing in plant diversity. Plant Soil 240:169-
456 180

457 Cesaro P, van Tuinen D, Copetta A, Chatagnier O, Berta G, Gianinazzi S, Lingua G (2008) Preferential colonization
458 of *Solanum tuberosum* L roots by the fungus *Glomus intraradices* in arable soil of a potato farming area. Appl
459 Environ Microbiol 74:5776-5783

460 Croll D, Wille L, Gamper HA, Mathimaran N, Lammers PJ, Corradi N, Sanders IR (2008) Genetic diversity and
461 host plant preferences revealed by simple sequence repeat and mitochondrial markers in a population of the
462 arbuscular mycorrhizal fungus *Glomus intraradices*. *New Phytol* 178:672-687

463 Davison J, Moora M, Jairus T, Vasar M, Öpik M, Zobel M (2016). Hierarchical assembly rules in arbuscular
464 mycorrhizal (AM) fungal communities. *Soil Biol Biochem* 97:63-70

465 Davison J, Moora M, Öpik M, Adholeya A, Ainsaar L, Bâ A, Burla S, Diedhiou AG, Hiiesalu I, Jairus T, Johnson
466 NC, Kane A, Koorem K, Kochar M, Ndiaye C, Pärtel M, Reier Ü, Saks Ü, Singh R, Vasar M, Zobel M (2015)
467 Global assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism. *Science* 349:970–
468 973

469 Davison J, Öpik M, Daniell TM, Moora M, Zobel M (2011) Arbuscular mycorrhizal fungal communities in plant
470 roots are not random assemblages. *FEMS Microbiol Ecol* 78:103-115

471 De Beenhouwer M, Van Geel M, Ceulemans T, Muleta D, Lievens B, Honnay O (2015) Changing soil
472 characteristics alter the arbuscular mycorrhizal fungi communities of Arabica coffee (*Coffea arabica*) in
473 Ethiopia across a management intensity gradient. *Soil Biol Biochem* 91:133-139

474 Feddermann N, Boller T, Salzer P, Elfstrand S, Wiemken A, Elfstrand M (2008) *Medicago truncatula* shows
475 distinct patterns of mycorrhizarelated gene expression after inoculation with three different arbuscular
476 mycorrhizal fungi. *Planta* 227:671-680

477 Gee GW, JW Bauder (1986) Particle-size analysis. In A Klute (Ed.) *Methods of Soil Analysis, Part 1. Physical and*
478 *Mineralogical Methods. Agronomy Monograph. 9 2nd edn, American Society of Agronomy/Soil Science*
479 *Society of America, Madison, WI, pp 383-411*

480 Gianinazzi S, Gollotte A, Binet MN, van Tuinen D, Redecker D, Wipf D (2010) Agroecology: the key role of
481 arbuscular mycorrhizas in ecosystem services. *Mycorrhiza* 20:519-530

482 Giovannetti M, Mosse B (1980) An evaluation of techniques for measuring vesicular arbuscular mycorrhizal
483 infection in roots. *New Phytol* 84:489-500

484 Gollotte A, van Tuinen D, Atkinson D (2004) Diversity of arbuscular mycorrhizal fungi colonising roots of the grass
485 species *Agrostis capillaris* and *Lolium perenne* in a field experiment. *Mycorrhiza* 14:111-117

486 Gustafson DJ, Casper BB (2006) Differential host plant performance as a function of soil arbuscular mycorrhizal
487 fungal communities: experimentally manipulating co-occurring *Glomus* species. *Plant Ecol* 183:257-263

488 Hart MM, Forsythe J, Oshowski B, Bücking H, Jansa J, Kiers ET (2013) Hiding in a crowd-does diversity facilitate
489 persistence of a low-quality fungal partner in the mycorrhizal symbiosis?. *Symbiosis* 59:47-56

490 Hazard C, Gosling P, van der Gast CJ, Mitchell DT, Doohan FM, Bending GD (2013) The role of local environment
491 and geographical distance in determining community composition of arbuscular mycorrhizal fungi at the
492 landscape scale. *ISME J* 7:498-508

493 Helgason T, Merryweather JW, Denison J, Wilson P, Young JPW, Fitter AH (2002) Selectivity and functional
494 diversity in arbuscular mycorrhizas of co-occurring fungi and plants from a temperate deciduous woodland. *J*
495 *Ecol* 90:371-384

496 Hildebrandt U, Ouziad F, Marnier F-JJ, Bothe H (2006) The bacterium *Paenibacillus validus* stimulates growth of
497 the arbuscular mycorrhizal fungus *Glomus intraradices* up to the formation of fertile spores. *FEMS Microbiol*
498 *Lett* 254:258-267

499 Hohnjec N, Vieweg ME, Pühler A, Becker A, Küster H (2005) Overlaps in the transcriptional profiles of *Medicago*
500 *truncatula* roots inoculated with two different *Glomus* fungi provide insights into the genetic program activated
501 during arbuscular mycorrhiza. *Plant Physiol* 137:1283-1301

502 Jansa J, Andrew SF, Smith SE (2008) Are there benefits of simultaneous root colonization by different arbuscular
503 mycorrhizal fungi? *New Phytol* 177:779-789

504 Jansa J, Erb A, Oberholzer H-R, Smilauer P, Egli S (2014) Soil and geography are more important determinants of
505 indigenous arbuscular mycorrhizal communities than management practices in Swiss agricultural soils. *Mol*
506 *Ecol* 23:2118-2135

507 Ji B, Bentivenga SP, Casper BB (2010) Evidence for ecological matching of whole AM fungal communities to the
508 local plant-soil environment. *Ecology* 91:3037-3046

509 Johnson D, Vandenkoornhuysen PJ, Leake JR, Gilbert L, Booth RE, Grime JP, Young JPW, Read DJ (2004) Plant
510 communities affect arbuscular mycorrhizal fungal diversity and community composition in grassland
511 microcosms. *New Phytol* 161:503-515

512 Jones Jr, Benton J, Wolf B, Mills HA (1991) *Plant analysis handbook - a practical sampling, preparation, analysis,*
513 *and interpretation guide.* Micro-Macro Publishing Inc, Athens, USA

514 Kiers ET, Duhamel M, Beesetty Y, Mensah JA, Franken O, Verbruggen E, Fellbaum CR, Kowalchuk GA, Hart
515 MM, Bago A, Palmer TM, West SA, Vandenkoornhuysen P, Jansa J, Bücking H (2011) Reciprocal rewards
516 stabilize cooperation in the mycorrhizal symbiosis. *Science* 333:880-882

517 Kivlin SN, Hawkes CV, Treseder KK (2011) Global diversity and distribution of arbuscular mycorrhizal fungi. *Soil*
518 *Biol Biochem* 43:2294-2303

519 Knecht B, Jansa J, Franken O, Engelmoer DJ, Werner GD, Bücking H, Kiers ET (2016) Host plant quality mediates
520 competition between arbuscular mycorrhizal fungi. *Fungal Ecol* 20:233-240.

521 Koch AM, Kuhn G, Fontanillas P, Fumagalli L, Goudet J, Sanders IR (2004) High genetic variability and low local
522 diversity in a population of arbuscular mycorrhizal fungi. *Proc Nat Acad Sci USA* 101:2369-2374

523 Koide RT (2000) Functional complementarity in the arbuscular mycorrhizal symbiosis. *New Phytol* 147:233-235

524 Koorem K, Tulva I, Davison J, Jairus T, Öpik M, Vasar M, Zobel M, Moora M (2017) Arbuscular mycorrhizal
525 fungal communities in forest plant roots are simultaneously shaped by host characteristics and canopy-
526 mediated light availability. *Plant Soil* 410:259-271

527 Lee J, Lee S, Young JPW (2008) Improved PCR primers for the detection and identification of arbuscular
528 mycorrhizal fungi. *FEMS Microbiol Ecol* 65:339-349

529 Maherali H, Klironomos JN (2007) Influence of phylogeny on fungal community assembly and ecosystem
530 functioning. *Science* 316:1746-1748

531 Massoumou M, van Tuinen D, Chatagnier O, Arnould C, Brechenmacher L, Sanchez L, Selim S, Gianinazzi S,
532 Gianinazzi-Pearson V (2007) *Medicago truncatula* gene responses specific to arbuscular mycorrhiza
533 interactions with different species and genera of Glomeromycota. *Mycorrhiza* 17:223-234

534 Moora M, Öpik M, Sen R, Zobel M (2004) Native arbuscular mycorrhizal fungal communities differentially
535 influence the seedling performance of rare and common *Pulsatilla* species. *Funct Ecol* 18:554-562

536 Morrison E, Lagos L, Al-Agely A, Glaab H, Johnson W, Jorquera MA, Ogram A (2017) Mycorrhizal inoculation
537 increases genes associated with nitrification and improved nutrient retention in soil. *Biol Fertil Soils* 53:275-
538 279

539 Mummey DL, Rillig MC (2008) Spatial characterization of arbuscular mycorrhizal fungal molecular diversity at the
540 submetre scale in a temperate grassland. *FEMS Microbiol Ecol* 64:260-270

541 Munkvold L, Kjoller R, Vestberg M, Rosendahl S, Jakobsen I (2004) High functional diversity within species of
542 arbuscular mycorrhizal fungi. *New Phytol* 164:357-364

543 Myers N, Mittermeier RA, Mittermeier CG, Da Fonseca GA, Kent J (2000) Biodiversity hotspots for conservation
544 priorities. *Nature* 403:853

545 Nelson DW, Sommers L (1982) Total carbon, organic carbon and organic matter. In: Page AL, Miller RH, Keeney
546 DR (Eds) *Methods of Soil Analysis. Part 2. Chemical and Microbiological Properties*, 2nd edn, Agronomy
547 Monograph 9. American Society of Agronomy, Madison, WI, pp 539-579

548 Njeru EM, Avio L, Bocci G, Sbrana C, Turrini A, Bärberi P, Giovannetti M, Oehl F (2015) Contrasting effects of
549 cover crops on 'hot spot' arbuscular mycorrhizal fungal communities in organic tomato. *Biol Fertil Soils*
550 51:151-166

551 Njeru EM, Bocci G, Avio L, Sbrana C, Turrini A, Giovannetti M, Bàrberi P (2017) Functional identity has a
552 stronger effect than diversity on mycorrhizal symbiosis and productivity of field grown organic tomato. Eur J
553 Agron 86:1-11

554 Oehl F, Laczko E, Bogenrieder A, Stahr K, Bösch R, van der Heijden M, Sieverding E (2010) Soil type and land use
555 intensity determine the composition of arbuscular mycorrhizal fungal communities. Soil Biol Biochem 42:724-
556 738

557 Oehl F, Laczko E, Oberholzer HR, Jansa J, Egli S (2017) Diversity and biogeography of arbuscular mycorrhizal
558 fungi in agricultural soils. Biol Fertil Soils. doi: 10.1007/s00374-017-1217-x

559 Oehl F, Sieverding E, Ineichen K, Mäder P, Boller T, Wiemken A (2003) Impact of land use intensity on the species
560 diversity of arbuscular mycorrhizal fungi in agroecosystems of Central Europe. Appl Environ Microbiol
561 69:2816-2824

562 Oehl F, Sieverding E, Ineichen K, Ris EA, Boller T, Wiemken A (2005) Community structure of arbuscular
563 mycorrhizal fungi at different soil depths in extensively and intensively managed agroecosystems. New Phytol
564 165:273-283

565 Ohsowski BM, Zaitsoff PD, Öpik M, Hart MM (2014) Where the wild things are: looking for uncultured
566 Glomeromycota. New Phytol 204:171-179

567 Olsen SR, Sommers LE (1982) Phosphorus. In: Page AL, Miller RH, Keeney DR (Eds) Methods of Soil Analysis.
568 Part 2. Chemical And Microbiological Properties. Agronomy Monograph 9, 2nd edn, American Society of
569 Agronomy, Madison, WI, pp 403-430

570 Öpik M, Moora M, Liira J, Zobel M (2006) Composition of root-colonizing arbuscular mycorrhizal fungal
571 communities in different ecosystems around the globe. J Ecol 94: 778-790

572 Pepe A, Sbrana C, Ferrol N, Giovannetti M (2017) An *in vivo* whole-plant experimental system for the analysis of
573 gene expression in extraradical mycorrhizal mycelium. Mycorrhiza. doi:10.1007/s00572-017-0779-7

574 Pivato B, Mazurier S, Lemanceau P, Siblot S, Berta G, Mougél C, van Tuinen D (2007) *Medicago* species affect the
575 community composition of arbuscular mycorrhizal fungi associated with roots. New Phytol 176:197-210

576 Pringle A, Bever JD (2002) Divergent phenologies may facilitate the coexistence of arbuscular mycorrhizal fungi in
577 a North Carolina grassland. Am J Bot 89:1439-1446

578 Read DJ (1991). Mycorrhizas in ecosystems. Experientia 47:376-391

579 Redecker D (2002) Molecular identification and phylogeny of arbuscular mycorrhizal fungi. Plant Soil 244:67-73

580 Redecker D, Hijri I, Wiemken A (2003) Molecular identification of arbuscular mycorrhizal fungi in roots:
581 perspectives and problems. Folia Geobot 38:113-124

582 Rodríguez-Echeverría S, Teixeira H, Correia M, Timóteo S, Heleno R, Öpik M, Moora M (2017) Arbuscular
583 mycorrhizal fungi communities from tropical Africa reveal strong ecological structure. *New Phytol* 213:380-
584 390.

585 Roger A, Colard A, Angelard C, Sanders IR (2013) Relatedness among arbuscular mycorrhizal fungi drives plant
586 growth and intraspecific fungal coexistence. *ISME J* 7:2137-2146

587 Rouphael Y, Franken P, Schneider C, Schwarz D, Giovannetti M, Agnolucci M, De Pascale S, Bonini F, Colla G
588 (2015) Arbuscular mycorrhizal fungi act as biostimulants in horticultural crops. *Sci Hort* 196:91-108

589 Sbrana C, Avio L, Giovannetti M (2014) Beneficial mycorrhizal symbionts affecting the production of health-
590 promoting phytochemicals. *Electrophoresis* 35:1535-1546

591 Scheublin TR, Ridgway KP, Young JPW, van der Heijden MGA (2004) Nonlegumes, legumes, and root nodules
592 harbor different arbuscular mycorrhizal fungal communities. *Appl Environ Microbiol* 70:6240-6246

593 Smith SE, Read DJ (2008) *Mycorrhizal symbiosis*, 3rd edn. Academic Press, London

594 Spatafora JW, Chang Y, Benny GL, Lazarus K, Smith ME, Berbee ML, Bonito G, Corradi N, Grigoriev I,
595 Gryganskyi A, James TY, O'Donnell K, Roberson RW, Taylor TN, Uehlin J, Vilgalys R, White MM, Stajich
596 JE (2016) A phylum-level phylogenetic classification of zygomycete fungi based on genome-scale data.
597 *Mycologia* 108:1028-1046

598 Stukenbrock EH, Rosendahl S (2005) Clonal diversity and population genetic structure of arbuscular mycorrhizal
599 fungi (*Glomus* spp.) studied by multilocus genotyping of single spores. *Mol Ecol* 14:743-752

600 Sumner ME, Miller WP (1996) Cation exchange capacity and exchange coefficients. In: Sparks DL (Ed) *Methods of*
601 *Soil Analysis. Part 3. Chemical Methods*. Soil Science Society of America, Madison, WI, pp. 1201-1229

602 Sýkorová Z, Wiemken A, Redecker A (2007) Co-occurring *Gentiana verna* and *Gentiana acaulis* and their
603 neighboring plants in two Swiss Upper Montane meadows harbor distinct arbuscular mycorrhizal fungal
604 communities. *Appl Environ Microbiol* 73:5426-5434

605 Treseder KK, Cross A (2006) Global distributions of arbuscular mycorrhizal fungi. *Ecosystems* 9:305-316

606 Turrini A, Agnolucci M, Palla M, Tomé E, Tagliavini M, Scandellari F, Giovannetti M (2017) Species diversity and
607 community composition of native arbuscular mycorrhizal fungi in apple roots are affected by site and orchard
608 management. *Appl Soil Ecol* 116:42-54

609 Turrini A, Avio L, Bedini S, Giovannetti M (2008) *In situ* collection of endangered arbuscular mycorrhizal fungi in
610 a Mediterranean UNESCO Biosphere Reserve. *Biodivers Conserv* 17:643-657

611 Turrini A, Giovannetti M (2012) Arbuscular mycorrhizal fungi in national parks, nature reserves and protected areas
612 worldwide: a strategic perspective for their *in situ* conservation. *Mycorrhiza* 22:81-97

613 Turrini A, Sbrana C, Avio L, Njeru EM, Bocci G, Bàrberi P, Giovannetti M (2016) Changes in the composition of
614 native root arbuscular mycorrhizal fungal communities during a short-term cover crop-maize succession. *Biol*
615 *Fertil Soils* 52:643-653

616 Uibopuu A, Moora M, Öpik M, Zobel M (2012) Temperate forest understorey species performance is altered by
617 local arbuscular mycorrhizal fungal communities from stands of different successional stages. *Plant Soil*
618 356:331-339

619 Vályi K, Rillig MC, Hempel S (2015) Land-use intensity and host plant identity interactively shape communities of
620 arbuscular mycorrhizal fungi in roots of grassland plants. *New Phytol* 205:1577-1586

621 Vandenkoornhuyse P, Ridgway KP, Watson IJ, Fitter AH, Young JPW (2003) Co-existing grass species have
622 distinctive arbuscular mycorrhizal communities. *Mol Ecol* 12:3085-3095

623 Van der Gast CJ, Gosling P, Tiwari B, Bending GD (2011) Spatial scaling of arbuscular mycorrhizal fungal
624 diversity is affected by farming practice. *Environ Microbiol* 13:241-249

625 van der Heijden MGA, Klironomos JN, Ursic M, Moutoglis P, Streitwolf-Engel R, Boller T, Wiemken A, Sanders
626 IR (1998) Mycorrhizal fungal diversity determines plant biodiversity, ecosystem variability and productivity.
627 *Nature* 396:69-72

628 Van Geel, M, Ceustermans A, Van Hemelrijck W, Lievens B, Honnay O (2015) Decrease in diversity and changes
629 in community composition of arbuscular mycorrhizal fungi in roots of apple trees with increasing orchard
630 management intensity across a regional scale. *Mol Ecol* 24:941-952

631 Vogelsang KM, Reynolds HL, Bever JD (2006) Mycorrhizal fungal identity and richness determine the diversity
632 and productivity of a tallgrass prairie system. *New Phytol* 172:554-562

633 Walder F, van der Heijden M.G (2015) Regulation of resource exchange in the arbuscular mycorrhizal symbiosis.
634 *Nature plants* 1:15159.

635

1. SOIL PHYSICAL-CHEMICAL CHARACTERIZATION OF 12



2. SELECTION OF 6 SIMILAR SOIL PLOTS FOR MICROCOSM EXPERIMENTS WITH 3 HOST PLANTS




3. MOLECULAR ANALYSES OF ROOT AMF COMMUNITIES FROM 2 NATIVE AMF TREATMENTS SHOWING THE MOST DIFFERENTIAL PLANT PERFORMANCES




18 AML1-AML2 libraries from 2 treatments, 3 host plants, 3 replicates


636
637 **Fig 1** Flow chart showing 1) the experimental field with the 12 plots, characterized for their soil physical and
638 chemical properties; 2) the six soil plots selected for uniform soil characteristics, used as native AMF treatments in
639 microcosm experiments aimed at evaluating plant growth and nutrition of three host plants; 3) the two native AMF
640 treatments, showing the most differential plant performance, selected for molecular analyses of AMF communities
641 occurring in the roots.



642
 643 **Fig 2** a) Radar graph representing the relative increases, for onion, pepper and lettuce, in shoot dry weight (SDW),
 644 nitrogen (N) and phosphorous (P) content of five selected native AMF treatment pairs; b) results of principal
 645 components analysis ordination of native AMF treatments, using scores of growth and nutrition variables for host
 646 plants. The first axis explained 37.5%, the second axis 22.1% of variation.
 647




649 **Fig 3** Neighbour-Joining phylogenetic tree of glomeromycotan sequences derived from host plants growing on the
650 native AMF treatment 2 and 22. Bootstrap values are shown when they exceed 65 % (1000 replications). The
651 analysis is based on partial nuclear small subunit ribosomal RNA gene sequences (SSU; ~800 bp; AML1/AML2
652 fragment) and involved 107 nucleotide sequences. Different sequence types are indicated in brackets and names are
653 reported in Table 4. AMF families are also reported. Sequences obtained in the present study are shown in bold and
654 their accession numbers are prefixed with plant species/native AMF treatment/ clone identifiers (O = onion; L=
655 lettuce; P = pepper; 2 and 22 = native AMF treatments). The tree is rooted with a reference sequence of
656 *Corallochytrium lymacisporum* (L42528).
657



659

660 **Fig 4** Relative abundance (%) of AMF OTUs detected in the roots of the different host plants (onion, pepper,

661 lettuce), growing in native AMF treatments 2 and 22.



662
 663 **Fig 5** Redundancy analysis (RDA) ordination biplot of AMF colonizing the three host species (onion, pepper and
 664 lettuce), growing in the AMF soil treatments 2 and 22. The inoculation treatment and the host plants were used as
 665 the explanatory variable, and the different OTUs were used as dependent variables. The names of the different
 666 OTUs are reported in Table 4. The first and second axes explain 51.2% of total variance.

667
 668

669 **Table 1** Descriptive statistics of soil chemical and physical properties of the 12 plots inside the “hot spot” field in
 670 the UNESCO Biosphere Reserve.

Soil property	pH	CEC (cmol kg⁻¹)	Total N (%)	Organic matter (%)	Olsen P (µg g⁻¹)	Clay (%)	Silt (%)	Sand (%)
Mean	7.43	12.71	1.70	2.58	6.13	15.4	16.7	67.9
Median	7.44	13.62	1.64	2.64	6.00	15.4	15.6	69.5
Minimum	6.90	10.15	1.32	1.63	5.11	11.9	13.4	57.5
Maximum	8.02	15.36	2.23	3.01	7.33	20.9	24.9	74.3
SE ^a	0.12	0.55	0.07	0.12	0.22	0.72	1.01	1.42
CV ^b	5.6%	15.0%	13.4%	15.5%	12.3%	16.1%	20.9%	7.2%

671 ^aSE (Standard Error); ^bCV (Coefficient of Variability)

672

673 **Table 1** Descriptive statistics of soil chemical and physical properties of the 12 plots inside the “hot spot” field in
 674 the UNESCO Biosphere Reserve.

Soil property	pH	CEC (cmol kg⁻¹)	Total N (%)	Organic matter (%)	Olsen P (µg g⁻¹)	Clay (%)	Silt (%)	Sand (%)
Mean	7.43	12.71	1.70	2.58	6.13	15.4	16.7	67.9
Median	7.44	13.62	1.64	2.64	6.00	15.4	15.6	69.5
Minimum	6.90	10.15	1.32	1.63	5.11	11.9	13.4	57.5
Maximum	8.02	15.36	2.23	3.01	7.33	20.9	24.9	74.3
SE ^a	0.12	0.55	0.07	0.12	0.22	0.72	1.01	1.42
CV ^b	5.6%	15.0%	13.4%	15.5%	12.3%	16.1%	20.9%	7.2%

675 ^aSE (Standard Error); ^bCV (Coefficient of Variability)

676

677 **Table 2** Summary of (a) two way ANOVA testing the effects of native AMF treatments and host species on root
 678 colonization and plant growth, and (b) one way ANOVA testing the effects of native AMF treatments on nutrient
 679 content of each host species.

a)	Colonized root length (%)			Shoot dry weight (mg plant⁻¹)	
	df	F	P	F	P
Source of variation	df	F	P	F	P
native AMF treatment	5,36	4.9	0.001	17.2	<0.001
host species	2,36	249.5	<0.001	24.9	<0.001
Interaction	10,36	3.6	0.002	2.9	0.008
b)	Shoot N content (mg plant⁻¹)			Shoot P content (mg plant⁻¹)	
	df	F	P	F	P
<i>Allium cepa</i>	5,12	0.725	0.618	1.5	0.273
<i>Capsicum annuum</i>	5,12	12.3	<0.001	4.9	0.011
<i>Lactuca sativa</i>	5,12	10.6	<0.001	10.1	0.001

680

681 **Table 3** Shoot dry matter, nutritional parameters and percentage of mycorrhizal colonization in the roots of *Allium*
682 *cepa*, *Capsicum annuum* and *Lactuca sativa*, grown in 6 microcosms using soils from different plots of the field site
683 within UNESCO Biosphere Reserve, showing the highest similarity in soil physical and chemical characteristics. In
684 columns, means (\pm standard error of the mean) within each plant species followed by different lowercase letters are
685 significantly different ($P < 0.05$).

Plant species	native AMF treatment	SDW ^a (mg plant ⁻¹)	Shoot N content (mg plant ⁻¹)	Shoot P content (mg plant ⁻¹)	Colonized root length (%)
<i>Allium cepa</i> (onion)	2	45.7 \pm 1.8 bA ^b	0.66 \pm 0.02 a	0.12 \pm 0.012 a	90.4 \pm 2.2 aA
	5	61.4 \pm 7.0 aA	0.72 \pm 0.06 a	0.18 \pm 0.019 a	86.7 \pm 2.6 aA
	9	54.5 \pm 1.2 abA	0.68 \pm 0.02 a	0.13 \pm 0.009 a	92.0 \pm 0.6 aA
	10	53.7 \pm 4.0 abA	0.72 \pm 0.06 a	0.13 \pm 0.032 a	89.8 \pm 0.8 aA
	14	49.2 \pm 3.4 abA	0.68 \pm 0.03 a	0.14 \pm 0.004 a	86.6 \pm 3.9 aA
	22	54.4 \pm 4.0 abB	0.78 \pm 0.07 a	0.15 \pm 0.002 a	90.0 \pm 3.6 aA
<i>Capsicum annuum</i> (pepper)	2	47.5 \pm 2.8 cA	0.76 \pm 0.04 b	0.07 \pm 0.007 c	55.6 \pm 4.6 abB
	5	56.5 \pm 2.3 bcA	0.77 \pm 0.05 b	0.12 \pm 0.009 a	66.8 \pm 3.3 aB
	9	61.8 \pm 3.9 abA	0.76 \pm 0.04 b	0.10 \pm 0.006 abc	38.9 \pm 1.3 bB
	10	54.7 \pm 3.7 bcA	0.74 \pm 0.08 b	0.07 \pm 0.015 bc	49.7 \pm 2.2 abB
	14	49.3 \pm 4.0 bcA	0.61 \pm 0.03 b	0.09 \pm 0.006 abc	41.4 \pm 4.7 bB
	22	74.5 \pm 3.7 aA	1.04 \pm 0.04 a	0.11 \pm 0.004 ab	39.5 \pm 1.2 bC
<i>Lactuca sativa</i> (lettuce)	2	25.2 \pm 1.0 cB	0.44 \pm 0.03 b	0.04 \pm 0.003 b	61.6 \pm 4.7 aB
	5	45.3 \pm 1.7 abB	0.52 \pm 0.09 b	0.08 \pm 0.011 ab	55.7 \pm 5.1 abB
	9	51.9 \pm 2.6 abA	0.61 \pm 0.05 b	0.11 \pm 0.010 a	41.2 \pm 5.7 bB
	10	48.3 \pm 2.4 abA	0.63 \pm 0.02 b	0.08 \pm 0.003 ab	45.5 \pm 5.9 abB
	14	41.5 \pm 0.3 bA	0.55 \pm 0.03 b	0.10 \pm 0.013 a	47.1 \pm 2.7 abB
	22	55.3 \pm 1.7 aB	0.87 \pm 0.02 a	0.11 \pm 0.005 a	57.9 \pm 2.4 abB

686 ^a SDW (Shoot Dry Matter)

687 ^b In columns, means for each native AMF treatment, followed by different uppercase letters, are significantly
688 different ($P < 0.05$).

689

690 **Table 4** Sequence types of arbuscular mycorrhizal fungi, identified using AML1-AML2 primers pair, in the roots of
691 onion, pepper and lettuce plants from treatments 2 and 22, using native soil collected from two different plots inside
692 the “hot spot” field within the UNESCO Biosphere Reserve “Selva Pisana”.

Sequence type	Identity (%)
Rh1	FR821553 (99%)
Rh2	JX296753 (99%)
Rh3	FR750222 (99%)
Rh4	JN791150 (98%)
Scle	KF386336 (98%)
Dom	HM153420 (99%)
Fun1	NG017178 (99%)
Fun2	AJ245637 (99%)
Sept1	KF386332 (99%)
Sept2	FR848639 (98%)
Rac	AJ306435 (100%)
Clar	AJ852598 (99%)
Par1	FN869853 (99%)
Par2	AM295493 (100%)

693 **Table 5** Diversity indices of AMF communities occurring in the roots of different plant species from treatments 2 and 22 (maens± SEM).

Treatment	Plant species	Taxa (S)	Simpson (1-D)	Shannon (H)	Evenness (e ^{H/S})	Chao
2	<i>Allium cepa</i>	2.67 ± 0.54a	0.26 ± 0.13a	0.49 ± 0.24a	0.65 ± 0.03a	2.67 ± 0.54a
	<i>Lactuca sativa</i>	3.00 ± 0.00a	0.25 ± 0.02a	0.47 ± 0.04a	0.54 ± 0.02a	3.33 ± 0.14a
	<i>Capsicum annuum</i>	2.00 ± 0.00a	0.25 ± 0.01a	0.39 ± 0.01a	0.76 ± 0.01a	2.00 ± 0.00a
	P*	0.135	0.874	0.874	0.427	0.118
22	<i>Allium cepa</i>	7.67 ± 0.54b	0.82 ± 0.00b	1.83 ± 0.02b	0.82 ± 0.04a	8.00 ± 0.82b
	<i>Lactuca sativa</i>	7.33 ± 0.27b	0.72 ± 0.03b	1.55 ± 0.07b	0.65 ± 0.02a	8.00 ± 0.61b
	<i>Capsicum annuum</i>	4.33 ± 0.27b	0.51 ± 0.02a	0.89 ± 0.03b	0.57 ± 0.02a	5.67 ± 0.95b
	P*	0.051	0.027	0.039	0.039	0.417

694 Values followed by different letters are significantly different (P≤0.05) among plant species within treatments after Kruskal Wallis test.

695 * P represents the significance level of the Kruskal Wallis test within each native AMF treatment.

696