

# Supplemental Information

**SUPPLEMENTAL TABLE 3** State- and Territory-Specific Changes in the Prevalence of High Weight-for-Length Among Infants in the WIC-PC Survey From 2000 Through 2004 and From 2010 Through 2014

State	2000 to 2004			2010 to 2014		
	<i>n</i> <sup>a</sup>	High WFL, %	Prevalence Change (95% CI) <sup>b</sup>	<i>n</i>	High WFL, %	Prevalence Change (95% CI) <sup>b</sup>
Alabama	31 267	12.5	0.7 (0.2 to 1.3)	43 957	12.9	0.2 (-0.2 to 0.7)
Alaska	5982	16.0	1.9 (0.5 to 3.4)	6 100	16.5	0.9 (-0.6 to 2.6)
Arizona	41 103	12.7	1.3 (0.8 to 1.8)	41 678	15.3	-2.8 (-3.2 to -2.3)
Arkansas	13 575	10.1	0.9 (0.2 to 1.7)	20 972	13.1	-2.0 (-2.6 to -1.4)
California	362 429	14.8	1.0 (0.8 to 1.2)	403 041	17.3	-3.6 (-3.7 to -3.4)
Colorado	15 053	6.7	0.4 (-0.2 to 1.0)	25 245	8.1	-1.1 (-1.5 to -0.7)
Connecticut	15 966	11.1	-0.7 (-1.4 to -0.0)	13 164	12.3	-1.7 (-2.5 to -0.9)
Delaware	3 587	14.6	-3.0 (-4.2 to -1.6)	5 201	14.0	0.1 (-1.2 to 1.4)
District of Columbia	3 598	10.2	1.2 (-0.2 to 2.8)	3 579	12.8	-1.7 (-3.1 to -0.1)
Florida	64 654	13.5	0.2 (-0.1 to 0.6)	115 168	14.2	-2.1 (-2.4 to -1.9)
Georgia	52 114	11.3	1.0 (0.6 to 1.4)	106 764	11.5	-1.7 (-2.0 to -1.5)
Hawaii	11 237	12.1	NA <sup>c</sup>	13 437	11.2	-1.3 (-2.0 to -0.6)
Idaho	11 020	8.2	0.2 (-0.5 to 1.0)	11 818	10.2	-1.0 (-1.7 to -0.2)
Illinois	81 315	13.5	3.8 (3.5 to 4.2)	100 445	12.5	-0.7 (-0.9 to -0.3)
Indiana	28 678	11.0	1.6 (1.1 to 2.2)	37 011	13.6	-1.9 (-2.4 to -1.4)
Iowa	17 017	13.0	0.8 (0.0 to 1.6)	25 237	13.4	0.4 (-0.2 to 1.0)
Kansas	12 936	12.0	4.6 (3.7 to 5.6)	25 911	12.6	-1.8 (-2.4 to -1.3)
Kentucky	23 691	17.8	1.2 (0.5 to 1.9)	27 129	19.6	-7.4 (-7.9 to -6.9)
Louisiana	21 918	13.3	2.7 (2.0 to 3.4)	33 380	16.8	-1.6 (-2.1 to -1.0)
Maine	4 557	14.2	0.2 (-1.1 to 1.7)	9 976	12.7	-2.3 (-3.1 to -1.4)
Maryland	18 556	14.5	-0.5 (-1.2 to 0.1)	31 121	14.7	-0.5 (-1.0 to 0.1)
Massachusetts	25 423	15.4	0.8 (0.1 to 1.4)	28 286	16.7	-2.4 (-3.0 to -1.9)
Michigan	74 870	10.6	1.3 (1.0 to 1.7)	51 570	12.9	-1.0 (-1.4 to -0.7)
Minnesota	17 774	13.3	0.5 (-0.3 to 1.0)	30 457	12.3	-1.2 (-1.7 to -0.7)
Mississippi	17 566	16.2	0.8 (0.0 to 1.6)	24 126	17.7	-2.4 (-3.0 to -1.7)
Missouri	41 369	11.6	2.0 (1.5 to 2.6)	46 897	12.4	-2.5 (-2.9 to -2.1)
Montana	4 375	12.1	0.8 (-0.5 to 2.3)	7 748	10.2	-0.9 (-1.8 to 0.0)
Nebraska	10 437	14.4	-0.0 (-0.9 to 0.9)	13 439	14.1	0.6 (-0.3 to 1.5)
Nevada	15 567	11.0	1.7 (1.0 to 2.5)	23 255	12.4	0.1 (-0.5 to 0.7)
New Hampshire	4 664	13.1	1.0 (-0.4 to 2.5)	6 411	13.0	-1.5 (-2.6 to -0.4)
New Jersey	39 922	15.1	1.7 (1.2 to 2.3)	53 110	15.1	-1.1 (-1.5 to -0.7)
New Mexico	17 331	9.3	0.6 (-0.0 to 1.3)	13 097	13.1	-2.6 (-3.3 to -1.8)
New York	113 640	12.9	0.2 (-0.1 to 0.5)	107 374	14.1	-2.2 (-2.4 to -1.9)
North Carolina	38 011	11.7	0.8 (0.4 to 1.3)	58 763	11.8	0.4 (0.1 to 0.8)
North Dakota	4 025	12.1	0.0 (-1.3 to 1.5)	5 065	12.3	-1.0 (-2.2 to 0.3)
Ohio	82 471	11.4	0.4 (0.1 to 0.7)	104 540	12.2	0.2 (-0.1 to 0.5)
Oklahoma	19 765	12.9	0.1 (-0.6 to 0.8)	21 964	12.8	-2.3 (-2.9 to -1.8)
Oregon	26 180	13.5	-2.4 (-2.9 to -1.9)	36 551	10.6	0.2 (-0.2 to 0.7)
Pennsylvania	51 506	13.4	-0.4 (-0.8 to -0.0)	57 842	13.6	-1.6 (-1.9 to -1.2)
Rhode Island	5 992	14.2	0.6 (-0.6 to 1.9)	5 764	15.9	-1.8 (-3.0 to -0.4)
South Carolina	22 248	12.2	3.5 (2.8 to 4.3)	27 838	13.7	-1.5 (-2.1 to -1.0)
South Dakota	5 666	11.3	3.5 (2.1 to 5.0)	4 973	16.2	-0.5 (-1.9 to 1.1)
Tennessee	32 139	13.4	-0.1 (-0.6 to 0.4)	36 840	15.0	-2.0 (-2.5 to -1.6)
Texas	139 778	16.4	0.9 (0.6 to 1.1)	221 750	16.2	-1.3 (-1.5 to -1.1)
Utah	18 279	9.7	1.4 (0.8 to 2.1)	24 944	11.3	-4.4 (-4.8 to -3.9)

TABLE 3 Continued

State	2000 to 2004			2010 to 2014		
	<i>r</i> <sup>a</sup>	High WFL, %	Prevalence Change (95% CI) <sup>b</sup>	<i>n</i>	High WFL, %	Prevalence Change (95% CI) <sup>b</sup>
Vermont	4848	9.2	0.8 (-0.3 to 2.1)	5183	9.0	0.6 (-0.6 to 1.9)
Virginia	35 149	13.6	8.2 (7.5 to 9.0)	33 245	24.0	-1.1 (-1.7 to -0.5)
Washington	53 153	10.6	1.5 (1.1 to 1.9)	64 628	12.8	-1.6 (-2.0 to -1.3)
West Virginia	10 027	10.6	0.4 (-0.4 to 1.2)	15 477	9.4	1.9 (1.1 to 2.8)
Wisconsin	21 815	12.8	1.9 (1.2 to 2.6)	28 409	15.5	-1.9 (-2.5 to -1.3)
Wyoming	3 492	7.3	-1.2 (-2.2 to 0.0)	2 997	9.8	-3.3 (-4.3 to -2.0)
Territory						
American Samoa	1 191	16.5	3.0 (0.1 to 6.6)	1 322	16.1	6.1 (2.9 to 9.8)
Guam	1 411	13.7	-0.1 (-2.2 to 2.6)	2 021	9.7	-1.7 (-3.2 to 0.1)
Northern Mariana Islands	NA <sup>c</sup>	NA <sup>c</sup>	NA <sup>c</sup>	1 022	13.1	-1.4 (-4.0 to 2.0)
Puerto Rico	45 134	18.8	-2.9 (-3.3 to -2.4)	51 259	19.1	-9.0 (-9.3 to -8.7)
Virgin Islands	1 853	11.7	-2.4 (-4.2 to -0.1)	1 231	11.1	0.7 (-1.8 to 3.8)

<sup>a</sup> For each time period, the columns represent the *n* and prevalence of high weight-for-length (WFL) at the beginning of the period. The prevalence ratio change is the prevalence at the end of the period minus the prevalence at the beginning. This change was calculated from the relative risk based on a Poisson regression model that included age and race as covariates.

<sup>b</sup> The prevalence at the beginning of the period and the prevalence ratio were used to estimate the adjusted percentage change.

<sup>c</sup> NA indicates that data were not available (missing). The 2002 and 2004 data for Hawaii were excluded because the prevalence estimates were inconsistent with other annual estimates from the state.