



PUBLISHED FOR SISSA BY SPRINGER

RECEIVED: August 2, 2012
ACCEPTED: October 15, 2012
PUBLISHED: October 30, 2012

4d $\mathcal{N} = 2$ gauge theories and quivers: the non-simply laced case

Sergio Cecotti and Michele Del Zotto

*Scuola Internazionale Superiore di Studi Avanzati (SISSA),
via Bonomea 265, I-34100 Trieste, Italy*

E-mail: cecotti@sisssa.it, eledelz@gmail.com

ABSTRACT: We construct the BPS quivers with superpotential for the 4d $\mathcal{N} = 2$ gauge theories with *non*-simply laced Lie groups (B_n , C_n , F_4 and G_2). The construction is inspired by the BIKMSV geometric engineering of these gauge groups as *non-split* singular elliptic fibrations. From the categorical viewpoint of [arXiv:1203.6734](https://arxiv.org/abs/1203.6734), the fibration of the light category $\mathcal{L}(\mathfrak{g})$ over the (degenerate) Gaiotto curve has a monodromy given by the action of the outer automorphism of the corresponding *unfolded* Lie algebra.

In view of the Katz-Vafa ‘matter from geometry’ mechanism, the monodromic idea may be extended to the construction of (Q, \mathcal{W}) for SYM coupled to higher matter representations. This is done through a construction we call *specialization*.

KEYWORDS: Supersymmetric gauge theory, Extended Supersymmetry, Nonperturbative Effects

ARXIV EPRINT: [1207.7205](https://arxiv.org/abs/1207.7205)

JHEP10(2012)190

Contents

1	Introduction	1
2	Geometric preliminaries	3
3	Q-systems and Dirac integrality conditions	6
4	Review of the light category $\mathcal{L} \subset \text{rep}(Q, \mathcal{W})$	7
5	The light category of non-simply laced $\mathcal{N} = 2$ SYM	11
5.1	Monodromic light categories	11
5.1.1	Preprojective algebras for non-simply laced (i.e. valued) graphs	11
5.1.2	The structure of $\text{mod } \mathcal{P}(\mathfrak{g})$	12
5.1.3	Example: B_2	13
5.1.4	Example: G_2	14
5.1.5	Example: F_4	15
5.1.6	Monodromy of $\mathcal{M}(\mathfrak{g})$	16
5.2	‘Auxiliary’ fields and the final form of the light category $\mathcal{L}(\mathfrak{g})$	17
6	The quiver superpotential \mathcal{W} for $BCFG$ SYM	19
7	Other applications of the monodromic construction	20
7.1	<i>Specialization</i> : new pairs (Q, \mathcal{W}) from old ones	21
7.1.1	The ‘heavy quark procedure’ revisited	21
7.1.2	More general (split) specializations: quiver gauge theories	22
7.2	Coupling matter in $\text{Sym}^n F_i$	23
8	Coupling $BCFG$ SYM to matter	24
A	Consistency with the perturbative Higgs mechanism	25
B	Consistency of $\text{rep}(Q, \mathcal{W}) \rightarrow \text{rep}(Q', \mathcal{W}')$	29
B.1	C_2	30
B.2	G_2	31

1 Introduction

In the last year there has been significant progress in the computation of BPS spectra of 4d $\mathcal{N} = 2$ supersymmetric gauge theories using the Representation Theory of quivers with superpotentials [1–7] (for previous work see [8–22]; for other approaches to the $\mathcal{N} = 2$ BPS

spectra see [23–28]). In the above references the discussion was always limited to theories whose gauge group G is a product of Lie groups of type ADE . This limitation arose from the lack of a construction of the quivers with superpotential for the $\mathcal{N} = 2$ models with gauge group of type $BCFG$. The goal of the present paper is to fill this gap.

The existence of a quiver description of the BPS spectrum for $\mathcal{N} = 2$ SYM with ADE gauge groups follows [1] from their geometric engineering in string theory as fibrations of ADE singularities. From this point of view, the strategy to extend the quiver description to the non-simply laced gauge groups appears obvious: in ref. [29] BIKMSV engineered all non-simply laced gauge groups by considering *non-split* singular elliptic fibrations (for a related construction see [30]). Roughly speaking, the dichotomy *simply-laced vs. non-simply-laced* corresponds geometrically to the dichotomy *split vs. non-split*. Thus one is naturally led to the question: ‘Is there a notion of split/non-split in the Representation Theory of a quiver with superpotential, (Q, \mathcal{W}) , which corresponds to the BIKMSV geometric notion?’ or, more directly, — ‘Which (Q, \mathcal{W}) ’s describe a non-split situation?’

The general (and rather abstract) theory developed in [6] has a simple answer to this question (see equation (2.9) below). Knowing that answer, one may write down the (Q, \mathcal{W}) ’s for all $BCFG$ pure SYM in a matter of few minutes. The resulting pairs (Q, \mathcal{W}) are guaranteed by (geometric) construction to have the physically right BPS spectrum in the weak coupling limit $g_{\text{YM}} \rightarrow 0$; in particular, in that limit the only stable particles with zero magnetic charge are vector-multiplets making precisely one copy of the adjoint of the gauge group G .

In the same vein, we take inspiration from the Katz-Vafa ‘matter from geometry’ construction [31] to get the pairs (Q, \mathcal{W}) for SYM coupled to matter. This allows to determine (Q, \mathcal{W}) for matter representations more general than those discussed in refs. [4, 6]. We call this method *specialization* of the quiver (with superpotential). Putting the two geometric constructions — monodromy and specialization — together, leads to the pairs (Q, \mathcal{W}) for $BCFG$ SYM coupled to matter.

The present paper is organized as follows. In section 2, following [29, 30], we describe the geometrical ideas on which our construction is based. In section 3 we determine the relevant quivers in two different ways: by relating them to certain integrable statistical systems, called Q -systems [32–34], as well as by using the Dirac integrality condition introduced in [6, 7]. In section 4 we review the notion of the light subcategory corresponding to the set of ‘perturbative’ states [6]. Section 5 is the heart of the paper: there it is shown how a light category with the BIKMSV monodromic properties arises naturally and uniquely from the special properties of the non-simply laced Dynkin graphs; the light BPS spectrum is computed and found to agree with the physically expected one. In section 6 we write down the superpotentials \mathcal{W} for all non-simply laced pure SYM theories. In section 7 we sketch other possible applications of the monodromic construction. In section 8 we present a preliminary discussion of the quivers for $BCFG$ SQCD with various matter content. Technical aspects and detailed computations are confined in the appendices.

2 Geometric preliminaries

We refer to [2–4, 6, 7] for general background on the correspondence between the Representation Theory of quivers with superpotential and the BPS spectra of 4d $\mathcal{N} = 2$ QFTs, including the notion of *stability* of a representation and the general properties of the stable representations.

The description of the BPS sector of a 4d $\mathcal{N} = 2$ gauge theory in terms of stable representations of a quiver Q , bounded by the Jacobian relations $\partial\mathcal{W} = 0$ arising from a superpotential \mathcal{W} , originates from the geometric construction of the gauge theory from string theory or M/F -theory. In this context the non-Abelian gauge groups arise from fibrations of ADE singularities [29]; *non*-simply laced gauge groups may also arise provided the fibration is *non-split*, as explained in [29]. This last case corresponds to the Slodowy generalization of the classical McKay correspondence to the non-simply laced case [36].

It is well-known [36, 37] that the Dynkin graph of a non-simply laced Lie group G arises by folding a parent simply-laced Dynkin graph G_{parent} along an automorphism. Specifically, the $G_{\text{parent}} \rightarrow G$ foldings are

$$D_{n+1} \longrightarrow B_n \tag{2.1}$$

$$A_{2n-1} \longrightarrow C_n \tag{2.2}$$

$$D_4 \longrightarrow G_2 \tag{2.3}$$

$$E_6 \longrightarrow F_4, \tag{2.4}$$

see figure 1. In all cases the folding corresponds to a \mathbb{Z}_2 automorphism of the parent graph, except for the G_2 case, where we may take it to be \mathbb{Z}_3 . To each node of the folded Dynkin diagrams in figure 1 there is attached an integer d_i , namely the number of nodes of the parent graph which were folded into it. This number corresponds to one-half the length-square¹ of the corresponding simple co-root α_i^\vee

$$d_i = \frac{1}{2}(\alpha_i^\vee, \alpha_i^\vee) \equiv \frac{2}{(\alpha_i, \alpha_i)} \quad i = 1, 2, \dots, r. \tag{2.5}$$

The BIKMSV [29] construction of a non-simply laced gauge group is based on a G_{parent} fibration having a non-trivial monodromy given by the action of the outer automorphism group \mathbb{Z}_2 (\mathbb{Z}_3) of the parent Lie algebra induced by the Dynkin diagram symmetry associated to the given folding (figure 1).

This monodromic construction may be imported in the context of Gaiotto engineering [24] of class- S 4d $\mathcal{N} = 2$ gauge theories from the 6d $(2, 0)$ theory of type $\mathfrak{g} = ADE$ as follows. In the Gaiotto framework, a weakly coupled SYM sector with simply-laced gauge group $G_{\text{s.l.}} = \exp[\mathfrak{g}]$ is realized by considering a long thin tube in the UV surface \mathcal{C} [24]. If the tube is very long, we may work locally in its bulk, and replace the surface \mathcal{C} by an infinite cylinder \mathbb{C}^\times ; this corresponds to ignoring the parts of the surface attached at the infinite ends of the tube which act as ‘matter’, whose global symmetry is gauged by the

¹The length of the short (long) co-roots (resp. roots) is normalized to 2.

given SYM sector. Neglecting the ‘matter’ is physically justified, since a very long tube corresponds to a very small YM coupling, and then the ‘matter’ asymptotically decouples.

Nothing prevents us from using a *twisted* version of this construction parallel to the BIKMSV *non-split* elliptic fibrations. (For a related construction in a slightly different setting see [35]). Indeed, considering the (2, 0) 6d theory of type, respectively,

$$\mathfrak{g} = D_{n+1}, A_{2n-1}, E_6, D_4$$

on such an infinite cylinder, \mathbb{C}^\times , times $\mathbb{R}^{1,2} \times S^1$, and reducing first on the S^1 factor, we are led [25] to solve the Hitchin equations for a $\mathfrak{g} \otimes K_{\mathbb{C}}$ valued Higgs field φ and the \mathfrak{g} -connection A satisfying a twisted periodic boundary conditions. Let $\theta = \arg z$ be the circle coordinate on $\mathbb{C}^\times \simeq S^1 \times \mathbb{R}$; we require

$$\left(A(\theta + 2\pi, r), \varphi(\theta + 2\pi, r) \right) = \left(\omega \cdot A(\theta, r), \omega \cdot \varphi(\theta, r) \right), \tag{2.6}$$

where ω is a non-trivial outer automorphism of \mathfrak{g} . Of course, this twisting is consistent only if we also twist the adjoint fields $A_\mu(x; \theta, r)$, $\psi(x; \theta, r)$ and $\Phi(x; \theta, r)$ which contain the 4d \mathfrak{g} -valued vector-multiplet. Then, from the 4d perspective, this ‘monodromic’ geometry corresponds to a weakly coupled $\mathcal{N} = 2$ SYM theory with the non-simply laced gauge group G whose Dynkin graph is obtained by folding the Dynkin diagram of \mathfrak{g} along the cyclic group $\{\omega^k\}$.

To make contact between this geometrical construction and the Representation Theory of quivers, we have to recall the categorical interpretation of the degeneration limit of the Gaiotto curve presented in ref. [6]. If the gauge group $G = \exp \mathfrak{g}$ is simple and simply-laced, the SYM light category² \mathcal{L} has the form³ [6]

$$\mathcal{L} = \bigvee_{\lambda \in \mathbb{P}^1} \mathcal{L}_\lambda, \tag{2.7}$$

where \mathcal{L}_λ are Abelian categories, (independent of λ up to equivalence). The bricks⁴ of \mathcal{L}_λ are rigid and their images in the lattice $K_0(\mathcal{L})$ are positive roots of \mathfrak{g} [6]. Physically, these two properties say that the BPs states at weak coupling with zero magnetic charge are vector-multiplets in the adjoint of the gauge group [6]. Cutting away the ends of the tube, corresponds to omitting the two fibers over the poles in the RHS of eq. (2.7), that is, to considering the ‘open’ subcategory

$$\bigvee_{\lambda \in \mathbb{C}^\times} \mathcal{L}_\lambda. \tag{2.8}$$

An outer automorphism ω of \mathfrak{g} induces an autoequivalence \mathcal{G} in each category \mathcal{L}_λ . The categorical analogue of the geometric twisting (2.6) is then

$$\mathcal{L}_{e^{2\pi i} \lambda} = \mathcal{G} \mathcal{L}_\lambda. \tag{2.9}$$

²The main properties of the light category \mathcal{L} are reviewed in section 4 below.

³Given two Krull-Schmidt additive \mathbb{C} -categories, \mathcal{A}, \mathcal{B} , we write $\mathcal{A} \vee \mathcal{B}$ for the category whose objects are the direct sums of objects of \mathcal{A} and \mathcal{B} . The equation in the text then says that an object in \mathcal{L} may be written as a (finite) direct sum $X_{\lambda_1} \oplus \dots \oplus X_{\lambda_\ell}$ with $X_{\lambda_s} \in \mathcal{L}_{\lambda_s}$.

⁴A representation X is called a *brick* iff $\text{End } X = \mathbb{C}$. All stable representations are, in particular, bricks.




Figure 1. Dynkin diagrams foldings. The graphs on the left are called the ‘parent graph’ G_{parent} of the graphs G on the right.

In other words: there is an equivalence of categories of the form

$$\gamma: \mathcal{L}(G)_\lambda \longrightarrow \mathcal{L}(G_{\text{par}})_\lambda \tag{2.10}$$

which associates to each object of the SYM light category for the non-simply laced gauge group G , $\mathcal{L}(G)_\lambda$, an object of the SYM light category with the simply-laced gauge group G_{par} , $\mathcal{L}(G_{\text{par}})_\lambda$. γ preserves all Hom spaces (in particular, it preserves the End rings). However, the equivalence γ is *not* canonical since it depends on the choice of a root of the algebraic equation

$$x^n = \lambda, \quad n \equiv \text{order } \omega. \tag{2.11}$$

Changing the root will replace the equivalence by making

$$\gamma \rightarrow \gamma \circ \mathcal{G}^k, \tag{2.12}$$

for some $k \in \mathbb{Z}$.

3 Q -systems and Dirac integrality conditions

We have very natural candidates for the quiver of all pure SYM with G non-simply laced. Indeed, for each finite-dimensional simple Lie algebra \mathfrak{g} there is a certain integrable statistical system — known as the Q -system of type \mathfrak{g} [32–34] — which is defined in terms of the cluster algebra associated to a certain mutation-class of quivers depending on \mathfrak{g} . For \mathfrak{g} simply-laced, the Q -system quivers coincide with those for pure \mathfrak{g} SYM as described in refs. [1, 4, 6]; it is natural to expect that the identification of the pure SYM quivers with the Q -system ones remains true also for \mathfrak{g} non-simply laced. Indeed, the (classical) Q -systems lead to recursion relations for characters of quantum groups [32] which, in turn, are crucial for the physical interpretation of $\text{rep}(Q, \mathcal{W})$ as the space of BPS of $\mathcal{N} = 2$ SYM with the given gauge group.

There is a general *first principle* strategy to find the quiver of a $\mathcal{N} = 2$ gauge theory (if it exists at all), namely the Dirac integrality conditions [6, 7]. As we are going to show, this method confirms that the Q -systems quivers are the correct ones for pure SYM even in the non-simply laced case.

Let C be the Cartan matrix of \mathfrak{g} ; then the exchange matrix of the (standard representative) quiver of the associated Q -system is the $2r \times 2r$ skew-symmetric matrix⁵

$$B = \left(\begin{array}{c|c} C - C^t & C^t \\ \hline -C & 0 \end{array} \right) \tag{3.1}$$

In other words, the quiver for \mathfrak{g} SYM is obtained by drawing $r = \text{rank } \mathfrak{g}$ vertical Kronecker subquivers $\uparrow\uparrow$ and then connecting the sink of the i -th Kronecker to the source of the j -th one by $|C_{ij}|$ arrows. In addition, one draws $|C_{ij} - C_{ji}|$ extra ‘diagonal’ arrows in such a way that the net number of arrows connecting any two vertical Kronecker subquivers is zero. The *net* number of arrows entering (leaving) the source (resp. the sink) of the i -th Kronecker subquiver \mathbf{Kr}_i is always equal to the integer d_i defined in eq. (2.5).

The extra ‘diagonal’ arrows guarantee that the indecomposable representations X_{α_i} having support in the i -th vertical Kronecker subquiver \mathbf{Kr}_i and dimension vector equal to the imaginary root δ_i of the \hat{A}_1 affine Dynkin graph underlying \mathbf{Kr}_i , are *mutually local*. This last property is crucial for the physical consistency, since the representation X_{α_i} corresponds to the α_i simple-root W -boson, and the W -bosons must be mutually-local.

The charges of the α_i W -boson with respect to the maximal torus $U(1)^r \subset G$ are given by

$$q_j(\alpha_i) \equiv \alpha_i(\alpha_j^\vee) \equiv \frac{2(\alpha_i, \alpha_j)}{(\alpha_j, \alpha_j)} \equiv C_{ij}. \tag{3.2}$$

Let Γ be the charge lattice of the 4d theory, which is identified with the dimension lattice of $\text{rep}(Q, \mathcal{W})$ [2–4]. From eq. (3.2) we see that the vector in $\Gamma \otimes \mathbb{Q}$ corresponding to the unit q_j charge is

$$\mathfrak{q}_j = (C^t)_{ji}^{-1} \dim X_{\alpha_i}. \tag{3.3}$$

⁵In the conventions of [32], this is the exchange matrix of the *opposite* quiver Q^{op} .

The magnetic weights $m_j(X)$ of the representation X are then defined through its Dirac electromagnetic pairing with the unit electric vectors \mathbf{q}_j [6, 7]

$$m_j(X) = -\langle \mathbf{q}_j, \dim X \rangle_{\text{Dirac}} \equiv -(C^t)_{ji}^{-1} \langle \mathbf{q}_j, \dim X \rangle_{\text{Dirac}} \quad (3.4)$$

In the basis (3.1), the charge vector of the W -boson associated to the α_i simple root of \mathfrak{g} is then

$$\dim X_{\alpha_i} \equiv \delta_i = \left(e_i \mid e_i \right) \quad (3.5)$$

where e_i stands for the row r -vector with 1 in position i and zero elsewhere. Hence

$$\langle \delta_i, \dim X \rangle_{\text{Dirac}} = \delta_i B (\dim X)^t = -(C^t \dim X)_{i,1} + (C^t \dim X)_{i,2}, \quad (3.6)$$

which gives the magnetic charges (cf. eq. (3.4))

$$m_j(X) = \dim X_{j,1} - \dim X_{j,2} \quad j = 1, 2, \dots, r, \quad (3.7)$$

which are consistent with Dirac integrality [6, 7].

The quivers for SYM with $\mathfrak{g} = BCFG$ are then as in figure 2.

It remains to determine the superpotential $\mathcal{W}(\mathfrak{g})$ such that the light subcategory $\mathcal{L}(\mathfrak{g}) \subset \text{rep}(Q(\mathfrak{g}), \mathcal{W}(\mathfrak{g}))$ has the required physical properties [6] and, moreover, it enjoys the monodromic properties discussed in the previous section, see eq. (2.9). Before doing that, we review some facts about such light subcategories in a form convenient for our present purposes.

4 Review of the light category $\mathcal{L} \subset \text{rep}(Q, \mathcal{W})$

Suppose we have a quiver Q which corresponds to a $\mathcal{N} = 2$ theory that, in some corner of its parameter space, is a weakly coupled gauge theory with gauge group G . If we know the representations of Q corresponding to the simple-roots α_i , we may define the magnetic charges $m_i(X) \in \mathbb{Z}^r$ of all representations of Q as in eqs. (3.6), (3.7). In the weak coupling limit $g_{\text{YM}} \rightarrow 0$, the central charge of the representation X behaves as [6]

$$Z(X) = -\frac{1}{g_{\text{YM}}^2} \sum_i C_i m_i(X) + O(1), \quad C_i > 0, \quad (4.1)$$

so that all states which have bounded mass as $g_{\text{YM}} \rightarrow 0$ must have⁶ $m_i(X) = 0$ for all i . A representation X with $m_i(X) = 0$ will be very unstable if it has a subrepresentation Y with $m_i(Y) > 0$ for some i . This (and some mathematical theory [6]) motivates the definition of the light subcategory \mathcal{L} of $\text{rep}(Q, \mathcal{W})$ as the full subcategory over the objects X satisfying the following two conditions:

$$(1) \quad m_i(X) = 0 \quad \text{for all } i = 1, 2, \dots, r, \quad (4.2)$$

$$(2) \quad \text{for all subobjects } Y \text{ of } X, \quad m(Y)_i \leq 0 \text{ for all } i. \quad (4.3)$$

⁶To avoid technical complications, one may take the real numbers C_i to be linearly independent over \mathbb{Q} .




Figure 2. The quivers of SYM with non-simply laced gauge group. The corresponding superpotentials are given in section 6.

One shows that \mathcal{L} is an exact Abelian subcategory which is closed under direct sum/summands and extensions.

The basic property of \mathcal{L} is that it contains all representations corresponding to the stable BPS particles which are light in the region $g_{\text{YM}} \sim 0$. If, in this regime, the theory has a weakly coupled Lagrangian description, the stable states of \mathcal{L} (at weak coupling) precisely correspond to the perturbative spectrum of the model. In this case, on physical grounds, the light stable states should consist in one vector-multiplet in the adjoint of \mathfrak{g} plus finitely many hypers in definite \mathfrak{g} -representations.

Such physical considerations imply other Representation Theoretical properties of the light category \mathcal{L} which may be rather surprising from the mathematical side. For instance, a perturbative state certainly cannot, at weak coupling, become non-perturbative by tuning the adjoint Higgs background to trigger the breaking $G \rightarrow \text{SU}(2) \times \text{U}(1)^{r-1}$. For a quiver as in figure 2, this means that⁷

$$X \in \mathcal{L} \implies X|_{\mathbf{Kr}_i} \in \mathcal{L}(\text{SU}(2)) \quad \text{for all } i = 1, 2, \dots, r, \quad (4.4)$$

where $\mathcal{L}(\text{SU}(2))$ denotes the light category for pure $\text{SU}(2)$ SYM, equal to the category of regular representations of the Kronecker quiver $\uparrow\uparrow$ [6, 38–40]. With some pain the property (4.4) may be proven directly from representation theory, see [6, 7] and appendix A.

As explained in refs. [6, 7], the property (4.4) implies a major simplification; indeed, it allows to identify the sink $i, 2$ and source $i, 1$ nodes of each \mathbf{Kr}_i subquiver by setting equal to 1 one of the arrows, say B_i , in each vertical Kronecker subquiver (cf. figure 2). In this way we get a smaller quiver Q' having r less nodes than the original one. Under this reduction, the superpotential \mathcal{W} induces an effective superpotential \mathcal{W}' for Q' [7] and one has simply

$$\mathcal{L} = \text{rep}(Q', \mathcal{W}').$$

Notice that in a theory with a Lagrangian description all states in \mathcal{L} are mutually local so that

$$\langle \cdot, \cdot \rangle_{\text{Dirac}}|_{\mathcal{L}} = 0, \quad (4.5)$$

and the net number of arrows between any two nodes of Q' must be zero

$$\#\{\text{arrows } i \rightarrow j \text{ in } Q'\} - \#\{\text{arrows } i \leftarrow j \text{ in } Q'\} = 0. \quad (4.6)$$

In particular, Q' , if connected, cannot be 2-acyclic as the typical quiver for a 4d $\mathcal{N} = 2$ theory. This is related to the fact that \mathcal{L} is not the non-perturbative Abelian category representing all BPS states of the theory, but rather it represent only a subsector (the perturbative one) which is not a full QFT. The complete spectrum also contain dyons with masses of order $O(1/g_{\text{YM}}^2)$. Lifting the perturbative category \mathcal{L} , which captures the classical physics, to its non-perturbative completion $\text{rep}(Q, \mathcal{W})$ is a sort of quantization procedure which we shall refer to as ‘categorical quantization’.

There is another property of the light category which is crucial for our purposes and was already mentioned in section 2. The pure $\text{SU}(2)$ light category, $\mathcal{L}(\text{SU}(2))$, has the following

⁷For the corresponding statement in more general situations, see [7].

structure [38–40]: there is a family of orthogonal subcategories $\mathcal{L}(\text{SU}(2))_\lambda \subset \mathcal{L}(\text{SU}(2))$, labelled by a point $\lambda \in \mathbb{P}^1$ such that each object $X \in \mathcal{L}(\text{SU}(2))$ is a (finite) direct sum $\oplus_\lambda X_\lambda$ with $X_\lambda \in \mathcal{L}(\text{SU}(2))_\lambda$. In fact, $\mathcal{L}(\text{SU}(2))_\lambda$ is the closure with respect to direct sums and extensions of the brick representation

$$\mathbb{C} \begin{array}{c} \xrightarrow{\lambda} \\ \xrightarrow{1} \end{array} \mathbb{C} \tag{4.7}$$

of dimension δ equal to the minimal imaginary root of \widehat{A}_1 , which is physically identified with the charge vector of the $\text{SU}(2)$ W -boson. In symbols these facts are expressed as [38]

$$\mathcal{L}(\text{SU}(2)) = \bigvee_{\lambda \in \mathbb{P}^1} \mathcal{L}(\text{SU}(2))_\lambda. \tag{4.8}$$

Physically, this equations just states that the W -boson is a vector-multiplet and that there are neither light hypermultiplets nor light higher-spin supermultiplets in the spectrum of pure $\text{SU}(2)$ SYM [6].

For general $\mathcal{N} = 2$ gauge theories eq. (4.8) generalizes as follows [6]: the light category has the form

$$\mathcal{L} = \bigvee_{\lambda \in N} \mathcal{L}_\lambda, \tag{4.9}$$

where N is a one-dimensional variety whose irreducible components N_a are in one-to-one correspondence with the simple factors in the gauge group G . Moreover, in class- S theories, N may be identified with the skeleton of the particular degeneration of the Gaiotto curve which is relevant for the S -duality frame specified by the given choice of magnetic charges $m_i(\cdot)$. Matter lives in the categories \mathcal{L}_{λ_s} where $\{\lambda_s\}$ is a finite set of points in N , and the matter in \mathcal{L}_{λ_s} may be charged only with respect to the gauge factor groups G_a whose irreducible component $N_a \ni \lambda_s$. Again, these statements just correspond to the fact that the gauge particles are vectors while the matter has only spin 0, 1/2 states. In the next section we shall show directly (and rigorously) that all these general statements hold true for the models (and categories) of interest in this paper.

The situations explicitly analyzed in [6] correspond to the *split*, i.e. ADE case. In that case, assuming the gauge group G to be simple, eq. (4.4) has a nice refinement: for all $\lambda \in \mathbb{P}^1$,

$$X \in \mathcal{L}_\lambda \implies X|_{\mathbf{K}r_i} \in \mathcal{L}(\text{SU}(2))_\lambda \quad \text{for all } i = 1, 2, \dots, r. \tag{4.10}$$

This last equation will not longer be true in the *non-split* case, and its failure is an alternative way of characterizing the non-split phenomenon.

It will be shown in the next two sections that the generalization of eq. (4.10) to the case of a gauge group G which is simple and *non*-simply laced is as follows. Let $d_i \in \mathbb{N}$ be the integers defined in eq. (2.5). Then eq. (4.10) is replaced by

$$X \in \mathcal{L}_\lambda \implies X|_{\mathbf{K}r_i} \in \mathcal{L}(\text{SU}(2))_{\lambda^{1/d_i}} \quad \text{for all } i = 1, 2, \dots, r. \tag{4.11}$$

The fact that in the RHS we have to choose a square root (or a cubic root in the G_2 case) of $\lambda \in \mathbb{C}^\times$ opens the possibility that going along a non-contractible loop in \mathbb{C}^\times we end up with a different choice for the root. This will produce a monodromy having the form predicted in eq. (2.9). More detailed statements below.

5 The light category of non-simply laced $\mathcal{N} = 2$ SYM

As in ref. [7], our strategy for fixing the superpotential $\mathcal{W}(\mathfrak{g})$ is to require that the corresponding category $\text{rep}(Q(\mathfrak{g}), \mathcal{W}(\mathfrak{g}))$ contains a light category $\mathcal{L}(\mathfrak{g})$ with the right physical properties as described in section 4, and, in particular, that the light BPS states at weak coupling consist of just one vector-multiplet in the adjoint of \mathfrak{g} . Contrary to ref. [7], however, we shall not put forward an ansatz for $\mathcal{W}(\mathfrak{g})$ and then check the properties of $\mathcal{L}(\mathfrak{g})$, but rather we shall directly construct the correct light category $\mathcal{L}(\mathfrak{g})$ from the monodromic properties it must enjoy (section 2), and then build the full non-perturbative category $\text{rep}(Q(\mathfrak{g}), \mathcal{W}(\mathfrak{g}))$ out of its light subcategory $\mathcal{L}(\mathfrak{g})$ by the process we have dubbed ‘categorical quantization’.

We start by constructing the non-simply laced monodromic counterparts to the usual *ADE* preprojective algebras [41–44] which played such a prominent role in refs. [6, 7].

5.1 Monodromic light categories

We start by considering certain monodromic module categories which play for the non-simply laced case a role somehow similar to that played, in the simply laced case [6], by the modules over the preprojective algebra $\mathcal{P}(G)$ (where the graph G is the Dynkin diagram of the gauge group).

5.1.1 Preprojective algebras for non-simply laced (i.e. valued) graphs

For our present purposes, a *valued* graph \mathfrak{g} is a *simply laced* graph together with the assignment of a pair (n_α, m_α) of positive integers to each link α of \mathfrak{g} ; at least one of the two numbers (n_α, m_α) must be 1, and if they are both 1 the link is said to have trivial valuation and is denoted by a plain edge. The Dynkin graph of a finite-dimensional (simple) Lie algebra \mathfrak{g} may be seen as a valued graph with valuation function

$$(n(\alpha), m(\alpha)) \equiv \left(|C_{s(\alpha), t(\alpha)}|, |C_{t(\alpha), s(\alpha)}| \right). \tag{5.1}$$

\mathfrak{g} is simply laced precisely iff all links have trivial valuation.

Given a valued graph \mathfrak{g} , we define its *$N = 2$ double* $\widehat{\mathfrak{g}}$ to be the quiver obtained by replacing each edge α of \mathfrak{g} with a pair of opposite oriented arrows⁸

$$\begin{array}{c} \xrightarrow{\alpha} \\ \xleftarrow{\alpha^*} \end{array}$$

In addition, the quiver $\widehat{\mathfrak{g}}$ also has an ‘adjoint’ loop A_i at each node i of \mathfrak{g} . For future convenience, between nodes i, j we also draw $(\max\{|C_{ij}|, |C_{ji}|\} - 1)$ *dashed* pairs of opposite arrows. We stress that the dashed arrows are NOT part of the quiver $\widehat{\mathfrak{g}}$, and most of the times we shall even *omit* to draw them; they will be needed only in section 5.2.

⁸We shall refer to the arrows α as the *direct* arrows, while the arrows α^* will be called *inverse* arrows.

For instance, for $\mathfrak{g} = F_4$ we get

$$\widehat{F}_4 : \quad A_1 \begin{array}{c} \circlearrowleft \\ \end{array} 1 \begin{array}{c} \xrightarrow{\alpha_1} \\ \xleftarrow{\alpha_1^*} \end{array} 2 \begin{array}{c} \xrightarrow{\alpha_2} \\ \xleftarrow{\alpha_2^*} \end{array} 3 \begin{array}{c} \xrightarrow{\alpha_3} \\ \xleftarrow{\alpha_3^*} \end{array} 4 \begin{array}{c} \circlearrowleft \\ \end{array} A_4 \quad (5.2)$$

We equip $\widehat{\mathfrak{g}}$ with the superpotential⁹

$$\mathcal{W} = \sum_{\alpha} \left(\alpha A_{s(\alpha)}^{n(\alpha)} \alpha^* - \alpha^* A_{t(\alpha)}^{m(\alpha)} \alpha \right), \quad (5.3)$$

where the sum is over the direct arrows only.

We define the algebra $\mathcal{P}(\mathfrak{g})$ to be the Jacobian algebra

$$\mathcal{P}(\mathfrak{g}) := \mathbb{C}\widehat{\mathfrak{g}}/(\partial\mathcal{W}). \quad (5.4)$$

In case \mathfrak{g} is simply laced (i.e. trivially valued), $\mathcal{P}(\mathfrak{g})$ is essentially the preprojective algebra of the graph [41–44]. We shall loosely refer to $\mathcal{P}(\mathfrak{g})$ as *the preprojective algebra* of \mathfrak{g} also in the non-simply laced case.

5.1.2 The structure of $\text{mod } \mathcal{P}(\mathfrak{g})$

From now on the valued graph \mathfrak{g} is the Dynkin diagram of a finite-dimensional simple Lie algebra.

We wish to prove the following two crucial facts:

- (1) *The Abelian category $\text{mod } \mathcal{P}(\mathfrak{g})$ has a decomposition into (exact, full, extension-closed) Abelian subcategories of the following form*

$$\text{mod } \mathcal{P}(\mathfrak{g}) \equiv \mathcal{M}(\mathfrak{g}) = \bigvee_{\lambda \in \mathbb{P}^1} \mathcal{M}(\mathfrak{g})_{\lambda}. \quad (5.5)$$

- (2) *The bricks of $\mathcal{P}(\mathfrak{g})$ belong to \mathbb{P}^1 families $X_{\lambda} \in \mathcal{M}(\mathfrak{g})_{\lambda}$ ($\lambda \in \mathbb{P}^1$) and their dimension vectors are positive roots of \mathfrak{g} .*

These results are known to hold in the simply-laced cases and, in that situation, they imply that in each weakly coupled chamber the light BPS spectrum consists precisely of vector-multiplets making one copy of the adjoint of \mathfrak{g} [6]. However, in the non-simply laced case, the corresponding representations will also have non-trivial monodromy, in agreement with the geometric discussion of section 2.

⁹Here and in the following we adopt the more concise mathematical notation for the superpotential [48]. Under the identification of the arrows of the quiver with the 1d ‘bifundamental’ Higgs fields, the physical superpotential is

$$W = \text{tr } \mathcal{W}.$$

Therefore, two words of $\mathbb{C}Q$, corresponding to an oriented cycle in Q , are considered equivalent as contributions to \mathcal{W} if they differ by a cyclic permutation of the arrows.

Again we write d_i ($i = 1, 2, \dots, r$) for the integers defined in eq. (2.5). The valuation of the edge $i \xrightarrow{\alpha} j$ of \mathfrak{g} is

$$(n(\alpha), m(\alpha)) = \left(\frac{d_i}{(d_i, d_j)}, \frac{d_j}{(d_i, d_j)} \right). \tag{5.6}$$

As a consequence of the relations $\partial \mathcal{W} = 0$, for all $X \in \text{mod } \mathcal{P}(\mathfrak{g})$ the set of linear maps

$$X_i \longmapsto A_i^{d_i} X_i \quad \text{for all } i, \tag{5.7}$$

gives an element of $\text{End } X$. Suppose X is *indecomposable*. Then $\text{End } X$ is a *local ring* [45], and hence there is a complex number $\lambda \in \mathbb{C}$ and integers f_i such that

$$(A_i^{d_i} - \lambda)^{f_i} = 0 \quad \text{for all } i = 1, 2, \dots, r. \tag{5.8}$$

We write $\mathcal{M}(\mathfrak{g})_\lambda$ for the Krull-Schmidt full subcategory of $\text{mod } \mathcal{P}(\mathfrak{g})$ whose objects are the direct sums of indecomposables satisfying (5.8) for the given $\lambda \in \mathbb{C}$. It is manifestly an Abelian category. This shows (1).

In the particular case that X is actually a brick we have the stronger statement

$$X \text{ is a brick} \implies A_i^{d_i} = \lambda \quad \text{for all } i = 1, 2, \dots, r. \tag{5.9}$$

Let us consider the case of λ generic (i.e. $\lambda \neq 0$). Then A_i is semisimple with eigenvalues $e^{2\pi i \ell_i / d_i} \lambda^{1/d_i}$, $\ell_i = 1, 2, \dots, d_i$. Let P_{i, ℓ_i} be the projector on the corresponding eigenspace, so that

$$X_i = P_{i,1} X_i \oplus \dots \oplus P_{i,d_i} X_i. \tag{5.10}$$

Thus we split each vector space X_i in as many vector spaces as the number of nodes of the parent graph $\mathfrak{g}_{\text{parent}}$ folded into the i -th node. By projecting the arrows in these eigenspaces, we get a representation X^b of the double quiver $\overline{\mathfrak{g}_{\text{parent}}}$ which satisfies the usual $\mathcal{P}(\mathfrak{g}_{\text{parent}})$ relations. It is elementary to check that X is a brick iff the $\mathcal{P}(\mathfrak{g}_{\text{parent}})$ module X^b is a brick. Then the statement reduces to the simply-laced case, which is known to hold ([6] and reference therein). Thus (2) is true.

A few explicit examples will make (1)(2) more concrete.

5.1.3 Example: B_2

In this case the quiver and superpotential are

$$A_1 \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} 1 \begin{array}{c} \xleftarrow{\alpha^*} \\ \xrightarrow{\alpha} \end{array} 2 \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} A_2 \tag{5.11}$$

$$\mathcal{W} = \alpha A_1 \alpha^* - \alpha \alpha^* A_2^2. \tag{5.12}$$

The relations are

$$\partial_\alpha \mathcal{W} \equiv A_1 \alpha^* - \alpha^* A_2^2 = 0 \tag{5.13}$$

$$\partial_{\alpha^*} \mathcal{W} \equiv \alpha A_1 - A_2^2 \alpha = 0 \tag{5.14}$$

$$\partial_{A_1} \mathcal{W} \equiv \alpha^* \alpha = 0 \tag{5.15}$$

$$\partial_{A_2} \mathcal{W} \equiv -\alpha \alpha^* A_2 - A_2 \alpha \alpha^* = 0. \tag{5.16}$$

from which we see that the map

$$\ell: (X_1, X_2) \mapsto (A_1 X_1, A_2^2 X_2), \quad (5.17)$$

is an element of $\text{End } X$ and hence a complex number $\lambda \in \mathbb{P}^1$ if X is a brick.

For λ generic (that is, $\lambda \neq 0, \infty$) A_2 may be written in the form

$$A_2 = \sqrt{\lambda} \left(\begin{array}{c|c} \mathbf{1}_{n \times n} & \mathbf{0}_{n \times m} \\ \hline \mathbf{0}_{m \times n} & -\mathbf{1}_{m \times m} \end{array} \right). \quad (5.18)$$

Let P_{\pm} the projectors to the two eigenspaces of A_2 . Clearly a brick X with $\lambda \neq 0, \infty$ induces a representation of the *unfolded* quiver

$$\begin{array}{ccccc} & & P_+ \alpha & & -\alpha^* P_- \\ & \swarrow & & \searrow & \\ P_+ X_2 & & & & P_- X_2 \\ & \nwarrow & & \nearrow & \\ & & \alpha^* P_+ & & P_- \alpha \end{array} \quad X_1 \quad (5.19)$$

which satisfies the relations

$$P_{\pm} \alpha \cdot \alpha^* P_{\pm} = 0 \quad (\text{by eq. (5.16)}) \quad (5.20)$$

$$\alpha^* P_+ \cdot P_+ \alpha - (-\alpha^* P_-) \cdot P_- \alpha = \alpha^* \alpha = 0, \quad (\text{by eq. (5.15)}) \quad (5.21)$$

which are precisely those of $\mathcal{P}(A_3)$. Hence the bricks of X are mapped into representations of $\mathcal{P}(A_3)$ which have dimension vectors equal to the positive roots of A_3 . By folding back the quiver, we get (essentially by definition) the positive roots of B_2 .

5.1.4 Example: G_2

The quiver is the same as in eq. (5.11) with a different superpotential

$$\mathscr{W} = \alpha A_1 \alpha^* - \alpha \alpha^* A_2^3. \quad (5.22)$$

The relations are

$$\partial_{\alpha} \mathscr{W} \equiv A_1 \alpha^* - \alpha^* A_2^3 = 0 \quad (5.23)$$

$$\partial_{\alpha^*} \mathscr{W} \equiv \alpha A_1 - A_2^3 \alpha = 0 \quad (5.24)$$

$$\partial_{A_1} \mathscr{W} \equiv \alpha^8 \psi = 0 \quad (5.25)$$

$$\partial_{A_2} \mathscr{W} \equiv -(A_2^2 \alpha \alpha^* + A_2 \alpha \alpha^* A_2 + \alpha \alpha^* A_2^2) = 0. \quad (5.26)$$

The map

$$(X_1, X_2) \mapsto (A_1 X_1, A_2^3 X_2) \quad (5.27)$$

is an element of $\text{End } X$, hence a complex number λ . For generic λ , A_2 has eigenvectors $\lambda e^{2\pi i k/3}$, with $k = 0, 1, 2$. Let P_0, P_1, P_2 be the projectors on the corresponding eigenspaces.

Then we get a representation of the quiver

$$\begin{array}{ccccc}
 & & P_0 \alpha & & \alpha^* P_2 \\
 & \swarrow & \curvearrowright & \searrow & \swarrow \\
 P_0 X_2 & & X_1 & & P_2 X_2 \\
 & \searrow & \curvearrowleft & \swarrow & \searrow \\
 & & P_1 \alpha & & P_2 \alpha \\
 & & P_1 X_2 & &
 \end{array}
 \quad (5.28)$$

which satisfies the relations of $\mathcal{P}(D_4)$ (cf. eqs. (5.26), (5.25))

$$P_k \alpha \cdot \alpha^* P_k = 0 \quad k = 0, 1, 2 \quad \sum_k (\alpha^* P_k) \cdot (P_k \alpha) = 0. \quad (5.29)$$

5.1.5 Example: F_4

The quiver and \mathscr{W} in this case are

$$\begin{array}{ccccccc}
 A_1 \circlearrowleft & 1 & \xleftarrow{\alpha_1^*} & 2 & \xleftarrow{\alpha_2^*} & 3 & \xleftarrow{\alpha_3^*} & 4 \circlearrowright A_4 \\
 & & & \curvearrowright A_2 & & \curvearrowright A_3 & &
 \end{array}
 \quad (5.30)$$

$$\mathscr{W} = A_1 \alpha_1^* \alpha_1 - A_2 \alpha_1 \alpha_1^* + A_2 \alpha_2^* \alpha_2 - A_3^2 \alpha_2 \alpha_2^* + A_3 \alpha_3^* \alpha_3 - A_4 \alpha_3 \alpha_3^*. \quad (5.31)$$

From the relations $\partial_{\alpha_i^*} \mathscr{W} = \partial_{\alpha_i} \mathscr{W} = 0$ it follows that the map

$$(X_1, X_2, X_3, X_4) \mapsto (A_1 X_1, A_2 X_2, A_3^2 X_3, A_4 X_4), \quad (5.32)$$

is an element of $\text{End } X$ and hence a number $\lambda \in \mathbb{C}$ if X is a brick. Then A_3 and A_4 have eigenvalues $\pm\sqrt{\lambda}$, and (for $\lambda \neq 0$) let P_{\pm} and Q_{\pm} be the corresponding projectors. We get the representation

$$\begin{array}{ccccc}
 & & & P_+ X_3 & \xleftrightarrow{Q_+ \alpha_3 P_+ / (2\sqrt{\lambda})} & Q_+ X_4 \\
 & & & \swarrow & \searrow & \\
 & & & \alpha_2^* P_+ & & \\
 & & & P_+ \alpha_2 & & \\
 X_1 & \xleftrightarrow{\alpha_1} & X_2 & & & \\
 & \swarrow & \searrow & & & \\
 & & & P_- \alpha_2 & & \\
 & & & \alpha_2^* P_- & & \\
 & & & P_- X_3 & \xleftrightarrow{-Q_- \alpha_3 P_- / (2\sqrt{\lambda})} & Q_- X_4 \\
 & & & & \swarrow & \searrow \\
 & & & & P_- \alpha_3^* Q_- &
 \end{array}
 \quad (5.33)$$

Notice that the relations imply

$$P_{\pm} \alpha_3^* Q_{\mp} = Q_{\pm} \alpha_3 P_{\mp} = 0, \quad (5.34)$$

so the arrows connect the various subspaces precisely as in the above quiver (5.33). The relations of the projective algebra $\mathcal{P}(E_6)$ are then satisfied; for instance

$$P_{\pm}\alpha_2 \cdot \alpha_2^*P_{\pm} - P_{\pm}\alpha_3^*Q_{\pm} \cdot \left(\pm \frac{1}{2\sqrt{\lambda}}Q_{\pm}\alpha_3P_{\pm} \right) = \mp \frac{1}{2\sqrt{\lambda}}P_{\pm} \partial_{A_3} \mathcal{W} P_{\pm} = 0. \quad (5.35)$$

Again our claim is verified.

5.1.6 Monodromy of $\mathcal{M}(\mathfrak{g})$

From the above examples, it is clear that there is an auto-equivalence

$$\mathcal{G}: \mathcal{M}(\mathfrak{g})_{\lambda} \rightarrow \mathcal{M}(\mathfrak{g})_{\lambda} \quad (5.36)$$

which acts by permuting the roots of the algebraic equation

$$\begin{aligned} x^2 &= \lambda && \text{for } \mathfrak{g} \neq G_2 \\ x^3 &= \lambda && \text{for } \mathfrak{g} = G_2. \end{aligned}$$

For instance, \mathcal{G} will act on a brick in $\mathcal{M}(G_2)_{\lambda}$ with dimension (1, 1) as

$$\lambda \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \mathbb{C} \begin{array}{c} \longleftarrow \\ \longrightarrow \end{array} \mathbb{C} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \lambda^{1/3} \xrightarrow{\mathcal{G}} \lambda \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} \mathbb{C} \begin{array}{c} \longleftarrow \\ \longrightarrow \end{array} \mathbb{C} \begin{array}{c} \circlearrowleft \\ \circlearrowright \end{array} e^{2\pi i/3} \lambda^{1/3} \quad (5.37)$$

which corresponds in terms of the covering $\mathcal{P}(D_4)$ representations to

$$\begin{array}{ccc} \begin{array}{c} \mathbb{C} \\ \swarrow \nearrow \\ \mathbb{C} \begin{array}{c} \longleftarrow \\ \longrightarrow \end{array} 0 \\ \swarrow \nearrow \\ 0 \end{array} & \longrightarrow & \begin{array}{c} 0 \\ \swarrow \nearrow \\ \mathbb{C} \begin{array}{c} \longleftarrow \\ \longrightarrow \end{array} \mathbb{C} \\ \swarrow \nearrow \\ 0 \end{array} \end{array} \quad (5.38)$$

Then the family of Abelian categories $\mathcal{M}(\mathfrak{g})_{\lambda}$ manifestly enjoys the monodromic properties discussed in section 2, namely

$$\mathcal{M}(\mathfrak{g})_{e^{2\pi i}\lambda} = \mathcal{G} \mathcal{M}(\mathfrak{g})_{\lambda}. \quad (5.39)$$

Note that the action of \mathcal{G} on the objects of $\mathcal{M}(\mathfrak{g})_{\lambda}$ (taken up to isomorphism) is not free. For instance, in the G_2 case the simple representation of dimension (1, 0) is fixed by the functor \mathcal{G} . More generally, under the lift

$$\mathcal{M}(\mathfrak{g}) \rightarrow \text{mod } \mathcal{P}(\mathfrak{g}_{\text{parent}})$$

the action of \mathcal{G} gets identified with the auto-equivalence ϖ associated to the automorphisms of the Dynkin graph \mathfrak{g} . Then the action of \mathcal{G} is not free precisely on the objects whose lift is ϖ -invariant.

Since the action of \mathcal{G} is not free, the present construction differs from the notion of a Galois cover of a quiver as defined by Gabriel [46, 47], which requires the group to act freely on the nodes of the quiver.

5.2 ‘Auxiliary’ fields and the final form of the light category $\mathcal{L}(\mathfrak{g})$

In the definition of the $N = 2$ double quiver $\widehat{\mathfrak{g}}$ in section 5.1.1 we added pairs $\langle \cdots \rangle$ of *dashed* arrows which we dropped in all subsequent discussions. Now it is time to put them to some good use.

The point is that, for \mathfrak{g} *non*-simply laced, the superpotential \mathcal{W} in eq. (5.3) is not in a form suitable for categorical quantization. In facts, it is not linear in the loops A_i — that is, in the physical language, is not of first order in the adjoint Higgs fields A_i . This makes 1d *extended* SUSY not manifest. There is a simple and well known remedy to this problem: We have to add to the 1d SQM massive ‘auxiliary’ Higgs in such a way that integrating them away will produce the higher order term in the superpotential. The dashed arrows in $\widehat{\mathfrak{g}}$ correspond to such ‘auxiliary’ Higgs fields. Then we shall describe the category $\mathcal{M}(\mathfrak{g})$, up to equivalence, in the form

$$\mathcal{M}(\mathfrak{g}) = \text{rep}(Q(\mathfrak{g})', \mathcal{W}(\mathfrak{g})') \equiv \mathcal{L}(\mathfrak{g}), \quad (5.40)$$

where the quiver $Q(\mathfrak{g})'$ is just $\widehat{\mathfrak{g}}$ with the dashed arrows made solid and $\mathcal{W}(\mathfrak{g})'$ is such that it becomes \mathcal{W} after integrating away the massive arrows, that is, in the mathematical language, $(\widehat{\mathfrak{g}}, \mathcal{W})$ is the *reduced* pair of $(Q(\mathfrak{g})', \mathcal{W}(\mathfrak{g})')$ in the sense of DWZ [48].

In the Dynkin graph of a non-simply laced finite-dimensional (simple) Lie algebra we have precisely *one* link with a non-trivial valuation (which is (1,2), except for G_2 which has a (1,3) edge). Therefore, in the sum in the RHS of eq. (5.3) there is precisely one term of higher order in the loop arrows A_i . This is the only term we have to rewrite in terms of lower order ones. Therefore we may focus on the full subquiver of $Q(\mathfrak{g})'$ over the two nodes connected by the link with the non-trivial valuation, since everything else will remain invariant.

We are reduced to consider two situations, corresponding to the two possible non-trivial link valuations. The relevant (sub)quiver in the two cases is, respectively,

$$\begin{array}{ccc}
 & \eta^* & \\
 & \curvearrowright & \\
 A_i \curvearrowright i & \xleftarrow{\alpha^*} & j \curvearrowleft A_j \\
 & \xrightarrow{\alpha} & \\
 & \curvearrowleft & \\
 & \eta &
 \end{array} \quad (5.41)$$

$$\begin{array}{ccc}
 & \xi^* & \\
 & \curvearrowright & \\
 A_i \curvearrowright i & \xleftarrow{\alpha^*} & j \curvearrowleft A_j \\
 & \xrightarrow{\alpha} & \\
 & \curvearrowleft & \\
 & \eta & \\
 & \xi &
 \end{array} \quad (5.42)$$

while the relevant terms in \mathcal{W} are (cf. eq. (5.3))

$$\mathcal{W} = A_i \alpha^* \alpha - A_j^2 \alpha \alpha^* + \cdots \quad (5.43)$$

$$\mathcal{W} = A_i \alpha^* \alpha - A_j^3 \alpha \alpha^* + \cdots \quad (5.44)$$

Using the auxiliary arrows η, η^*, ξ, ξ^* , these terms are rewritten, respectively, as

$$\mathcal{W}' = A_i \alpha^* \alpha + A_j \alpha \eta^* + A_j \eta \alpha^* + \eta \eta^* + \dots \tag{5.45}$$

$$\mathcal{W}' = A_i \alpha^* \alpha - A_j \alpha \xi^* - A_j \xi \eta^* - A_j \eta \alpha^* + \xi \xi^* + \eta \eta^* + \dots \tag{5.46}$$

It is easy to check that integrating out the (massive) auxiliary arrows gives back (5.43), (5.44). Using this rule, we construct $\mathcal{W}(\mathfrak{g})'$ for arbitrary (non-simply laced) \mathfrak{g} .

The conclusion of this section is that we have an equivalence between the light category for \mathfrak{g} $\mathcal{N} = 2$ SYM which (for the moment) we just *define* to be

$$\mathcal{L}(\mathfrak{g}) \equiv \text{rep}(Q(\mathfrak{g})', \mathcal{W}(\mathfrak{g})') \tag{5.47}$$

and the module category $\mathcal{M}(\mathfrak{g})$ we introduced previously. Then, using results (1)(2) of section 5.1.2, we deduce the following crucial properties:

$$(1) \quad (\mathbb{P}^1\text{-family}) \quad \mathcal{L}(\mathfrak{g}) = \bigvee_{\lambda \in \mathbb{P}^1} \mathcal{L}(\mathfrak{g})_\lambda \tag{5.48}$$

$$(2) \quad (\text{non-split monodromy}) \quad \mathcal{L}(\mathfrak{g})_{e^{2\pi i} \lambda} = \mathcal{G} \mathcal{L}(\mathfrak{g})_\lambda \tag{5.49}$$

$$(3) \quad (\text{bricks \& positive roots}) \quad X \text{ is a brick} \Rightarrow \dim X \in \Delta^+(\mathfrak{g}). \tag{5.50}$$

In facts, (3) is slightly stronger than what we have actually proven, since we have shown it only for $\lambda \neq 0, \infty$. It is easy to see that the result extends to $\lambda = 0$. For instance, in the example of section 5.1.3, at $\lambda = 0$, $\alpha^* \alpha = 0$ while the matrices $\gamma_1 = \alpha \alpha^*$ and $\gamma_2 = A_2$ satisfy the Clifford algebra of the trivial quadratic form

$$\gamma_1^2 = \gamma_2^2 = \gamma_1 \gamma_2 + \gamma_2 \gamma_1 = 0, \tag{5.51}$$

which, by the universality of the Clifford algebra, means that the representations of the quiver

$$1 \begin{array}{c} \xleftarrow{\alpha^*} \\ \xrightarrow{\alpha} \end{array} 2 \begin{array}{c} \circlearrowleft \\ \gamma_2 \end{array} \tag{5.52}$$

which are not direct sums of representations with either γ_1 or γ_2 vanishing, must contain a direct summand of the form

$$\alpha = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \otimes \mathbf{1}_2, \quad \alpha^* = \begin{pmatrix} 0 & 1 \end{pmatrix} \otimes \mathbf{1}_2, \quad A_2 = \sigma_3 \otimes \sigma^+, \tag{5.53}$$

whose endomorphism ring contains the nilpotent $\mathbf{1}_2 \otimes \sigma^+$. Hence, if X is a brick, either $\alpha \alpha^*$ or A_2 is zero. If $A_2 = 0$, we are reduced to a string algebra [49] whose strings have dimensions in $\Delta^+(B_2)$. In the other case, $\alpha \alpha^* = \alpha^* \alpha = 0$, and we again reduce to a string algebra whose strings have dimensions in $\Delta^+(B_2)$ or equal to $(2, 2)$, and in the last case $\text{End } X$ necessarily contains nilpotent elements. In conclusion, all bricks in $\mathcal{L}(B_2)_0$ have dimension vectors in $\Delta^+(B_2)$. The opposite case $\lambda = \infty$ will be briefly commented in the next section.

From the physical point of view the fact that (3) extends to $\lambda = 0, \infty$ is often obvious. Indeed, the existence of bricks at $\lambda = 0, \infty$ with dimensions $\notin \Delta^+(\mathfrak{g})$ would be interpreted as the coupling to SYM of matter which must be Lagrangian and carry no flavor charge, that is, to consist of *half*-hypermultiplets. For most non-simply laced \mathfrak{g} there is just no such matter systems.

6 The quiver superpotential \mathcal{W} for $BCFG$ SYM

To complete the construction, we have to endow the quivers in figure 2 with superpotentials $\mathcal{W}(\mathfrak{g})$, then determine the corresponding light categories $\mathcal{L}(\mathfrak{g})$ — which are well-defined since we know the magnetic charges, see section 3 — and finally check that we obtain the same light categories as defined in eq. (5.47). After that, we are guaranteed that our pairs $(Q(\mathfrak{g}), \mathcal{W}(\mathfrak{g}))$ predict the physically correct weak coupling spectrum, namely just one light BPS vector-multiplet in the adjoint of \mathfrak{g} , and hence the pair $(Q(\mathfrak{g}), \mathcal{W}(\mathfrak{g}))$ is consistently identified with the quiver with superpotential for pure $\mathcal{N} = 2$ \mathfrak{g} SYM. On the other hand, these pairs precisely correspond to the geometrical split/non-split realization of ADE vs. $BCFG$ gauge symmetries, so if the ADE pairs are correctly identified (and they are!!) so are the $BCFG$ ones.

The quivers in figure 2 by construction satisfy the conditions that the X_{α_i} have Kronecker supports; in facts they are precisely the representations with support in the vertical Kroneckers $\uparrow\uparrow$ (resp. $\downarrow\downarrow$) and dimension equal to the corresponding minimal imaginary root. Then the inverse process from the light effective superpotential \mathcal{W}' to the full one \mathcal{W} consists in just reinserting the arrows B_i (cf. figure 2) which were set to 1 in the reduction from the full non-perturbative module category to its light subcategory. There is a unique gauge-invariant way of doing this, so \mathcal{W} is uniquely recovered.

With reference to the labeling of arrows in figure 2, the superpotentials are

$$\begin{aligned} \mathcal{W}(B_r) = & A_2 \alpha_1^* B_1 \alpha_1 + A_1 \alpha_1 B_2 \eta^* + A_1 \eta \alpha_1^* + B_1 \eta \eta^* + \\ & + \sum_{s=2}^{r-1} (A_{s+1} \alpha_s^* B_s \alpha_s - B_{s+1} \alpha_s^* A_s \alpha_s) \end{aligned} \quad (6.1)$$

$$\begin{aligned} \mathcal{W}(C_r) = & A_1 \alpha_1^* B_2 \alpha_1 + A_2 \alpha_1 B_1 \eta^* + A_2 \eta \alpha_1^* + B_2 \eta \eta^* + \\ & + \sum_{s=2}^{r-1} (A_s \alpha_s^* B_{s+1} \alpha_s - B_s \alpha_s^* A_{s+1} \alpha_s) \end{aligned} \quad (6.2)$$

$$\mathcal{W}(G_2) = A_1 \alpha^* B_2 \alpha - A_2 \alpha B_1 \xi^* - A_2 \xi \eta^* - A_2 \eta \alpha^* + B_2 \xi \xi^* + B_2 \eta \eta^* \quad (6.3)$$

$$\begin{aligned} \mathcal{W}(F_4) = & A_1 \alpha_1^* B_2 \alpha_1 - B_1 \alpha_1^* A_2 \alpha_1 + A_2 \alpha_2^* B_3 \alpha_2 + A_3 \alpha_2 B_2 \eta^* + \\ & + A_3 \eta \alpha_2^* + B_3 \eta \eta^* + A_3 \alpha_3^* B_4 \alpha_3 - B_3 \alpha_3^* A_4 \alpha_3. \end{aligned} \quad (6.4)$$

The only thing that remains to be done is to check the consistency of the reduction

$$\text{rep}(Q, \mathcal{W}) \rightarrow \text{rep}(Q', \mathcal{W}') \equiv \mathcal{L}. \quad (6.5)$$

This requires two properties to hold. First the non-perturbative category $\text{rep}(Q, \mathcal{W})$ should be consistent with the perturbative Higgs mechanism [6, 7] with requires

$$X \in \mathcal{L} \quad \Rightarrow \quad X|_{\mathbf{Kr}_i} \in \mathcal{L}(\text{SU}(2)). \quad (6.6)$$


This property, common to all quivers of a $\mathcal{N} = 2$ gauge theory, implies the quiver reduction $Q \rightarrow Q'$ for \mathcal{L} . The proof of (6.6) is by now standard [6, 7] and it is spelled out for the C_2 example in appendix A. Then one has to show that all representations of Q' which satisfy

the Jacobian relations $\partial\mathcal{W}' = 0$ also satisfy the original relations $\partial\mathcal{W} = 0$ with the B_i 's set to 1. In other words, one has to check that the relations

$$\partial_{B_i} \mathcal{W} \Big|_{B_i=1} = 0 \tag{6.7}$$

are automatically satisfied. The check is presented in appendix B.

$\lambda = \infty$ corresponds to setting $A_i = 1$, $B_i = 0$. E.g. for B_2 we reduce to the representations of the quiver



subjected to the relations

$$\eta^* \alpha_1 = \eta = \alpha_1^* = 0. \tag{6.9}$$

Deleting the zero arrows, we get a string algebra whose strings have, again, dimensions in $\Delta^+(B_2)$.

7 Other applications of the monodromic construction

The monodromic quiver construction has also other applications besides the quivers for $\mathcal{N} = 2$ super-Yang Mills with non-simply laced gauge groups. We stress that most considerations in this section are *purely formal* (they may or may not correspond to consistent QFTs), and should be taken with a grain of salt.

The methods of refs. [4, 6] give the pair (Q, \mathcal{W}) for $G = ADE$ SQCD with quarks in representations which are direct sums of the irrepr. F_i whose Dynkin labels have the form

$$F_i \equiv [0, 0, \dots, 0, 1, 0, \dots, 0], \quad 1 \text{ in the } i\text{-th position, } i = 1, 2, \dots, r$$

(these r irrepr. are called *fundamental* representations of G). At the formal level, the methods of [4, 6] may be extended to *reducible* representations of the form [4]

$$\overbrace{F_i \otimes F_i \otimes F_i \otimes \dots \otimes F_i}^{n \text{ times}}, \tag{7.1}$$

and their direct sums. For instance, the usual quiver for $SU(2)$ $\mathcal{N} = 2^*$ corresponds in facts to $SU(2)$ SYM coupled to a hypermultiplet in the reducible representation [4, 6]

$$\mathbf{2} \otimes \mathbf{2} = \mathbf{3} \oplus \mathbf{1}. \tag{7.2}$$

Indeed, the corresponding light subcategory contains an extra singlet hypermultiplet.

It would be desirable to have a more general construction which allows to construct (Q, \mathcal{W}) for all matter representations and, in particular, for the symmetric products

$$\text{Sym}^n F_i \equiv [0, \dots, 0, n, 0, \dots, 0]. \tag{7.3}$$

The monodromic construction allows one to construct natural pairs (Q, \mathcal{W}) whose light subcategories contain precisely the *ADE* W -bosons plus a hyper in the $\text{Sym}^n F_i$. Basically, making $\lambda \rightarrow e^{2\pi i} \lambda$, induces a cyclic permutation of the direct factors in eq. (7.1). (Of course, the matter should be located in an Abelian subcategory \mathcal{L}_{λ_0} over a point $\lambda_0 \in \mathbb{P}^1$ which is not a fixed point of the $U(1)$ action, that is, $\lambda_0 \neq 0, \infty$).

There is a special reason to be interested in such extensions. In [6] it was noticed the following remarkable fact: *$\mathcal{N} = 2$ SYM with an ADE gauge group G coupled to a hypermultiplet in the F_i is asymptotically free if and only if the augmented graph $G[i]$ (obtained by adding a node to the Dynkin graph of G connected by a single edge to the i -th node) is again an ADE Dynkin graph.* This does not reproduce the full list of asymptotically-free ADE SQCDs because there are matter representations not of the form F_i which are consistent with asymptotic freedom; in facts, for $\mathfrak{g} = \text{ADE}$ there is just one infinite series of such non-fundamental asymptotically-free representations, namely $SU(N)$ coupled to the symmetric two-index representation of dimension $(\mathbf{N} + 1)\mathbf{N}/2$. (This representation is not in the list of the ‘nice’ ones admitting a Type IIB geometric engineering [51].) In [6] it was suggested that, maybe, this last case is also consistent with the correspondence

$$(\text{augmented Dynkin graphs}) \longleftrightarrow (\text{asymptotically free } \mathcal{N} = 2 \text{ SQCD})$$

provided one identifies the relevant augmented Dynkin graph with the C_N one. This proposal was not explored in [6] since, at the time, the non-symply case was not yet understood.

Now we are in a position to answer that question. Before doing that, however, it is better to introduce yet another ‘geometric’ tool to construct (Q, \mathcal{W}) pairs for 4d $\mathcal{N} = 2$ theories, namely quiver *specialization*.

7.1 Specialization: new pairs (Q, \mathcal{W}) from old ones

Our construction of the *BCFG* SYM quivers was modeled on the BIKMSV construction of non-simply laced gauge symmetries in string/ F -theory [29]. That construction has been extended by Katz and Vafa to cover the matter sector [31]. In F -theory the gauge and matter sectors arise from fibrations of the same kind of singularities, except that matter is related to singularities in codimension one relatively to the ones responsible for the gauge interaction. In order to construct (Q, \mathcal{W}) for various matter contents, we mimic this ‘matter from geometry’ idea through the procedure we shall call *specialization*.

7.1.1 The ‘heavy quark procedure’ revisited

We start with an alternative viewpoint on the ‘heavy quark procedure’ [1, 4, 6, 7] for the construction of the quiver for $\mathcal{N} = 2$ SYM with gauge group $G = \text{ADE}$ coupled to a hypermultiplet in a fundamental representation F_i of G . One starts by constructing the augmented graph $G[i]$ and defines its ‘Cartan’ matrix as the $(r + 1) \times (r + 1)$ matrix $C[i] \equiv 2 - I$, I being the incidence matrix of the augmented graph $G[i]$. Then one considers the *fake* SYM quiver with exchange matrix


$$B = C[i] \otimes S, \tag{7.4}$$

equipped with the standard square-tensor-product $G[i] \square \widehat{A}(1,1)$ superpotential [6, 50]

$$\mathcal{W}_{G[i] \square \widehat{A}(1,1)} = \sum_{\substack{\alpha \text{ link} \\ \text{of } G[i]}} \left(A_{t(\alpha)} \alpha B_{s(\alpha)} \alpha^* - B_{t(\alpha)} \alpha A_{s(\alpha)} \alpha^* \right) \quad (7.5)$$

Notice that (7.4), (7.5) is not, in general, an ADE SYM quiver pair, since $G[i]$ needs not to be a Dynkin graph, and therefore $C[i]$ is not necessarily a Cartan matrix of ADE Lie algebra. According to [6], B would be a *bona fide* ADE SYM quiver if and only if the coupled model is asymptotically free.

The pair (Q, \mathcal{W}) for G SYM coupled to a hypermultiplet in the representation F_i is obtained from the above fake SYM one by setting the two arrows A_0, B_0 of the vertical Kronecker subquiver over the augmentation node 0 of $G[i]$ equal to two fixed complex numbers $(a_0, b_0) \equiv \lambda_0 \in \mathbb{P}^1$. The sink and source nodes of \mathbf{Kr}_0 are then identified *via* the isomorphism B_0 ,



so that the resulting quiver has $2r + 1$ nodes and the exchange matrix has one zero eigenvalue, corresponding to the flavor symmetry f of the matter sector. The associated superpotential is

$$\mathcal{W} = \mathcal{W}_{G[i] \square \widehat{A}(1,1)} \Big|_{(A_0, B_0) = (\lambda_0, 1)} \quad (7.7)$$

Thus the SQCD pair (Q, \mathcal{W}) is just a *specialization* at the augmented node of the fake SYM pair. This implies that the generic light category \mathcal{L}_λ is equivalent to the pure G SYM one, while the *specialized* category $\mathcal{L}_{(a_0: b_0)}$ coincides with the one for ‘pure $G[i]$ fake SYM’. Of course the last category is well behaved iff $G[i]$ is a Dynkin quiver [6]. (Physically, specialization corresponds to replacing the would be simple-root W -boson vector associated to the augmentation node with a scalar).

7.1.2 More general (split) specializations: quiver gauge theories

In section 7.1.1 we specialized the augmented square-tensor-products of the form $G[i] \square \widehat{A}(1,1)$ at the augmentation Kronecker subquiver $\uparrow\uparrow$. We could as well have specialized at any other Kronecker subquiver, or even at several such subquivers.

A first generalization of the ‘heavy quark procedure’ may be described as follows. Let P be any alternating quiver (recall that a quiver is *alternating* if all its nodes are either sinks or sources; all connected trees admit two alternating orientations). Then the square-tensor-product $P \square \widehat{A}(1,1)$ is a well defined [50] pair (Q_P, \mathcal{W}_P) with a vertical Kronecker subquiver for each node in P . If $I \subset P_0$ is a subset of the nodes of P , the specialization at I corresponds to specializing

$$(A_i, B_i) \rightsquigarrow (a_i, b_i) \in \mathbb{P}^1 \quad \text{for all } i \in I, \quad (7.8)$$

while identifying in pairs the sources and sinks of the Kronecker subquivers \mathbf{Kr}_i for $i \in I$.

For instance, we can specialize $A_N \square \widehat{A}(1, 1)$ at the K -th Kronecker subquiver. The resulting pair (Q, \mathcal{W}) corresponds to the quiver gauge theory with gauge quiver

$$\bigcirc \longrightarrow \bigcirc \tag{7.9}$$

where the nodes stand, respectively, for $SU(K)$ and $SU(N - K)$ gauge groups and the arrow to a $(N - K, \overline{K})$ bifundamental hypermultiplet.

From this example it is pretty obvious that we may construct the pair (Q, \mathcal{W}) of any quiver (in the gauge sense) $\mathcal{N} = 2$ model with ADE gauge groups simply by specializing a suitable $P \square \widehat{A}(1, 1)$ at the appropriate set I of Kronecker subquivers.

The specializations of ADE SYM at *one* Kronecker precisely correspond to the list of cases explicitly discussed in ref. [31], namely

$$\begin{aligned} A_n &\rightarrow A_{n-k} \times A_{k-1} \\ D_n &\rightarrow D_{n-1}, A_{n-1}, D_{n-r} \times A_{r-1} \\ E_6 &\rightarrow D_5, A_5 \\ E_7 &\rightarrow D_6, E_6, A_6 \\ E_8 &\rightarrow E_7. \end{aligned} \tag{7.10}$$

In all instances the matter representations found in [31] using geometric methods agree with those found in [6] using the Representation Theory of the associated quivers with superpotential. In facts, in both approaches one is effectively reduced to a breaking of the adjoint representation of the Lie groups in the left part of eq. (7.10) to the Lie groups on the right *times* $U(1)_f$, where — after specialization — $U(1)_f$ is interpreted as the *global* flavor symmetry of the matter sector. (Note that specialization at k vertical Kronecker subquivers reduces the numbers of nodes of the quiver by k ; in particular, the exchange matrices of one-Kronecker specializations have odd rank, so that $\det B = 0$, consistent with the fact that there is a flavor $U(1)_f$ charge).

For more general 4d $\mathcal{N} = 2$ gauge models we have to combine specialization with the monodromic constructions discussed in the first part of the present paper.

7.2 Coupling matter in $\text{Sym}^n F_i$

The above construction by specialization may be generalized to the case in which the link between the augmentation node 0 and the i -th node of G has a non trivial value $(1, n)$, with $n \in \mathbb{N}$. We claim that, after the specialization

$$(A_0, B_0) = (a_0, b_0) \in \mathbb{P}^1,$$

this construction corresponds to coupling to G SYM a hypermultiplet in the representation $\text{Sym}^n F_i$ of G .

The physical case of this construction is $G = A_{N-1}$ coupled to $\text{Sym}^2 \mathbf{N} = (\mathbf{N} + \mathbf{1})\mathbf{N}/\mathbf{2}$. The corresponding augmented valued graph is the C_N Dynkin diagram. The generic category \mathcal{L}_λ has bricks with charges equal to the positive roots of $SU(N)$, while the bricks of the specialized category \mathcal{L}_{λ_0} have charges corresponding to the positive roots of C_N . Since

$$\text{AdSp}(N) = \text{AdSU}(N) \oplus (\mathbf{N} + \mathbf{1})\mathbf{N}/\mathbf{2} \oplus \overline{(\mathbf{N} + \mathbf{1})\mathbf{N}/\mathbf{2}} \oplus \mathbf{1}, \tag{7.11}$$

we get the claim for this example. In particular, this shows that the conjecture of [6] is true.

The Langlands dual construction in which one interchanges the role of long and short roots in the augmented valued graph, with the effect of replacing C_N with B_N , corresponds to a non-generic (in the sense of section 8.4 of [6]) construction of the pair (Q, \mathcal{W}) for the coupling to $SU(N)$ SYM to a hypermultiplet in the reducible representation

$$\mathbf{N} \oplus \mathbf{N}(\mathbf{N} - \mathbf{1})/\mathbf{N}. \quad (7.12)$$

in agreement with the group-theoretical branch rule

$$\text{Ad SO}(2N + 1) = \text{Ad SU}(N) \oplus \mathbf{N}(\mathbf{N} - \mathbf{1})/2 \oplus \overline{\mathbf{N}(\mathbf{N} - \mathbf{1})/2} \oplus \mathbf{N} \oplus \overline{\mathbf{N}} \oplus \mathbf{1}, \quad (7.13)$$

The specialization of G_2 SYM leads to the quiver for $SU(2)$ coupled to a hypermultiplet of isospin $3/2$ already discussed in [1] (see also [6]). The corresponding QFT is not UV complete, and, while the light subcategory \mathcal{L} is perfectly nice, the full non-perturbative category $\text{rep}(Q, \mathcal{W})$ is expected to be quite wild [6] (it *is* wild in the RT technical sense).


8 Coupling $BCFG$ SYM to matter

A detailed analysis of all models arising from the coupling of $BCFG$ $\mathcal{N} = 2$ SYM to matter is beyond the scope of the present paper. Here we limit ourselves to a sketchy discussion of some simple models, leaving a more systematic treatment to future work.

We focus on the non-simply laced analogues of the ‘matter from geometry’ models in eq. (7.10), namely one-Kronecker specializations of $BCFG$ SYM quivers (different from the ones corresponding to ADE SYM coupled to $\text{Sym}^n F_i$ matter already discussed in section 7.2). The non-simply laced counterpart to eq. (7.10) is¹⁰

$\mathfrak{g}_{\text{geometry}} \rightarrow \mathfrak{g}_{\text{gauge}}$	matter representations	
$B_n \rightarrow A_k \times B_{n-k-1}$	$(\mathbf{k} + \mathbf{1}, \mathbf{2}(\mathbf{n} - \mathbf{k}) - \mathbf{1})$	(8.1)
$C_n \rightarrow A_k \times C_{n-k-1}$	$(\mathbf{k} + \mathbf{1}, \mathbf{2}(\mathbf{n} - \mathbf{k} - \mathbf{1}))$	
$F_4 \rightarrow B_3$	$\mathbf{8} \oplus \mathbf{7}$	
$F_4 \rightarrow C_3$	$\mathbf{14} \equiv [0, 0, 1]$	

where the matter consists of one hypermultiplet in the indicated representation of $\mathfrak{g}_{\text{gauge}}$. The quivers and superpotentials for the various models are obtained straightforwardly from the specialization procedure. For instance, in the $A_k \times B_{n-k-1}$ case, the quiver is



¹⁰We omit the gauge singlets.

where the vertical Kronecker subquiver \mathbf{Kr}_{n-k} in the first quiver of figure 2 has been contracted to a single node attached by oriented triangles [2] to the rest of the quiver. Correspondingly, the superpotential is simply a specialization of the pure $\mathrm{SO}(2n+1)$ SYM one

$$\mathcal{W}_{\mathrm{SU}(k+1) \times \mathrm{SO}(2n-2k-1)} = \mathcal{W}_{\mathrm{SO}(2n+1)} \Big|_{(A_{n-k}, B_{n-k}) = (\lambda_0, 1)}. \quad (8.2)$$

As explained in [4, 6], if we have N_f flavors, to get the pair (Q, \mathcal{W}) we just reiterate the same construction N_f times.

In particular, taking $k = 0$ in the first two rows of eq. (8.1), we get the pairs (Q, \mathcal{W}) for $\mathrm{SO}(2m+1)$ and $\mathrm{Sp}(m)$ SYM coupled to N_f fundamentals.

Acknowledgments

We have greatly benefited of discussions with Murad Alim, Clay Córdova, Sam Espahbodi, Ashwin Rastogi and especially Cumrun Vafa. We thank them all. SC thanks the Simons Center for Geometry and Physics, where this work was completed, for hospitality.

A Consistency with the perturbative Higgs mechanism

The consistency of our pairs (Q, \mathcal{W}) with the perturbative Higgs mechanism requires the following mathematical **Fact** to hold

Fact. *If $X \in \mathcal{L}$, then*

$$X \Big|_{\mathbf{Kr}_i} \in \mathcal{L}(\mathrm{SU}(2)) \equiv \mathcal{T}_{\mathbf{Kr}} \quad \forall i. \quad (A.1)$$

Here $\mathcal{T}_{\mathbf{Kr}}$ stands for the Abelian category of the regular representations of the Kronecker quiver [38–40]. The strategy of the proof of this kind of results is rather standard see the appendices of ref. [6, 7]. Here we limit ourselves to present the details for the B_2 example.

Proof (for $G = B_2$). The B_2 quiver and superpotential are

$$\begin{array}{ccc}
 2, 1 & \xleftarrow{\alpha^*} & 1, 2 \\
 \downarrow B_2 & \xleftarrow{\eta^*} & \uparrow B_1 \\
 & \searrow \eta & \\
 & & 1, 1 \\
 \uparrow A_2 & \xrightarrow{\alpha} & \downarrow A_1
 \end{array} \quad (A.2)$$

$$\mathcal{W} = A_2 \alpha^* B_1 \alpha + A_1 \alpha B_2 \eta^* + A_1 \eta \alpha^* + B_1 \eta \eta^*.$$

Since $m_i(X) = 0$, if $X|_{\mathbf{Kr}} \notin \mathcal{T}$ it must have a preinjective summand. Consider first \mathbf{Kr}_1 . If $X|_{\mathbf{Kr}_1}$ has a preinjective summand, there are vectors — not *all* zero ([6] appendix) — $v_1, \dots, v_\ell \in X_{1,1}$ such that

$$A_1 v_1 = 0, \quad A_1 v_{a+1} = B_1 v_a, \quad \text{for } a = 1, \dots, \ell - 1 \quad B_1 v_\ell = 0. \quad (A.3)$$

Let us prove by induction on ℓ that, if this is the case, then $X \notin \mathcal{L}(B_2)$. The case $\ell = 1$ is trivial: if $A_1 v = 0 = B_1 v$, then $\mathbb{C}v$ is subrepresentation with positive m_1 magnetic charge and is not in \mathcal{L} . Assuming that it is true for $\ell - 1$ vectors, let us now prove that it is true also for ℓ of them. By the two equations

$$\partial_\eta \mathcal{W} = \alpha^* A_1 + \eta^* B_1 = 0 \quad (\text{A.4})$$

$$\partial_\alpha \mathcal{W} = A_2 \alpha^* B_1 + B_2 \eta^* A_1 = 0, \quad (\text{A.5})$$

we have that there exist $\ell - 1$ vectors

$$w_i \equiv \eta^* B_1 v_{i+1} \in X_{2,1} \quad i = 1, \dots, \ell - 1 \quad (\text{A.6})$$

that are such that $w_0 = 0$, $w_{\ell-1} = 0$, and

$$\begin{aligned} A_2 w_a &\equiv A_2 \eta^* B_1 v_{a+1} \\ &\stackrel{\text{eq. (A.4)}}{=} -A_2 \alpha^* A_1 v_{a+1} = \\ &\stackrel{\text{eq. (A.3)}}{=} -A_2 \alpha^* B_1 v_a = \\ &\stackrel{\text{eq. (A.5)}}{=} B_2 \eta^* A_1 v_a = \\ &\stackrel{\text{eq. (A.3)}}{=} B_2 \eta^* B_1 v_{a-1} = \\ &= B_2 w_{a-2}. \end{aligned} \quad (\text{A.7})$$

In particular, this implies that

$$A_2 w_1 = A_2 w_2 = 0 = B_2 w_{\ell-2}. \quad (\text{A.8})$$

Notice that if the vectors w_i are all equal to zero, then we have that $X|_{\mathbf{K}r_1}$ is a subrepresentation of X that being a preinjective summand of $X|_{\mathbf{K}r_1}$, has positive m_1 charge. Now, let us define

$$z_i \equiv \eta w_i + \alpha B_2 w_{i-1} \in X_{1,1} \quad i = 1, \dots, \ell - 1. \quad (\text{A.9})$$

If all the z_a are equal to zero, then by (A.7),

$$\eta w_a = -\alpha B_2 w_{a-1} = -\alpha A_2 w_{a+1}. \quad (\text{A.10})$$

Then, if also $\eta w_a \equiv 0$, eq. (A.4)(A.5), and

$$\partial_{A_1} \mathcal{W} = \alpha B_2 \eta^* + \eta \alpha^* = 0 \quad (\text{A.11})$$

give the additional relations

$$\begin{aligned} 0 \equiv \eta w_a &\equiv \eta \eta^* B_1 v_{a+1} \stackrel{\text{eq. (A.4)}}{=} -\eta \alpha^* A_1 v_{a+1} \\ &\stackrel{\text{eq. (A.11)}}{=} \alpha B_2 \eta^* A_1 v_{a+1} \stackrel{\text{eq. (A.5)}}{=} -\alpha A_2 \alpha^* B_1 v_{a+1}. \end{aligned} \quad (\text{A.12})$$

Now, for each node of $Q(B_2)$, consider the following subspaces:

$$\begin{aligned} V_{1,1} &\equiv \text{span}(v_1, \dots, v_\ell) \subset X_{1,1} \\ V_{1,2} &\equiv B_1 V_{1,1} = A_1 V_{1,1} \subset X_{1,2} \\ V_{2,1} &\equiv \eta^* V_{1,2} = -\alpha^* V_{1,2} \subset X_{2,1} \\ V_{2,2} &\equiv A_2 V_{2,1} = B_2 V_{2,1} \subset X_{2,2}. \end{aligned} \quad (\text{A.13})$$

The relations we have just obtained in (A.12) entail that

$$\eta V_{2,1} \equiv 0 \quad \text{and} \quad \alpha V_{2,2} \equiv 0, \quad (\text{A.14})$$

therefore, if $\eta w_a \equiv 0$, the module X has the following submodule (arrows are defined by restriction)

$$V \equiv (V_{1,1}, V_{1,2}, V_{2,1}, V_{2,2}) \subset X. \quad (\text{A.15})$$

Since

$$V|_{\mathbf{Kr}_1} \text{ is preinjective} \Rightarrow m_1(V) > 0. \quad (\text{A.16})$$

So, we have shown that X has a submodule with positive m_1 magnetic charge: This proves that if X is such that $\eta w_a \equiv 0$, $X \notin \mathcal{L}(B_2)$. Thus, assume that ηw_a are not all zero. If this is the case, but we still have that $z_a \equiv 0$, composing B_1 to (A.10) on the left, we have that

$$B_1 \eta w_a = -B_1 \alpha A_2 w_{a+1}, \quad (\text{A.17})$$

but by

$$\partial_{\alpha^*} \mathcal{W} = B_1 \alpha A_2 + A_1 \eta = 0 \quad (\text{A.18})$$

we have that

$$A_1 \eta w_a = -B_1 \alpha A_2 w_a \quad (\text{A.19})$$

Thus

$$A_1 \eta w_a = B_1 \eta w_{a-1}. \quad (\text{A.20})$$

Moreover, by (A.3), (A.4), and (A.11) we obtain

$$\begin{aligned} A_1 \eta w_1 &\equiv A_1 \eta \eta^* B_1 v_2 \\ &\stackrel{\text{eq. (A.4)}}{=} -A_1 \eta \alpha^* A_1 v_2 \\ &\stackrel{\text{eq. (A.3)}}{=} -A_1 \eta \alpha^* B_1 v_1 \\ &\stackrel{\text{eq. (A.11)}}{=} A_1 \alpha B_2 \eta^* B_1 v_1 \\ &\stackrel{\text{eq. (A.4)}}{=} -A_1 \alpha B_2 \alpha^* A_1 v_1 \\ &= 0. \end{aligned} \quad (\text{A.21})$$

That, together with

$$B_1 \eta w_{\ell-1} = 0, \quad (\text{A.22})$$

that holds by definition, entail that there are $\ell - 1$ vectors (not all zero), namely

$$\eta w_a \in X_{1,1} \quad a = 1, \dots, \ell - 1, \quad (\text{A.23})$$

such that (A.3) holds: Thus by the inductive hypothesis, $X \notin \mathcal{L}(B_2)$. Therefore, we can safely assume that the z_a 's are not all zero: By the equations of motion (A.18) and

$$\partial_{\eta^*} \mathcal{W} = A_1 \alpha B_2 + B_1 \eta = 0 \quad (\text{A.24})$$

the vectors z_a are such that, for $a = 2, \dots, \ell - 1$

$$\begin{aligned}
 A_1 z_a &\equiv A_1 \eta w_a + A_1 \alpha B_2 w_{a-1} \\
 &\stackrel{\text{eqs. (A.18), (A.24)}}{=} -B_1 \alpha A_2 w_a - B_1 \eta w_{a-1} \\
 &\stackrel{\text{eq. (A.7)}}{=} -B_1 \alpha B_2 w_{a-2} - B_1 \eta w_{a-1} \\
 &= -B_1 z_{a-1}.
 \end{aligned} \tag{A.25}$$

Then, since $w_0 = 0$,

$$A_1 z_1 \equiv A_1 \eta w_1 \stackrel{\text{eq. (A.21)}}{=} 0 \tag{A.26}$$

and, since $w_{\ell-1} = 0$,

$$\begin{aligned}
 B_1 z_{\ell-1} &\equiv \alpha B_2 w_{\ell-2} \\
 &\equiv \alpha B_2 \eta^* B_1 v_{\ell-1} \\
 &\stackrel{\text{eq. (A.3)}}{=} \alpha B_2 \eta^* A_1 v_\ell \\
 &\stackrel{\text{eq. (A.11)}}{=} -\eta \alpha^* A_1 v_\ell \\
 &\stackrel{\text{eq. (A.4)}}{=} \eta \eta^* B_1 v_\ell \\
 &= 0.
 \end{aligned} \tag{A.27}$$

So we have shown that if there is a family of ℓ vectors of type (A.3), then either X has a positive magnetic-charge submodule or there is an analogous family of $\ell - 1$ vectors (not all zero), and therefore $X|_{\mathbf{K}\mathbf{r}_1}$ cannot have preinjective summands, by induction, and it must be regular. Now, consider $X|_{\mathbf{K}\mathbf{r}_2}$. If $X|_{\mathbf{K}\mathbf{r}_2}$ has a preinjective summand then there are v_1, \dots, v_ℓ vectors in $X_{2,1}$ such that

$$A_2 v_1 = 0, \quad A_2 v_{a+1} = B_2 v_a, \text{ for } a = 1, \dots, \ell - 1 \quad B_2 v_\ell = 0. \tag{A.28}$$

Again we will proceed by induction on ℓ . The case $\ell = 1$ is trivial: we have $A_2 v = 0 = B_2 v$ and so that by eqs. (A.18), (A.24)

$$A_1 \eta v = 0 = B_1 \eta v. \tag{A.29}$$

If $\eta v = 0$ the $\mathbb{C}v \subset X_{2,1}$ is a destabilizing subrep, if not, then $\mathbb{C}\eta v \subset X_{1,1}$ is one. Now define

$$u_a \equiv \eta^* B_1 \alpha B_2 v_a \in X_{2,1} \quad i = 1, \dots, \ell - 1. \tag{A.30}$$

Now, if the u_a are zero, then, by

$$\partial_{\eta^*} \mathcal{W} = A_1 \alpha B_2 + B_1 \eta = 0 \tag{A.31}$$

we have the relations

$$\begin{aligned}
 0 &= \eta^* A_1 \eta v_{a+1} \\
 &\stackrel{\text{eq. (A.18)}}{=} \eta^* B_1 \alpha A_2 v_{a+1} \\
 &\stackrel{\text{eq. (A.28)}}{=} \eta^* B_1 \alpha B_2 v_a \\
 &\stackrel{\text{eq. (A.4)}}{=} \alpha^* A_1 \alpha B_2 v_a \stackrel{\text{eq. (A.28)}}{=} \alpha^* A_1 \alpha A_2 v_{a+1} \\
 &\stackrel{\text{eq. (A.31)}}{=} \alpha^* B_1 \eta v_a.
 \end{aligned} \tag{A.32}$$

that, together with

$$\begin{aligned}
 \partial_{A_2} \mathcal{W} &= \alpha^* B_1 \alpha = 0 \\
 \partial_{B_2} \mathcal{W} &= \eta^* A_1 \alpha = 0 \\
 \eta^* B_1 \eta &=_{\text{eq. (A.24)}} \eta^* A_1 \alpha B_2 = 0 \\
 \alpha^* A_1 \eta &=_{\text{eq. (A.18)}} \alpha^* B_1 \alpha A_2 = 0.
 \end{aligned} \tag{A.33}$$

entail, proceeding as above, that, the following subspaces

$$\begin{aligned}
 V'_{2,1} &\equiv \text{span}(v_1, \dots, v_\ell) \subset X_{2,1} \\
 V'_{2,2} &\equiv B_2 V'_{2,1} = A_2 V'_{2,1} \subset X_{2,2} \\
 V'_{1,1} &\equiv \alpha V'_{2,2} + \eta V'_{2,2} \subset X_{1,1} \\
 V'_{1,2} &\equiv A_1 V'_{1,1} + B_1 V'_{1,1} \subset X_{1,2}
 \end{aligned} \tag{A.34}$$

are such that

$$\alpha^* V'_{1,2} = 0 = \eta^* V'_{1,2}. \tag{A.35}$$

Therefore, if the module X is such that $u_a \equiv 0$, such a module has a submodule (arrows are defined by restriction)

$$V' \equiv (V'_{1,1}, V'_{1,2}, V'_{2,1}, V'_{2,2}) \subset X. \tag{A.36}$$

But

$$V'|_{\mathbf{Kr}_2} \text{ is preinjective} \Rightarrow m_2(V') > 0, \tag{A.37}$$

therefore X has a submodule of positive magnetic charge and it cannot belong to the light subcategory. This shows that we can assume that the u_a are not all zero. Now, by (A.4), (A.5), (A.18), (A.24), and (A.28)

$$\begin{aligned}
 A_2 u_a &\equiv A_2 \eta^* B_1 \alpha B_2 v_a \\
 &=_{\text{eq. (A.4)}} -A_2 \alpha^* A_1 \alpha B_2 v_a \\
 &=_{\text{eq. (A.24)}} A_2 \alpha^* B_1 \eta v_a \\
 &=_{\text{eq. (A.5)}} B_2 \eta^* A_1 \eta v_a \\
 &=_{\text{eq. (A.18)}} B_2 \eta^* B_1 \alpha A_2 v_a \\
 &=_{\text{eq. (A.28)}} B_2 \eta^* B_1 \alpha B_2 v_{a-1} \\
 &= B_2 u_{a-1}.
 \end{aligned} \tag{A.38}$$

This also shows that $A_2 u_1 = 0$ and that $B_2 u_{\ell-1} = 0$. So, by the inductive hypothesis on \mathbf{Kr}_2 , $X \notin \mathcal{L}(B_2)$. \square

B Consistency of $\text{rep}(Q, \mathcal{W}) \rightarrow \text{rep}(Q', \mathcal{W}')$

We need to check eq. (6.7) only at the Kronecker subquivers associated to nodes of the Dynkin graph which are connected by a link with a non-trivial value since at all other nodes we reduce to the corresponding statement for ADE SYM. In other words, we need to check only the two cases C_2 and G_2 .

B.1 C_2

 Let us recall (Q, \mathcal{W}) for C_2 SYM

$$(B.1)$$

$$\mathcal{W} = \alpha A_1 \alpha^* B_2 + \alpha B_1 \eta^* A_2 + \eta(\eta^* B_2 + \alpha^* A_2) \quad (B.2)$$

 Assuming the usual lemma [6, 7] on the restriction of $X \in \mathcal{L}$ on the Kronecker subquivers to hold, we set $B_1, B_2 = 1$, and consistency of this restrictions requires the relations

$$\partial_{B_2} \mathcal{W} \equiv \alpha A_1 \alpha^* + \eta \eta^* = 0 \quad (B.3)$$

$$\partial_{B_1} \mathcal{W} \equiv \eta^* A_2 \alpha = 0 \quad (B.4)$$

 to hold identically for all elements of $\text{rep}(Q', \mathcal{W}')$ where

$$(B.5)$$

$$\mathcal{W}' \equiv \mathcal{W}|_{B_1, B_2=1} = \alpha A_1 \alpha^* + \alpha \eta^* A_2 + \eta \eta^* + \eta \alpha^* A_2. \quad (B.6)$$

 Here we check that eqs. (B.3), (B.4) are indeed automatically satisfied. As in the main body of the paper, we may integrate out the massive fields η, η^* using their equations of motion

$$\eta^* = -\alpha^* A_2 \quad (B.7)$$

$$\eta = -A_2 \alpha. \quad (B.8)$$

 In this way we are reduced to the category $\text{rep}(Q'', \mathcal{W}'')$ where

$$(B.9)$$

$$\mathcal{W}'' = \alpha A_1 \alpha^* - \alpha \alpha^* A_2^2. \quad (B.10)$$

The relations of this last Jacobian algebra are

$$\partial_\alpha \mathcal{W}'' \equiv A_1 \alpha^* - \alpha^* A_2^2 = 0 \quad (\text{B.11})$$

$$\partial_{\alpha^*} \mathcal{W}'' \equiv \alpha A_1 - A_2^2 \alpha = 0 \quad (\text{B.12})$$

$$\partial_{A_1} \mathcal{W}'' \equiv \alpha^* \alpha = 0 \quad (\text{B.13})$$

$$\partial_{A_2} \mathcal{W}'' \equiv -\alpha \alpha^* A_2 - A_2 \alpha \alpha^* = 0. \quad (\text{B.14})$$

Let us check that eqs. (B.3), (B.4) are consequences of these relations. Indeed, using (B.7), (B.8), we get

$$\alpha A_1 \alpha^* + \eta \eta^* = \alpha A_1 \alpha^* + A_2 \alpha \alpha^* A_2 = \quad (\text{B.15})$$

$$= \alpha A_1 \alpha^* - \alpha \alpha^* A_2^2 + (\partial_{A_2} \mathcal{W}'') A_2 = \alpha (\partial_\alpha \mathcal{W}'') + (\partial_{A_2} \mathcal{W}'') A_2 = 0$$

$$\eta^* A_2 \alpha = -\alpha^* (A_2^2 \alpha) = -\alpha^* \alpha A_1 + \alpha^* (\partial_{\alpha^*} \mathcal{W}'') = -(\partial_{A_1} \mathcal{W}'') A_1 + \alpha^* (\partial_{\alpha^*} \mathcal{W}'') = 0. \quad (\text{B.16})$$


B.2 G_2

With reference to the G_2 SYM quiver in figure 2, the superpotential is given by eq. (6.3). Again, to restrict to the light category we set $B_1, B_2 = 1$ and identify the source and target nodes of the vertical Kronecker subquivers. The relations which should automatically hold for consistency are

$$\partial_{B_1} \mathcal{W}|_{B_1, B_2=1} \equiv \xi^* A_2 \alpha = 0 \quad (\text{B.17})$$

$$\partial_{B_2} \mathcal{W}|_{B_1, B_2=1} \equiv \alpha A_1 \alpha^* + \xi \xi^* + \eta \eta^* = 0. \quad (\text{B.18})$$

The light category is then given in terms of (Q', \mathcal{W}') where



$$\mathcal{W}' = \alpha A_1 \alpha^* - \xi^* A_2 \alpha - \eta^* A_2 \xi + \xi \xi^* - \alpha^* A_2 \eta + \eta \eta^*. \quad (\text{B.20})$$

We integrate out the massive fields η, ξ, η^*, ξ^* using their equations of motion

$$\xi = A_2 \alpha \quad (\text{B.21})$$

$$\xi^* = \eta^* A_2 \quad (\text{B.22})$$

$$\eta = A_2 \xi = A_2^2 \alpha \quad (\text{B.23})$$

$$\eta^* = \alpha^* A_2 \quad (\text{B.24})$$

$$\Rightarrow \xi^* = \alpha^* A_2^2. \quad (\text{B.25})$$

and we remain with the pair (Q'', \mathcal{W}'')

$$Q'' \quad A_1 \begin{array}{c} \curvearrowright \\ \curvearrowleft \end{array} 1 \begin{array}{c} \xleftarrow{\alpha^*} \\ \xrightarrow{\alpha} \end{array} 2 \begin{array}{c} \curvearrowleft \\ \curvearrowright \end{array} A_2 \quad (B.26)$$

$$\mathcal{W}'' = \alpha A_1 \alpha^* - \alpha \alpha^* A_2^3. \quad (B.27)$$

The corresponding Jacobian relations are

$$\partial_\alpha \mathcal{W}'' \equiv A_1 \alpha^* - \alpha^* A_2^3 = 0 \quad (B.28)$$

$$\partial_{\alpha^*} \mathcal{W}'' \equiv \alpha A_1 - A_2^3 \alpha = 0 \quad (B.29)$$

$$\partial_{A_1} \mathcal{W}'' \equiv \alpha^* \alpha = 0 \quad (B.30)$$

$$\partial_{A_2} \mathcal{W}'' \equiv -(A_2^2 \alpha \alpha^* + A_2 \alpha \alpha^* A_2 + \alpha \alpha^* A_2^2) = 0. \quad (B.31)$$

Let us check that these relations imply eqs. (B.17), (B.18). Indeed, in view of eqs. (B.21)–(B.25),

$$\xi^* A_2 \alpha = \alpha^* A_2^3 \alpha = \alpha^* (\alpha A_1) = (\partial_{A_1} \mathcal{W}'') A_1 = 0, \quad (B.32)$$

$$\begin{aligned} \alpha A_1 \alpha^* + \xi \xi^* + \eta \eta^* &= \alpha A_1 \alpha^* + A_2 \alpha \alpha^* A_2^2 + A_2^2 \alpha \alpha^* A_2 = \\ &= \alpha A_1 \alpha^* - A_2 (\partial_{A_2} \mathcal{W}'') - A_2^3 \alpha \alpha^* = (\partial_{\alpha^*} \mathcal{W}'') \alpha^* - A_2 (\partial_{A_2} \mathcal{W}'') = 0. \end{aligned} \quad (B.33)$$

References

- [1] S. Cecotti, A. Neitzke and C. Vafa, *R-Twisting and 4d/2d Correspondences*, [arXiv:1006.3435](#) [INSPIRE].
- [2] S. Cecotti and C. Vafa, *Classification of complete N = 2 supersymmetric theories in 4 dimensions*, [arXiv:1103.5832](#) [INSPIRE].
- [3] M. Alim, S. Cecotti, C. Cordova, S. Espahbodi, A. Rastogi and C. Vafa, *BPS Quivers and Spectra of Complete N = 2 Quantum Field Theories*, [arXiv:1109.4941](#) [INSPIRE].
- [4] M. Alim, S. Cecotti, C. Cordova, S. Espahbodi, A. Rastogi and C. Vafa, *N = 2 Quantum Field Theories and Their BPS Quivers*, [arXiv:1112.3984](#) [INSPIRE].
- [5] S. Cecotti and M. Del Zotto, *On Arnold's 14 'exceptional' N = 2 superconformal gauge theories*, *JHEP* **10** (2011) 099 [[arXiv:1107.5747](#)] [INSPIRE].
- [6] S. Cecotti, *Categorical tinkertoys for N = 2 gauge theories*, [arXiv:1203.6734](#) [INSPIRE].
- [7] S. Cecotti and M. Del Zotto, *Half-hypers and quivers*, *JHEP* **09** (2012) 135 [[arXiv:1207.2275](#)] [INSPIRE].
- [8] F. Denef, *Supergravity flows and D-brane stability*, *JHEP* **08** (2000) 050 [[hep-th/0005049](#)] [INSPIRE].
- [9] F. Denef and G.W. Moore, *Split states, entropy enigmas, holes and halos*, *JHEP* **11** (2011) 129 [[hep-th/0702146](#)] [INSPIRE].
- [10] M.R. Douglas and G.W. Moore, *D-branes, quivers and ALE instantons*, [hep-th/9603167](#) [INSPIRE].

- [11] D.-E. Diaconescu and J. Gomis, *Fractional branes and boundary states in orbifold theories*, *JHEP* **10** (2000) 001 [[hep-th/9906242](#)] [[INSPIRE](#)].
- [12] M.R. Douglas, B. Fiol and C. Romelsberger, *Stability and BPS branes*, *JHEP* **09** (2005) 006 [[hep-th/0002037](#)] [[INSPIRE](#)].
- [13] M.R. Douglas, B. Fiol and C. Romelsberger, *The spectrum of BPS branes on a noncompact Calabi-Yau*, *JHEP* **09** (2005) 057 [[hep-th/0003263](#)] [[INSPIRE](#)].
- [14] B. Fiol and M. Mariño, *BPS states and algebras from quivers*, *JHEP* **07** (2000) 031 [[hep-th/0006189](#)] [[INSPIRE](#)].
- [15] B. Fiol, *The BPS spectrum of $N = 2$ $SU(N)$ SYM and parton branes*, [hep-th/0012079](#) [[INSPIRE](#)].
- [16] F. Denef, *Quantum quivers and Hall/hole halos*, *JHEP* **10** (2002) 023 [[hep-th/0206072](#)] [[INSPIRE](#)].
- [17] B. Feng, A. Hanany, Y.H. He and A. Iqbal, *Quiver theories, soliton spectra and Picard-Lefschetz transformations*, *JHEP* **02** (2003) 056 [[hep-th/0206152](#)] [[INSPIRE](#)].
- [18] B. Feng, A. Hanany and Y.-H. He, *D-brane gauge theories from toric singularities and toric duality*, *Nucl. Phys. B* **595** (2001) 165 [[hep-th/0003085](#)] [[INSPIRE](#)].
- [19] B. Feng, A. Hanany and Y.-H. He, *Phase structure of D-brane gauge theories and toric duality*, *JHEP* **08** (2001) 040 [[hep-th/0104259](#)] [[INSPIRE](#)].
- [20] A. Hanany and K.D. Kennaway, *Dimer models and toric diagrams*, [hep-th/0503149](#) [[INSPIRE](#)].
- [21] S. Franco, A. Hanany, K.D. Kennaway, D. Vegh and B. Wecht, *Brane dimers and quiver gauge theories*, *JHEP* **01** (2006) 096 [[hep-th/0504110](#)] [[INSPIRE](#)].
- [22] B. Feng, Y.-H. He, K.D. Kennaway and C. Vafa, *Dimer models from mirror symmetry and quivering amoebae*, *Adv. Theor. Math. Phys.* **12** (2008) 489 [[hep-th/0511287](#)] [[INSPIRE](#)].
- [23] A.D. Shapere and C. Vafa, *BPS structure of Argyres-Douglas superconformal theories*, [hep-th/9910182](#) [[INSPIRE](#)].
- [24] D. Gaiotto, *$N = 2$ dualities*, *JHEP* **08** (2012) 034 [[arXiv:0904.2715](#)] [[INSPIRE](#)].
- [25] D. Gaiotto, G.W. Moore and A. Neitzke, *Wall-crossing, Hitchin Systems and the WKB Approximation*, [arXiv:0907.3987](#) [[INSPIRE](#)].
- [26] D. Gaiotto, G.W. Moore and A. Neitzke, *Framed BPS States*, [arXiv:1006.0146](#) [[INSPIRE](#)].
- [27] D. Gaiotto, G.W. Moore and A. Neitzke, *Wall-Crossing in Coupled 2d-4d Systems*, [arXiv:1103.2598](#) [[INSPIRE](#)].
- [28] D. Gaiotto, G.W. Moore and A. Neitzke, *Spectral networks*, [arXiv:1204.4824](#) [[INSPIRE](#)].
- [29] M. Bershadsky, K.A. Intriligator, S. Kachru, D.R. Morrison, V. Sadov and C. Vafa, *Geometric singularities and enhanced gauge symmetries*, *Nucl. Phys. B* **481** (1996) 215 [[hep-th/9605200](#)] [[INSPIRE](#)].
- [30] C. Vafa, *Geometric origin of Montonen-Olive duality*, *Adv. Theor. Math. Phys.* **1** (1998) 158 [[hep-th/9707131](#)] [[INSPIRE](#)].
- [31] S.H. Katz and C. Vafa, *Matter from geometry*, *Nucl. Phys. B* **497** (1997) 146 [[hep-th/9606086](#)] [[INSPIRE](#)].
- [32] P. Di Francesco and R. Kedem, *Q -systems as cluster algebras II: Cartan matrix of finite type and the polynomial property*, *Lett. Math. Phys.* **89** (2009) 183 [[arXiv:0803.0362](#)] [[INSPIRE](#)].

- [33] P.D. Francesco and R. Kedem, *Q-systems, heaps, paths and cluster positivity*, *Commun. Math. Phys.* **293** (2010) 727 [[arXiv:0811.3027](#)].
- [34] P. di Francesco and R. Kedem, *Q-system cluster algebras, paths and total positivity*, *SIGMA* **6** (2010) 14 [[arXiv:0906.3421](#)].
- [35] Y. Tachikawa, *On S-duality of 5d super Yang-Mills on S^1* , *JHEP* **11** (2011) 123 [[arXiv:1110.0531](#)] [[INSPIRE](#)].
- [36] R. Stekolshchik, *Springer Monographs in Mathematics: Notes on Coxeter Transformations and the McKay Correspondence*, Springer, Berlin Germany (2008).
- [37] D. Bump, *Graduate Text in Mathematics. Vol. 225: Lie groups*, Springer, Berlin Germany (2004).
- [38] C.M. Ringel, *Lecture Notes in Mathematics. Vol. 1099: Tame Algebras and Integral Quadratic Forms*, Springer, Berlin Germany (1984).
- [39] D. Simson and A. Skowronski, *London Mathematical Society Student Texts. Vol. 71: Elements of the Representation Theory of Associative Algebras. 2: Tubes and Concealed Algebras of Euclidean type*, Cambridge University Press, Cambridge U.K. (2007).
- [40] W. Crawley-Boevey, *Lectures on Representations of Quivers*, available on line at <http://www1.maths.leeds.ac.uk/~pmtwc/quivlecs.pdf>.
- [41] I.M. Gelfand and V.A. Ponomarev, *Model algebras and representations of graphs*, *Funkts. Anal. Prilozh.* **13** (1979) 1.
- [42] C.M. Ringel, *The preprojective algebra of a quiver*, in *Proceedings of Algebras and modules, II*, Geiranger Norway (1996), pg. 467 [*CMS Conf. Proc.* **24**, Amer. Math. Soc., Providence U.S.A. (1998)].
- [43] W. Crawley-Boevey, *On the exceptional fibres of Kleinian singularities*, *Amer. J. Math.* **122** (2000) 1027.
- [44] C. Geiss, B. Leclerc and J. Schröer, *Kac-Moody groups and cluster algebras*, [arXiv:1001.3545](#).
- [45] I. Assem, D. Simson and A. Skowronski, *London Mathematical Society Student Texts. Vol. 65: Elements of the Representation Theory of Associative Algebras. 1: Techniques of Representation Theory*, Cambridge University Press, Cambridge U.K. (2006).
- [46] P. Gabriel, *The universal cover of a representation-finite algebra*, in *Lectures Notes in Mathematics. Vol. 903: Representation Theory I*, Springer, Heidelberg Germany (1981), pg. 68.
- [47] R. Martinez-Villa and J.A. de la Peña, *The universal cover of a quiver with relations*, *J. Pure Appl. Algebra* **30** (1983) 277.
- [48] H. Derksen, J. Weyman and A. Zelevinsky, *Quivers with potentials and their representations I: Mutations*, *Selecta Math.* **14** (2008) 59 [[arXiv:0704.0649](#)].
- [49] M.C.R. Butler and C.M. Ringel, *Auslander-Reiten sequences with few middle terms and applications to string algebras*, *Comm. Algebra* **15** (1987) 145.
- [50] B. Keller, *The periodicity conjecture for pairs of Dynkin diagrams*, [arXiv:1001.1531](#) [[INSPIRE](#)].
- [51] Y. Tachikawa and S. Terashima, *Seiberg-Witten geometries revisited*, *JHEP* **09** (2011) 010 [[arXiv:1108.2315](#)] [[INSPIRE](#)].