

Constraints on the primordial power spectrum from high-resolution Lyman α forest spectra and *WMAP*

Matteo Viel,^{*} Jochen Weller and Martin G. Haehnelt

Institute of Astronomy, Madingley Road, Cambridge CB3 0HA

Accepted 2004 October 13. Received 2004 September 17; in original form 2004 July 19

ABSTRACT

The combined analysis of the cosmic microwave background on large scales and Lyman α forest on small scales provides a sufficiently long lever arm to obtain strong constraints on the slope and curvature of the power spectrum of primordial density fluctuations. We present results from the combination of the first year *WMAP* data and the dark matter power spectrum inferred by Viel, Haehnelt & Springel for two different sets of high-resolution and high signal-to-noise ratio quasar absorption spectra: the Croft et al. sample with a median redshift $z = 2.72$ and the LUQAS sample with a median redshift $z = 2.125$. The best-fitting value for the rms fluctuation amplitude of matter fluctuations is $\sigma_8 = 0.94 \pm 0.08$ and $n_s = 0.99 \pm 0.03$, if we do not include running of the spectral index. The best-fitting model with a running spectral index has parameters $n_s = 0.959 \pm 0.036$ and $n_{\text{run}} = -0.033 \pm 0.025$. The data are thus consistent with a scale-free primordial power spectrum with no running of the spectral index. We further include tensor modes and constrain the slow-roll parameters of inflation.

Key words: quasars: absorption lines – cosmic microwave background – cosmology: observations – cosmology: theory.

1 INTRODUCTION

The *Wilkinson Microwave Anisotropy Probe* team (*WMAP*, Bennett et al. 2003; Spergel et al. 2003) has presented impressive confirmation for what is now considered the standard cosmological model: a flat universe composed of matter, baryons and vacuum energy with a nearly scale-invariant spectrum of primordial fluctuations. One of the surprising results of the *WMAP* team's combined analysis of cosmic microwave background (CMB), galaxy redshift survey and Lyman α forest data was the evidence for a curvature of the power spectrum of primordial density fluctuations. This was expressed by the *WMAP* team in terms of a non-vanishing derivative $n_{\text{run}} = dn_s/d \ln k$ of the spectral index n_s . In inflationary models the shape of the primordial power spectrum is directly related to the potential of the inflaton field. For slow-roll inflation simple relations exist between the slow-roll parameters and the ratio of tensor to scalar fluctuation, the power-law index and the derivative of the power-law index of primordial fluctuations (see Liddle & Lyth 2000 for a review). Constraints on the power spectrum of primordial density fluctuations can thus at least in principle constrain inflationary models (Hannestad et al. 2002; Peiris et al. 2003; Leach & Liddle 2003; Kinney et al. 2004; Tegmark et al. 2004).

The Lyman α forest data constrain the dark matter (DM) power spectrum on scales of $\sim 4\text{--}40 h^{-1}$ Mpc in the redshift range $2 < z < 4$ [Croft et al. 1998, 1999; Hui 1999; McDonald et al. 2000;

Hui et al. 2001; Croft et al. 2002 (C02); McDonald 2003; Viel et al. 2003, 2004 (VHS)]. Owing to the long lever arm between the Lyman α forest on small scales and the CMB on large scales, the combined analysis of these data sets alone can provide strong constraints on the slope and curvature of the primordial power spectrum in the context of CDM-like models. Croft et al. (1999) inferred an amplitude and slope that was consistent with a *COBE* normalized Λ CDM model with a primordial scale-invariant fluctuation spectrum (Phillips et al. 2001). McDonald et al. (2000) and C02, using a larger sample of better quality data, found a somewhat shallower slope and smaller fluctuation amplitude. These latter data were part of the *WMAP* team's analysis mentioned above, which gave evidence for a tilted primordial CMB-normalized fluctuation spectrum ($n_s < 1$) and/or a running spectral index (Bennett et al. 2003; Spergel et al. 2003; Verde et al. 2003). There are, however, a number of systematic uncertainties in the amplitude of the DM power spectrum inferred from the Lyman α forest data (C02, Zaldarriaga, Hui & Tegmark 2001; Gnedin & Hamilton 2002; Seljak, McDonald & Makarov 2003; Zaldarriaga, Scoccimarro & Hui 2003; VHS).

VHS have recently used the method developed by C02 together with a large suite of new high-resolution numerical hydro-simulations to obtain an improved estimate of the DM power spectrum from the Lyman α forest. Two large samples of high-resolution spectra were used for this analysis, the sample of HIRES/LRIS spectra in the original analysis of C02 and the Large Sample of UVES QSO Absorption Spectra (LUQAS) of Kim et al. (2004). VHS demonstrated that the inferred rms fluctuation amplitude of the matter density is about 20 per cent higher than that inferred by C02 if the best

^{*}E-mail: viel@ast.cam.ac.uk

possible estimate for the effective optical depth, as obtained from high-resolution absorption spectra, is used. VHS further found that with this effective optical depth the inferred DM spectrum is consistent with a scale-invariant *COBE*-normalized DM power spectrum thus confirming the findings of Seljak et al. (2003). The running spectral index model advocated by the *WMAP* team falls significantly below the DM spectrum inferred from the Lyman α forest data (VHS).

In this paper, we will perform a combined analysis of the results of VHS based on the two independent data sets of C02 and K04 together with the *WMAP* data. The plan of the paper is as follows. In Section 2 we present the two data sets and briefly outline the method used to infer the linear dark matter power spectrum. Section 3 contains a description of the method used to estimate rms fluctuation amplitude σ_8 , the spectral index n_s , a possible running of the spectral index and the parameters of slow-roll inflationary models. In Section 4 we summarize and discuss our results.

2 THE *WMAP* AND LYMAN α FOREST DATA

2.1 The *WMAP* data

The CMB power spectrum has been measured by the *WMAP* team over a large range of multipoles ($l < 800$) to unprecedented precision on the full sky (Hinshaw et al. 2003; Kogut et al. 2003). For the analysis we use the one-year release of the *WMAP* temperature and temperature–polarization cross-correlation power spectrum.¹ The CMB power spectrum depends on the underlying primordial fluctuation via the photon transfer functions. The angular power spectrum in the temperature exhibits acoustic peaks at $l = 220$ with an amplitude of $\Delta T \approx 75 \mu\text{K}$ and $l = 540$ with $\Delta T \approx 49 \mu\text{K}$ (Page et al. 2003). The *WMAP* team (Spergel et al. 2003) showed that the measured CMB power spectrum is in excellent agreement with a flat universe and throughout this paper we will adopt this as a prior. In order to estimate parameters from the CMB we exploit for *COSMOMC*² an adapted version of the CMB likelihood code provided by the *WMAP* team (Verde et al. 2003; Hinshaw et al. 2003; Kogut et al. 2003).

2.2 The Lyman α forest data

The power spectrum of the observed flux in high-resolution Lyman α forest data constrains the DM power spectrum on scales of $0.003 \text{ s km}^{-1} < k < 0.03 \text{ s km}^{-1}$. At larger scales the errors due to uncertainties in fitting a continuum to the absorption spectra become large while at smaller scales the contribution of metal absorption systems becomes dominant (see e.g. Kim et al. 2004 and McDonald et al. 2004 for discussions). We will use here the DM power spectrum that VHS inferred for the C02 sample (Croft et al. 2002) and the LUQAS sample (Kim et al. 2004). The C02 sample consist of 30 Keck HIRES spectra and 23 Keck LRIS spectra and has a median redshift of $z = 2.72$. The LUQAS sample contains 27 spectra taken with the UVES spectrograph and has a median redshift of $z = 2.125$. The resolution of the spectra is 6 km s^{-1} , 8 km s^{-1} and 130 km s^{-1} for the UVES, HIRES and LRIS spectra, respectively. The signal-to-noise ratio (S/N) per resolution element is typically 30–50.

¹ <http://lambda.gsfc.nasa.gov>

² <http://www.cosmologist.info>

The method used to infer the dark matter power spectrum from the observed flux has been proposed by Croft et al. (1998) and has been improved by C02, Gnedin & Hamilton (2002) and VHS. The method relies on hydrodynamical simulations to calibrate a bias function between flux and matter power spectrum: $P_F(k) = b^2(k) P(k)$, on the range of scales of interest, and has been found to be robust (Gnedin & Hamilton 2002).

There are a number of systematic uncertainties and statistical errors that affect the inferred power spectrum, and an extensive discussion can be found in C02, Gnedin & Hamilton (2002), Zaldarriaga et al. (2003) and VHS. VHS estimated the uncertainty of the rms fluctuation amplitude of matter fluctuation to be 14.5 per cent. There is a wide range of factors contributing to this uncertainty. In the following we present a brief summary. The effective optical depth that is used to calibrate the numerical simulation and has to be determined separately from the absorption spectra is a major systematic uncertainty. Unfortunately, as discussed in VHS there is a considerable spread in the measurement of the effective optical depth in the literature. Determinations from low-resolution low S/N spectra give systematically higher values than those from high-resolution high S/N spectra. There is little doubt that the lower values from high-resolution high S/N spectra are appropriate and the range suggested in VHS leads to a 8 per cent uncertainty in the rms fluctuation amplitude of the matter density field (see table 5 in VHS). Other uncertainties are the slope and normalization of the temperature–density relation of the absorbing gas, which is usually parametrized as $T = T_0(1 + \delta)^{\gamma-1}$. T_0 and γ together contribute up to 5 per cent to the error of the inferred fluctuation amplitude. Our estimate of the uncertainties arising from the method was also 5. Somewhat surprisingly VHS found substantial differences between numerical simulations of different authors and assigned a corresponding uncertainty of 8 per cent. Other possible systematic errors include fluctuations of the UV background, temperature fluctuation in the intergalactic medium (IGM) owing to late helium reionization and the effect of galactic winds on the surrounding IGM. In VHS we had assigned an uncertainty of 5 per cent for these. Summed in quadrature this led to the estimate of an overall uncertainty of 14.5 per cent in the rms fluctuation amplitude of the matter density field.

For our analysis we take the inferred DM power spectrum in the range $0.003 \text{ s km}^{-1} < k < 0.03 \text{ s km}^{-1}$ as given in table 4 of VHS. Note that we have reduced the values by 7 per cent to mimic a model with $\gamma = 1.3$, the middle of the plausible range for γ (Ricotti, Gnedin & Shull 2000; Schaye et al. 2000). In the next section we will also show results for the DM power spectrum inferred by C02 in their original analysis (their table 3).

3 CONSTRAINING THE PRIMORDIAL POWER SPECTRUM AND INFLATION

3.1 The power spectrum

We will first attempt to constrain the primordial power spectrum of scalar fluctuations. We parametrize the power spectrum by

$$P(k) = P(k_0) \left(\frac{k}{k_0} \right)^{n_s(k)}, \quad (1)$$

where we choose the pivot scale to be $k_0 = 0.05 \text{ Mpc}^{-1}$ for the analysis in this subsection. The simplest case is a scale-invariant power spectrum with $n_s = 1$ constant. In a more general context,

and assuming $d^2 n_s / d \ln k^2 = 0$, the spectral index is defined by

$$n_s(k) = n_s(k_0) + \frac{dn_s}{d \ln k} \ln \left(\frac{k}{k_0} \right), \quad (2)$$

with $n_{\text{run}} \equiv dn_s / d \ln k$ parametrizing the evolution of the spectral index to first order.

We extended the *COSMOMC* package (Lewis & Bridle 2002) to include the Lyman α data as discussed in Section 2. As mentioned before, the main uncertainty of the Lyman α data is its relative calibration with respect to the underlying dark matter power spectrum. This is mainly influenced by the effective optical depth τ_{eff} , the normalization and slope of the temperature–density relation, T_0 and γ and the other uncertainties described in Section 2.2. We describe this uncertainty with an effective calibration amplitude A and obtain for the log-likelihood

$$-\ln \mathcal{L}_{\text{Ly}\alpha} \propto \sum_k \frac{[A P_{\text{dat}}(k) - P_{\text{th}}(k)]^2}{\sigma_p^2}. \quad (3)$$

We introduce the calibration A as an extra parameter into the Markov chain and multiply the likelihood with the prior

$$P_{\text{prior}} \propto \exp \left[-\frac{1}{2} \frac{(A - \bar{A})^2}{\sigma_A^2} \right]. \quad (4)$$

In order to obtain constraints on the cosmological parameters we will marginalize over the calibration parameter A . Note that this differs from the method described in Verde et al. (2003). As a standard prior on the mean we choose $\bar{A} = 1$.

The cosmological parameters that are varied for the combined analysis are the physical baryon density $\Omega_b h^2$, the physical dark matter density $\Omega_c h^2$, the angular diameter distance to last scattering θ , the optical depth τ owing to instantaneous complete reionization, the amplitude A_s of primordial perturbations and as mentioned above the spectral index of scalar perturbations, either running or constant. Note that we restrict the analysis to a flat universe with a cosmological constant, where we neglect the contribution from neutrinos and tensor modes. We further restrict the analysis to optical depths $\tau < 0.3$ as suggested by the *WMAP* team (Spergel et al. 2003; Verde et al. 2003). The amplitude of the matter fluctuations in a $8 h^{-1}$ Mpc sphere, σ_8 , is a derived quantity for this analysis.

We first discuss the results when we assume a constant spectral index n_s ($n_{\text{run}} = 0$).

In Fig. 1 we show the joint likelihood for the scalar spectral index n_s and σ_8 . On the left is the result for *WMAP* only. Note that we *only* include the *WMAP* data set and *not* the extended CMB data set in

order to concentrate on the constraints that are obtained by adding the Lyman α forest. The *WMAP* data alone give only very weak limits on the amplitude of the matter fluctuations and the spectral index. For a marginalized likelihood the spectral index is $n_s = 0.990 \pm 0.039$, consistent with a scale-invariant power spectrum as found by the *WMAP* team and other authors. In the middle panel we show the result if we combine the *WMAP* data with Lyman α data from C02 as compiled by Gnedin & Hamilton (2002). Note that we impose an uncertainty on the calibration of $\sigma_A = 0.25$ if $A < 1$ and $\sigma_A = 0.29$ if $A > 1$ (Verde et al. 2003). The spectral index differs now from a scale-invariant $n_s = 1$ primordial spectrum at the 2σ level. The marginalized best-fitting value is $n_s = 0.955 \pm 0.023$. The right panel shows the result for the *WMAP* data combined with the DM power spectrum as inferred by VHS. For these data we use a prior with $\sigma_A = 0.29$ throughout, which is twice the uncertainty of the rms fluctuation amplitude as discussed in Section 2. As discussed in detail by VHS, the inferred amplitude is about 20 per cent larger compared with that inferred by the C02 data with the lower effective optical depth. The spectral index is consistent with a scale-invariant spectrum with $n_s = 0.992 \pm 0.032$. There is *no* significant evidence for a non-scale-invariant spectrum, in agreement with the simpler analysis in VHS which obtained $n_s = 1.01 \pm 0.02(\text{stat.}) \pm 0.06(\text{syst.})$ and $\sigma_8 = 0.93 \pm 0.03(\text{stat.}) \pm 0.09(\text{syst.})$.

In Fig. 2 we show the dependence of the σ_8 inferred from the combined analysis on the amplitude of the power spectrum inferred from the Lyman α forest (parametrized as A , equation 3). The dependence around the best-fitting value is quite close to linearity ($\sigma_8^2 \approx A$). This plot was obtained with a prior on A given by equation (3). The contours close at small and large values owing to the constraints from the *WMAP* data alone. On the same axis we show explicitly the dependence on the effective optical depth (relative to the assumed best-fitting value). The degeneracy of the inferred amplitude with the assumed effective optical depth already discussed by C02 and Seljak et al. (2003) is clearly seen. For the relation between amplitude A and effective optical depth we assumed $A \propto \tau_{\text{eff}}^{-1.4}$ as found by VHS.

We will now discuss the constraints that we obtain if we include a running spectral index. In Fig. 3 we show the joint 1σ and 2σ likelihoods in the $n_{\text{run}}-n_s$ plane marginalized over all other parameters. In the left panel we show the constraints from *WMAP* only. Again there is no significant evidence for a deviation from a simple scale-invariant power spectrum, although the marginalized best-fitting values are $n_s = 0.922 \pm 0.062$ and $n_{\text{run}} = -0.050 \pm 0.037$. In the middle we show the result from combining *WMAP* with the C02 data. The best-fitting values are $n_s = 0.924 \pm 0.030$ and




Figure 1. Joint 1- and 2σ likelihoods on σ_8 and n_s . In the left panel the constraints are from *WMAP* only, in the middle panel from *WMAP* combined with the Lyman α power spectrum of C02, for a $\tau_{\text{eff}} = 0.349$. On the right the constraints are from *WMAP* combined with VHS, for a $\tau_{\text{eff}}(\zeta = 2.125) = 0.17$ and $\tau_{\text{eff}}(\zeta = 2.72) = 0.305$. The dashed lines indicate the spectral index for a scale-invariant $n_s = 1$ primordial spectrum.




Figure 2. Joint 1- and 2σ likelihoods on σ_8 as a function of the amplitude of the power spectrum A and of the effective optical depth.

$n_{\text{run}} = -0.052 \pm 0.016$. This would rule out the most simple scale-invariant power spectrum on the 3σ level. This is consistent with the findings of the *WMAP* team but somewhat unusual for slow-roll inflationary models, which require n_{run} to be of the order of $(|n_s - 1|)^2$ (Kosowsky & Turner 1995; but see Dodelson & Stewart 2002). If we instead combine the VHS data with *WMAP* (right panel) the contours shift again towards a scale-invariant spectrum with no running of the spectral index. The marginalized one-dimensional errors are $n_s = 0.959 \pm 0.035$ and $n_{\text{run}} = -0.033 \pm 0.025$. This makes the difference to a simple scale-invariant primordial power spectrum statistically insignificant.

In Table 1 we show the constraints on all the cosmological parameters for the combined *WMAP* and Lyman α analysis with and without a running spectral index.

3.2 Constraining slow-roll inflation

In order to obtain model independent constraints on inflation we concentrate on slow roll models. Since we are using a running of the spectral index of the scalar perturbations over a wide range in wavenumbers, we use a second-order extension of the standard slow-roll approximation (Liddle & Lyth 1993). This is described by

the parameters (Martin & Schwarz 2000; Leach et al. 2002)

$$\epsilon_1 \simeq \frac{M_{\text{Pl}}^2}{2} \left(\frac{V'}{V} \right)^2, \quad (5)$$

$$\epsilon_2 \simeq 2M_{\text{Pl}}^2 \left[\left(\frac{V'}{V} \right)^2 - \frac{V''}{V} \right], \quad (6)$$

$$\epsilon_2 \epsilon_3 \simeq 2M_{\text{Pl}}^4 \left[\frac{V''' V'}{V^2} - 3 \frac{V''}{V} \left(\frac{V'}{V} \right)^2 + 2 \left(\frac{V'}{V} \right)^4 \right], \quad (7)$$

where primes denote derivatives with respect to the field and $M_{\text{Pl}} = (8\pi G)^{-1/2}$ is the reduced Planck mass. Note that ϵ_1 , ϵ_2 , ϵ_3 correspond to the conventional slow-roll parameters by $\epsilon_1 = \epsilon_V$, $\epsilon_2 = 2\epsilon_V - 2\eta_V$ and $\epsilon_2 \epsilon_3 = 4\epsilon_V^2 - 6\epsilon_V \eta_V + 2\xi_V$. If we calculate the spectrum of primordial perturbations in the inflaton field we can identify the slow-roll parameters with parameters of the power spectrum. In general inflation also predicts tensor fluctuations or gravitational waves. We neglect here the running of the tensor index which is given by $n_t^{\text{run}} = \epsilon_1 \epsilon_2$. With the slow-roll conditions we obtain

$$n_s - 1 \approx -2\epsilon_1 - \epsilon_2 - 2\epsilon_1^2 - C\epsilon_2 \epsilon_3, \quad (8)$$

$$r \approx 16\epsilon_1, \quad (9)$$

$$n_{\text{run}} \approx -\epsilon_2 \epsilon_3, \quad (10)$$

with $C \approx 0.73$. We then obtain as a ‘consistency’ relation between the scalar to tensor ratio r and the tensor spectral index

$$n_t = -\frac{r}{8} - \frac{r^2}{128}. \quad (11)$$

Note that this is only true if we neglect the running of the tensor spectral index. We will calculate the slow-roll quantities from the Markov chains generated in the parameters n_s , n_{run} and r . In this case we chose as the pivot scale $k_0 = 0.002 \text{ Mpc}^{-1}$ (Peiris et al. 2003).

We perform a parameter analysis as described in Section 3.1 at this pivot scale with the addition of tensor modes with a free amplitude ratio $r < 1$ and the constraint in equation (11). In Fig. 4 we show the results of this analysis. The plot shows the results in the $\eta_V - \xi_V$ plane. The models are colour-coded according to the third slow-roll parameter ϵ_V . From equation (9) we see that $0 < \epsilon_V < 0.0625$, if we




Figure 3. Joint 1- and 2σ likelihoods on n_{run} and n_s . In the left panel the constraints are from the *WMAP* only, in the middle panel from *WMAP* combined with the Lyman α power spectrum of C02, for a $\tau_{\text{eff}}(z = 2.72) = 0.349$. On the right the constraints are from *WMAP* combined with the results of VHS, for a $\tau_{\text{eff}}(z = 2.125) = 0.17$ and $\tau_{\text{eff}}(z = 2.72) = 0.305$. The dashed lines indicate a scale-invariant $n_s = 1$ and $n_{\text{run}} = 0$ primordial spectrum.

Table 1. The marginalized results on cosmological parameters. The quoted values correspond to the peaks of the marginalized probability distributions. Errors are the 68 per cent confidence limits.

	WMAP+C02 (power law)	WMAP+VHS (power law)	WMAP+C02 (running)	WMAP+VHS (running)
$\Omega_c h^2$	0.109 ± 0.010	0.124 ± 0.013	0.124 ± 0.013	0.132 ± 0.015
$\Omega_b h^2$	0.022 ± 0.009	0.024 ± 0.001	0.022 ± 0.001	0.023 ± 0.001
h	0.721 ± 0.040	0.704 ± 0.047	0.683 ± 0.048	0.677 ± 0.047
τ	0.115 ± 0.056	0.157 ± 0.068	0.192 ± 0.065	0.182 ± 0.069
σ_8	0.794 ± 0.052	0.936 ± 0.079	0.921 ± 0.071	0.978 ± 0.090
n_s	0.955 ± 0.023	0.992 ± 0.032	0.924 ± 0.030	0.959 ± 0.036
n_{run}	–	–	-0.052 ± 0.016	-0.033 ± 0.025




Figure 4. Joint likelihood in the η_V - ξ_V plane, where the colour coding of the dots is scaled with ϵ_V . The contours are the 68 and 95 per cent confidence levels.

restrict the analysis to $r < 1$. For most models of inflation there is a negligible contribution of gravitational waves to the perturbations, i.e. $r \sim \epsilon_V \approx 0$ (Liddle & Lyth 2000). In this case the likelihood contour in Fig. 4 is restricted to the area depicted by the dark points, towards the lower left of the figure. η_V and ξ_V are thus relatively small, which according to equations (8) and (10) corresponds to a spectrum very close to scale-invariant scalar fluctuations. This is consistent with our findings in Section 3.1. However, with the current data, models with a large tensor contribution are also valid, which correspond to the light coloured points in Fig. 4, in the top right corner. The marginalized best-fitting values for the slow-roll parameters are $\epsilon_V = 0.032 \pm 0.018$, $\eta_V = 0.020 \pm 0.025$ and $\xi_V = 0.015 \pm 0.014$. The constraint on the tensor to scalar ratio is $r = 0.499 \pm 0.296$.

4 CONCLUSION

We have combined the linear dark matter power spectrum inferred from the Lyman α forest with the first-year *WMAP* data to obtain constraints on the rms fluctuation amplitude of matter fluctuations σ_8 and on the slope and shape parameter of the primordial density power spectrum n_s and n_{run} . We have restricted our analysis to spatially flat models with a low gravitational wave contribution to the CMB fluctuations. The *WMAP* data alone provide only weak

constraints on σ_8 , n_s and n_{run} . Adding the Lyman α forest significantly tightens the constraints because of the resulting long lever arm, which spans more than 3 orders of magnitude in wavenumber. The *WMAP* data combined with the DM power spectrum inferred by Croft et al. (2002) from their Lyman α forest data require a tilt ($n_s < 1$) and/or running of the spectral index ($n_{\text{run}} < 0$) to fit the data which is significant at the 3σ level. If we combine instead the *WMAP* data with the DM power spectrum inferred by VHS, a smaller tilt and running of the spectral index are required which are not statistically significant. The best-fitting value for the rms fluctuation amplitude of matter fluctuations is $\sigma_8 = 0.94 \pm 0.08$ and $n_s = 0.99 \pm 0.03$, in good agreement with the simpler analysis presented by VHS.

The difference of the analysis for the DM power spectrum of C02 and VHS is mainly due to the more appropriate smaller effective optical depth obtained from high resolution, high S/N QSO absorption spectra that was assumed in VHS. We further included tensor modes and put constraints on the slow-roll parameters. At present the limits on ϵ_V , ξ_V and η_V are not very tight, but future CMB missions such as *Planck*, together with better understanding of the systematic uncertainties of the Lyman α forest data, will improve the constraints.

ACKNOWLEDGMENTS

This work is supported by the European Community Research and Training Network ‘The Physics of the Intergalactic Medium’. The simulations were performed at the UK National Cosmology Supercomputer Center funded by PPARC, HEFCE and Silicon Graphics/Cray Research. We thank Sarah Bridle, George Efstathiou, Sabino Matarrese, Hiranya Peiris for useful discussions, Anthony Lewis for technical help and useful discussions and Dominik Schwarz for helpful comments on the manuscript.

REFERENCES

- Bennett C. L. et al., 2003, *ApJS*, 148, 1
- Croft R. A. C., Weinberg D. H., Katz N., Hernquist L., 1998, *ApJ*, 495, 44
- Croft R. A. C., Weinberg D. H., Pettini M., Hernquist L., Katz N., 1999, *ApJ*, 520, 1
- Croft R. A. C., Weinberg D. H., Bolte M., Burles S., Hernquist L., Katz N., Kirkman D., Tytler D., 2002, *ApJ*, 581, 20 (C02)
- Dodelson S., Stewart E., 2002, *Phys. Rev. D*, 65, 101301
- Gnedin N. Y., Hamilton A. J. S., 2002, *MNRAS*, 334, 107
- Hannestad S., Hansen S. H., Villante F. L., Hamilton A. J. S., 2002, *Astropart. Phys.*, 17, 375
- Hinshaw G. et al., 2003, *ApJS*, 148, 135
- Hui L., 1999, *ApJ*, 516, 519
- Hui L., Burles S., Seljak U., Rutledge R. E., Magnier E., Tytler D., 2001, *ApJ*, 552, 15

- Kim T.-S., Viel M., Haehnelt M. G., Carswell R. F., Cristiani S., 2004, *MNRAS*, 347, 355 (K04)
- Kinney W. H., Kolb E. W., Melchiorri A., Riotto A., 2004, *Phys. Rev. D*, 69, 103516
- Kogut A. et al., 2003, *ApJS*, 148, 161
- Kosowsky A., Turner S. M., 1995, *Phys. Rev. D*, 52, 1739
- Leach S. M., Liddle A. R., Martin J., Schwarz D. J., 2002, *Phys. Rev. D*, 66, 023515
- Leach S. M., Liddle A. R., 2003, *Phys. Rev. D*, 68, 123508
- Lewis A., Bridle S., 2002, *Phys. Rev. D*, 66, 103511
- Liddle A. R., Lyth D. H., 1993, *Phys. Rep.*, 231, 1
- Liddle A. R., Lyth D. H., 2000, *Cosmological Inflation and Large-Scale Structure*. Cambridge Univ. Press, Cambridge
- Martin J., Schwarz D., 2000, *Phys. Rev. D*, 62, 103520
- McDonald P., Miralda-Escudé J., Rauch M., Sargent W. L., Barlow T. A., Cen R., Ostriker J. P., 2000, *ApJ*, 543, 1
- McDonald P., 2003, *ApJ*, 585, 34
- McDonald P. et al., 2004, *ApJ*, submitted (astro-ph/0405013)
- Page L. et al., 2003, *ApJS*, 148, 233
- Peiris H. et al., 2003, 148, 213
- Phillips J., Weinberg D. H., Croft R. A. C., Hernquist L., Katz N., Pettini M., 2001, 560, 15
- Ricotti M., Gnedin N., Shull M., 2000, *ApJ*, 534, 41
- Schaye J., Theuns T., Rauch M., Efstathiou G., Sargent W. L. W., 2000, *MNRAS*, 318, 817
- Seljak U., McDonald P., Makarov A., 2003, *MNRAS*, 342, L79
- Spergel D. N. et al., 2003, *ApJS*, 148, 175
- Tegmark M. et al., 2004, *ApJ*, 606, 702
- Verde L. et al., 2003, *ApJS*, 148, 195
- Viel M., Matarrese S., Theuns T., Munshi D., Wang Y., 2003, *MNRAS*, 340, L47
- Viel M., Haehnelt M. G., Springel V., 2004, *MNRAS*, 354, 684 (VHS)
- Zaldarriaga M., Hui L., Tegmark M., 2001, *ApJ*, 557, 519
- Zaldarriaga M., Scoccimarro R., Hui L., 2003, *ApJ*, 590, 1

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.