

## NOTES ON GEOGRAPHIC DISTRIBUTION

### Mammalia, Chiroptera, Vespertilionidae, *Myotis albescens*: New occurrence site in the state of Rio Grande do Sul, Brazil.

Fernando Marques Quintela<sup>1</sup>  
Rafael Almeida Porciúncula<sup>1</sup>  
Susi Missel Pacheco<sup>2</sup>


<sup>1</sup> Fundação Universidade Federal do Rio Grande, Departamento de Ciências Morfo-Biológicas.  
Avenida Itália km 8. Caixa Postal 474, CEP 96201-900, Rio Grande, RS, Brazil. E-mail: boiruna@yahoo.com.br

<sup>2</sup> Instituto Sauver. Rua Dr. Paulo Franco dos Reis 40. CEP 90480-090, Porto Alegre, RS, Brazil.

*Myotis albescens* (E. Geoffroy, 1806) is a medium-sized bat (Chiroptera: Vespertilionidae) characterized by a dark-chocolate dorsum and a whitish belly. Its urogenital and perianal regions have white fur from the base to the top, what makes it impossible to mix it and *M. levis* up (Barquez et al. 1993). Besides, its femoral membrane has a pigmented edge with fur. *Myotis albescens* feeds on insects and may form large colonies, or even cohabitiate with other bat species (Achaval et al. 2004). Some records from Paraguay, Uruguay, and Venezuela suggest that this species occurs mainly in rural areas and, eventually, in human constructions (González 1973; Laval 1973; Einsenberg and Redford 1991; Pacheco and Freitas 2003). Peracchi et al. (2006) and Tavares et al. (in press) reported the occurrence of this species in all Brazilian states; Miranda et al. (2007)

recorded *M. albescens* in Paraná and Rio Grande do Sul states. Voss (1973) and Pacheco and Marques (2006) observed that this species is most commonly observed in savannas and pioneer vegetations and may use either inhabited or abandoned houses.

According to Simmons (2005), the occurrence area of *M. albescens* extends all over Latin America, from southern Vera Cruz in Mexico to Uruguay and northern Argentina. Pacheco et al. (2007) recorded the species in five sub-basins: Camaquã, Piratini-Mirim-São Gonçalo, Caí, Santa Maria, Apuaê-Inhandava, that include the cities of Bagé, São Lourenço do Sul, Maquiné, Barracão, Itaara, Santa Maria, and Frederico Westphalen. We present a new occurrence site of *M. albescens* for Rio Grande do Sul (Figure 1).



**Figure 1.** A, South America showing the state of Rio Grande do Sul, Brazil (red); B, sites where specimens of *Myotis albescens* were collected in the state (dark points, previous records; red point, present record).

## NOTES ON GEOGRAPHIC DISTRIBUTION

The city of Rio Grande ( $31^{\circ}01'40''$  S,  $52^{\circ}05'40''$  W) is located in the southern region of Rio Grande do Sul coastal plain on the shore of Lagoa dos Patos, at sea level. During the field work of a project on the city's bat fauna, a male individual of *M. albescens* (Figure 2) was captured in a mist net positioned over a streamlet along the bridge near the salt marsh of Lagoa dos Patos ( $31^{\circ}59.247'$  S,

$52^{\circ}14.636'$  W) on 31 January 2006, around 22:30 h. The predominant vegetation of this area is composed of grassweeds (*Scirpus* sp.), rushes (*Juncus* sp.), ferns (*Blechnum* sp.), and some sparse trees. The measurements of the collected specimen (Table 1) and its hair color and robust feet confirm the identity (Barquez et al. 1999).



**Figure 2.** General view of a *Myotis albescens* specimen collected in the municipality of Rio Grande, southern Brazil. Photo by F. M. Quintela.

**Table 1.** Specimens of *Myotis albescens* deposited in scientific collections at Rio Grande do Sul, Brazil.

Measurements (mm)	Specimens			
	MA 1	MA 2	MA 3	MA 4
E	12.2	15.0	-	-
FA	37.0	38.4	34.3	-
FT	9.8	8.0	-	-
HB	44.0	49.0	-	-
LT	78.6	91.0	-	-
T	32.2	42.0	-	-
GSL	14.4	-	14.3	13.4
ZB	-	-	-	-
BCH	5.8	-	5.8	5.8
BB	7.4	-	-	7.3
MTR	5.1	-	5.1	5.1
PW	4.1	-	4.0	4.1
UMR	5.3	-	5.1	5.6
IW	4.9	-	-	5.0
BMt	7.5	-	7.5	7.50
DL	10.1	-	-	10.0
LTR	5.3	-	-	5.2
C1-C1	3.8	-	3.8	3.7

The individual was preserved in alcohol (70 %) and deposited in the mastozoological collection of the *Museu de Ciências e Tecnologia* at *Pontifícia Universidade Católica do Rio Grande do Sul* (MCT 1721).

The present record contributes to the knowledge on the geographic distribution of *M. albescens*. Because there are too few data on its distribution in southern Brazil, the conservation status of this species, in Rio Grande do Sul, has been considered as Data Deficient (DD) according to Pacheco and Freitas (2003).

### Acknowledgements

We are grateful to Ana C. Oliveira, Josiane A. S. Silva, Renato Lopes, Marco A. R. Mello, and Cleber P. Silva for their support; Rafael Pinheiro for helping during the field work, and Thasiane Alves for helping to identify plants.

## NOTES ON GEOGRAPHIC DISTRIBUTION

### Literature cited

- Achaval, F., M. Clara, and A. Olmos. 2004. Mamíferos de la República Oriental del Uruguay. Montevideo, Imprimex. 176 p.
- Barquez, R. M., N. P. Giannini, and M. A. Mares. 1993. Guide to the bats of Argentina. Norman, Oklahoma Museum of Natural History, 119 p.
- Bárquez, R. M., M. A. Mares, and J. K. Braun. 1999. The bats of Argentina. Special Publication of the Museum of Texas Tech University 42, 273 p.
- Eisenberg, J. F. and K. H. Redford. 1999. Mammals of the Neotropics. Vol. 3. The Central Neotropics: Ecuador, Peru, Bolivia, Brazil. Chicago, University of Chicago Press, 609 p.
- González, J. C. 1973. Observaciones sobre algunos mamíferos de Bopicuá (Dpto. de Rio Negro, Uruguay). Comunicaciones del Museo Municipal de Historia Natural del Rio Negro 1:1-14.
- Laval, R. K. 1973. A revision of the neotropical bats of the genus *Myotis*. Science Bulletin Natural History Museum Los Angeles County 15: 1-53.
- Miranda, J. M. D, A. P. Leite, I. P. Bernardi, and F. C. Passos. 2007. Primeiro registro de *Myotis albescens* (E. Geoffroy, 1806) (Chiroptera, Vespertilionidae) para o Estado do Paraná, Brasil. Biota Neotropica 7 (1): 13-15.
- Pacheco, S. M. and T. R. O. Freitas. 2003. Quirópteros, p. 493-497 In C. S. Fontana, G. A. Bencke, and R. E. Reis (ed.). Livro vermelho da fauna ameaçada de extinção no Rio Grande do Sul. Porto Alegre, Edipucrs.
- Pacheco, S. M. and R. V. Marques. 2006. Conservação de morcegos no Rio Grande do Sul, p. 91-106 In T. R. O. Freitas, E. Vieira, S. M. Pacheco, and A. Christoff (org.). Mamíferos brasileiros: sistemática, genética, ecologia e conservação. Rio de Janeiro, Sociedade Brasileira de Genética. 170 p.
- Pacheco, S. M., M. L. Sekiama, K. P. A. Oliveira, F. Quintela, M. M. Weber, R. V. Marques, D. Geiger, and D. D. Silveira. 2007. Biogeografia de Quirópteros da Região Sul. Ciência & Ambiente 35: 181-202.
- Peracchi, A. L., I. P. Lima, N. R. Reis, M. R. Nogueira, and H. Ortêncio-Filho. 2006. Ordem Chiroptera, p. 150-226 In N. R. Reis, A. L. Peracchi, W. A. Pedro, and I. Passos (org.). Mamíferos do Brasil. Londrina, Editora da Universidade Estadual de Londrina.
- Simmons, N. B. 2005. Order Chiroptera, p. 312-529 In D. E. Wilson and D. M. Reeder (ed.). Mammal species of the World: a taxonomic and geographic reference, 3<sup>rd</sup> edition. Baltimore, Johns Hopkins University Press.
- Tavares, V. C., R. Gregorin, and A. L. Peracchi. In Press. Sistemática: a diversidade de morcegos no Brasil, In S. M. Pacheco, R. V. Marques, and C. E. L. Esbérard. Morcegos no Brasil: biologia, sistemática, ecologia e conservação.
- Voss, W. A. 1973. Ensaio de lista sistemática dos mamíferos do Rio Grande do Sul, Brasil. Revista da Faculdade de Filosofia, Ciências e Letras de São Leopoldo 25: 1-36.

Received January 2007

Accepted March 2008

Published online March 2008