



INDO AMERICAN JOURNAL OF PHARMACEUTICAL RESEARCH



A REVIEW ON PHARMACO KINETIC DRUG INTERACTIONS OF STATINS

A Satishchandra*, P Sureshkumar, V Radhika, T Anitha Chowdary, V. Jagannath Patro, A Sireesha

Department of Pharmacology, Browns College of Pharmacy, Khammam, India.

ARTICLE INFO

Article history

Received 12/01/2017

Available online

31/01/2017

Keywords

Statins,
Drug Interactions,
Cytochrome P450 (CYP)
Enzymes,
Drug Transporters,
Adverse Effects.

ABSTRACT

The 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibitors (statins) are generally well tolerated as monotherapy. Statins are associated with two important adverse effects, asymptomatic elevation in liver enzymes and myopathy. Myopathy is most likely to occur when statins are administered with other drugs. Statins are substrates of multiple drug transporters (including OAT- -P1B1, BCRP and MDR1) and several cytochrome P450 (CYP) enzymes (including CYP3A4, CYP2C8, CYP2C19, and CYP2C9). Possible adverse effects of statins can occur due to interactions in concomitant use of drugs that substantially inhibit or induce their metabolic pathway. This review aim is to summarize the most important interactions of statins.

Corresponding author

Mr. A. Satish Chandra

Department of Pharmacology

Browns college of Pharmacy

Khammam, Telangana

9441125557

satishchandranano.1@gmail.com

Please cite this article in press as **Mr. A. Satish Chandra** et al. A Review on Pharmacokinetic Drug Interactions of Statins. *Indo American Journal of Pharmaceutical Research*.2017;7(01).

Copy right © 2017 This is an Open Access article distributed under the terms of the Indo American journal of Pharmaceutical Research, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Statins (or HMG-CoA reductase inhibitors) represent a class of drugs used to lower cholesterol levels by inhibiting the enzyme HMG-CoA reductase, which plays a central role in the production of cholesterol in the liver. As of 2010, a number of statins have been on the market: atorvastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin. Statins are associated with two uncommon, but important, adverse effects, asymptomatic elevation in liver enzyme activity and myopathy. (1, 2).

At the pharmacodynamic level (i.e., at their site of action), statins are not prone to interfere with other drugs. However, at the pharmacokinetic level (i.e., absorption, distribution, metabolism and excretion of a given drug), the available statins show notable differences, including half-life, systemic exposure, maximum plasma concentration (C_{max}), bioavailability, protein binding, lipophilicity, metabolism, presence of active metabolites, and excretion routes that is explained below with examples for mechanism of drug interactions.

Mechanisms

There are several mechanisms by which drugs may interact (3-5), and most of these mechanisms can be categorized as pharmacokinetic (involving intestinal absorption, distribution, metabolism, and elimination) or as pharmacodynamic, or as additive toxicity, respectively. Pharmacokinetic interactions: the interaction in intestinal absorption is best illustrated by an example: tetracyclines and other broad-spectrum antibiotics may impair the absorption of oral contraceptives (in particular those with low-dose progestogens and/or estrogens) and hence render contraception unsafe. Several drugs are subject to inactivation via metabolic degradation in the liver, catalysed by various liver enzymes.

The formation of these enzymes can be induced or enhanced by drugs such as rifampicine, griseofulvine, and several anti-epileptics (carbamazepin, phenytoine, phenobarbital), but also by regular alcohol consumption.

This process, which requires several weeks of treatment and which is indicated as enzyme induction, enhances the metabolic degradation of several drugs.

In practice, enzyme induction may play a relevant role for oral anticoagulants (coumarin type), corticosteroids (glucocorticoids), oral contraceptives, or quinidine. Accordingly, these categories of drugs are metabolized/inactivated more rapidly and their doses should therefore be increased.

A comparable but opposite problem is the inhibition of liver enzymes involved in the biotransformation by a variety of drugs, such as cimetidine, erythromycin, metronidazole, tricyclic antidepressants, phenothiazine-neuroleptics, and sulphonamides (also in cotrimoxazole). Enzyme inhibitors of this type impair the biodegradation of certain drugs and hence increase their effects.

A well-known problem is the enhanced effect of anticoagulants (as reflected by bleeding) induced by additional treatment with cotrimoxazole. Certain drugs may impair the renal excretion (3-5) of other agents, usually at the renal tubular level.

A well-known relevant example is the rise in the plasma level and toxicity of digoxin, provoked by verapamil, amiodarone, or quinidine. Similarly, thiazide diuretics may decelerate the renal elimination of lithium salts and hence reinforce their toxicity. A beneficial effect of such an interaction is the impaired excretion of penicillin antibiotics induced by simultaneously administered probenecide.

Pharmacodynamic interactions and additive toxicity (3-5): Pharmacodynamic interactions between similarly acting drugs may lead to additive or even over-additive effects (potentiation). A well-known example is the combination of i.v. verapamil and a β -blocker, which may cause additive impairment of cardiac A-V conduction and the risk of A-V block.

Another possibility is the inhibition of the therapeutic effect of a drug by an additional agent. Over-additive adverse reactions are illustrated by the following example: a most important interaction, probably caused by non-specific mechanisms, is the mutual enhancement of the central nervous depressant effects of all drugs that are known to dampen the activity of the central nervous system.

This interaction holds for hypnotics, anxiolytics (minor tranquilizers), antipsychotics (neuroleptics, major tranquilizers), anti-epileptics, and opioids, but also for drugs with central nervous depressant adverse reactions, such as antihistamines, centrally acting antitussives (codeine, etc.), and scopolamine (3-5, 6).

Furthermore, alcohol enhances the central nervous depressant effects of all of the aforementioned therapeutics. Accordingly, enhanced sedation, impaired psychomotor skills (driving), but also respiratory depression may occur. The major drug interactions of statins tabulated below table.

Interacting Drug	Findings and Mechanism of Interaction	Recommendations/Comments
Amiodarone (e.g., Cordarone, generics)	<ul style="list-style-type: none"> Increased risk for myopathy/rhabdomyolysis due to decreased metabolism of atorvastatin, lovastatin, and simvastatin. Amiodarone is a CYP3A4 and CYP2C9 (moderate) inhibitor.(2) 	<ol style="list-style-type: none"> Do not exceed 40 mg lovastatin or 20 mg simvastatin.^{7,9,11} Consider limiting atorvastatin dose.¹¹ Fluvastatin is primarily metabolized by cyp3a4
Azole Antifungals Fluconazole (Diflucan, generics) Itraconazole (Sporanox) Ketoconazole (Nizoral, generics) Posaconazole (Noxafil [U.S.], Posanol [Canada]) Voriconazole (Vfend)	<ul style="list-style-type: none"> Increased risk for myopathy/rhabdomyolysis due to decreased metabolism of atorvastatin, fluvastatin, lovastatin, and simvastatin. Itraconazole, ketoconazole, posaconazole, and voriconazole are strong CYP3A4 inhibitors.⁶⁴ Fluconazole is a moderate CYP3A4 and CYP2C9 inhibitor.⁶⁴ 	<ol style="list-style-type: none"> Hold lovastatin and simvastatin during the course of itraconazole, ketoconazole, or posaconazole treatment.^{7,9} Consider reduced lovastatin or simvastatin dose when used with voriconazole.^{7,9,16} Hold atorvastatin during the course of itraconazole treatment.²⁵ Consider reduced atorvastatin dose if used with ketoconazole, posaconazole, or voriconazole.²⁶ Use caution when coadministering fluconazole with atorvastatin, lovastatin, simvastatin, or fluvastatin.^{18,19} Fifty percent statin dose reduction suggested.¹⁸ Limit fluvastatin dose to 20 mg twice daily.²¹ Consider fluvastatin, rosuvastatin, or pravastatin rather than lovastatin, simvastatin, or atorvastatin with itraconazole or ketoconazole.^{30,31}
Bile Acid Sequestrants Cholestyramine (e.g., Prevalite [U.S.], Olestyr [Canada], generics) Colestipol (Colestid, generics [U.S.])	Decreased bioavailability of statins due to the drugs binding in the intestine.	<ol style="list-style-type: none"> Administer statins 1 hr before or at least 4 hrs after cholestyramine or colestipol.²⁵⁻²⁷ Some Canadian statin monographs (atorvastatin, lovastatin, and simvastatin) recommend at least 2 hours elapse between statin administration and administration of bile acid sequestrants.^{12,14,24}
Calcium Channel Blockers (Amlodipine, Diltiazem, Verapamil) Grapefruit/Grapefruit Juice	<p>Amlodipine: Increased risk for myopathy/rhabdomyolysis due to decreased metabolism of simvastatin</p> <p>Increased risk for myopathy/rhabdomyolysis due to decreased metabolism of atorvastatin, lovastatin, and simvastatin. Grapefruit juice inhibits CYP3A4 and Pglycoprotein.</p>	<ol style="list-style-type: none"> Do not exceed lovastatin 20 mg or simvastatin 10 mg with verapamil or diltiazem.^{11,13} Experts suggest avoiding grapefruit with atorvastatin, simvastatin, and lovastatin.³³ Consider using pravastatin, rosuvastatin, or fluvastatin.
Macrolide Antibiotics Clarithromycin (e.g., <i>Biaxin</i> , generics) Erythromycin	Increased risk for myopathy/rhabdomyolysis due to decreased metabolism of atorvastatin, lovastatin, and simvastatin. These macrolides are CYP3A4 inhibitors. ⁶⁴	<ol style="list-style-type: none"> Lovastatin and simvastatin should be held during treatment with these macrolides.^{11,15} Do not exceed atorvastatin 20 mg with clarithromycin. Also consider cautious dosing with erythromycin.²⁴ Use azithromycin if treatment

Warfarin	Potential increase in INR due to decreased warfarin metabolism and displacement of warfarin from protein binding sites.	<p>with a macrolide antibiotic is unavoidable.</p> <ol style="list-style-type: none"> 4. Do not exceed pravastatin 40 mg with clarithromycin.²⁹ 5. Use rosuvastatin or fluvastatin in patients who require frequent or prolonged treatment with clarithromycin or erythromycin. 6. • Limit pitavastatin to 1 mg with erythromycin. <ol style="list-style-type: none"> 1. Monitor INR closely when initiating, stopping, or changing the statin dose 2. Atorvastatin may be less likely to interact.^{24,29}
----------	---	---

CONCLUSION

Statins have proven to be safe in numerous clinical trials. However, studies have shown that they can interact with other co-administered medicines. The different pharmacokinetic profiles of different statins should be carefully examined in order to understand the different spectra of drug interactions. These interactions are important determinants of safety in patients with hypercholesterolemia, especially those requiring long-term therapy with drugs that are well-known CYP3A4 inhibitors/inducers

REFERENCES

1. T. Uno, K. Sugimoto, K. Sugawara and T. Tateishi, The role of cytochrome P2C19 in R-warfarin pharmacokinetics and its interaction with omeprazol, *Ther. Drug Monit.* 30 (2008) 276–281; DOI: 10.1097/FTD.0b013e31816e2d8e.
2. C. Bolego, R. Baetta, S. Bellosta, A. Corsini and R. Paoletti, Safety considerations for statins, *Curr. Opin. Lipidol.* 13 (2002) 637–644.
3. Hansten PhD. Important drug interactions. In: Katzung BG (Ed). *Basic and clinical pharmacology*. Prentice-Hall Int, Englewood Cliffs NJ, USA, 5th Ed, 1992; pp 931 - 42.
4. Stockley IH. *Drug interactions*. Pharmaceutical Press, London, 5th Ed, 1999.
5. Opie LH. Cardiovascular drug interactions. In: Messerli FH (Ed). *Cardiovascular drug therapy*. W.B. Saunders Company, Philadelphia, USA, 2nd Ed, 1996; pp 347 - 53.
6. Van Zwielen PA, Eijssman L. *Drug therapy in cardio-thoracic surgery*. Van Zuiden Communications, Alphen a/d Rijn, The Netherlands, 2nd Ed, 2001; pp 262 – 9
7. Product information for *Mevacor*. Merck & Co., Inc. Whitehouse Station, NJ 08889. February 2012.
8. Product monograph for *Mevacor*. Merck Frosst Canada Ltd. Kirkland, QC H9H 3L1. March 2011.
9. Product information for *Zocor*. Merck & Co., Inc. Whitehouse Station, NJ 08889. February 2012.
10. Product monograph for *Zocor*. Merck Canada, Inc. Kirkland, QC H9H 4M7. October 2011.
11. Product information for *Cordarone*. Wyeth Pharmaceuticals Inc. Philadelphia, PA 19101. October 2011.
12. Kahri J, Valkonen M, Backlund T, et al. Rhabdomyolysis in a patient receiving atorvastatin and fluconazole. *Eur J Clin Pharmacol* 2005;60:905-7.
13. Tufan A, Dede DS, Cavus S, et al. Rhabdomyolysis in a patient treated with colchicine and atorvastatin. *Ann Pharmacother* 2006;40:1466-9.
14. Shaikat A, Benekli M, Vladutiu GD, Slack JL, Wetzler M, Baer MR. Simvastatin/fluconazole causing rhabdomyolysis. *Ann Pharmacother* 2003;37:1032-5.
15. Product information for *Vfend*. Pfizer Inc. New York, NY 10017. November 2011.
16. Product monograph for *Multaq*. Sanofi-Aventis Canada Inc. Laval, QC H7L 4A8. November 2011.
17. Product information for *Lescol* and *Lescol XL*. Novartis Pharmaceuticals Corporation. East Hanover, NJ 07936. February 2012.
18. Product monograph for *Lescol* and *Lescol XL*. Novartis Pharmaceuticals Canada, Inc. Dorval, QC H9S 1A9. March 2012.
19. Neuvonen PJ, Niemi M, Backman JT. Drug interactions with lipid-lowering drugs: mechanisms and clinical relevance. *Clin Pharmacol Ther* 2006;80:565-81.
20. Product monograph for *Lipitor*. Pfizer Canada Inc. Kirkland, QC H9J 2M5. March 2011.
21. Product information for cholestyramine. Upsher-Smith Laboratories, Inc. Minneapolis, MN 55447. September 2011
22. Product information for *Colestid* tablets. Pfizer Inc. New York, NY 10017. June 2006.
23. Product monograph for *Colestid* granules. Pfizer Canada Inc. Kirkland, QC H9J 2M5. September 2003.
24. Choi DH, Choi J, Li C, Choi JS. Effects of simvastatin on the pharmacokinetics of diltiazem and its main metabolite, desacetyldiltiazem, after oral and intravenous administration in rats: possible role of p-glycoprotein and CYP3A4 inhibition by simvastatin. *Pharmacol Rep* 2011;63:1574-82.
25. Product information for *Lipitor*. Pfizer Inc. New York, NY 10017. February 2012.
26. Product information for *Crestor*. AstraZeneca Pharmaceuticals LP. Wilmington, DE 19850. February 2012.

27. Product monograph for *Crestor*. AstraZenecaCanada Inc. Mississauga, ON L4Y 1M4. April 2012.
28. Pasternak RC, Smith SC Jr, Bairey-Merz CN, et al. ACC/AHA/NHLBI clinical advisory on the use and safety of statins. *J Am Coll Cardiol* 2002;40:567-72.
29. Bailey DG, Dresser GK. Interactions between grapefruit juice and cardiovascular drugs. *Am J Cardiovasc Drugs* 2004;4:281-97.
30. Strandell J, Bate A, Hagg S, Edwards IR. Rhabdomyolysis a result of azithromycin and statins: an unrecognized interaction. *Br J Clin Pharmacol* 2009;68:427-34.
31. . Product monograph for *Pravachol*. Bristol-Myers Squibb Canada. Montreal, QC H4S 0A4. October 2010.



54878478451161107



Submit your next manuscript to **IAJPR** and take advantage of:

Convenient online manuscript submission

Access Online first

Double blind peer review policy

International recognition

No space constraints or color figure charges

Immediate publication on acceptance

Inclusion in **ScopeMed** and other full-text repositories

Redistributing your research freely

Submit your manuscript at: editorinchief@iajpr.com

